Journal Pre-proof

Decoding the role of the cerebellum in the early stages of reading acquisition

Hehui Li, Olga Kepinska, Jocelyn N. Caballero, Leo Zekelman, Rebecca A. Marks, Yuuko Uchikoshi, Ioulia Kovelman, Fumiko Hoeft

PII: S0010-9452(21)00168-4

DOI: <https://doi.org/10.1016/j.cortex.2021.02.033>

Reference: CORTEX 3233

To appear in: *Cortex*

Received Date: 19 June 2020

Revised Date: 3 January 2021

Accepted Date: 8 February 2021

Please cite this article as: Li H, Kepinska O, Caballero JN, Zekelman L, Marks RA, Uchikoshi Y, Kovelman I, Hoeft F, Decoding the role of the cerebellum in the early stages of reading acquisition, *CORTEX*, <https://doi.org/10.1016/j.cortex.2021.02.033>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Published by Elsevier Ltd.

Decoding the role of the cerebellum in the early stages of reading acquisition

Hehui Li^{1,2}, Olga Kepinska^{2,3,4}, Jocelyn N. Caballero², Leo Zekelman^{2,5}, Rebecca A. Marks⁶, Yuuko Uchikoshi⁷, Ioulia Kovelman⁶, Fumiko Hoeft^{2,4,8,9*}

¹Center for Brain Disorders and Cognitive Science, Shenzhen University, Shenzhen, 518060, PR China

²Department of Psychiatry and Behavioral Sciences, and Weill Institute for Neurosciences, University of California San Francisco, San Francisco, CA, 94143, USA

³Department of Cognition, Emotion, and Methods in Psychology, Faculty of Psychology, University of Vienna, Liebiggasse 5, 1010 Vienna, Austria

⁴Department of Psychological Sciences, University of Connecticut, Storrs, CT, 06269, USA

⁵Speech and Hearing Bioscience and Technology, Harvard University, Cambridge, MA 02138, USA

⁶Department of Psychology, University of Michigan, Ann Arbor, MI, 48105, USA

⁷School of Education, University of California Davis, Davis, CA, 95616, USA

⁸Brain Imaging Research Center (BIRC), University of Connecticut, Storrs, CT 06269, USA

⁹Haskins Laboratories, New Haven, CT, 06511, USA

* corresponding author: Fumiko Hoeft, fumiko.hoeft@uconn.edu

Decoding the role of the cerebellum in the early stages of reading acquisition

Hehui Li^{1,2}, Olga Kepinska^{2,3,4}, Jocelyn N. Caballero², Leo Zekelman^{2,5}, Rebecca A. Marks⁶, Yuuko Uchikoshi⁷, Ioulia Kovelman⁶, Fumiko Hoeft^{2,4,8,9*}

¹ Center for Brain Disorders and Cognitive Sciences, Shenzhen University, Shenzhen, 518060, PR China

² Department of Psychiatry and Behavioral Sciences, and Weill Institute for Neurosciences, University of California San Francisco, San Francisco, CA, 94143, USA

³ Department of Cognition, Emotion, and Methods in Psychology, Faculty of Psychology, University of Vienna, Liebiggasse 5, 1010 Vienna, Austria

⁴ Department of Psychological Sciences, University of Connecticut, Storrs, CT, 06269, USA

⁵ Speech and Hearing Bioscience and Technology, Harvard University, Cambridge, MA 02138, USA

⁶ Department of Psychology, University of Michigan, Ann Arbor, MI, 48105, USA

⁷ School of Education, University of California Davis, Davis, CA, 95616, USA

⁸ Brain Imaging Research Center (BIRC), University of Connecticut, Storrs, CT 06269, USA

⁹ Haskins Laboratories, New Haven, CT, 06511, USA

* corresponding author: Fumiko Hoeft, fumiko.hoeft@uconn.edu

ABSTRACT. Numerous studies have consistently reported functional activation of the cerebellum during reading tasks, especially in the right cerebellar hemisphere. However, it remains unclear whether this region is also involved in reading in the earliest stages of reading acquisition. Here, we investigated whether and how the cerebellum contributes to reading acquisition. We tested 80 5-6-year-old kindergarteners, who performed a visual word matching task during which functional MRI (fMRI) data were collected. We found that bilateral cerebellar hemispheres were significantly activated during visual word processing. Moreover, activation of left cerebellar lobule VII extended to lobule VIII negatively and significantly correlated with current reading ability, whereas activation of right cerebellar lobule VII extended to lobule VIII significantly and positively correlated with future reading ability. Functional decoding via functional connectivity patterns further revealed that left and right cerebellar lobules connected with different cerebral cortex. Our results suggest a division of labor between the left and right cerebellar lobules in beginning readers.

Keywords: cerebellum, reading, longitudinal, beginning readers, division of labor.

1 Introduction

Successful reading requires the integration of different neural systems, such as the attention, articulatory, phonological, and semantic networks (Achal, Hoeft, & Bray, 2015). Previous studies have mainly focused on how these systems work with each other during reading in the cerebrum (Cao, 2016; D'Mello & Gabrieli, 2018). Recently, extensive studies have also observed cerebellar activation during reading or reading-related tasks, especially in the right cerebellar hemisphere (Cattinelli, Borghese, Gallucci, & Paulesu, 2013; Martin, Schurz, Kronbichler, & Richlan, 2015; Meng et al., 2016; Peng et al., 2003; Tan, Laird, Li, & Fox, 2005). In addition, these regions are known to be functionally connected with regions within the reading network established in the cerebrum (Alvarez & Fiez, 2018; Booth, Cho, Burman, & Bitan, 2007). Functional or structural deficits in right cerebellar lobules VI and VII are also associated with reading impairment (Eckert et al., 2016; Feng et al., 2017; Hancock, Richlan, & Hoeft, 2017;

Linkersdörfer, Lonnemann, Lindberg, Hasselhorn, & Fiebach, 2012). Overall, these results suggest that the right cerebellar hemisphere, in particular, lobules VI and VII play important roles in reading. An intriguing issue concerns whether these regions are also involved in reading in the early stages of reading acquisition.

As to the cerebellar contributions to early reading, previous reviews provide some clues. For example, the Cerebellar Deficit Hypothesis proposed that the cerebellum supports pre-reading skills, such as motor skills or articulatory/phonological skills (Nicolson et al., 2001). Moreover, this hypothesis also suggests that the cerebellum has a longitudinal influence on reading during reading acquisition. Deficits in the cerebellum in the early stage of reading may lead to reading impairment. This idea is consistent with a consensus paper on language and the cerebellum (Mariën et al., 2014). Recently there have been two studies by the same research group that reported the longitudinal effect of the cerebellum on reading, which showed that the left inferior cerebellar peduncle (white matter to the brain stem) at age six could predict an oral reading outcome that required reading comprehension at age eight (Borchers et al., 2019; Bruckert et al., 2019). However, the relation between microstructural properties of white matter or gray matter in the cerebellum and subsequent reading abilities (e.g., comprehension or single word reading) has not been found (or at least reported) by any other research groups (for gray or white matter volume that predicted reading outcome in cerebral regions, see (Linkersdörfer et al., 2014; Myers et al., 2014). On one hand, this divergence in whether the cerebellar structure may be associated with reading outcome may be due to different brain indices (whiter matter integrity or gray/white matter volume), sample sizes, or reading measurements, used across these studies. On the other hand, this inconsistency also indicates that it is still an open question about whether the development of the cerebellum in the early stages of reading contributes to future reading. In addition, these longitudinal studies were mainly focused on anatomical properties, which might not determine whether the cerebellum is an active participant during reading acquisition. Moreover, these studies mainly focused on relations between the cerebellum and reading without quantitatively decoding the role of the cerebellum. Therefore, the role of the cerebellum in reading remains elusive.

In the current study, we focused on three issues: (1) whether cerebellar lobule VI and/or VII are already associated with reading at the beginning stages of reading acquisition. (2) whether the functional status of these reading related regions in early reading influences future reading ability; and (3) what the functional contributions of the cerebellum to reading are. Reasons to focus on the cerebellar lobules VI and VII are twofold. First, these regions are frequently reported in reading research (Ang et al., 2020; Hancock et al., 2017; Li et al., 2020; Norton et al., 2014; Peng et al., 2003). Second, a recent study parcellated the cerebellum into ten functional subareas with multiple functional MRI (fMRI) tasks. The cerebellar regions engaged in language were mainly located at the right lobules VI and VII (King, Hernandez-Castillo, Poldrack, Ivry, & Diedrichsen, 2019);

To address these issues, kindergarten children who were just beginning to learn to read (Time 1), were instructed to perform a reading-related task during fMRI. Additionally, participants completed a series of behavioral assessments outside of the scanner to measure their reading ability or reading related skills (Time 1). One year later, after receiving formal reading instruction, reading ability was tested again (Time 2). Using data collected at time 1, we conducted voxel-wise activation analysis and brain-behavior (i.e., the cerebellar activation &

reading ability) correlation analysis to identify regions in the cerebellum associated with early reading. Then, a similar correlation analysis was further conducted between activation at Time 1 and reading ability at Time 2 (one year later) to examine whether the reading related regions identified at Time 1 would predict later reading ability at Time 2. We hypothesized that multiple regions in the cerebellum are related to early reading, especially right cerebellar lobule VI and/or VII, and functional activity in these regions might predict future reading ability. An emerging view suggests that the role of a brain region is best determined by its functional connectivity (FC) pattern with other regions (Achal et al., 2015; Alvarez & Fiez, 2018; Buckner, Krienen, Castellanos, Diaz, & Yeo, 2011; Johnson, 2000; Li et al., 2020). We then inferred the fine-grained role of reading related regions in the cerebellum by its connectivity patterns to cerebral regions (Buckner et al., 2011) with the current imaging data and a bigger database integrated into Neurosynth, i.e., F1000 (De La Vega, Chang, Banich, Wager, & Yarkoni, 2016; Rubin et al., 2017; Yarkoni, Poldrack, Nichols, Van Essen, & Wager, 2011).

2 Materials and Methods

2.1 Participants

Eighty beginning readers (mean age = 5.8 years old, range: 5.6-6.6 years old; boy/girl: 39/41) were included in the current study as part of a larger project (NIH/NICHD R01HD078351). Two prior studies overlapped partially with the current sample (Haft, Kepinska, Caballero, Carreiras, & Hoeft, 2019; Marks et al., 2019). All participants were native English speakers, and attended kindergartens in Northern Californian public schools in the United States. Among these children, eight participants were left-handed, three ambidextrous and the remaining sixty-nine participants were right-handed.

2.2 Behavioral measures obtained outside the scanner

2.2.1 Baseline behavioral measures (Time 1)

Kaufman Brief Intelligence Test (KBIT) was used to measure nonverbal intelligence (Kaufman & Kaufman, 2004). Children with typical intelligence were included in this study, quantified by a standard score above or equal to 85. Children's reading abilities were measured by the Woodcock-Johnson IV Letter Word Identification subtest (WJ LWID; Schrank, Mather, & McGrew, 2014). Initial items (from 1 to 13) of this subtest were designed to identify letters and words. From item 14, participants were required to read words arranged in order of difficulty.

Eight tests tapping into processes such as language skills related to reading, were also administered at Time 1. The Woodcock-Johnson IV Word Attack subtest (WJ Word Attack; Schrank et al., 2014) and the Test of Orthographic Competence Homophone Choice subtest (TOC Homophone Choice; Mather, Roberts, Hammill, & Allen, 2008) were used to measure phonological decoding ability. Three subtests from the Comprehensive Test of Phonological Processing (CTOPP-2): Elision, Blending Words, and Sound Matching, were used to assess phonological manipulation ability (Wagner, Torgesen, Rashotte, & Pearson, 2013). The Peabody Picture Vocabulary Test (PPVT-IV) was used to measure children's receptive vocabulary level (Dunn & Dunn, 2007). The Woodcock-Johnson IV Picture Vocabulary (WJ Picture Vocabulary)

and the Woodcock-Johnson IV Oral Comprehension (WJ Oral Comprehension) were used to assess their oral language ability (Schrack et al., 2014).

2.2.2 Behavioral measures at follow-up (Time 2)

One year later, 43 out of 80 participants returned for a follow-up behavioral assessment (mean age = 7.1 years old, range: 6.5-7.5 years old; boy/girl: 22/21). The mean duration between Time 1 and Time 2 testing was 1.3 years with a range of 1 year to 1.5 years. Letter-word identification (LWID) was assessed again among the 43 participants.

2.3 MRI data acquisition and analyses

2.3.1 Visual word matching task during fMRI

Given that the participants were all beginning readers with limited reading skills, we used a relatively easy task, a visual word matching task, to ensure the task was accessible to them. Visual word matching has been widely used as an implicit reading task (Feng et al., 2017; Gao et al., 2015). During the task, 24 pairs of words were presented in sequence with each trial lasting for 6000 ms. The first word was presented above a fixation cross for 2000 ms, followed by the second word under the fixation cross for another 2000 ms. Immediately after, a question mark was presented for another 2000 ms replacing the fixation cross. Participants were required to indicate whether the two words were the same or not (“bear” – “shark” = no; “wolf” – “wolf” = yes). There were 6 24-s task blocks (12 matching and 12 unmatching) that consisted of 4 trials as well as 6 12-s resting blocks after each task block. The whole run lasted approximately 4 minutes (see also Marks et al., 2019).

Participants in the current study are from the same dataset as Marks et al. (2019). Because they were beginning readers, the stimuli used in the current study were all high-frequency nouns, taken from pictures of kindergarten classrooms with high-frequency words on the wall (e.g., pot, boat, dog, or green; Table S1) and from publicly available 1st and 2nd-grade spelling lists to ensure their familiarity. According to two age of acquisition indices, these words were typically acquired before age five (Giholly & Logie, 1980; Kuperman, Stadthagen-Gonzalez, & Brysbaert, 2012). All words had one or two syllables, with an average of 4.74 letters, 3.71 phonemes, and 1.24 syllables. For each pair, stimuli were matched for the number of phonemes and syllables. By and large, words were also matched for the number of letters, although three out of 24 word pairs had words that differed in length by one letter (e.g., bear – shark). There were no significant differences observed between the 12 matching (i.e., same words) and 12 non-matching (i.e., different words) pairs in terms of phoneme length, age of acquisition, familiarity, written frequency, or imageability.

A sensitivity index (d') was used to estimate children’s performance on this task. This score was calculated by subtracting the z-transformed false alarm rate from the z-transformed hit rate. Hit rate was defined as the proportion of trials that participants correctly responded as “yes” relative to all “yes” trials. False alarm rate was defined as the proportion of trials where participants incorrectly responded as “yes” relative to all “no” trials. Children with d' larger than zero were included in the analyses, which means that they could differentiate

between “same” and “different” trials. The larger the d' score, the better the subject's sensitivity. The mean d' score was 2.37, with a standard deviation of 1.12 ($n = 80$).

2.3.2 MRI data acquisition

Data was acquired using a 3-T Siemens Prisma Fit MRI scanner equipped with a 64-channel head coil. Whole-brain functional images were acquired using a gradient-echo echo-planar pulse sequence [repetition time (TR) = 1250 ms, echo time (TE) = 33.40 ms, flip angle (FA) = 45°, field of view (FOV) = 220 mm, voxel size = 2.2 mm³, 64 contiguous 2.20-mm axial slices, 0-mm inter-slice gap]. Parameters for high resolution T1-weighted anatomical images were as follows: matrix size 256*256; 160 contiguous axial slices; voxel resolution 1 mm, TR = 2300 ms, TE = 2.98 ms, T1 = 900 ms; and FA = 9°.

2.3.3 Pre-processing of MRI data

Imaging data were processed using FSL software Version 6.0.0 (FMRIB's Software Library, FMRIB, Functional Magnetic Resonance Imaging of the Brain, www.fmrib.ox.ac.uk/fsl; Jenkinson et al., 2012). Pre-processing was carried out using FMRI Expert Analysis Tool (FEAT) Version 6.00. The pre-statistics processing steps included: removal of the first 11 volumes for signal equilibration (173 volumes retained); motion correction using MCFLIRT (Jenkinson, Bannister, Brady, & Smith, 2002); non-brain removal using BET (Smith, 2002); grand-mean intensity normalization of the entire 4D dataset by a single multiplicative factor; spatial smoothing using a Gaussian kernel of 5 mm full width at half maximum (FWHM); and B0 unwarping using BBR (Boundary-Based Registration; https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/FLIRT_BBR; Jenkinson et al., 2002; Jenkinson and Smith, 2001), which also performs simultaneous registration to the high resolution T1-image (rigid body, 6 degrees of freedom). After this, T1-images were registered to MNI-152 standard space using 12-parameter affine transformation and non-linear registration with a warp resolution of 10 mm in FNIRT (Andersson, Jenkinson, & Smith, 2007a, 2007b).

After conducting standard data preprocessing, we performed ICA-AROMA (Independent Component Analysis-based Automatic Removal of Motion Artifacts; <https://github.com/maartenmennes/ICA-AROMA>; Pruim et al., 2015) to identify and remove motion-related artifacts on data that passed motion quality control (i.e., data where the absolute mean displacement during the task was less than 5 mm). The de-noised data were then high-pass filtered with a cutoff of 36 s (0.036 Hz) and the registration parameters were reapplied.

The design matrix for each participant included task blocks as events of interest, which was convolved with the Hemodynamic Response function (HRF). In addition, signals from white matter and cerebrospinal fluid were included as nuisance regressors. For the FC analysis, only the time series during the task were extracted. Given the effect of the hemodynamic response, we delayed the onset time of each task block to 4TRs (each TR lasts 1.25s; 4TRs equal to 5 s in total).

2.3.4 MRI data analyses

2.3.4.1 The validity of the current task in eliciting reading-related activation

In the first stage of data analyses, we aimed to examine the validity of the visual word matching fMRI task in eliciting reading-related activation. To this end, two analyses were performed:

- (1) First, we compared areas showing significant activation during this task with the canonical reading network generated via Neurosynth. Areas showing significant activation were obtained by voxel-based whole brain activation analyses by contrasting task to fixation conditions. The threshold for significant activation of the statistical images was set non-parametrically using clusters determined by $Z > 2.3$ and a (corrected) cluster significance threshold of $p = .05$. The classical reading network was constructed via Neurosynth. Neurosynth is a platform for automated synthesis of large-scale fMRI data (<http://www.neurosynth.org/>; Yarkoni, Poldrack, Nichols, Van Essen, and Wager, 2011). The database consists of 507,891 activations reported in 14,371 studies and meta-maps for 1,335 keywords. When building the meta-map of reading, “reading” was used as a keyword and an association test map was automatically generated, which was corrected by false discovery rate (FDR of $\alpha = .05$), indicating regions more likely to activate in studies containing “reading” (i.e., the occurrence frequency of reading was more than 1/1,000 words) compared to studies that do not (i.e., occurrence was equal to or less than 1/1,000 words). We examined overlap in areas showing significant activation and the canonical reading network by overlaying these two maps.
- (2) Second, we performed reverse inferencing to qualitatively identify potential cognitive states associated with brain activity pattern induced by the current task using Neurosynth Image Decoder (<http://www.neurosynth.org/decode/>; Poldrack et al., 2011; Yarkoni et al., 2011), wherein Pearson’s correlation analyses were automatically performed between the unthresholded activation map from the current study and the unthresholded meta-maps of 1,335 keywords in the Neurosynth database. Correlation coefficients were then used to generate a ranking of the keywords associated with the current image of interest (whole-brain activation pattern).

2.3.4.2 Concurrent relationship between cerebellar activation (at Time 1) and reading ability at Time 1

Voxel-based correlation analyses were further conducted with FSL, during which general linear models were built to examine whether and identify which regions in the cerebellum were associated with reading. Statistical images were thresholded using a non-parametric cluster threshold determined by $Z > 2.3$ at the voxel-level and a (corrected) cluster significance threshold of $p = .05$. To control for the effect of variables of no-interest, age, gender, and handedness were used as covariates.

2.3.4.3 Longitudinal influence of cerebellar activation (at Time 1) on future reading (at Time 2)

Further correlation analyses between reading ability at Time 2 or reading gain (differences between LWID raw scores at Time 2 and Time 1) and activation in the cerebellum at Time 1 were conducted to investigate whether activation in the cerebellum could predict future reading ability. The threshold for statistical image was the same as above. Gender and handedness were again used as covariates.

2.3.4.4 Deciphering the function of the cerebellum by examining the FC pattern between the cerebellum and cerebral regions

An emerging view suggests that the role of a brain region is determined by its FC pattern with other regions (Johnson, 2000). Therefore, in the following analysis, we aimed to use FC to infer the role of the cerebellum.

- (1) We further examined the FC between cerebellar regions and regions within a “canonical reading network.” The cerebellar regions were defined as regions in the cerebellum that correlated with reading ability at Time 1 or Time 2. The coordinates of the “canonical reading network” that we used were those reported in a recent meta-analysis on single word reading (Murphy, Jógia, & Talcott, 2019). 11 spheres were created (regions of interest, ROIs) with a radius of 5 mm and coordinates in Table 1 as the center.

Table 1. Central coordinates of regions related to single word reading in the cerebral cortex (based on Murphy, Jógia, & Talcott, 2019).

ROIs	BA	MNI coordinates		
		X	Y	Z
L.PostG	4	-50	-8	44
L.MFG _a	9	-40	28	24
L.SPL	7	-22	-68	48
R.IFG	9	48	12	24
L.IOG	17	-24	-98	-4
L.MFG _b	6	-4	-2	56
R.SPL	7	34	-56	50
L.FG _a	37	-50	-48	-8
L.FG _b	37	-42	-54	-18
L.IFG	44	-44	6	26
L.STG	41	-54	-16	8

Then, FC between the cerebellar regions and these 11 cerebral ROIs within “canonical reading network” were calculated. To determine functional connections between the cerebellum and those ROIs, the first eigenvariate of the time series of each region was extracted from the fMRI data (Thurston et al., 2015), which reflects the time course of the first component that contributes most to a region’s response (Hanson & Bunzl, 2010; Saxe, Brett, & Kanwisher, 2006). For each participant, we calculated the correlation coefficient between the first eigenvariate of each ROI and the cerebellar regions where activity correlated with reading ability. The values were then transformed to Z-scores via Fisher z transformation, indicating the strength of FC. We focused on FC of the time-series only during the task condition. Given the effect of

the hemodynamic response, we delayed the onset time of each task block to 4TR (each TR lasts 1.25 s, 4TR are equal to 5 s). For each FC value between the cerebellum and an ROI, we subtracted the mean FC of the cerebellum from the remaining ROIs to indicate the specific FC with that particular ROI. This index indicates a cerebellar cluster to be relatively more connected to a given cerebral reading ROI relative to the remaining reading ROIs. One sample t-tests were conducted to estimate the significance of each specific FC value, which were corrected for multiple comparisons at an FDR of $\alpha = 0.05$. The role of the cerebellum was inferred by the reading-related cerebral ROI that showed the most significant specific FC with cerebellar areas.

To supplement this result, we used the F1000 data in Neurosynth comprised of 1000 adults because of the availability of the large sample and the fact that the connectivity pattern would most likely reflect the mature brain network associated with reading and the cerebellum (Buckner et al., 2011). The peak of each reading-related region in the cerebellum described above was used as a seed region separately to pinpoint its resulting target cerebral regions (<https://www.neurosynth.org/locations/>). Threshold to identify these regions was set as voxel-level correlation coefficient larger than 0.2, cluster volume larger than 200 mm³.

- (2) Finally, we used Neurosynth's "location based analysis" feature (<https://www.neurosynth.org/locations/>) to decode the cognitive terms associated with cerebral ROIs that connected with the cerebellar clusters. Based on the three-dimensional coordinates of the central voxel of a given ROI (Table 1), an association table with the keywords ranked by its relations with this location was generated using Neurosynth. The role of each region was inferred via the top 12 cognitive terms (Alvarez & Fiez, 2018).

3 Results

3.1 Demographic information

For behavioral performance on the LWID at Time 1, only 21 children scored under 14, indicating that they were still in the alphabetic phase of reading acquisition and could not identify words (Figure 1). The remaining children could all identify at least one high frequency word. We compared the behavioral performance on all tests measured at Time 1 between children who returned ($n = 43$) and did not return one year later ($n = 37$). However, no significant group differences were observed except that children who did not return had a greater LWID score and behavioral performance (d' score) in the scanner in kindergarten than children who returned for Time 2 testing ($t(78) = 2.8, p = .02$; Table 2).

All children who returned one year later ($n = 43$) scored 14 or more on LWID at Time 2 (Figure 1). Moreover, LWID scores of these children significantly increased from Time 1 to Time 2 ($Mean_{Time1} = 16.4, SD_{Time1} = 7.2$; $Mean_{Time2} = 41.8, SD_{Time2} = 11.1, t(42) = -17.6, p < .001$).

Figure 1. The distribution of reading performance at Time 1 and Time 2. The dotted line represents LWID scores equal to 14. Scores under 14 indicate that the children were still in the alphabetic phase of reading acquisition and could not identify words. Blue dots represent children who returned for Time 2, red dots represent children who did not return for Time 2.

Table 2. Demographic information of participants

	Whole sample			Return group		Non-return group		T (Return & Non-return)	p
	Mean	SD	Range	Mean	SD	Mean	SD		
Sample size	80			43		37			
Gender (boy/girl)	39/41			22/21		17/20		$\chi^2=0.6$	0.7
Age	5.8	0.4	5.1-6.5	5.8	0.4	5.7	0.4	1.9	0.06
Behavioral performance outside the scanner									
KBIT Non-verbal IQ ⁵	105.2	14.1	85.0-141.0	104.0	11.8	106.5	16.4	0.1	0.92
WJ Word Attack subtest	7.9	4.0	0-18	7.3	3.7	8.7	4.2	-1.53	0.13
TOC Homophone Choice	5.6	4.2	0-22	5.0	3.8	6.4	4.5	-1.49	0.14
CTOPP Blending	14.4	4.8	5-24	14.0	4.6	15.0	5.0	-1.00	0.32
CTOPP Elision	13.2	5.8	0-30	12.7	4.9	13.7	6.7	-0.83	0.41
CTOPP Sound Matching	15.3	6.5	1-25	14.8	6.3	15.9	6.9	-0.76	0.45
Peabody Picture Vocabulary Test	115.3	20.1	63-152	118.7	18.3	111.4	21.7	1.63	0.11
WJ Picture Vocabulary	25.1	3.4	17-33	25.4	3.2	24.9	3.6	0.67	0.51
WJ Oral Comprehension	12.5	3.9	1-20	13.1	3.7	11.9	4.1	1.37	0.18
WJ Letter Word Identification subtest	18.5	8.9	2-48	16.4	7.2	21.0	10.2	-2.4	0.02
Behavioral performance inside the scanner									
d' score	2.4	1.1	0.3-5.4	2.1	1.0	2.6	1.2	-1.9	0.05

Note: ⁵ standard score. Others were raw scores. SD = standard deviation.

3.2 Activation pattern during the visual word matching task

Voxel-wise whole brain activation analysis showed that bilateral frontal gyri, bilateral superior parietal lobule extending to left inferior parietal lobule, and bilateral inferior occipital gyri were

significantly activated during the visual word matching task (Table 3, Figure 2A, regions with red/yellow spectral color). In addition to these regions, bilateral cerebellar hemispheres were also significantly activated, even though no peak was reported in this region (Figure 2A, regions with red/yellow spectral color). Notably, no significant difference in activation was observed for participants who did and did not return for Time 2 testing.

Further analyses were conducted to examine whether this activation pattern was related to reading. First, we used "reading" as a keyword and generated a reading-related network from the Neurosynth database (Yarkoni, Poldrack, Nichols, Van Essen, & Wager, 2011; Figure 2A, lime region). The results showed that regions that showed significant activation in the visual word matching task largely overlapped with the reading-related network defined by Neurosynth (Figure 2A). The overlap was observed both in the cerebrum and cerebellum.

The *Neurosynth Image Decoder* was further used to decode the function of the activated regions, where the correlation between all term-based association test maps in the Neurosynth database and the unthresholded current activation map (β map) were calculated (Chang, Yarkoni, Khaw, & Sanfey, 2012; Rubin et al., 2017). Based on these correlations, we observed that the whole brain activation pattern induced by the current task had a greater correlation with reading (Figure 2B), task processing, goal, face, and visual processing compared to other Neurosynth terms.

Together, these results suggest that the visual word matching fMRI task that we used in the current study appropriately induced brain activation within the reading network including bilateral cerebellar hemispheres.

Figure 2. Whole brain activation pattern and its relations with reading. (A) The topographic relations between activation regions in the current study (lime color), the reading network established by Neurosynth (blue color), and the overlapping areas between the two (red). (B) The first 25 cognitive terms showing the greatest correlation with the whole brain activation map. The size of the font represents the strength of the correlation between the current activation map and term-based meta-maps generated by Neurosynth. The first five terms were colored orange and the remaining terms were colored black.

Table 3. Coordinates of peak activation during the visual word matching task

Cluster	Cluster Size	Z	x	y	z	
Left inferior occipital gyrus	56680	8.91	-42	-82	-10	Inferior occipital gyrus
		8.89	-38	-86	-10	
		8.84	-6	16	46	Supplementary motor area
		8.55	-32	-90	-4	Middle temporal gyrus
		8.53	-8	8	54	Supplementary motor area
Left superior parietal lobule	2150	6.79	-24	-62	44	Superior parietal lobule
		6.27	-28	-56	46	
		5.97	-26	-54	40	
		5.21	-28	-70	24	
		4.92	-48	-40	42	Inferior parietal lobule
		4.17	-48	-44	52	
Right middle frontal gyrus	365	4.64	24	44	-16	Middle frontal gyrus
		3.15	16	52	-22	Superior frontal gyrus
		2.67	32	64	-6	

3.3 Correlations between the cerebellum and reading ability at Time 1

We next examined the correlation between the cerebellar activation at Time 1 and reading ability at Time 1 (performance on LWID). We limited our correlation analysis to the voxels within the cerebellum. The results showed that activation in left cerebellar Crus II extending to left cerebellar lobule VIII showed a significant negative correlation with reading ability at Time 1 (as measured by LWID). From hereon we refer to this cluster with peak coordinates: $x = -6$, $y = -84$, $z = -30$, $p = .02$, cluster size = 250, $Z_{\max} = 3.91$ as *Left T1CRBLM* (Figure 3A, B). This result suggests that greater reading skill was associated with less left cerebellar activity. In addition, the *Left T1CRBLM* largely overlapped with the regions that showed significant activation during the visual word matching task but was more focal (Figure 3C).

Notably, we used a visual word matching task, which required participants to judge whether the words were identical or not. It might be argued that the performance on this task could also reflect visual perceptual abilities. In other words, the variation of activation in the *Left T1CRBLM* might be accounted for by the visual perceptual ability, which in turn correlates with reading ability (d' score in the scanner significantly and positively correlated with LWID at Time 1, $r = 0.31$, $p = .006$). To address this concern, participants' d' score on the visual word matching task were used as a covariate in a follow-up brain-behavior correlation analysis. The result showed that left cerebellar lobule VII was still significantly and negatively correlated with children's performance on LWID at Time 1 (peak coordinates: $x = -6$, $y = -84$, $z = -30$, $p = .02$, cluster size = 240, $Z_{\max} = 4.25$), thus pointing to the discrimination ability between the items in the scanner task not being the only driving force behind the significant negative relationship between the activation in the *Left T1CRBLM* cluster and reading ability.

On the other hand, if variation in the *Left T1CRBLM* activation was due to visual perceptual ability, then a similar effect should be observed in cerebral regions responsible for visual processing (e.g., bilateral occipital gyrus). However, the results showed that reading ability at Time 1 negatively and significantly correlated with activation in left middle temporal gyrus, right supramarginal gyrus, and right inferior frontal gyrus. Activation in these regions

together shows a pattern that is consistent with semantic processing, phonological processing, or social cognitive processes (Hartwigsen et al., 2010; Hartwigsen, Neef, Camilleri, Margulies, & Eickhoff, 2019; Weiss, Cweigenberg, & Booth, 2018), none of which are typically associated with low-level visual processing. These two complementary analyses suggest that signal change in the *Left T1CRBLM* might be associated with the difference in reading abilities rather than visual perceptual abilities.

We also examined the negative brain-behavior relationship in the smaller sample of 43 participants that returned for Time 2 testing. The results showed that activation in the left cerebellum was still negatively and significantly correlated with reading ability at Time 1 (peak coordinates: $x = -14$, $y = -72$, $z = -36$, $p < .001$, cluster size = 553, $Z_{\max} = 3.69$). This region overlapped with *Left T1CRBLM* (Figure 4A).

3.4 Longitudinal influence of the cerebellum on future reading ability

To examine whether cerebellar activation during the early stages of reading acquisition predicts future reading, using the 43 data-sets of participants who returned for the follow-up behavioral testing one year later, we conducted a similar voxel-based brain and behavior correlation analysis. Interestingly, we did not observe any significant correlation between activation in the left cerebellum at Time 1 and reading ability at Time 2. Instead, we observed that activation in right cerebellar lobule VII extending to right cerebellar VIII significantly and positively correlated with reading ability at Time 2. From hereon we call this cluster with peak coordinates: $x = 14$, $y = -76$, $z = -42$, $p = .04$, cluster size = 221, $Z_{\max} = 3.95$, *Right T2CRBLM* (Figure 3A, D). Even after we excluded an extreme outlier (with 4 standard deviations from the group mean; $x = 14$, $y = -76$, $z = -42$, $p = .005$ cluster size = 221, $Z_{\max} = 4.16$, Figure 3E) or regressed out reading ability at Time 1 ($x = 14$, $y = -76$, $z = -40$, $p = .046$, cluster size = 217, $Z_{\max} = 3.78$), this result remained significant.

Notably, as we presented before, the list of items is not homogeneous in LWID. The first 13 items require children to point to spoken letters (from item 1 to item 6), to name letters (from item 7 to item 10), and to point to spoken words (from item 10 to item 13), whereas the remaining items require word reading (from item 14). Accordingly, the participants who scored under 14 could only point to or name letters while those scoring at or above 14 were able to read at least some frequent words. In the present study (Figure 1), 21 out of 80 children could only identify letters at Time 1. In contrast, no child read fewer than 3 words at Time 2. Therefore, reading performance at Time 1 refers to either letter or word reading while reading performance at Time 2 refers exclusively to word reading. As a result, the positive relationship between activation in right cerebellar lobule VII and reading ability at Time 2 might be a reflection of the word reading effect rather than a longitudinal effect. To address this issue, the whole sample was split into two groups with a cut-off score of 14: the first one consisting of 21 children who mainly processed letters, and the second one including 59 children who were able to process words at Time 1. If the negative correlation between *Left T1CRBLM* and T1 reading represents a letter reading effect, and the positive correlation between *Right T2CRBLM* and T2 reading represents a word reading effect, then we should observe a negative relationship between T1 reading ability and left cerebellar activity among pre-readers, and a positive relationship between reading ability and right cerebellar activation among more advanced

readers. However, this expected effect was absent, suggesting that the observed hemispheric dissociation in cerebellar activation was not due to differences in the process measured (the left cluster, *Left T1CRBLM*, primarily due to letter, and the right, *Right T2CRBLM*, primarily due to word processing), but rather due to developmental shifts.

As a complementary analysis, we also examined the relationship between activation in the cerebellum and reading gain across time (difference of raw scores between LWID at Time 2 and that at Time 1) with age, gender, and handedness as covariates. If the positive correlation between Time 2 reading and activation in the *Right T2CRBLM* represents a longitudinal effect, we should observe a similar effect between activation in the right cerebellum and reading gain. The result showed that activation in the right cerebellum was indeed positively correlated with reading gain ($x = 14$, $y = -76$, $z = -40$, $p < .001$, cluster size = 625, $Z_{\max} = 3.91$; $x = -22$, $y = -80$, $z = -48$, $p < .005$, cluster size = 357, $Z_{\max} = 3.81$; $x = 16$, $y = -52$, $z = -12$, $p = .045$, cluster size = 211, $Z_{\max} = 4.39$), and these clusters partially overlapped with the *Right T2CRBLM* (Figure 4B).

These results together support the idea that brain activity in the right cerebellum in the early stages of reading may be critical for future reading ability. More importantly, these results suggest that the left and right cerebellum differentially contribute to predicting children's concurrent as opposed to future reading ability. Specifically, activation in the right cerebellar lobule VII at baseline was associated with future reading outcomes, approximately one-year after the start of formal reading acquisition.

Table 4. Cerebro-cerebellar connections decoded by Neurosynth.

Peaks	MNI coordinates			L/R	FCs with the cerebrum (<i>r</i> > 0.2)
	x	y	z		
Regions in the left cerebellum that negatively and significantly correlated with LWID at Time 1 (<i>Left T1CRBLM</i>):					
First peak (located in left cerebellar VII)	-6	-84	-30	R	Angular gyrus; inferior temporal gyrus; middle frontal gyrus; inferior frontal gyrus; inferior frontal sulcus; superior frontal gyrus.
Second peak (located in left cerebellar lobule VIII)	-20	-68	-42		---
Regions in the right cerebellum that positively and significantly correlated with LWID at Time 2 (<i>Right T2CRBLM</i>):					
First peak (located in right cerebellar lobule VII)	14	-76	-42		---
Second peak (located in right cerebellar lobule VIII)	22	-72	-46	L	Inferior frontal gyrus; prefrontal gyrus; superior temporal gyrus

Figure 3. Brain (Mean % signal change) and behavior (LWID) relationship. LWID = Woodcock-Johnson letter word identification subtest. *Left T1CRBLM* = regions in the left cerebellum that negatively and significantly correlated with reading ability at Time 1; *Right T2CRBLM* = regions in the right cerebellum that positively correlated with reading ability at Time 2. (A) locations of *Left T1CRBLM* and *Right T2CRBLM*. (B) Scatter plot to visualize the significant negative relationship between the mean % signal change of *Left T1CRBLM* and LWID score at Time 1 ($n = 80$, corresponding to statistical analysis in section 3.3). (C) Topographic relationship between the regions that significantly activated during the reading fMRI task (yellow), *Left T1CRBLM* (blue; overlap with activation shown in light blue), and *Right T2CRBLM* (red; overlap with activation shown in light red). (D) Scatter plot to visualize the significant positive relationship between the mean % signal change of *Right T2CRBLM* and LWID score at Time 2 ($n = 43$, corresponding to statistical analysis in section 3.4). Dot in the black circle was an extreme value with almost 4 standard deviations from the group mean.

Figure 4: Topographic relationship between different analyses. (A) Regions in the cerebellum that show negative correlations with reading ability at Time 1 within the whole sample ($n = 80$) are shown in red, and within children who returned at Time 2 are shown in blue ($n = 43$). Overlapping regions are shown in pink. (B) Regions in the cerebellum that show positive correlations with reading ability at Time 2 are shown in red, and with reading gain across time are shown in blue. Overlapping regions are shown in pink.

3.5 Deciphering the function of the cerebellum using FC

3.5.1 FC between the cerebellum and cerebral regions based on task imaging data

Differential relationships with concurrent and future reading abilities suggest unique functional roles of cerebellar hemispheres in reading. Here, we further inferred the roles of *Left T1CRBLM* and *Right T2CRBLM* by estimating their FC patterns with the cerebral regions based on task imaging data. We focused on 11 cerebral ROIs within the reading network (see Table 1). The role of the cerebellum was presumed by the ROI that the cerebellum showed relatively greater connectivity bias.

We found that *Left T1CRBLM* was significantly functionally connected with the R.SPL ($t(79) = 7.8$, $p < .001$; FDR corrected), the L.SPL ($t(79) = 5.1$, $p < .001$; FDR corrected), and the L.FG (L.FG_b, $t(79) = 4.0$, $p = .001$; FDR corrected), see Figure 5B. We also observed that *Right T2CRBLM* showed significant bias in FC with the R.SPL ($t(42) = 2.6$, $p = .012$; FDR corrected), and the L.SPL ($t(42) = 3.2$, $p = .002$; FDR corrected), see Figure 5B.

When comparing the relative FCs indices seeded from *Left T1CRBLM* and *Right T2CRBLM*, we found that R.SPL ($t(42) = -3.6$, $p < .001$; FDR corrected) and L.IOG ($t(42) = -2.4$, $p = .019$; FDR corrected) showed greater connectivity with *Left T1CRBLM* compared to *Right T2CRBLM* (Figure 5B). On the other hand, L.STG ($t(42) = 2.4$, $p = .007$; FDR corrected), L.IFG ($t(42) = 3.3$, $p = .002$; FDR corrected), and L.MFG ($t(42) = 2.5$, $p = .018$; FDR corrected) had significantly higher connectivity with *Right T2CRBLM* than with *Left T1CRBLM* (Figure 5B).

Figure 5. FC patterns during the task. FC = functional connectivity. L.MFG = left middle frontal gyrus, L.IFG = left inferior frontal gyrus, L.PostG = left postcentral gyrus; L.STG = left superior temporal gyrus; L.SPL = left superior parietal lobule; L.FG = left fusiform gyrus; L.IOG = left inferior occipital gyrus; R.SPL = right superior parietal lobule. (A) ROIs associated with reading based on Murphy et al., (2019). Blue dots represent regions that had greater FC with *Left T1CRBLM*. Red dots represent regions that have greater FC with *Right T2CRBLM*. (B) Visualization of the strength of FC between *Left T1CRBLM* and with *Right T2CRBLM* and regions within the reading network. The value in each unit represents the difference between the FC from the cerebellum to a given ROI and the mean FC from the cerebellum to the remaining ROIs. This index indicates a cerebellar cluster to be relatively more connected to a given cerebral reading ROI relative to the remaining reading ROIs. Dots represent regions that showed significantly different FC with *Left T1CRBLM* and with *Right T2CRBLM*. *** $p < .001$, FDR corrected; ** $p < .01$, FDR corrected; * $p < .05$, FDR corrected.

3.5.2 FC between the cerebellum and the cerebrum based on the Neurosynth

To further estimate the role of the left and right cerebellum in reading, we used the Neurosynth to decode the function of *Left T1CRBLM* (blue color cluster in Figure 3A, Result Section 3.2) and *Right T2CRBLM* (red color cluster in Figure 3A and Result Section 3.3), respectively. For each of the two clusters, we extracted the location of their first two peaks. The locations were for the *Left T1CRBLM*: cerebellar lobule VII (Table 4, Figure 6, blue dot, L1), and cerebellar lobule VIII (Table 4, Figure 6, blue dot, L2); and for the *Right T2CRBLM*: cerebellar lobule VII (Table 4, Figure 6, red dot, R1), and cerebellar lobule VIII (Table 4, Figure 6, red dot, R2).

The four peaks identified above in *Left T1CRBLM* and *Right T2CRBLM* were further used as seeds to identify cerebral regions that were functionally connected with these peaks using F1000 data in Neurosynth. Results showed that the first peak (L1) in the left cerebellum (that predicted Time 1 reading) positively correlated with right angular gyrus, right frontal regions, and right inferior temporal gyrus (Figure 6). We did not observe significant FC between the second peak (L2) in the left cerebellum cluster and any cerebral regions.

In the right cerebellum (that predicted Time 2 reading), we found the second peak (R2) to correlate with phonological regions: including left inferior frontal gyrus, left precentral gyrus/inferior frontal gyrus, and left supramarginal gyrus (Figure 6).

Notably, FC patterns decoded by Neurosynth (Figure 6) were consistent with FC patterns observed during tasks (Figure 5). Specifically, based on Neurosynth, *Left T1CRBLM* connected significantly to the right angular gyrus, which overlapped with R.SPL where activation was observed during our fMRI task (Figure 7). On the other hand, *Right T2CRBLM* connected significantly to the left precentral gyrus in Neurosynth, and largely corresponded to L.IFG observed during our fMRI task (Figure 7)

3.5.3 Functional attributions of R.SPL and L.IFG via Neurosynth

Given that left and right cerebellar lobules VII showed a biased functional connection pattern with R.SPL and L.IFG, respectively, which were supported by results from Neurosynth, we then inferred the roles of left and right cerebellar lobule VII by decoding the role of R.SPL and L.IFG with the Neurosynth “location based analysis” feature. The central coordinates of R.SPL ($x = 34$, $y = -56$, $z = 50$, Table 1), as defined by the Murphy et al. (2019) meta-analysis, were first entered into Neurosynth. The closest available location ($x = 34$, $y = -56$, $z = 48$, Neurosynth coordinates) to these central coordinates offered the following top 12 cognitive terms: “task”, “working memory”, “symbolic”, “calculation”, “arithmetic”, “task difficulty”, “preparation”, “interference”, “memory”, “navigation”, “verbal”, and “maintenance”. A similar method was applied to L.IFG. The first 12 cognitive terms associated with the closest available location ($x = -44$, $y = 6$, $z = 24$) in Neurosynth to the central coordinates of left inferior frontal gyrus as defined by Murphy et al. (2019) ($x = -44$, $y = 6$, $z = 26$, Table 1), included “phonological”, “orthographic”, “lexical”, “reading”, “tasks”, “words”, “demands”, “word”, “task”, “fluency”, “semantic”, and “language”.

Thus, the results suggest the left and right cerebellar regions connected with R.SPL and L.IFG may play differential roles during reading acquisition and influence reading via different neural circuits.

Figure 6. FC patterns decoded by Neurosynth. Regions in the cerebral cortex that functionally connected with the peaks in the left and right cerebellum clusters, which correlated with reading ability at Time 1 and Time 2, respectively. R.AG = right angular gyrus; R.ITG = right inferior temporal gyrus; R.MFG = right middle frontal gyrus; R.IFG = right inferior frontal gyrus; R.IFS = right

inferior frontal sulcus; R.SFG = right superior frontal gyrus. Threshold to identify these regions was set as voxel-level correlation coefficient larger than 0.2, cluster volume larger than 200 mm³.

Figure 7. Overlap in FC patterns. FC = functional connectivity; Yellow regions in the cerebellum represent *Left T1CRBLM*, which positively correlated with the region represented by the yellow dot (R.SPL) in the cerebrum during the task. Green regions in the cerebellum represent *Right T2CRBLM*, which positively correlated with the region presented by the green dot (L.IFG) in the cerebrum during the task. The blue dot in the cerebellum represents the first peak of *Left T1CRBLM*, which demonstrated FC with blue regions in the cerebrum as decoded by Neurosynth. The red dot in the cerebellum represents the second peak of *Right T2CRBLM*, which demonstrated FC with the red region in the cerebrum as decoded by Neurosynth. R.SPL = right superior parietal lobule; L.IFG = left inferior frontal gyrus.

4 Discussion

In the current study, we examined how the cerebellum contributes to reading in the early stages of reading acquisition. To address this issue, we tested 80 kindergartners who participated in an fMRI experiment, during which the children performed a visual word matching task. One year after their first visit, the reading level of 43 out of the 80 children was tested again. First, we observed significant bilateral cerebellar activation during this task. Activation in left cerebellar lobule VII significantly and negatively predicted reading ability at Time 1, whereas activation in right cerebellar lobule VII significantly and positively predicted reading ability at Time 2. These results suggest that bilateral cerebellar lobules VII play differential roles in early reading. In order to further examine how the left and right cerebellar lobules VII differentially contribute to reading, we investigated the FC pattern with the cerebral cortex using the current task data and resting state data implemented in Neurosynth. We found that left and right cerebellar lobules VII had greater FC with the R.SPL and L.IFG both in task state and resting state. Brain-behavior correlation analysis and FC pattern with the cerebral cortex jointly suggest a division of labor of bilateral cerebellum in the earliest stage of reading acquisition. These results offer insights into how the cerebellum is engaged in the early stages of reading development. In the following sections, we elaborate on three critical findings.

4.1 Functional dissociation of the left and right cerebellar lobules VII in reading

Our study found that the left and right cerebellar hemispheres were involved in reading differently. We observed significant bilateral cerebellar lobules VI/VII activation in the visual word matching task during voxel-based activation analyses. This is consistent with previous findings that the cerebellar VI/VII is responsive to reading tasks (Gao et al., 2015; Martin et al., 2015; Peng et al., 2003). However, we further observed that activation in left cerebellar lobule VII negatively and significantly correlated with current reading ability, measured at the beginning of kindergarten, whereas right cerebellar lobule VII positively and significantly predicted future reading ability, measured one year later. These results suggest that beginning readers in our study might have tended to reduce the reliance on the left cerebellum and might have depended more on the right cerebellum during reading acquisition.

Different brain and behavior relationships suggest that bilateral cerebellar lobules VII play differential roles in reading. To identify the role of the cerebellum, we investigated the cerebro-cerebellar connection patterns. To estimate these, for each cerebral reading ROI, FC from cerebellar clusters (mean percent signal change of all voxels) to all other cerebral reading ROIs (weighted for the number of ROIs) were subtracted from cerebellar FC to the given reading ROI, and examined for statistical significance. This index indicates a cerebellar cluster to be relatively more connected to a given cerebral reading ROI relative to the remaining reading ROIs. The results showed that both left and right cerebellar clusters were significantly more connected with the right superior parietal lobule (R.SPL) and left occipital gyrus, relative to other cerebral areas. The left inferior occipital gyrus is a region known for lower-level visual processing, such as extracting elementary lines, curves, or surfaces from the input word (Dehaene, 2009). This likely explains why bilateral cerebellar hemispheres showed significantly greater connectivity with this region in a visual word matching task. Another brain region where we found significantly greater connectivity with bilateral cerebellar clusters was the R.SPL, also known to be important during visual matching tasks that require participants to judge whether two successive strings (e.g., F T S H G and S T F H G) are identical or not (Reilhac, Peyrin, Démonet, & Valdois, 2013). This region was considered as part of the fronto-parietal attentional network, which allocates attentional resources to the current task (Maidan, Jacob, Giladi, Hausdorff, & Mirelman, 2019; Moore, Porter, & Weissman, 2009). Studies have shown that dyslexics with deficits in visual attention span show reduced activity in this region (Lobier, Peyrin, Pichat, Le Bas, & Valdois, 2014). As a complementary analysis, we performed a data driven decoding via Neurosynth's "location based analysis" feature, which provides relative quantitative functional attributions of locations of interest based on a large database with a sample size of 1000 adults. The closest available Neurosynth coordinates to the central coordinates of R.SPL were associated with terms related to "mental effort", such as "working memory", "task difficulty", "preparation", "interference", and "maintenance", thus suggesting that functional attribution of R.SPL to reading may largely be attributable to mental effort. This was consistent with a previous study, which showed that bilateral SPLs are more strongly activated when processing multiple characters (such as letters, digits, pseudo-letters, Hiragana) (e.g., D F K M V) compared to single characters (which were surrounded by four pound signs to match in length) (e.g., # # D # #) (Lobier, Peyrin, Le Bas, & Valdois, 2012). In the current study, since all participants were beginning readers, they may have required more mental effort during a word processing task.

In line with this interpretation, a study reported SPL activation at the very beginning of literacy acquisition (Braga et al., 2017). Together, stronger connectivity with left occipital gyrus and R.SPL compared to other reading ROIs suggests the cerebellum to be functionally connected to cerebral cortical regions that are likely involved in lower-level visual processing of input words and mental effort required in early reading acquisition.

We then compared the relative FC index associated with each cerebral reading ROI seeded from the left and right cerebellar lobules VII. Interestingly, we observed that compared to right cerebellar lobule VII, the left cerebellar lobule VII had significantly greater FC with R.SPL. This finding was further verified in the F1000 functional connectome analysis (Buckner, Krienen, Castellanos, Diaz, & Yeo, 2011), which is integrated into Neurosynth. This analysis showed that one peak in the left cerebellar cluster functionally connected with right angular gyrus within the R.SPL. These results further indicate that the left cerebellar cluster was more likely to be associated with mental effort during visual word processing. In contrast, compared to left cerebellar lobule VII, right cerebellar lobule VII showed significantly increased FC with L.IFG, which overlapped with the left precentral gyrus reported by Neurosynth. While the role of the inferior frontal gyrus in language processing is complex, it is known to be involved with phonological manipulation (Meng et al., 2016), articulation (Hancock, Richlan, & Hoeft, 2017), semantic and syntactic processing (Hagoort, 2016), and cognitive control (Fedorenko, Duncan, & Kanwisher, 2012). To quantitatively decode the role of the particular cluster in the inferior frontal region, we also used the “location based analysis” feature in Neurosynth. The results showed that the top 12 cognitive terms associated with the available location in Neurosynth closest to the central coordinates of L.IFG were related broadly to reading (e.g., “phonological”, “orthographic”, “lexical”, “reading”, “words”, “fluency”, “semantic”, and “language”). This result indicates that the L.IFG observed in the current study may be more likely to be associated with reading. A stronger connection to this region suggests the FC from right cerebellar lobule VII to cerebral cortical regions reflect reading-related (e.g., phonological) processes.

Functional differentiation in the bilateral cerebellum observed in the current study was consistent with recent functional parcellation analyses of the cerebellum (Guell, Schmahmann, Gabrieli, & Ghosh, 2018; King et al., 2019). King et al. (2019) and Guell et al. (2018) used different tasks to delineate functional subregions in the cerebellum. The left cerebellar region whose activity negatively and significantly correlated with reading ability at Time 1 in our study overlapped with the functionally defined regions from Guell et al.’s parcellation involved in social, motor, and working memory tasks, and from King et al.’s parcellation that implicated attention, emotion, and language processing. In contrast, the right Time 2 reading-related region in our study overlapped with the regions from Guell et al.’s functional subdivisions associated with language and social processing, and from King et al.’s parcellation that implicated word comprehension and verbal fluency.

In short, FC patterns suggest that left and right cerebellar lobules VII might be associated with mental effort and reading-related processes (e.g., phonological processing), respectively. The division of labor of bilateral cerebellar hemispheres during reading contributes to the interpretation of reversed brain and behavior relationships observed in the left and right cerebellum. As reading abilities increase, regions responsible for mental effort (i.e., left cerebellar lobule VII) might decrease their activity during reading. Whereas the functional properties of regions responsible for reading-related processes (i.e., right cerebellar lobule VII)

in the early reading stages might positively contribute to future reading. Together, these results indicate that the left cerebellum may perform a scaffolding function in the early stage of reading acquisition, whereas the right cerebellum could be a contributor to future reading achievements.

4.2 Lateralization pattern during reading in the cerebellum

Previous studies showed a right lateralization pattern of language processing in the cerebellum (Hubrich-Ungureanu, Kaemmerer, Henn, & Braus, 2002; Jansen et al., 2005; Sokolov Arseny, Miall R, & Ivry Richard, 2017). For example, the right cerebellum was engaged in language processing in verb generation (Frings et al., 2006). In addition, the right cerebellum showed consistent activation in phonological processing and connected with the phonological processing network (Alvarez & Fiez, 2018; Tan et al., 2005). More importantly, reading impairment was associated with functional and structural abnormality in the right cerebellum (Hancock et al., 2017; Pernet, Poline, Demonet, & Rousselet, 2009).

In the current study, we observed bilateral cerebellar activation during visual word processing in beginning readers. However, with increase in reading ability, suppression of the left cerebellar activity (i.e., a significantly negative relationship between the left cerebellar lobule VII and reading ability), and increase in right cerebellar activity (i.e., a positive relationship between the right cerebellar lobule VII and future reading ability) may facilitate the reading network shifting from bilateral to right-lateralized (in the cerebellum).

This gradual establishment of hemispheric dominance for language processing has also been reported in the cerebrum. Different from the cerebellum, language processing is characterized by leftward asymmetry in the cerebrum (Cattinelli et al., 2013; Houdé, Rossi, Lubin, & Joliot, 2010; Knecht et al., 2000; Murphy et al., 2019). However, early in development, language processing has been shown to be bilateral (Holland et al., 2001). Everts et al. (2009) showed that functional lateralization strengthens over time and pointed to the dynamics of the process: emerging from an initially bilateral pattern, the cognitive functions in the language domain were shown to develop toward a specialized unilateral network. Similarly, increasing language lateralization during childhood was reported by Szaflarski, Holland, Schmithorst, and Byars (2006), and Ressel, Wilke, Lidzba, Lutzenberger, and Krägeloh-Mann (2008). Activation patterns including the right-hemisphere have been shown to be beneficial for the early stages of language learning even among adults (Kepinska, de Rover, Caspers, & Schiller, 2017; Prat, Keller, & Just, 2007). Our data seem to be consistent with the findings from the cerebrum that language processing in the beginning stages is bilateral in both cerebellum and cerebrum, but will shift to the right cerebellum and left cerebrum with increasing proficiency.

4.3 Implications for reading impairment

There has been a long debate about whether abnormality in the cerebellum was a cause or a consequence of reading disorders (Bishop, 2002). Some studies suggest that deficit in the cerebellum is the cause of reading impairment (Nicolson et al., 2001). As proposed by the Cerebellar Deficit Hypothesis, deficits in the cerebellum before reading acquisition lead to future reading impairment via articulation and automatization (Nicolson et al., 2011). However, this hypothesis is inconsistent with Zeffiro and Eden (2001), who argued that if the cerebellar deficits result in reading impairment, reading disorders should show classical cerebellar clinical

syndrome, such as disorders of the rate or regularity of voluntary movement. Additionally, why did patients with cerebellar damage not show disorders in reading or phonological processing? As an alternative, they proposed that deficits in the cerebellum could only be a bystander in developmental dyslexia (Zeffiro & Eden, 2001).

In recent years, a mounting number of studies comparing brain activation patterns between dyslexic and normal readers have been conducted aiming to investigate the role of the cerebellum in reading impairment, during which both hypo-activation (Cullum, Hodgetts, Milburn, & Cummine, 2019; Meng et al., 2016; Nicolson et al., 1999) and hyper-activation (Feng et al., 2017; Hancock et al., 2017; Patael et al., 2018; Yang, Bi, Long, & Tao, 2013) in the cerebellum were reported in dyslexics. However, these studies can hardly answer the causality issue (Li, Tao, Peng, & Ding, 2017; Xia, Hancock, & Hoeft, 2017).

One possible way to address this issue is to consider pre-readers. If a deficit in the cerebellum is observed in these children who have not received formal reading instruction, it might suggest that an abnormality in the cerebellum is the cause rather than the consequence of reading disorders (Ozernov-Palchik & Gaab, 2016a, 2016b; Raschle, Chang, & Gaab, 2011). Raschle, Zuk, and Gaab (2012) compared children with a family risk for dyslexia (FHD⁺) and without a family risk for dyslexia (FHD⁻), and found that FHD⁻ had greater activation in the bilateral cerebellar regions compared to FHD⁺, suggesting that the cerebellar deficits could be a cause for future reading deficits. Another way is to conduct a longitudinal study. Longitudinal effects of the cerebellum on reading might also be evidence for causal relationships. Studies found that the integrity of the left inferior cerebellar peduncle at age six could predict reading ability at age eight (Borchers et al., 2019; Bruckert et al., 2019). In addition, reading interventions also significantly increased the gray matter volume of the right cerebellum (Krafnick, Flowers, Napoliello, & Eden, 2011).

In the current study, we also used a longitudinal design, but we focused on brain activity during reading. We found that activation in the right cerebellum positively and significantly predicted reading performance after receiving a one-year formal reading instruction. Overall, our study supports the causal account assuming that the functional properties of the cerebellum might contribute to future reading achievement. Our study offers direct evidence that a functional deficit in the cerebellum might impair future reading ability.

4.4 Limitations

Although informative as to the role of the cerebellum in early reading, the present study has two limitations. This first limitation has to do with the reading ROIs, which were not based on pediatric studies. To define ROIs, meta-analytic studies have advantages in providing a consistent and reliable location for reading related regions. However, until now, few meta-analyses have ever focused on beginning readers, especially for children from 5 to 7 years old. Martin et al. (2015) investigated the reading network associated with children reading, but the age range, from 7 to 12 years old, was still inconsistent with the current study. As an alternative, we selected ROIs from a meta-analytic study conducted by Murphy et al. (2019). The reasons are twofold. First, it is a very recent study focusing on single word reading, which is similar to our task. Second, this study focused on native English speakers which were the same as the current study. However, this meta-analysis was not specific for beginning reading. Similar to this limitation, we used FC patterns generated by Neurosynth to supplement our results, which

pattern was based on 1000 adult readers, rather than children's imaging data. The use of pediatric meta-analyses of reading in future studies will therefore be an appropriate extension of the current results.

The second limitation was the visual word matching task that we used in the scanner, which served as an implicit reading task rather than an explicit reading task. We performed this task to ensure that it was accessible to beginning readers. In addition, implicit word reading tasks are frequently used to detect reading related brain activity in beginning readers. For example, Dehaene-Lambertz, Monzalvo, and Dehaene (2018) used a passive word view task to investigate the emergence of visual word form area in children of ages 5 to 6 years old, which was similar to Feng et al. (2020) in examining neural correlates of dyslexic children (8 to 12 years of age) across languages. One recent study used a task to detect whether a given word has a tall feature (e.g., "eaten" has one letter that is tall, i.e., a tall feature, whereas "manor" has no such feature), which was also considered as an implicit reading task (Ashburn, Flowers, Napoliello, & Eden, 2020), and has been shown to elicit activation in phonological areas, such as left superior temporal cortex (Evans, Flowers, Luetje, Napoliello, & Eden, 2016; Turkeltaub, Gareau, Flowers, Zeffiro, & Eden, 2003). In the current study, we also observed activation in this region in the visual word matching task. However, for non-readers, they may perform these tasks simply using visual perception. Future studies with explicit reading tasks are required to verify the current results.

5 Conclusion

In the current study, we investigated the roles of the cerebellum in the early stages of reading acquisition. We found that bilateral cerebellar hemispheres are involved in reading at the onset of schooling. Moreover, the cerebellum might have a longitudinal influence on reading development. Interestingly, we observed functional differentiation of the left and right cerebellar lobules VII in reading. Specifically, the left cerebellar lobule VII negatively and significantly correlated with reading ability and had greater functional connectivity with the right parietal lobule possibly underlying mental effort. In contrast, right cerebellar lobule VII predicted reading ability over time and had stronger connectivity with regions responsible for phonological processing, suggesting that right cerebellar lobule VII lays the foundation for reading development and is associated with reading related processing (e.g., phonological processing). This study is the first to test whether and how the cerebellum contributes to early reading. More importantly, this study lays the foundation to advance future research that specifically investigates the functional differentiation of the cerebellum as one learns to read.

Acknowledgments

This work was supported by grants from the National Institutes of Health/National Institute of Child Health and Human Development (NIH/NICHHD R01HD078351). It was also supported by a grant from the China Scholarship Council (CSC, No. 201806040171) and by Project funded by China Postdoctoral Science Foundation (Grant No. 2020M682846). We are grateful to two anonymous reviewers for their helpful and insightful feedback on an earlier version of the manuscript.

Open practice

No part of the study procedures or the study analyses was pre-registered prior to the research being conducted. We report how we determined our sample size, all data exclusions (if any), all inclusion/exclusion criteria, whether inclusion/exclusion criteria were established prior to data analysis, all manipulations, and all measures in the study. Public archiving of anonymised study data is not permitted by the ethics approval. Readers seeking access to the data should contact the corresponding author via email (fumiko.hoeft@uconn.edu). Data will be released to researchers on the following conditions: approval from the local research ethics committee and with appropriate safeguards to protect from the identification of individuals as well as completion of a request form (<https://drive.google.com/file/d/1kbYYfdomHHrPH6RbE4kV4WDVx1aAhNFn/view?usp=sharing>). Experiment presentation code could be found at [OSF | task materials for CORTEX-D-20-00596R1](#). No custom analysis code was used. Legal copyright restrictions prevent public archiving of the various assessment instruments used in the study (see section 2.2.1). These can be obtained from the copyright holders in the cited references accompanying each test.

Declarations of interest

The authors have no conflicts of interest to declare.

6 References

- Achal, S., Hoeft, F., & Bray, S. (2015). Individual Differences in Adult Reading Are Associated with Left Temporo-parietal to Dorsal Striatal Functional Connectivity. *Cerebral Cortex*, 26(10), 4069-4081. doi: <https://doi.org/10.1093/cercor/bhv214>
- Alvarez, T. A., & Fiez, J. A. (2018). Current Perspectives on the Cerebellum and Reading Development. *Neuroscience & Biobehavioral Reviews*, 92, 55-66. doi: <https://doi.org/10.1016/j.neubiorev.2018.05.006>
- Andersson, J. L., Jenkinson, M., & Smith, S. (2007a). Non-linear registration aka Spatial normalisation FMRIB Technical Report TR07JA2. *FMRIB Analysis Group of the University of Oxford*.
- Andersson, J. L., Jenkinson, M., & Smith, S. (2007b). Non-linear registration, aka spatial normalisation *FMRIB technical report TR07JA2*: Oxford, United Kingdom FMRIB Centre.
- Ang, C., Zhang, J., Chu, M., Li, H., Tian, M., Feng, X., . . . Ding, G. (2020). Intrinsic Cerebro-Cerebellar Functional Connectivity Reveals the Function of Cerebellum VI in Reading-Related Skills. *Frontiers in psychology*, 11, 420.
- Ashburn, S. M., Flowers, D. L., Napoliello, E. M., & Eden, G. F. (2020). Cerebellar function in children with and without dyslexia during single word processing. *Human brain mapping*, 41(1), 120-138. doi: <https://doi.org/10.1002/hbm.24792>
- Beckmann, C. F., & Smith, S. M. (2004). Probabilistic independent component analysis for functional magnetic resonance imaging. *IEEE transactions on medical imaging*, 23(2), 137-152. doi: 10.1109/TMI.2003.822821
- Bishop, D. (2002). Cerebellar abnormalities in developmental dyslexia: cause, correlate or consequence. *cortex*, 38, 491-498. doi: [https://doi.org/10.1016/S0010-9452\(08\)70018-2](https://doi.org/10.1016/S0010-9452(08)70018-2)

- Booth, J. R., Cho, S., Burman, D. D., & Bitan, T. (2007). Neural correlates of mapping from phonology to orthography in children performing an auditory spelling task. *Developmental science*, 10(4), 441-451. doi: <https://doi.org/10.1111/j.14677687.2007.00598.x>
- Borchers, L. R., Bruckert, L., Dodson, C. K., Travis, K. E., Marchman, V. A., Ben-Shachar, M., & Feldman, H. M. (2019). Microstructural properties of white matter pathways in relation to subsequent reading abilities in children: a longitudinal analysis. *Brain Structure and Function*, 224(2), 891-905. doi: <https://doi.org/10.1007/s00429-018-1813-z>
- Braga, L. W., Amemiya, E., Tauil, A., Suguieda, D., Lacerda, C., Klein, E., . . . Dehaene, S. (2017). Tracking adult literacy acquisition with functional MRI: A single-case study. *Mind, Brain, and Education*, 11(3), 121-132.
- Bruckert, L., Borchers, L. R., Dodson, C. K., Marchman, V. A., Travis, K. E., Ben-Shachar, M., & Feldman, H. M. (2019). White matter plasticity in reading-related pathways differs in children born preterm and at term: a longitudinal analysis. *Frontiers in human neuroscience*, 13. doi: <https://doi.org/10.3389/fnhum.2019.00139>
- Buckner, R. L., Krienen, F. M., Castellanos, A., Diaz, J. C., & Yeo, B. T. (2011). The organization of the human cerebellum estimated by intrinsic functional connectivity. *American Journal of Physiology-Heart and Circulatory Physiology*, 106(5), 2322-2345. doi: <https://doi.org/10.1152/jn.00339.2011>
- Cao, F. (2016). Neuroimaging studies of reading in bilinguals. *Bilingualism: Language and Cognition*, 19(4), 683-688. doi: <https://doi.org/10.1017/S1366728915000656>
- Cattinelli, I., Borghese, N. A., Gallucci, M., & Paulesu, E. (2013). Reading the reading brain: A new meta-analysis of functional imaging data on reading. *Journal of Neurolinguistics*, 26(1), 214-238. doi: <https://doi.org/10.1016/j.jneuroling.2012.08.001>
- Chang, L. J., Yarkoni, T., Khaw, M. W., & Sanfey, A. G. (2012). Decoding the role of the insula in human cognition: functional parcellation and large-scale reverse inference. *Cerebral cortex*, 23(3), 739-749. doi: <https://doi.org/10.1093/cercor/bhs065>
- Clark, K. A., Helland, T., Specht, K., Narr, K. L., Manis, F. R., Toga, A. W., & Hugdahl, K. (2014). Neuroanatomical precursors of dyslexia identified from pre-reading through to age 11. *Brain*, 137(12), 3136 - 3141.
- Cullum, A., Hodgetts, W. E., Milburn, T. F., & Cummine, J. (2019). Cerebellar Activation During Reading Tasks: Exploring the Dichotomy Between Motor vs. Language Functions in Adults of Varying Reading Proficiency. *The Cerebellum*, 1-17. doi: <https://doi.org/10.1007/s12311-019-01024-6>
- D'Mello, A. M., & Gabrieli, J. D. (2018). Cognitive neuroscience of dyslexia. *Language, speech, and hearing services in schools*, 49(4), 798-809. doi: https://doi.org/10.1044/2018_LSHSS-DYSLC-18-0020
- Danelli, L., Berlinger, M., Bottini, G., Ferri, F., Vacchi, L., Sberna, M., & Paulesu, E. (2013). Neural intersections of the phonological, visual magnocellular and motor/cerebellar systems in normal readers: Implications for imaging studies on dyslexia. *Human brain mapping*, 34(10), 2669-2687. doi: <https://doi.org/10.1002/hbm.22098>
- De La Vega, A., Chang, L. J., Banich, M. T., Wager, T. D., & Yarkoni, T. (2016). Large-scale meta-analysis of human medial frontal cortex reveals tripartite functional organization.

- Journal of Neuroscience*, 36(24), 6553-6562. doi: <https://doi.org/10.1523/JNEUROSCI.4402-15.2016>
- Dehaene-Lambertz, G., Monzalvo, K., & Dehaene, S. (2018). The emergence of the visual word form: Longitudinal evolution of category-specific ventral visual areas during reading acquisition. *PLoS biology*, 16(3), e2004103.
- Dunn, L. M., & Dunn, D. M. (2007). Peabody Picture Vocabulary Test: Manual (Fourth Ed.). Minneapolis: Pearson.
- Eckert, M. A., Berninger, V. W., Vaden, K. I., Gebregziabher, M., Tsu, L., & Consortium, D. D. (2016). Gray Matter Features of Reading Disability: A Combined Meta-Analytic and Direct Analysis Approach. *eneuro*, 3(1), ENEURO.0103-0115.2015. doi: [10.1523/ENEURO.0103-15.2015](https://doi.org/10.1523/ENEURO.0103-15.2015)
- Evans, T. M., Flowers, D. L., Luetje, M. M., Napoliello, E., & Eden, G. F. (2016). Functional neuroanatomy of arithmetic and word reading and its relationship to age. *Neuroimage*, 143, 304-315.
- Everts, R., Lidzba, K., Wilke, M., Kiefer, C., Mordasini, M., Schroth, G., . . . Steinlin, M. (2009). Strengthening of laterality of verbal and visuospatial functions during childhood and adolescence. *Human brain mapping*, 30(2), 473-483. doi: <https://doi.org/10.1002/hbm.20523>
- Fedorenko, E., Duncan, J., & Kanwisher, N. (2012). Language-selective and domain-general regions lie side by side within Broca's area. *Current Biology*, 22(21), 2059-2062.
- Feng, X., Altarelli, I., Monzalvo, K., Ding, G., Ramus, F., Shu, H., . . . Dehaene-Lambertz, G. (2020). A universal reading network and its modulation by writing system and reading ability in French and Chinese children. *Elife*, 9, e54591.
- Feng, X., Li, L., Zhang, M. L., Yang, X. J., Tian, M. Y., Xie, W. Y., . . . Ding, G. S. (2017). Dyslexic Children Show Atypical Cerebellar Activation and Cerebro-Cerebellar Functional Connectivity in Orthographic and Phonological Processing. *Cerebellum*, 16(2), 496-507. doi: [10.1007/s12311-016-0829-2](https://doi.org/10.1007/s12311-016-0829-2)
- Frings, M., Dimitrova, A., Schorn, C. F., Elles, H.-G., Hein-Kropp, C., Gizewski, E. R., . . . Timmann, D. (2006). Cerebellar involvement in verb generation: an fMRI study. *Neuroscience letters*, 409(1), 19-23. doi: <https://doi.org/10.1016/j.neulet.2006.08.058>
- Gao, y., Wei, N., Wang, Z., Jian, J., Ding, G., Meng, X., & Liu, L. (2015). Interaction between native and second language processing: Evidence from a functional magnetic resonance imaging study of Chinese-English bilingual children. *Acta Psychologica Sinica*, 47(12), 1419-1432.
- Guell, X., Schmahmann, J. D., Gabrieli, J. D., & Ghosh, S. S. (2018). Functional gradients of the cerebellum: a fundamental movement-to-thought principle. *bioRxiv*, 254326.
- Haft, S. L., Kepinska, O., Caballero, J. N., Carreiras, M., & Hoeft, F. (2019). Attentional Fluctuations, Cognitive Flexibility, and Bilingualism in Kindergarteners. *Behavioral Sciences*, 9(5), 58. doi: <https://doi.org/10.3390/bs9050058>
- Hagoort, P. (2016). MUC (Memory, Unification, Control): A model on the neurobiology of language beyond single word processing *Neurobiology of language* (pp. 339-347): Elsevier.

- Hancock, R., Richlan, F., & Hoefft, F. (2017). Possible roles for fronto-striatal circuits in reading disorder. *Neuroscience & Biobehavioral Reviews*, 72, 243-260. doi: <https://doi.org/10.1016/j.neubiorev.2016.10.025>
- Hanson, S. J., & Bunzl, M. (2010). *Foundational issues in human brain mapping*: MIT Press.
- Hartwigsen, G., Baumgaertner, A., Price, C. J., Koehnke, M., Ulmer, S., & Siebner, H. R. (2010). Phonological decisions require both the left and right supramarginal gyri. *Proceedings of the National Academy of Sciences*, 107(38), 16494-16499.
- Hartwigsen, G., Neef, N. E., Camilleri, J. A., Margulies, D. S., & Eickhoff, S. B. (2019). Functional segregation of the right inferior frontal gyrus: evidence from coactivation-based parcellation. *Cerebral Cortex*, 29(4), 1532-1546.
- Holland, S. K., Plante, E., Byars, A. W., Strawsburg, R. H., Schmithorst, V. J., & Ball Jr, W. S. (2001). Normal fMRI brain activation patterns in children performing a verb generation task. *Neuroimage*, 14(4), 837-843.
- Houdé, O., Rossi, S., Lubin, A., & Joliot, M. (2010). Mapping numerical processing, reading, and executive functions in the developing brain: an fMRI meta-analysis of 52 studies including 842 children. *Developmental science*, 13(6), 876-885. doi: <https://doi.org/10.1111/j.1467-7687.2009.00938.x>
- Hubrich-Ungureanu, P., Kaemmerer, N., Henn, F. A., & Braus, D. F. (2002). Lateralized organization of the cerebellum in a silent verbal fluency task: a functional magnetic resonance imaging study in healthy volunteers. *Neuroscience letters*, 319(2), 91-94. doi: [https://doi.org/10.1016/S0304-3940\(01\)02566-6](https://doi.org/10.1016/S0304-3940(01)02566-6)
- Jansen, A., Flöel, A., Van Randenborgh, J., Konrad, C., Rotte, M., Förster, A. F., . . . Knecht, S. (2005). Crossed cerebro-cerebellar language dominance. *Human brain mapping*, 24(3), 165-172. doi: <https://doi.org/10.1002/hbm.20077>
- Jenkinson, M., Bannister, P., Brady, M., & Smith, S. (2002). Improved optimization for the robust and accurate linear registration and motion correction of brain images. *Neuroimage*, 17(2), 825-841.
- Jenkinson, M., & Smith, S. (2001). A global optimisation method for robust affine registration of brain images. *Medical image analysis*, 5(2), 143-156. doi: [https://doi.org/10.1016/S1361-8415\(01\)00036-6](https://doi.org/10.1016/S1361-8415(01)00036-6)
- Johnson, M. H. (2000). Functional brain development in infants: Elements of an interactive specialization framework. *Child development*, 71(1), 75-81. doi: <https://doi.org/10.1111/1467-8624.00120>
- Kaufman, A. S., & Kaufman, N. L. (2004). Kaufman Assessment Battery for Children, Second Edition. *Bloomington, MN: Pearson*.
- Kepinska, O., de Rover, M., Caspers, J., & Schiller, N. O. (2017). On neural correlates of individual differences in novel grammar learning: an fMRI study. *Neuropsychologia*, 98, 156-168. doi: <https://doi.org/10.1016/j.neuropsychologia.2016.06.014>
- King, M., Hernandez-Castillo, C., Poldrack, R., Ivry, R., & Diedrichsen, J. (2019). Functional boundaries in the human cerebellum revealed by a multi-domain task battery. *Nature neuroscience*. doi: <https://doi.org/10.1038/s41593-019-0436-x>
- Knecht, S., Deppe, M., Dräger, B., Bobe, L., Lohmann, H., Ringelstein, E.-B., & Henningsen, H. (2000). Language lateralization in healthy right-handers. *Brain*, 123(1), 74-81. doi: <https://doi.org/10.1093/brain/123.1.74>

- Krafnick, A. J., Flowers, D. L., Napoliello, E. M., & Eden, G. F. (2011). Gray matter volume changes following reading intervention in dyslexic children. *Neuroimage*, 57(3), 733 - 741. doi: [org/10.1016/j.neuroimage.2010.10.062](https://doi.org/10.1016/j.neuroimage.2010.10.062)
- Li, H., Booth, J. R., Feng, X., Wei, N., Zhang, M., Zhang, J., . . . Meng, X. (2020). Functional parcellation of the right cerebellar lobule VI in children with normal or impaired reading. *Neuropsychologia*, 107630.
- Li, H., Tao, W., Peng, D., & Ding, G. (2017). The Causal Relationship between Developmental Dyslexia and Abnormal Brain Regions: Research Paradigms and the Relevant Findings. *Psychological Development and Education*, 33(5), 631-640.
- Linkersdörfer, J., Jurcoane, A., Lindberg, S., Kaiser, J., Hasselhorn, M., Fiebach, C. J., & Lonnemann, J. (2014). The association between gray matter volume and reading proficiency: a longitudinal study of beginning readers. *Journal of Cognitive Neuroscience*, 27(2), 308-318.
- Linkersdörfer, J., Lonnemann, J., Lindberg, S., Hasselhorn, M., & Fiebach, C. J. (2012). Grey matter alterations co-localize with functional abnormalities in developmental dyslexia: an ALE meta-analysis. *PloS One*, 7(8), e43122. doi: <https://doi.org/10.1371/journal.pone.0043122>
- Lobier, M., Peyrin, C., Le Bas, J.-F., & Valdois, S. (2012). Pre-orthographic character string processing and parietal cortex: A role for visual attention in reading? *Neuropsychologia*, 50(9), 2195-2204.
- Lobier, M., Peyrin, C., Pichat, C., Le Bas, J.-F., & Valdois, S. (2014). Visual processing of multiple elements in the dyslexic brain: evidence for a superior parietal dysfunction. *Frontiers in human neuroscience*, 8, 479.
- Maidan, I., Jacob, Y., Giladi, N., Hausdorff, J. M., & Mirelman, A. (2019). Altered organization of the dorsal attention network is associated with freezing of gait in Parkinson's disease. *Parkinsonism & related disorders*, 63, 77-82.
- Mariën, P., Ackermann, H., Adamaszek, M., Barwood, C. H., Beaton, A., Desmond, J., . . . Küper, M. (2014). Consensus paper: language and the cerebellum: an ongoing enigma. *The Cerebellum*, 13(3), 386-410. doi: <https://doi.org/10.1007/s12311-013-0540-5>
- Marks, R. A., Kovelman, I., Kepinska, O., Oliver, M., Xia, Z., Haft, S. L., . . . Hancock, R. (2019). Spoken language proficiency predicts print-speech convergence in beginning readers. *NeuroImage*, 201, 116021. doi: <https://doi.org/10.1016/j.neuroimage.2019.116021>
- Martin, A., Schurz, M., Kronbichler, M., & Richlan, F. (2015). Reading in the brain of children and adults: A meta-analysis of 40 functional magnetic resonance imaging studies. *Human brain mapping*, 36(5), 1963-1981. doi: <https://doi.org/10.1002/hbm.22749>
- Mather, N., Roberts, R., Hammill, D. D., & Allen, E. A. (2008). TOC Examiner's Manual. Austin, Texas: PRO-ED.
- Meng, X. Z., You, H. L., Song, M. X., Desroches, A. S., Wang, Z. K., Wei, N., . . . Ding, G. S. (2016). Neural deficits in auditory phonological processing in Chinese children with English reading impairment. *Bilingualism-Language and Cognition*, 19(2), 331-346. doi: <https://doi.org/10.1017/S1366728915000073>
- Moore, K. S., Porter, C. B., & Weissman, D. H. (2009). Made you look! Consciously perceived, irrelevant instructional cues can hijack the attentional network. *NeuroImage*, 46(1), 270-279.

- Murphy, K., Jogia, J., & Talcott, J. (2019). On the neural basis of word reading: A meta-analysis of fMRI evidence using activation likelihood estimation. *Journal of Neurolinguistics*, 49, 71-83. doi: <https://doi.org/10.1016/j.jneuroling.2018.08.005>
- Myers, C. A., Vandermosten, M., Farris, E. A., Hancock, R., Gimenez, P., Black, J. M., . . . Hendren, R. L. (2014). White matter morphometric changes uniquely predict children's reading acquisition. *Psychological science*, 25(10), 1870 - 1883.
- Nicolson, R. I., Fawcett, A. J., Berry, E. L., Jenkins, I. H., Dean, P., & Brooks, D. J. (1999). Association of abnormal cerebellar activation with motor learning difficulties in dyslexic adults. *The Lancet*, 353(9165), 1662-1667. doi: [https://doi.org/10.1016/S0140-6736\(98\)09165-X](https://doi.org/10.1016/S0140-6736(98)09165-X)
- Nicolson, R. I., Fawcett, A. J., & Dean, P. (2001). Developmental dyslexia: the cerebellar deficit hypothesis. *Trends in neurosciences*, 24(9), 508-511. doi: [https://doi.org/10.1016/S0166-2236\(00\)01896-8](https://doi.org/10.1016/S0166-2236(00)01896-8)
- Norton, E. S., Black, J. M., Stanley, L. M., Tanaka, H., Gabrieli, J., D. E., Sawyer, C., & Hoeft, F. (2014). Functional neuroanatomical evidence for the double-deficit hypothesis of developmental dyslexia. *Neuropsychologia*, 61, 235-246. doi: [10.1016/j.neuropsychologia.2014.06.015](https://doi.org/10.1016/j.neuropsychologia.2014.06.015)
- Ozernov-Palchik, O., & Gaab, N. (2016a). Tackling the 'dyslexia paradox': reading brain and behavior for early markers of developmental dyslexia. *Wiley Interdisciplinary Reviews: Cognitive Science*, 7(2), 156-176. doi: <https://doi.org/10.1002/wcs.1383>
- Ozernov-Palchik, O., & Gaab, N. (2016b). Tackling the Early Identification of Dyslexia with the Help of Neuroimaging. *Perspectives on Language and Literacy*, 42(1), 11-17.
- Patael, S. Z., Farris, E. A., Black, J. M., Hancock, R., Gabrieli, J. D., Cutting, L. E., & Hoeft, F. (2018). Brain basis of cognitive resilience: Prefrontal cortex predicts better reading comprehension in relation to decoding. *PloS one*, 13(6), e0198791.
- Peng, D. I., Xu, D., Jin, Z., Luo, Q., Ding, G. s., Perry, C., . . . Liu, Y. (2003). Neural basis of the non-attentional processing of briefly presented words. *Human Brain Mapping*, 18(3), 215-221. doi: <https://doi.org/10.1002/hbm.10096>
- Perfetti, C., Cao, F., & Booth, J. (2013). Specialization and universals in the development of reading skill: How Chinese research informs a universal science of reading. *Scientific Studies of Reading*, 17(1), 5-21. doi: <https://doi.org/10.1080/10888438.2012.689786>
- Pernet, C. R., Poline, J. B., Demonet, J. F., & Rousselet, G. A. (2009). Brain classification reveals the right cerebellum as the best biomarker of dyslexia. *BMC neuroscience*, 10(1), 67. doi: <https://doi.org/10.1186/1471-2202-10-67>
- Prat, C. S., Keller, T. A., & Just, M. A. (2007). Individual differences in sentence comprehension: a functional magnetic resonance imaging investigation of syntactic and lexical processing demands. *J. Cogn. Neurosci.*, 19, 1950-1963. doi: [org/10.1162/jocn.2007.19.12.1950](https://doi.org/10.1162/jocn.2007.19.12.1950)
- Pruim, R. H., Mennes, M., van Rooij, D., Llera, A., Buitelaar, J. K., & Beckmann, C. F. (2015). ICA-AROMA: A robust ICA-based strategy for removing motion artifacts from fMRI data. *Neuroimage*, 112, 267-277. doi: <https://doi.org/10.1016/j.neuroimage.2015.02.064>
- Raschle, N. M., Chang, M., & Gaab, N. (2011). Structural brain alterations associated with dyslexia predate reading onset. *Neuroimage*, 57(3), 742 - 749. doi: <https://doi.org/10.1016/j.neuroimage.2010.09.055>

- Raschle, N. M., Zuk, J., & Gaab, N. (2012). Functional characteristics of developmental dyslexia in left-hemispheric posterior brain regions predate reading onset. *Proceedings of the National Academy of Sciences*, 109(6), 2156-2161. doi: <https://doi.org/10.1073/pnas.1107721109>
- Reilhac, C., Peyrin, C., Démonet, J.-F., & Valdois, S. (2013). Role of the superior parietal lobules in letter-identity processing within strings: fMRI evidence from skilled and dyslexic readers. *Neuropsychologia*, 51(4), 601-612.
- Ressel, V., Wilke, M., Lidzba, K., Lutzenberger, W., & Krägeloh-Mann, I. (2008). Increases in language lateralization in normal children as observed using magnetoencephalography. *Brain and language*, 106(3), 167-176. doi: <https://doi.org/10.1016/j.bandl.2008.01.004>
- Rubin, T. N., Koyejo, O., Gorgolewski, K. J., Jones, M. N., Poldrack, R. A., & Yarkoni, T. (2017). Decoding brain activity using a large-scale probabilistic functional-anatomical atlas of human cognition. *PLoS computational biology*, 13(10), e1005649. doi: <https://doi.org/10.1371/journal.pcbi.1005649>
- Sörös, P., Hoxhaj, E., Borel, P., Sadohara, C., Feige, B., Matthies, S., . . . Philipsen, A. (2019). Hyperactivity/restlessness is associated with increased functional connectivity in adults with ADHD: a dimensional analysis of resting state fMRI. *BMC psychiatry*, 19(1), 43. doi: <https://doi.org/10.1186/ISRCTN12722296>
- Saxe, R., Brett, M., & Kanwisher, N. (2006). Divide and conquer: a defense of functional localizers. *Neuroimage*, 30(4), 1088-1096. doi: <https://doi.org/10.1016/j.neuroimage.2005.12.062>
- Schrank, F., Mather, N., & McGrew, K. (2014). Woodcock-Johnson IV Tests of Oral Language. *Rolling Meadows, IL: Riverside*.
- Smith, S. M. (2002). Fast robust automated brain extraction. *Human brain mapping*, 17(3), 143-155.
- Sokolov Arseny, A., Miall R, C., & Ivry Richard, B. (2017). The Cerebellum: Adaptive Prediction for Movement and Cognition. *Trends in Cognitive Sciences*, 21(5), 313-332. doi: <https://doi.org/10.1016/j.tics.2017.02.005>
- Szaflarski, J. P., Holland, S. K., Schmithorst, V. J., & Byars, A. W. (2006). fMRI study of language lateralization in children and adults. *Human brain mapping*, 27(3), 202-212. doi: <https://doi.org/10.1002/hbm.20177>
- Tan, L. H., Laird, A. R., Li, K., & Fox, P. T. (2005). Neuroanatomical correlates of phonological processing of Chinese characters and alphabetic words: A meta-analysis. *Human brain mapping*, 25(1), 83-91. doi: <https://doi.org/10.1002/hbm.20134>
- Thurston, R. C., Maki, P. M., Derby, C. A., Sejdić, E., & Aizenstein, H. J. (2015). Menopausal hot flashes and the default mode network. *Fertility and sterility*, 103(6), 1572-1578. e1571. doi: <https://doi.org/10.1016/j.fertnstert.2015.03.008>
- Turkeltaub, P. E., Gareau, L., Flowers, D. L., Zeffiro, T. A., & Eden, G. F. (2003). Development of neural mechanisms for reading. *Nature neuroscience*, 6(7), 767-773.
- Wagner, R. K., Torgesen, J. K., Rashotte, C. A., & Pearson, N. A. (2013). Comprehensive Test of Phonological Processing (CTOPP-2): Examiner's Manual (Second). Austin, Texas: PRO-ED.
- Weiss, Y., Cweigenberg, H. G., & Booth, J. R. (2018). Neural specialization of phonological and semantic processing in young children. *Human brain mapping*.

- Xia, Z., Hancock, R., & Hoeft, F. (2017). Neurobiological bases of reading disorder Part I: Etiological investigations. *Language and Linguistics Compass*, 11(4). doi: 10.1111/lnc3.12239
- Yang, Y., Bi, H.-Y., Long, Z.-Y., & Tao, S. (2013). Evidence for cerebellar dysfunction in Chinese children with developmental dyslexia: an fMRI study. *International Journal of Neuroscience*, 123(5), 300-310. doi: <https://doi.org/10.3109/00207454.2012.756484>
- Yarkoni, T., Poldrack, R. A., Nichols, T. E., Van Essen, D. C., & Wager, T. D. (2011). Large-scale automated synthesis of human functional neuroimaging data. *Nature methods*, 8(8), 665. doi: <https://doi.org/10.1038/nmeth.1635>
- Zeffiro, T., & Eden, G. (2001). The cerebellum and dyslexia: perpetrator or innocent bystander? *Trends in Neurosciences*, 24, 512-513.

Credit author statement

Hehui Li: Conceptualization, Methodology, Formal analysis, Writing - Original Draft, Funding acquisition.

Olga Kepinska: Conceptualization, Methodology, Formal analysis, Investigation, Writing - Original Draft, Writing - Review & Editing, Supervision.

Jocelyn N. Caballero: Investigation, Writing - Review & Editing.

Leo Zekelman: Investigation, Writing - Review & Editing.

Rebecca A. Marks: Investigation, Writing - Review & Editing.

Yuuko Uchikoshi: Investigation, Writing - Review & Editing, Project administration.

Ioulia Kovelman: Investigation, Writing - Review & Editing, Project administration.

Fumiko Hoeft: Conceptualization, Methodology, Investigation, Writing - Original Draft, Writing - Review & Editing, Project administration, Supervision, Funding acquisition.