1 **ORIGINAL ARTICLE**

2
3
4 **Visual discrimination of primate species based on faces in**
5 **chimpanzees**
6

7 **Duncan A. Wilson · Masaki Tomonaga**

8
9
10 D. A. Wilson (✉)

11 Primate Research Institute, Kyoto University, Inuyama, Aichi, 484-8506, Japan

12 E-mail: duncan.wilson.76x@st.kyoto-u.ac.jp

13 Phone: (+81) 0568-63-0567

14 Fax: (+81) 0568-63-0085
15

16 M. Tomonaga

17 Primate Research Institute, Kyoto University, Inuyama, Aichi, 484-8506, Japan
18
19

20 © Japan Monkey Centre and Springer Japan KK, part of Springer Nature 2018. This
21 paper is not the copy of record and may not exactly replicate the final, authoritative
22 version of the article. Please do not copy or cite without authors' permission. The final
23 article is available via its DOI: 10.1007/s10329-018-0649-8

Abstract

Many primate studies have investigated discrimination of individual faces within the same species. However, few studies have looked at discrimination between primate species faces at the categorical level. This study systematically examined the factors important for visual discrimination between primate species faces in chimpanzees, including: colour, orientation, familiarity and perceptual similarity. Five adult female chimpanzees were tested on their ability to discriminate identical and categorical (non-identical) images of different primate species faces in a series of touchscreen matching-to-sample experiments. Discrimination performance for chimpanzee, gorilla and orangutan faces was better in colour than in greyscale. An inversion effect was also found, with higher accuracy for upright than inverted faces. Discrimination performance for unfamiliar (baboon and capuchin monkey) and highly familiar (chimpanzee and human) but perceptually different species was equally high. After excluding effects of colour and familiarity, difficulty in discriminating between different species faces can be best explained by their perceptual similarity to each other. Categorical discrimination performance for unfamiliar, perceptually similar faces (gorilla and orangutan) was significantly worse than unfamiliar, perceptually different faces (baboon and capuchin monkey). Moreover, Multidimensional Scaling analysis of the image similarity data based on local feature matching revealed greater similarity between chimpanzee, gorilla and orangutan faces than between human, baboon and capuchin monkey faces. We conclude our chimpanzees appear to perceive similarity in primate faces in a similar way to humans. Information about perceptual similarity is likely prioritised over the potential influence of previous experience or a conceptual representation of species for categorical discrimination between species faces.

- 48 **Keywords** Species discrimination · Colour · Familiarity · Perceptual similarity ·
- 49 Chimpanzees

Introduction

For primates the most important visual cue for gathering social information about another individual is the face (Fujita 1993a; Spencer et al. 1997; Burton et al. 1999). Information about identity, age, gender and emotion can be extracted from the face and facial information leads to the fastest and most accurate identification of an individual (Dahl et al. 2013a; Kano and Tomonaga 2009; Parr 2011; Pascalis and Wirth 2011). The face is also used to discriminate between individuals within the same species and other prey or predator species (Pascalis and Wirth 2011).

Exactly how facial features are processed in human and non-human animals has been a topic of intense study for decades (Bruce and Young 1986; Ge et al. 2009; Valentine 1991). Diamond and Carey (1986) propose that faces are processed according to first and second-order relational features. First-order relational features refer to the eyes, nose and mouth and their fixed position relative to each other, used to discriminate between face and non-face stimuli. Second-order relational features refer to the relative spatial distances between facial features, used to discriminate between the faces of different individuals. Interestingly, when faces are inverted judging distances between their second-order relational features becomes more difficult compared to non-face stimuli. This is known as the ‘inversion effect’ (Yin 1969). The inversion effect is consistently found in humans (Goldstein 1965; Scapinello and Yarmey 1970; Yin 1969) and chimpanzees (Dahl et al. 2013c; Parr et al. 1998; Tomonaga 1999, 2007). Faces are thought to be processed configurally. Configural processing refers to “the emergent features of a face that only become apparent when two or more of its basic features (e.g., the eyes, nose, or mouth) are processed at the same time.” (Piepers and Robbins

2012, p. 2). The integration of this information into a perceptual whole allows individual recognition and discrimination within and between other species (Parr 2011).

A number of primate studies have investigated visual discrimination of species based on whole body images, including the face. Humphrey (1974) conducted one of the earliest experiments on species discrimination in rhesus monkeys using a habituation task. The monkeys could distinguish individuals within their own species but not between species. Fujita (1987, 1990, 1993b) and Fujita and Watanabe (1995) found longer looking durations for own-species images in Japanese macaques and rhesus monkeys using a sensory reinforcement procedure. In addition, Demaria and Thierry (1988) observed stump-tailed macaques looked for longer at own-species images compared to other macaque species images. These experiments show a consistent advantage for processing own-species faces in monkeys.

In contrast, in chimpanzees Tanaka (2003, 2007) found a visual preference for human stimuli over other primate species in free-choice tasks. Tanaka (2003) presented five adult chimpanzees with images of humans, chimpanzees, gorillas, orangutans and other monkey species. Human images were touched more than any other species and preference strength did not differ with phylogenetic distance from chimpanzees. Later, Tanaka (2007) tested eight adult and three infant chimpanzees using the same paradigm and species categories. Consistent with the previous study, adult chimpanzees showed a preference for humans, whilst infants showed a preference for either chimpanzees or no species preference. These differences may be explained by social experience during infancy; the adult chimpanzees were mainly raised by humans, whereas the infants were raised by their mothers. Conspecific social experience during infancy may also account for the own-species preferences found in earlier monkey studies (Fujita 1990, 1993b).

Several studies have also investigated visual discrimination of species based on stimuli featuring only faces. Pascalis and Bachevalier (1998) used a visual paired-comparison (VPC) task to investigate species preference in humans and macaques. In the VPC task a preference for looking at new stimuli is measured after a period of familiarisation with another stimulus. Looking longer at the new stimulus indicates recognition of the familiarised stimulus. An own-species effect across species was found; humans showed a novelty preference for human faces but not monkey faces, whereas monkeys showed the opposite preference. Dufour et al. (2006) examined recognition of different species faces using the VPC task in humans, Tonkean macaques and brown capuchin monkeys and found recognition was limited to own-species faces. Dahl et al. (2007) used an adaptation paradigm with rhesus monkeys and found they were better at identifying individuals within their own species compared to other species and processed the faces configurally. Gothard et al. (2009) found macaques can recognise individual faces of both macaques and humans. However, recognising own-species faces involved configural and feature-based processing, whilst recognising human faces mainly involved feature-based processing. Consistent with discrimination studies involving the whole body and face, own-species effects are also observed with stimuli featuring only the face.

Related to the own-species effect, other studies have focused on the influence of experience and familiarity. An interesting study in chimpanzees by Martin-Malivel and Okada (2007) used a matching-to-sample task (MTS) to investigate the influence of familiarity on categorical perception (CP) of morphed human and chimpanzee faces. CP “occurs when members of a class of stimuli which vary in their sensory characteristics are nevertheless processed as if they are equivalent.” (Campbell et al. 1997, p. 1429).

They found one group of chimpanzees more exposed to human faces than chimpanzee faces showed better discrimination of human than chimpanzee faces, and a CP effect for human faces. In another group of chimpanzees, familiar with both chimpanzees and humans, no discrimination advantage or CP effects for human or chimpanzee faces were found. Also in chimpanzees, Dahl et al. (2013b) investigated the effects of lifetime experience on discriminating individual chimpanzee and human faces using a delayed MTS task. Infant chimpanzees were better at discriminating chimpanzee faces and older chimpanzees were better at discriminating human faces. These results support Tanaka (2003, 2007) in demonstrating the importance of developmental experience in species discrimination in the same group of chimpanzees.

In addition to experience and familiarity, perceptual features are also important for individual and species discrimination. For example, Quinn and Eimas (1996) found three to four month old human infants used the internal features and outer edges of the face to discriminate between cat and dog images. Roberts and Mazmanian (1988) showed humans had greater difficulty in discriminating between pictures of kingfishers and other birds in greyscale compared to colour. Parr et al. (2000) demonstrated chimpanzees were worse at discriminating conspecific faces with masked eyes compared to control stimuli, but rhesus monkeys were worse when both the eyes and mouth were masked. Marsh and MacDonald (2008) studied the classification of great ape, gibbon and monkey, and prosimian faces by orangutans using a two-choice touch screen procedure. Discrimination performance was best for stimuli in colour and featuring the face only, and worst for stimuli in greyscale and with modified eyes, i.e. removed eyes or inserted infant eyes. Together, such visual cues help primates to discriminate between faces.

Although many studies have investigated discrimination of individual faces within the same species, few studies have investigated discrimination between primate species faces at the categorical level, especially in chimpanzees. The aim of the current study was to systematically examine the factors important for visual discrimination between different species faces in chimpanzees using a MTS task. Examining how chimpanzees, our closest living relatives, visually discriminate between primate species faces may give us greater insight into how the human visual system has evolved to perceive similarities and differences in categories of faces. We predicted discrimination accuracy would be higher for faces in colour than greyscale, upright than inverted orientation, highly familiar than unfamiliar species faces, and perceptually different than perceptually similar species faces. In addition, higher discrimination accuracy was predicted for identical compared to categorical (non-identical) faces within the same species.

Methods

Participants and housing

Five adult female chimpanzees (*Pan troglodytes*) participated in the study at the Primate Research Institute, Kyoto University (KUPRI), Japan (Table 1) (Watanuki et al. 2014). The chimpanzees were members of a social group of 12 individuals living in an environmentally enriched facility consisting of two outdoor enclosures (250 m² and 280 m²), an openair outdoor enclosure (700 m²) and indoor living rooms linked to testing rooms. The open air outdoor enclosure was equipped with 15 m high climbing frames and included streams and trees (Matsuzawa et al. 2006; Yamanashi and Hayashi 2011). No food or water deprivation was used in the study. The experimental protocol was

approved by the Animal Welfare and Care Committee of the KUPRI and the Animal Research Committee of Kyoto University and followed the *Guidelines for the Care and Use of Laboratory Primates* of the KUPRI (Version 3, 2010). The chimpanzees had extensive experience participating in perceptual and cognitive tasks using a touchscreen, including MTS tasks (e.g. Dahl et al. 2013b; Tomonaga 1999).

Apparatus

Experiments were conducted in an experimental booth ($1.80 \times 2.15 \times 1.75$ m) inside a testing room. Each chimpanzee voluntarily walked to the booth through an overhead walkway connected to the living rooms. Two 17-inch touch-sensitive LCD monitors (1280×1024 pixels) encased inside Plexiglas were used to present visual stimuli at approximately 40 cm distance. Food rewards (eight-millimetre apple cubes) were delivered via a universal feeder device. All experimental events were controlled by a PC and the computer task was programmed using Microsoft Visual Basic 2010 (Express Edition).

Stimuli

Photographs (200×225 pixels) of six species of primate faces including; chimpanzee (*Pan troglodytes*), western lowland gorilla (*Gorilla gorilla*), orangutan (*Pongo pygmaeus* and *Pongo abelii*) (20 images each), Japanese human (*Homo sapiens*), olive baboon (*Papio anubis*) and white-headed capuchin monkey (*Cebus capucinus*) faces (10 images each) were obtained from the internet and colleagues (for examples see Figure 1). All images were cropped and featured unfamiliar faces with a neutral closed-mouth expression against a variety of naturalistic backgrounds. The images were prepared in colour, greyscale, upright and inverted format and controlled for luminance

and contrast using Adobe Photoshop CS2 (Version 9.0). In addition, for conditions presenting faces of two primate species, 10 ‘random dot’ control images (five images for each species) were composed by randomly assigning each pixel of the original primate face images to a new position. This procedure ensured content-related information was removed while average brightness levels were maintained.

Procedure

The chimpanzees participated in a series of zero-delay MTS tasks (Figure 1). To begin each trial the chimpanzees touched a blue start key at the bottom-center of the screen, followed by a sample stimulus directly above the start key position. When the sample was touched it disappeared and three comparison stimuli appeared in the middle of the screen. When a correct choice was made (the comparison touched matched the sample) a chime sound was played and a food reward given, when an incorrect choice was made (the comparison did not match the sample) a buzzer sound was played and no food reward given. The inter-trial interval was two seconds. The number of correct choices and response times (ms) were recorded by a PC.

Conditions

The study consisted of five conditions to investigate the influence of colour, orientation, familiarity, and perceptual similarity on the ability to discriminate primate faces under identical and categorical matching formats. On identical matching trials, the sample and correct comparison stimuli were identical to each other, whilst on categorical matching trials, the images were different from each other but within the same species category. Identical and categorical matching was alternated on each trial in each of the conditions. For the orientation condition only identical matching was presented. The chimpanzees

completed a total of 864 trials (72 trials $\times$ 12 sessions) for each condition. Trial order was randomised within and across sessions.

In Condition 1 the chimpanzees matched chimpanzee, gorilla and orangutan faces in colour and greyscale. In Condition 2 they matched the same species in upright and inverted orientation in greyscale. Both the sample and comparison images were presented in the same orientation (Tomonaga 1999; Dahl et al. 2013c). In Conditions 3, 4 and 5 they matched greyscale baboon and capuchin monkey faces (unfamiliar and perceptually different), chimpanzee and human faces (highly familiar and perceptually different), and orangutan and gorilla faces (unfamiliar and perceptually similar) respectively. Perceptual similarity was defined as the relative similarity between faces of two different species as perceived by the experimenters. In Conditions 3, 4 and 5 a third ‘random dot’ control image was added to the two comparison face images to ensure an equal number of choices across all conditions. Ten novel chimpanzee, gorilla and orangutan faces were used in Conditions 4 and 5 to reduce the possible influence of learning effects.

Image similarity analysis

Similarity between chimpanzee, orangutan, and gorilla faces (20 images each) and baboon, capuchin monkey and human faces (10 images each) in greyscale was calculated. First, the faces were aligned vertically and eyes positioned centrally. Using a customised programme written using Python and OpenCV commands AKAZE (Accelerated KAZE) local features were then extracted (cf. Alcantarilla, Nuevo and Bartoli 2011). These features were analysed using the Brute-Force Matcher method and “cv2.bfmatcher” function. The customised programme generated similarity values for each face pair, where lower values indicate higher similarity based on local feature

matching (OpenCV Development Team, 2014). This similarity-matrix data was used for metric Multidimensional Analysis (MDS) using the “cmdscale” function in the library “stats” of R version. 3.2.2 (R Core Team, 2015). This produced spatial representations of similarities between stimuli. We adopted a two-dimensional solution for the present analysis. To evaluate goodness of fit, the “cmdscale” function generated a GOF (Goodness of Fit) value. The higher the GOF value the better the spatial solution.

Statistical analysis

The data were analysed using SPSS (version 23). Paired-comparison *t*-tests and repeated measures analysis of variance (ANOVAs) were used to analyse accuracy (the mean number of correct choices) and response times (on correct trials). One-sample *t*-tests were used to analyse difference in accuracy from chance level. Trials with response times of five seconds or longer after the sample image presentation disappeared may not have been retained and so were excluded from the analysis. In Conditions 3, 4 and 5 the third random dot control image was never touched and so chance level was 50%.

Results

Condition 1: Colour effects

Figure 2. shows the mean percentage of correct choices for identical and categorical matching in colour and greyscale. One-sample *t*-tests found accuracy was above chance level (33.33%) in all conditions; identical colour ($M = 95\%$ correct, $t_{(4)} = 42.59$, $p < .001$), categorical colour ($M = 59\%$ correct, $t_{(4)} = 16.92$, $p < .001$), identical greyscale ($M = 87\%$ correct, $t_{(4)} = 20.49$, $p < .001$) and categorical greyscale ($M = 44\%$ correct, $t_{(4)}$

= 3.30, $p = .030$). A two-way ANOVA, with colour type (colour and greyscale) and matching type (identical and categorical) as independent variables was conducted. A main effect of colour type on accuracy was found, with higher accuracy for colour ($M = 77\%$ correct) than greyscale ($M = 65\%$ correct) matching ($F_{(1,4)} = 172.19, p < .001$). A main effect of matching type was also found, with higher accuracy for identical ($M = 91\%$ correct) than categorical ($M = 52\%$ correct) matching ($F_{(1,4)} = 166.38, p < .001$). No interactions were found. For response times on correct trials, a two-way ANOVA found a main effect of colour type, with faster response times for colour ($M = 768$ ms) than greyscale ($M = 938$ ms) matching ($F_{(1,4)} = 25.46, p = .007$). A main effect of matching type was found, with faster response times for identical ($M = 696$ ms) than categorical ($M = 1010$ ms) matching ($F_{(1,4)} = 27.29, p = .006$). No interactions were found.

Condition 2: Orientation effects

Figure 3. shows the mean percentage of correct choices for identical matching in upright and inverted orientation in greyscale. One sample t -tests found accuracy was above chance level (33.33%) in both orientations; upright ($t_{(4)} = 8.91, p = .001$) and inverted ($t_{(4)} = 8.84, p = .001$). A paired comparison t -test found accuracy was higher for upright ($M = 75\%$ correct) than inverted ($M = 66\%$ correct) matching ($t_{(4)} = 3.06, p = .038$). A paired comparison t -test found no difference in response times on correct trials between upright ($M = 889$ ms) and inverted ($M = 882$ ms) orientation.

Conditions 3, 4 and 5: Familiarity and perceptual similarity effects

Figure 4. shows the mean percentage of correct choices for identical and categorical matching of baboon and capuchin monkey (Condition 3), chimpanzee and human

(Condition 4), and gorilla and orangutan (Condition 5) faces in greyscale. One-sample t -tests found accuracy was above chance level (50%) in all conditions; baboon and capuchin monkey (identical: $M = 91\%$ correct, $t_{(4)} = 20.67$, $p < .001$), (categorical: $M = 82\%$ correct, $t_{(4)} = 15.43$, $p < .001$), chimpanzee and human (identical: $M = 95\%$ correct, $t_{(4)} = 23.79$, $p < .001$), (categorical: $M = 87\%$ correct, $t_{(4)} = 10.72$, $p < .001$) and gorilla and orangutan (identical: $M = 83\%$ correct, $t_{(4)} = 14.03$, $p < .001$), (categorical: $M = 58\%$, $t_{(4)} = 8.77$, $p = .001$). A 3 x 2 ANOVA with stimulus type (baboon and capuchin monkey, chimpanzee and human, and gorilla and orangutan) and matching type (identical and categorical) as independent variables was conducted. A main effect of stimulus type was found with higher accuracy for matching baboon and capuchin monkey ($M = 87\%$) and chimpanzee and human ($M = 91\%$) than gorilla and orangutan ($M = 70\%$) faces ($F_{(2,8)} = 26.48$, $p < .001$). A main effect of matching type was found, with higher accuracy for identical ($M = 90\%$) than categorical ($M = 76\%$) matching ($F_{(1,4)} = 78.30$, $p < .001$). An interaction between stimulus type and matching type was found, with a sharp decline in accuracy for categorical gorilla and orangutan face matching ($F_{(1,4)} = 51.80$, $p < .001$). For response times on correct trials a 3 x 2 ANOVA was conducted. A main effect of stimulus type was found ($F_{(1,4)} = 4.60$, $p = .047$) with longer response times for matching gorilla and orangutan (772 ms) than baboon and capuchin monkey (698 ms) or chimpanzee and human (668 ms) faces. A main effect of matching type was found with faster response times for identical (669 ms) than categorical (756 ms) matching ($F_{(1,4)} = 48.93$, $p = .002$). No interactions were found.

Image similarity analysis

Figure 5. shows the two-dimensional representation of the Multidimensional Scaling analysis of the image similarity data based on local feature matching. The GOF value is

0.207. The chimpanzee, gorilla and orangutan features were grouped more closely together indicating greater image similarity, and the human, baboon and capuchin monkey features were grouped further away indicating less similarity.

Discussion

This study focused on the visual elements important for categorical discrimination between species faces. As predicted, discrimination performance for great ape faces was better in colour than in greyscale. This is consistent with the findings of Roberts and Mazmanian (1988) in humans and Marsh and MacDonald (2008) in orangutans, that pictures of animals and faces are easier to discriminate in colour than greyscale. An inversion effect for discriminating great ape faces was found, with lower accuracy for discriminating inverted faces, suggesting the stimuli were perceived as faces and processed configurally. Previous MTS and visual search studies with our chimpanzees (Tomonaga 1999, 2007) have found inversion effects for images of faces but not houses, chairs or hands as control stimuli, suggesting our results are also face-specific.

Interestingly, although discrimination performance was high for both unfamiliar (baboon and capuchin monkey) and highly familiar (chimpanzee and human) perceptually different species faces, it was not significantly better for highly familiar species faces as predicted. Two studies in chimpanzees have examined the influence of familiarity on discrimination of individual human and chimpanzee faces using the MTS task (Dahl et al. 2013b; Martin-Malivel and Okada 2007). Both studies found discrimination accuracy was highest for species to which the chimpanzees had the most real-life previous exposure. Given that previous experience enhances discrimination of

individual faces within species, why did previous experience appear to have no influence on discrimination between species at the categorical level in the current study?

One explanation may be from an evolutionary perspective, discrimination of individuals within one's own species is especially important for the social lives of primates. In chimpanzees, higher-order features (a combination of first and second-order features) influence individual face discrimination within species (Dahl et al. 2013b; Martin-Malivel and Okada 2007). However, discrimination between individuals of other species, or between species at the categorical level, is less socially relevant and likely to involve fewer processing resources allocated to recognition (Pascalis and Wirth 2011). Studies examining the other-race effect have found humans are better at recognising faces of individuals within their own race and better at categorising faces of other races (Ge et al. 2009; Levin 1996, 2000). This may be due to greater processing resources allocated to individuating information of own-race faces we are experienced with, and to categorical information of other-race faces we are less experienced with (Ge et al. 2009). Similarly, when discriminating between categories of primate species faces, rather than within own-species faces, our chimpanzees may have paid more attention to information important for categorisation than recognition.

Another explanation is the perceived similarity between species faces. We chose primate faces representing a range of phylogenetic groups including humans, great apes, old world monkeys and new world monkeys, to examine the potential relationship between phylogenetic distance and discrimination ability. Performance for categorical discrimination of unfamiliar, perceptually similar and phylogenetically closer faces (gorilla and orangutan) was significantly worse than for unfamiliar, perceptually different and phylogenetically more distant faces (baboon and capuchin monkey). In

support of this finding, Campbell et al. (1997) found humans were better able to discriminate between physically morphed faces of monkeys and cows, and humans and cows, compared to humans and monkeys. Categorical perception was sharpest for monkey and cow and human and cow faces, suggesting the phylogenetically and morphologically closer human and monkey faces were perceived as more similar to each other. In our chimpanzees, Tanaka (2003, 2007) did not find a relationship between visual preference of primate species and phylogenetic distance from chimpanzees, although they were not required to directly discriminate between species.

After excluding effects of colour and familiarity, we propose difficulty in discriminating between primate species faces can be best explained by their perceived similarity to each other. This conclusion is supported by both the behavioural data; categorical discrimination performance for perceptually similar primate faces was significantly worse than for perceptually different faces, and the Multidimensional Scaling analysis of the stimuli; greater local feature similarity between chimpanzee, gorilla and orangutan faces than between human, baboon and capuchin monkey faces. Greater perceptual similarity between great ape faces likely explains the relatively poor discrimination performance in the absence of colour cues, and greater perceptual difference explains the high discrimination performance for baboon and capuchin monkey, and chimpanzee and human faces. Although we do not exclude the possibility previous experience has some influence on categorical discrimination between species faces, information about perceptual similarity appears to be prioritised. This could have masked the potential influence of previous experience or a conceptual representation of species (cf. Martin-Malivel et al. 2006).

In this study the experimenters made subjective judgements about the relative perceptual similarity of primate species faces. This was subsequently supported by the objective Multidimensional Scaling analysis of the stimuli. However, a useful future extension to this study would be to test whether our findings generalise across primate species, by testing them under the same experimental conditions as our chimpanzees. If the same pattern of responses are found, we could more concretely conclude non-human primates perceive similarity in different species faces like humans. In addition, although the Multidimensional Scaling analysis revealed the relative similarities between faces by local feature matching, it does not tell us which specific features, e.g. external face contours or internal features such as the eyes, nose, and mouth may be responsible for discrimination between species. Although the inversion effect we observed provides evidence of configural processing, further analysis of part-based processing by extracting curvature information about the eyes, nose and mouth image surfaces (e.g. Dahl et al. 2014a, b) or systematic removal of different facial features (e.g. Parr et al. 2000; Quinn and Eimas 1996; Tomonaga and Imura 2015) may help to reveal features most important for between species discrimination. Finally, we acknowledge some of the general limitations of our study including the small sample size, the limited stimuli number and species used for each phylogenetic group, and the fact sex differences could not be examined as all the chimpanzees were female.

We conclude our chimpanzees appear to perceive similarity in primate faces in a similar way to humans. Information about perceptual similarity is likely prioritised over the potential influence of previous experience or a conceptual representation of species for categorical discrimination between species faces.

Acknowledgements

We sincerely thank Prof. Tetsuro Matsuzawa and Drs. Ikuma Adachi, Yuko Hattori and Misato Hayashi of the Language and Intelligence Section, Primate Research Institute, Kyoto University, for their helpful comments and generous support of this research. We also thank Dr. Fumito Kawakami, Mr. Yoshiki Kurosawa and Ms. Etsuko Ichino of the Language and Intelligence Section, Primate Research Institute, Kyoto University, for assistance with conducting the experiments and staff at the Center for Human Evolution Modelling Research, Primate Research Institute, Kyoto University, for their daily care of the chimpanzees. In addition, we thank Drs. Yumi Yamanashi, Yoko Sakuraba and Renata Mendonça for permission to use their photographs of chimpanzees, gorillas and orangutans. This research was financially supported by the MEXT (#150528), JSPS-MEXT KAKENHI (#15H05709, #16H06283, #20002001, #23220006, #24000001), JSPS-CCSN and the JSPS Leading Graduate Program in Primatology and Wildlife Science (U04) at Kyoto University.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Ethical approval This study was approved by the Animal Welfare and Care Committee of the Primate Research Institute, Kyoto University and the Animal Research Committee of Kyoto University, and followed the Guidelines for the Care and Use of Laboratory Primates of the Primate Research Institute, Kyoto University (Version 3, 2010).

References

- Alcantarilla PF, Nuevo J, Bartoli A (2011) Fast explicit diffusion for accelerated features in nonlinear scale spaces. *IEEE Trans Patt Anal Mach Intell* 34(7):1281-98
- Bruce V, Young A (1986) Understanding face recognition. *Br J Psychol* 77(3):305–327
- Burton AM, Wilson S, Cowan M, Bruce V (1999) Research Article Evidence From Security Surveillance. *Psychol Sci* 10:243–248
- Campbell R, Pascalis O, Coleman M, Wallace SB, Benson PJ (1997) Are faces of different species perceived categorically by human observers? *Proc Biol Sci* 264:1429–1434
- Dahl CD, Chen C-C, Rasch MJ (2014a) Own-race and own-species advantages in face perception: a computational view. *Sci Rep* 4:6654
- Dahl CD, Logothetis NK, Hoffman KL (2007) Individuation and holistic processing of faces in rhesus monkeys. *Proc R Soc London Ser B-Biological Sci* 274(1622):2069–76
- Dahl CD, Rasch MJ, Chen CC (2014b) The other-race and other-species effects in face perception - A subordinate-level analysis. *Front Psychol* 5:1–9
- Dahl CD, Rasch MJ, Tomonaga M, Adachi I (2013a) Laterality Effect for Faces in Chimpanzees (*Pan troglodytes*). *J Neurosci* 33:13344–13349

439 Dahl CD, Rasch MJ, Tomonaga M, Adachi I (2013b) Developmental processes in face
 440 perception. *Sci Rep* 3:1044

441 Dahl CD, Rasch MJ, Tomonaga M, Adachi I (2013c) The face inversion effect in non-
 442 human primates revisited - an investigation in chimpanzees (*Pan troglodytes*).
 443 *Sci Rep* 3(2):2504

444 Demaria C, Thierry B (1988) Responses to animal stimulus photographs in stump-tailed
 445 macaques (*Macaca arctoides*). *Primates* 29:237–244

446 Diamond R, Carey S (1986) Why faces are and are not special: an effect of expertise. *J*
 447 *Exp Psychol Gen* 115(2):107–17

448 Dufour V, Pascalis O, Petit O (2006) Face processing limitation to own species in
 449 primates: A comparative study in brown capuchins, Tonkean macaques and
 450 humans. *Behav Processes* 73(1):107–113

451 Fujita K (1987) Species recognition by five macaque monkeys. *Primates* 28(3):353–366

452 Fujita K (1990) Species preference by infant macaques with controlled social
 453 experience. *Int J Primatol* 11:553–573

454 Fujita K (1993a) Role of some physical characteristics in species recognition by pigtail
 455 monkeys. *Primates* 34(2):133–140

456 Fujita K (1993b) Development of visual preference for closely related species by infant
 457 and juvenile macaques with restricted social experience. *Primates* 34(2):141–
 458 150

- 459 Fujita K, Watanabe K (1995) Visual preference for closely related species by Sulawesi
460 macaques. *Am J Primatol* 37:253–261
- 461 Ge L, Zhang H, Wang Z, Quinn PC, Pascalis O, Kelly D, Slater A, Tian J, Lee K (2009)
462 Two faces of the other-race effect: Recognition and categorisation of Caucasian
463 and Chinese faces. *Perception* 38(8):1199–1210
- 464 Goldstein AG (1965) Learning of inverted and normally oriented faces in children and
465 adults. *Psychon Sci* 3:447–448
- 466 Gothard KM, Brooks KN, Peterson MA (2009) Multiple perceptual strategies used by
467 macaque monkeys for face recognition. *Anim Cogn* 12:155–167
- 468 Humphrey N (1974) Species and individuals in the perceptual world of monkeys.
469 3:105–114
- 470 Kano F, Tomonaga M (2009) How chimpanzees look at pictures: a comparative eye-
471 tracking study *Proc Royal Soc B* 276(1664):1949-1955
- 472 Levin DT (1996) Classifying faces by race: The structure of face categories. *J Exp*
473 *Psychol Learn Mem Cogn* 22: 1364-1382
- 474 Levin DT (2000) Race as a visual feature: Using visual search and perceptual
475 discrimination tasks to understand face categories and the cross-race recognition
476 deficit. *J Exp Psychol Gen* 129: 559-574
- 477 Marsh HL, MacDonald SE (2008) The use of perceptual features in categorization by
478 orangutans (*Pongo abelli*). *Anim Cogn* 11(4):569–585

479 Martin-Malivel J, Mangini MC, Fagot J, Biederman I (2006) Do Humans and Baboons
 480 Use the Same Information When Categorizing Human and Baboon Faces?
 481 Psychol Sci 17:599–607

482 Martin-Malivel J, Okada K (2007) Human and chimpanzee face recognition in
 483 chimpanzees (*Pan troglodytes*): role of exposure and impact on categorical
 484 perception. Behav Neurosci 121(6):1145–1155

485 Matsuzawa T (2006) Sociocognitive development in chimpanzees: A synthesis of
 486 laboratory work and field work. In: Matsuzawa T, Tomonaga M, Tanaka M
 487 (eds.) Cognitive Development in Chimpanzees. Springer-Verlag, Tokyo

488 OpenCV Development Team (2014) Feature matching. URL
 489 [https://docs.opencv.org/3.0-](https://docs.opencv.org/3.0-beta/doc/py_tutorials/py_feature2d/py_matcher/py_matcher.html)
 490 [beta/doc/py_tutorials/py_feature2d/py_matcher/py_matcher.html](https://docs.opencv.org/3.0-beta/doc/py_tutorials/py_feature2d/py_matcher/py_matcher.html). Accessed 21
 491 November 2017

492 Parr LA (2011) The evolution of face processing in primates. Philos Trans R Soc Lond
 493 B Biol Sci 366(1571):1764–1777

494 Parr LA, Dove T, Hopkins WD (1998) Why faces may be special: evidence of the
 495 inversion effect in chimpanzees. J Cogn Neurosci 10(5):615–622

496 Parr LA, Winslow JT, Hopkins WD, de Waal FBM (2000) Recognizing Facial Cues:
 497 Individual Discrimination by Chimpanzees (*Pan troglodytes*) and Rhesus
 498 Monkeys (*Macaca mulatta*). J Comp Psychol 114(1):47–60

- 499 Pascalis O, Bachevalier J (1998) Face recognition in primates: a cross-species study.
500 Behav Process 43(1):87-96
- 501 Pascalis O, Wirth S (2011) Recognizing the Faces of Other Species: What Can a
502 Limited Skill Tell Us About Face Processing? In: Calder AJ, Rhodes G, Johnson
503 MH, Haxby JV (eds.) The Oxford Handbook of Face Perception, Oxford
504 University Press Inc., New York
- 505 Piepers DW, Robbins RA (2012) A review and clarification of the terms “holistic,”
506 “configural,” and “relational” in the face perception literature. Front Psychol
507 3(559):1–11
- 508 Quinn PC, Eimas PD (1996) Perceptual cues that permit categorical differentiation of
509 animal species by infants. J Exp Child Psychol 63:189–211
- 510 R Core Team (2015) R: a language and environment for statistical computing. R
511 Foundation for Statistical Computing, Vienna, Austria. URL [https://www.R-](https://www.R-project.org/)
512 project.org/. Accessed 21 November 2017
- 513 Roberts WA, Mazmanian DS (1988) Concept learning at different levels of abstraction
514 by pigeons, monkeys, and people. J Exp Psychol Anim Behav Process 14:247–
515 260
- 516 Scapinello KF, Yarmey AD (1970) The role of familiarity and orientation in immediate
517 and delayed recognition of pictorial stimuli. Psychon Sci 21:329–330

518 Spencer J, Quinn PC, Johnson MH, Karmiloff-Smith A (1997) Heads you win, tails you
 519 lose: evidence for young infants categorizing mammals by head and facial
 520 attributes. *Early Dev Parent* 6:113–126

521 Tanaka M (2003) Visual preference by chimpanzees (*Pan troglodytes*) for photos of
 522 primates measured by a free choice-order task: implication for influence of
 523 social experience. *Primates* 44(2):157–165

524 Tanaka M (2007) Development of the visual preference of chimpanzees (*Pan*
 525 *troglodytes*) for photographs of primates: Effect of social experience. *Primates*
 526 48(4):303–309

527 Tomonaga M (1999) Inversion Effect in Perception of Human Faces in a Chimpanzee
 528 (*Pan troglodytes*). *Primates* 40(3):417–438

529 Tomonaga M (2007) Visual search for orientation of faces by a chimpanzee (*Pan*
 530 *troglodytes*): Face-specific upright superiority and the role of facial configural
 531 properties. *Primates* 48(1):1–12

532 Tomonaga M, Imura T (2015) Efficient search for a face by chimpanzees (*Pan*
 533 *troglodytes*). *Sci Rep* 5:11437

534 Valentine T (1991) A unified account of the effects of distinctiveness, inversion, and
 535 race in face recognition. *Q J Exp Psychol Sect A Hum Exp Psychol* 43(2):161–
 536 204

- 537 Watanuki K, Ochiai T, Hirata S, Morimura N, Tomonaga M, Idani G, Matsuzawa T
538 (2014) Review and long-term survey of the status of captive chimpanzees in
539 Japan in 1926-2013. *Primate Res* 30:147-156
- 540 Yamanashi Y, Hayashi M (2011) Assessing the effects of cognitive experiments on the
541 welfare of captive chimpanzees (*Pan troglodytes*) by direct comparison of
542 activity budget between wild and captive chimpanzees. *Am J Primatol* 73:1231–
543 1238
- 544 Yin R (1969) Looking at upside-down faces. *J Exp Psychol Gen* 81:141–145

Tables

Table 1. Basic information about the five chimpanzees

Name	GAIN ID number ^a	Sex	Age (at study start)
Ai	0434	Female	40
Chloe	0441	Female	36
Cleo	0609	Female	16
Pal	0611	Female	16
Pendensa	0095	Female	38

^aIdentification number (ID) for each chimpanzee listed in the database of the Great Ape Information Network (GAIN): <https://shigen.nig.ac.jp/gain/>

Table 1. Basic information about the five chimpanzees. ^a Identification number (ID) for each chimpanzee listed in the database of the Great Ape Information Network (GAIN): <https://shigen.nig.ac.jp/gain/>

Figures

Fig. 1 Chimpanzee 'Pal' choosing an orangutan face in the colour condition.

Fig. 2 Mean percentage of correct choices for identical and categorical matching of chimpanzee, gorilla and orangutan faces in colour and greyscale. The solid horizontal line indicates a main effect of colour on accuracy. The dashed horizontal line represents 33.33% chance level. Error bars represent the standard error of the mean.

Fig. 3 Mean percentage of correct choices for identical matching of chimpanzee, gorilla and orangutan faces in upright and inverted orientation in greyscale. The solid horizontal line indicates a significant difference in accuracy between orientation presentations. The dashed line represents 33.33% chance level. Error bars represent the standard error of the mean.

Fig. 4 Mean percentage of correct choices for identical and categorical matching of baboon and capuchin monkey, chimpanzee and human, and gorilla and orangutan faces in greyscale. The solid horizontal line indicates a main effect of stimulus type on accuracy. The dashed line represents 50% chance level. Error bars represent the standard error of the mean.

● ● Baboon ▨ ▨ Human ● ● Capuchin monkey
 ● ● Chimpanzee ▤ ▤ Orangutan ● ● Gorilla

575

576 **Fig. 5** Two-dimensional representation of the Multidimensional Scaling analysis of the
 577 image similarity data based on local feature matching. Large circles represent average
 578 values and small circles represent individual values. Chimpanzee, gorilla and orangutan
 579 faces consisted of 20 images each, and human, baboon and capuchin monkey faces
 580 consisted of 10 images each.