**The structure of psychopathology and association with poor sleep, self-harm,
suicidality, risky sexual behaviour, and low self-esteem in a population sample of
adolescents**

Matthew Sunderland¹, Miriam K. Forbes², Louise Mewton³, Andrew Baillie⁴, Natacha Carragher^{5,6}, Samantha J. Lynch¹, Philip J. Batterham⁷, Alison L. Calear⁷, Cath Chapman¹,
Nicola C. Newton¹, Maree Teesson¹, Tim Slade¹.

1. The Matilda Centre for Research in Mental Health and Substance Use, Sydney Medical School, Faculty of Medicine and Health, The University of Sydney, Australia.
2. Centre for Emotional Health, Department of Psychology, Macquarie University, Australia.
3. Centre for Healthy Brain Aging, School of Psychiatry, UNSW Sydney, Australia.
4. Faculty of Health Sciences, The University of Sydney, Australia.
5. World Health Organisation, Geneva, Switzerland.
6. Office of Medical Education, UNSW Sydney, Australia.
7. Centre for Mental Health Research, Australian National University, Canberra, Australia.

Address for correspondence: Dr Matthew Sunderland, The Matilda Centre for Research in Mental Health and Substance Use, Level 6, Jane Foss Russell Building (G02), University of Sydney, Sydney, NSW 2006, Australia. Email: matthew.sunderland@sydney.edu.au

ABSTRACT

There is a growing body of evidence highlighting the presence of a single general dimension of psychopathology that can account for multiple associations across mental and substance use disorders. However, relatively little evidence has emerged regarding the validity and utility of this model with respect to a range of factors that have been previously implicated across multiple disorders. The current study utilised a cross-sectional population survey of adolescents (n=2003) to examine the extent that broad psychopathology factors account for specific associations between psychopathology and poor sleep, self-harm, suicidality, risky sexual behaviour, and low self-esteem. Confirmatory factor models, latent class models, and factor mixture models were estimated to identify the best structure of psychopathology. Structural equation models were then estimated to examine the broad and specific associations between each psychopathology indicator and the validity indicators. A confirmatory factor model with three lower order factors, representing internalizing, externalizing, and thought disorder, and a single higher order factor evidenced the best fit. The associations between manifest indicators of psychopathology and validity indicators were largely non-specific. However, significant direct effects were found between several important pairwise associations. These findings have implications for the identification of potential targets for intervention and/or tailoring of prevention programs.

Keywords: psychopathology, confirmatory factor analysis, validity, adolescents.

INTRODUCTION

Mental disorders affect one in seven young Australians in any given year and are responsible for the largest burden of disease in this age group (Erskine et al., 2015). Epidemiological data on population samples has consistently identified a high rate of comorbidity (or co-occurring disorders) between mood, anxiety, psychotic, and substance use disorders with the presence of any disorder associated with increased odds of experiencing another disorder (Slade, McEvoy, Chapman, Grove, & Teesson, 2015). Higher rates of comorbidity have also been linked to increased levels of distress, higher disorder severity, service use, and poor treatment response (Lubman, Allen, Rogers, Cementon, & Bonomo, 2007; Teesson, Slade, & Mills, 2009). In younger populations, comorbidity is the rule rather than the exception and many disorders first manifest as transdiagnostic indicators or common symptoms that further develop into more defined and recognisable disorders (McElroy, Belsky, Carragher, Fearon, & Patalay, 2018). These findings have led researchers to speculate that there might be several underlying common or transdiagnostic factors that better account for co-occurring multiple disorders and their association with other clinically relevant factors or outcomes, such as suicide or treatment response (Conway et al., 2019). More recently, the literature has placed considerable attention on identifying an underlying “p-factor” that accounts for the broad tendency to experience any psychopathology, which might manifest in numerous specific ways (Caspi et al., 2014). Indeed, identifying and targeting broad levels of the p-factor might result in more efficient treatment and prevention efforts to reduce the sizable burden of psychopathology (Caspi & Moffitt, 2018).

Primarily based on empirical data, the Hierarchical Taxonomy of Psychopathology (HiTOP) consortium has proposed that psychopathology might be more efficiently captured and organized through a system of continuous latent factors that are structured within a larger hierarchy, ranging from specific symptom clusters at the lowest level through to a broad *p*-

factor at the superspectrum level (Kotov et al., 2017). Previous studies have identified and validated various levels of the HiTOP hierarchy and employed bi-factor models to identify general and specific factors related to adolescent psychopathology (Afzali, Sunderland, Carragher, & Conrod, 2017; Carragher et al., 2016; Castellanos-Ryan et al., 2016; Haltigan et al., 2018; Laceulle, Vollebergh, & Ormel, 2015; Tackett, 2006; Waldman, Poore, van Hulle, Rathouz, & Lahey, 2016). However, these models assume the structure underlying psychopathology is inherently continuous and normally distributed, relying on the use of variable-centred latent modelling. Other potential models can relax and test these strict assumptions by fitting person-centred categorical latent variables (in a latent class modelling framework) or a mixture of continuous and categorical latent variables (factor mixture models) (Lubke & Muthén, 2005). Additional insights into the nature of psychopathology can be obtained via detailed model comparisons, for example whether the latent distribution might be considered continuous but non-normally distributed (Masyn, Henderson, & Greenbaum, 2010). Among adult samples, comprehensive model comparisons have indicated that the structure of symptom-based data across multiple mental and substance use disorders is best represented by continuous and normally distributed factors (Wright et al., 2013).

In addition to identifying the latent structure of psychopathology, studies have demonstrated the various degrees of validity and utility associated with latent models when capturing the association between multiple individual mental disorders (i.e., comorbidity) and other key indicators of interest that have been hypothesized to be associated with multiple disorders (Conway et al., 2019). If the common or broad latent factors (either categorical, continuous, or both) adequately capture the total observed variance between individual indicators of psychopathology and other key validity indicators, it may be inferred that the association is primarily driven by commonalities across multiple disorders and/or symptoms rather than information that is specific to individual disorders and/or symptoms. Among adult

samples, the shared and specific associations between psychopathology and numerous additional clinical validators have been investigated, including: suicidality, treatment seeking, functional impairment/disability, traumatic events, intergenerational transmission, and adverse childhood events (Albott, Forbes, & Anker, 2018; Keyes et al., 2012; Starr, Conway, Hammen, & Brennan, 2014; Sunderland et al., 2016; Sunderland & Slade, 2014). These studies show that broad latent factors, such as internalizing and externalizing, can adequately capture multiple associations between additional clinical indicators and individual disorder indicators that comprise the latent factors. Moreover, these studies provide further evidence for the utility of the HiTOP model as an organisational structure for psychopathology. Yet much of this structural work has been confined to adults. Adolescence is a developmental period marked by significant neurobiological, behavioural, and social changes. It is also a time when substance use disorders, psychosis, personality disorders, and eating disorder first emerge (Paus, Keshavan, & Giedd, 2008), with critical implications for adult development and health.

In adolescents, there are several key validity indicators, including suicidality, self-harm, sleep, self-esteem and risky sexual behaviours, that have been previously associated with numerous mental health and substance use conditions. Moreover, these indicators have been linked to increased levels of distress, hospitalization, mortality, and in the case of risky sexual behaviour, sexually transmitted infections and unwanted pregnancies (Borschmann, Stark, Prakash, & Sawyer, 2018; Prendergast, Toumbourou, McMorris, & Catalano, 2019). Often these indicators further contribute to and exacerbate psychopathology and negative outcomes (Baglioni et al., 2016; Daraganova, 2017; Mars et al., 2014; Masselink, Van Roekel, & Oldehinkel, 2018). Further empirical evidence is required to better understand the shared and specific relationships that might exist between psychopathology and these key validity indicators. Indeed, specific associations would indicate there is something unique

about a pairwise association between a symptom domain/disorder and a key validity indicator, with implications for the development of transdiagnostic and/or tailored interventions that seek to reduce these harmful and inter-related behaviours.

As such, the current study: 1) examined the fit and compared a range of alternative and theoretically plausible models of psychopathology from a large sample of Australian adolescents; and 2) investigated if these broad models of psychopathology captured the total association among specific indicators of psychopathology (e.g., major depression) and other hypothesized key validity indicators implicated across multiple disorders (e.g. poor sleep, self-harm, suicidal ideation, suicide attempts, multiple sexual partners, condom use, and low self-esteem).

METHODS

Sample

Data for the current study were drawn from the Second Australian Child and Adolescent Survey of Mental Health and Wellbeing (Young Minds Matter; YMM) (Hafekost, Johnson, et al., 2016), a stratified, multistage area probability sample of Australian households with at least one child aged 4-17 years. The survey was administered by the Telethon Kids Institute at the University of Western Australia in partnership with Roy Morgan Research. The survey received ethical approval from the Australian Government Department of Health. Moreover, parents or carers and young people provided informed consent prior to participating in the survey and the self-report measures. The survey data have been weighted to represent the population of all Australian young people aged 11-17 years old. The clustered nature of the sample design was incorporated into the analyses when estimating standard errors and confidence intervals. The current study restricted the analysis to adolescents aged 14 years or older given those younger than 14 were not administered

detailed self-report questions on alcohol, tobacco, drug use, risky sexual behaviour, and psychotic like experiences. More detail on the survey methods and technical details can be found elsewhere (Hafekost, Lawrence, et al., 2016).

Measures

Indicators of Psychopathology

All diagnostic variables included in the analyses were assessed using the parent-reported Diagnostic Interview Schedule for Children version IV (DISC-IV) developed by the US National Institute of Mental Health (Shaffer, Fisher, Lucas, Dulcan, & Schwab-Stone, 2000). The parent version of the DISC-IV has previously demonstrated acceptable reliability compared to the self-report version (Shaffer, Fisher, Lucas, Hilsenroth, & Segal, 2004). DSM-IV criteria were assessed in the past 12 months prior to the survey and diagnostic variables were generated for eight disorders: separation anxiety disorder (SD), social anxiety disorder (SAD), generalised anxiety disorder (GAD), major depressive disorder (MDD), attention deficit hyperactivity disorder (ADHD), obsessive-compulsive disorder (OCD), conduct disorder (CD), and oppositional defiance disorder (ODD). Preliminary inspection of the data indicated zero cells in the bivariate correlation matrix when running the latent variables models. As such, the diagnoses of ODD and OCD were removed from the analysis.

Indicators of substance use were measured using a combination of self-report items (e.g., frequency and quantity of alcohol use) and single use questions (e.g., have you ever smoked cigarettes daily). More specifically, binge drinking in the past 30 days was measured by calculating the frequency of single occasion drinking with bingeing classified as greater than or equal to 4 standard drinks in a row (using the Australian definition of a standard drink equivalent to 10 grams of alcohol). Tobacco use was determined by asking if the respondent had ever smoked cigarettes daily. Cannabis use and other illegal drug use were assessed by

combining multiple binary variables about whether the respondent had ever used cannabis or other illegal drugs, including sniffing petrol, glue, aerosols, solvents, paints, or nitrous.

Psychotic like experiences were measured using a five-item self-report screening questionnaire that obtained information on feelings of paranoia (“In the last year, have you ever believed that people were spying on you?”), unusual beliefs (“In the last year, have you ever believed that you were being sent special messages through the TV or radio or that a program had been arranged for you alone and no one else?”), “In the last year, have you ever believed that someone was using special powers to read your mind?”), visual hallucinations (“In the last year, have you ever seen something or someone that other people who were present could not see, that is had a vision when you were completely awake?”), or auditory hallucinations (“In the last year, have you heard things other people could not hear, such as a voice?”).

Key validity indicators

Poor sleep was measured using two self-report questions to generate the average number of hours of sleep obtained by the respondent on weeknights (Sunday-Thursday) and weekend nights (Friday-Saturday). An average for the whole week was calculated using both variables and dichotomised into poor sleep (<7 hours or >11 hours) or adequate sleep (between 7-11 hours) based on established sleep guidelines (Hirshkowitz et al., 2015). Self-harm was measured by a single dichotomous (yes/no) self-report question “Have you deliberately harmed or injured yourself without intending to end your own life during the past 12 months?”. Suicidal ideation and suicide attempts were measured using two dichotomous (yes/no) self-report questions: “During the past 12 months, did you ever seriously consider attempting suicide?” and “Did you attempt suicide during the past 12 months?”. Risky sexual behaviour was measured by two indicators representing 1) multiple sexual partners

dichotomised into those who have had 4 or more sexual partners across their lifetime and those with less than 4 sexual partners and 2) condom use by a single dichotomous (yes/no) question “Have you ever had sexual intercourse without using a condom?”. Finally, low self-esteem was measured using the Adolescent Self-esteem Questionnaire (ASQ) developed specifically for the Young Minds Matter survey (Hafekost, Boterhoven de Haan, Lawrence, Sawyer, & Zubrick, 2017). Scores were dichotomised to indicate the presence of low self-esteem (or not) based on a previously validated cut-point (Hafekost et al., 2017).

Statistical analysis

Delineating the Structure of Psychopathology

To determine the structure underlying the selected indicators of psychopathology in the current study, a series of latent variable models were estimated with different underlying assumptions regarding the distribution of the latent factors. Confirmatory factor analysis (CFA) was used to estimate latent models with continuous and normally distributed latent variables. Latent class analysis (LCA) was used to estimate latent models with categorical latent variables with an increasing number of classes. Finally, factor mixture models (FMM) were used to estimate latent models with non-normal continuously distributed latent variables. All models were estimated using data weighted to the sex and age characteristics of the young Australian population and standard errors were adjusted for the clustered nature of the sampling design (Muthen & Muthen, 2015).

The current study proceeded by fitting four CFA models: 1) a one factor model with all indicators loading on a single latent factor representing general psychopathology; 2) a correlated trait model with three factors representing internalizing (MDD, SD, SAD, GAD, ADHD), externalizing (binge drinking, daily smoking, cannabis use, illegal drug use, and conduct disorder), and psychotic like experiences (paranoia, receiving special messages,

believing people can read your mind, visual hallucinations, and auditory hallucinations); 3) a bifactor model with all indicators of psychopathology loading on a single factor representing general psychopathology as well as indicators of internalizing, externalizing, and psychotic like experiences (as described in the correlated latent trait model); and 4) a higher order model with three lower order factors representing internalizing, externalizing, and psychotic like experiences, and a single higher order factor representing general psychopathology that accounts for correlations among lower order factors. All models were estimated using a robust maximum likelihood estimator (MLR) with numerical integration, suitable for binary indicators, as implemented in Mplus (Muthen & Muthen, 2015). Models were compared based on the Bayesian Information Criterion (BIC) and the sample size adjusted Bayesian Information Criterion (ssaBIC), with lower values indicating better model fit (Raftery, 1995). Additional bifactor model fit indices were calculated including the Explained Common Variance (ECV), Percent of Uncontaminated Correlations (PUC), and the OmegaH coefficient (derived from standardized factor loadings) (Rodriguez, Reise, & Haviland, 2016). Moreover, we compared the bifactor and correlated factor models using the *H* coefficient given recent simulations demonstrating that model fit statistics like BIC and ssaBIC tend to be biased towards selection of bifactor models (Greene et al., 2019). The *H* coefficient provides an indication of the replicability of the latent factors with values ranging from 0 to 1 and estimates >0.8 indicating well-defined latent variables (Rodriguez et al., 2016). Finally, the CFA models were re-run using a weighted least squares estimator (WLSMV) to generate the Comparative Fit Index (CFI), Tucker-Lewis Fit Index (TLI), and the root mean square error of approximation (RMSEA). Prior empirically derived cut-points were used to determine excellent fit (CFI and TLI >0.95 and RMSEA <0.05) (Hu & Bentler, 1999).

A series of latent class models were then estimated on the indicators of psychopathology with an increasing number of latent classes included until either there was no decrease in BIC values between successive models or there was insufficient information to fit additional classes. The models were fit using the MLR estimator with increasing the number of random starting values to ensure multiple replications of the best log-likelihood value. The LCA models were compared using BIC and ssaBIC values as well as the Lo-Mendell-Rubin log-likelihood difference test used to compare a model with k classes with a model with $k-1$ classes (Nylund, Asparouhov, & Muthén, 2007). A non-significant result indicates that the k class model does not significantly improve model fit over the $k-1$ class model. An entropy value was calculated to determine the classification accuracy of the latent classes with scores ranging from 0 to 1 and higher scores indicating greater classification accuracy.

Finally, a series of FMMs that contain both categorical and continuous latent variables were estimated based on the results of the CFA and increasing the number of latent classes until BIC values no longer decreased or there was insufficient information to model more classes. In the current study, the FMMs were estimated in which the only parameters of the factor model that varied across classes were the factor means. Item thresholds and loadings were held invariant across classes and the factor variance and covariance matrix was fixed to zero (Clark et al., 2013). This model can be interpreted as a non-linear factor analysis with differences across latent classes representing differences in the severity location on the underlying dimensional continuum of each factor. FMMs were compared using BIC and ssaBIC as well as Lo-Mendell-Rubin log-likelihood difference test and entropy. The best fitting model across CFA, LCA, and FMMs was selected by comparing BIC and ssaBIC values, with the lowest value indicating better fit.

Associations between psychopathology and key validity indicators

The best fitting model of psychopathology (identified using the method described above) was then used to examine the association between broad latent variables of psychopathology (that account for the shared variance across multiple indicators) and the key validity indicators of interest. The results of the structural modelling dictated the specific analytic approach utilised to address the current aim. As such, the current study applied latent variable indirect effect models to decompose the total effect associated with each indicator of psychopathology and key validity indicators into direct effects (e.g., the association remaining after partialling out the variance explained by the higher and lower order latent factors) and indirect effects (the association explained via the higher order and lower order latent factors). The models can be interpreted in a similar manner to a measurement invariance analysis, in that the indirect effect provides an indication of the association between indicators of psychopathology (e.g., depression) and key validity indicators (e.g., self-harm) assumed by the structural model. Whereas, significant direct effects indicate the need for additional information specific to the indicators to either amplify or reduce the assumed association via the higher order factors. Significant and large direct effects relative to the indirect effects suggest that the structural model does not adequately capture the association (i.e. measurement non-invariance), whereas non-significant and small direct effects relative to indirect effects suggest that the model does adequately capture the association and therefore might provide a highly useful and parsimonious model that accounts for multiple pairwise associations (i.e., measurement invariance).

The indirect effect models estimated in the current study are described further in Figure 1. Each of the observed factor indicators and the lower-order factors (internalizing, externalizing, psychotic like experiences) are regressed on the key validity indicators whilst controlling for age, sex and the higher order factors. For example, the path labelled *cI* estimates the total effect of the covariate “poor sleep” on mean latent general

psychopathology scores (estimated with mean 0 and standard deviation 1) controlling for age and sex. The path labelled $c2$ represents the total effect of “poor sleep” on mean latent internalizing scores (estimated with mean 0 and standard deviation 1) whereas the path $c2'$ represents the direct effect of “poor sleep” on mean latent internalizing after subtracting the indirect association via the higher order factor (paths $a1 \times b1$) from the total effect. Finally, the path labelled $c3$ represents the total effect of “poor sleep” on the probability of major depression whereas the path $c3'$ represents the direct effect of “poor sleep” on the probability of major depression after subtracting the indirect association via the higher order latent factors (paths $(a1 \times b1 \times b2) + (a2 \times b2)$). This set of models enables the examination of potential specific effects between key validity indicators and observed indicators (e.g., major depression) within lower-order factors (e.g., internalizing) as well as specific effects between key validity indicators and lower-order factors (e.g., internalizing) within higher-order factors (e.g., general psychopathology). Significant regression path estimates were determined using a more conservative $p < 0.01$ given the large sample size and number of statistical tests.

RESULTS

Descriptive statistics

Of the 2,003 young Australians aged 14-17 years included in the survey, 1029 (51.4%, SE=1.19) were male with an average age of 15.5 (SE=0.03) years old. The unweighted frequencies, weighted percentages, and standard errors associated with the indicators of psychopathology and the key validity indicators are provided in Table 1. Poor sleep was experienced by 21.4% (SE=1.05) and 3.0% (SE=0.40) reported ever attempting suicide. The indicators of psychopathology ranged from 1.9% (SE=0.4) for CD through to 15.9% (0.91) for auditory hallucinations. Tetrachoric correlations for all the binary indicators of psychopathology are provided in Table 2 with results indicating moderate to high

correlations for all conceptually related indicators. There was evidence of a moderate correlation between ADHD and CD ($r=0.539$), however ADHD exhibited low correlations ($r_s<0.268$) with other externalizing indicators.

Structural analysis

The model fit statistics generated by the CFA, LCA, and FMMS are provided in Table 3. The best fitting CFA model, according to BIC and ssaBIC, was the correlated three factor model and therefore by extension the higher order model with 3 lower order factors and 1 higher order factor. The path diagram for the best fitting model and standardized factor loadings are provided in Figure 2. Loadings ranged from 0.44 (INT by ADHD) to 0.90 (EXT by Cannabis) and all loadings were significant at the $p<0.01$ level. The bifactor model generated excellent model fit according to the CFI, TLI, and RMSEA values but was considered worse than the correlated three factor/higher order model according to BIC and ssaBIC. For the general factor, ECV=0.366, PUC=0.714, and OmegaH=0.59 suggested that the presence of multidimensionality in the specific factors was severe enough to disqualify the interpretation of psychopathology as primarily unidimensional (Rodriguez et al., 2016). Moreover, the H coefficients for the bifactor model (General factor = 0.87, Internalizing=0.81, Externalizing=0.78, Psychotic like experiences=0.82) were acceptable but slightly lower than the H coefficients for the correlated factor model (Internalizing=0.90, Externalizing=0.93, Psychotic like experiences=0.90).

The best fitting LCA model contained 5 classes according to the BIC and ssaBIC values with good classification accuracy (entropy=0.88). However, the 3-factor 7-class FMM evidenced substantially lower BIC and ssaBIC values indicating superior fit relative to all LCA models. The predicted class probabilities for the 3-factor 7-class FMM are provided in Figure 3. The largest class representing 74% of the sample (Class 4) were relatively free from

psychopathology across the three factors. The second largest class representing 9% of the sample (Class 3) had relatively moderate levels of externalizing, low internalizing and low psychotic like experiences. Class 7 comprised 8% of the sample and were relatively high in psychotic like experiences but low on internalizing and externalizing. Class 6 comprised 4% of the sample and demonstrated relatively high levels of internalizing, low externalizing, and low psychotic like experiences. Class 1 comprised 2% of the sample and demonstrated high levels of comorbidity between externalizing and psychotic like experiences but low levels of internalizing. Finally, Classes 5 and 2 both comprised 1% of the population and represented relatively moderate to high levels of psychopathology across the three factors. However, Classes 1, 5, and 2 comprised relatively low proportions ($\leq 2\%$) and suggests that a more parsimonious solution might be considered, e.g. the 3 factor 4 class solution. Moreover, inspection of the BIC and ssaBIC values over all models indicates the three-factor correlated model/higher order model provided better fit relative to all FMMs.

Latent variable indirect effects models

The standardized path coefficients for the latent variable indirect effect models (i.e. total effects, total indirect effects, and direct effects) are provided in the online supplementary material. The ratio of the absolute direct effect and absolute total indirect effect was calculated for each model separately for the key validity indicators and presented in Figure 4.

For sleep, the association with each indicator of psychopathology was accounted for by the higher order latent variables with direct effects reflecting approximately half or less than that of the total indirect effects. However, the general p factor did not account for the total association between internalizing and poor sleep with a ratio of 2.05 for the direct effect relative to indirect effect. This indicates that adolescents reporting poor sleep patterns had

significantly higher internalizing scores than those with good sleep patterns ($b=0.40$, $p<0.01$), holding underlying levels of the p-factor constant.

For self-harm, the associations with 12 out of 15 observable indicators of psychopathology were accounted for by the higher order latent factors, as the direct effects were less than half that of the indirect effects. Notably, the higher order factors failed to account for the total association between MDD and self-harm with a significant direct effect ($b=1.39$, $p<0.01$) indicating that adolescents who self-harmed were approximately 4 times more likely to experience MDD than those who did not self-harm, holding underlying levels of internalizing and p-factor constant. Similarly, the p-factor was not able to fully account for the association between self-harm and internalizing with the size of the direct effect representing 80% of the indirect effect.

For suicidality (suicidal ideation and suicide attempt), the associations with 14 out of 15 observable indicators of psychopathology were accounted for by the higher order latent factors. The exception in both cases was MDD with the direct effects representing 79% and 74% of the total indirect effects for suicidal ideation and suicide attempt, respectively. Moreover, there were relatively large direct effects for the association between internalizing and suicidality in comparison to the indirect effect via the p-factor, particularly for suicide attempt given the direct effect was 1.87 times that of the indirect effect.

For risky sexual behaviour, the association between MDD and condom use was not captured by the higher order latent factors with the direct effects approximately 1.18 times that of the indirect effects, respectively. In addition, the direct association between ADHD and multiple sexual partners was 1.09 times that of the indirect effect suggesting that those with multiple sexual partners were less likely (but not statistically significant) to experience ADHD ($b=-0.74$, $p=0.154$), holding levels of internalizing and p-factor constant. There were

also relatively large, positive, and significant direct effects between externalizing and risky sexual behaviours ($b=1.00$, $p<0.01$ and $b=0.90$, $p<0.01$, respectively), holding levels of the p-factor constant.

Finally, for self-esteem there were relatively large but non-significant direct effects associated with self-esteem and major depression, ADHD, binge drinking, and mind reading, holding the corresponding higher order factors constant. Importantly, there were relatively large, positive, and significant direct effects associated with internalizing and self-esteem ($b=0.75$, $p<0.01$), holding levels of the p-factor constant.

DISCUSSION

The current study utilised a large population-based sample of Australian adolescents aged 14 to 17 years old to: 1) compare alternative and theoretically plausible models of psychopathology; and 2) investigate the shared and specific associations among multiple indicators of psychopathology (e.g., major depression, internalizing, etc.) and other key validity indicators: poor sleep, self-harm, suicidal ideation, suicide attempts, multiple sexual partners, condom use, and low self-esteem. The study found evidence that the best fitting model among this sample comprised three latent factors representing internalizing, externalizing and psychotic-like experiences. The correlations between these three factors might be represented by a single higher order factor indexing the overall liability to experience psychopathology. Moreover, the best fitting CFA model provided substantially better fit according to BIC and ssaBIC values in comparison to the best fitting FMM. This provides some evidence that the latent factors underlying psychopathology could be sufficiently modelled as continuous and normally distributed in this population.

The study indicated that the hierarchical model sufficiently accounted for the majority of the pairwise associations between psychopathology and the key validity indicators,

specifically between individual psychopathology indicators and sleep, self-harm, and suicidal ideation. However, there were a few notable exceptions, for example: MDD and condom use, ADHD and multiple sexual partners, and ADHD and self-esteem. Importantly, looking at the associations between first order latent factors and the key validity indicators, the overall *p*-factor could not sufficiently explain the total association between internalizing and self-esteem, sleep, and suicide attempts, as well as externalizing and condom use, and multiple sexual partners. Borrowing terminology from the field of psychometrics, these direct effects provide some indication of *measurement non-invariance* of the higher-order psychopathology model in relation to the validity indicators (Vandenberg & Lance, 2000). For example, the estimated association via the hypothesized model does not adequately capture the total observed association between each psychopathology indicator and the validity indicator. There are potentially additional factors unique to the identified psychopathology indicators (e.g., internalizing) that either increase or decrease the observed association with the validity indicators (e.g., self-esteem) than would be expected if estimated via the higher order latent factors alone (e.g., *p*-factor). Indeed, these results also provide evidence of the incremental validity of the lower-order factors over and above the general *p*-factor with respect to these validity indicators. Additional research is needed to focus on identifying the specific mechanisms driving these direct associations.

The results of the current study have several implications regarding the meaning and interpretation of the higher order latent factors, particularly the *p*-factor and its relative importance with lower order factors and specific indicators of psychopathology. For example, the relatively high direct associations between internalizing and self-esteem, poor sleep, and suicide attempts indicates that knowledge of an individual's proclivity to experience psychopathology in general (e.g., scores on the *p*-factor) will not provide enough detail regarding their likelihood of also experiencing low self-esteem, poor sleep, or suicide

attempts. This also suggests that the internalizing factor might be specifically characterized and differentiated from general psychopathology by associations with low self-esteem, poor sleep, and suicide attempts. Similarly, the study demonstrated that risky sexual behaviours were associated with higher externalizing scores over and above general psychopathology, reflecting and supporting previous findings that externalizing might be differentiated from general psychopathology by increased rates of sensation seeking and the tendency to engage in risky or impulsive behaviours (Carragher et al., 2016). However, it was interesting that at the individual indicator level, the associations between externalizing indicators and all seven of the validity indicators was adequately captured by the externalizing factor, which suggests these indicators have relatively homogeneous associations with the key validity indicators included in the current study. Importantly, these findings suggest that additional validity indicators or external risk factors of psychopathology might also be organised into a hierarchy that either account for commonalities or specificities when considered in conjunction with the HiTOP model (Forbes, Rapee, & Krueger, 2019). Moreover, the findings further contribute to the utility of specifying multiple levels of the psychopathology hierarchy simultaneously rather than focusing research on aspects of psychopathology that could potentially be too broad or too narrow, missing part of the picture. Finally, the fact that the current study demonstrates these outcomes among adolescents, a time when symptoms of psychopathology first emerge, is a considerable strength of the current study and offers several avenues for clinical application.

Regarding clinical implications, by estimating and comparing associations with key validity indicators at varying levels of the psychopathology hierarchy the current study provides some guidance regarding the development of broad or targeted prevention or early intervention programs. For example, interventions that broadly target internalizing disorders may also improve sleep, and self-esteem, and reduce suicide attempts in a more efficient and

effective manner than an intervention that broadly targets all psychopathology (i.e., p-factor, including externalizing and psychotic like experiences). In contrast, an intervention that narrowly focuses on features specific to MDD may not efficiently or effectively address levels of poor sleep in comparison to a transdiagnostic intervention that targets commonalities across all internalizing disorders (including anxiety and pathological worry). Conversely, interventions to improve sleep and self-esteem may have broad positive impacts on internalizing disorders, but with less impact on externalizing or psychotic disorders. Indeed, these findings highlight the potential of efficient stepped or staged care models for the early intervention or prevention of psychopathology in adolescents based on the hierarchical organisation proposed in the HiTOP model (Forbes et al., 2019). Such interventions might begin with broad components that adequately address commonalities associated with general psychopathology before stepping up into more focused interventions for those with higher internalizing to further address factors that have additional specific associations with those conditions (e.g., poor sleep, self-esteem, etc).

There are several limitations of the current study that should be considered when interpreting the results. First, the indicators of psychopathology used in the current study represent a mix of self-report and parent-reported indicators as well as a mix of different reference timeframes (e.g., 30 day/12-month/no reference). Moreover, there was a lack of robust indicators available in the survey to measure thought disorder and externalizing, with the current study relying on the use of screening questions to infer the presence of more severe psychopathology rather than clinically assessed symptoms of psychopathology. The high prevalence of auditory hallucinations observed in the current study might suggest that the question “hearing voices no one else can hear” is too ambiguous or interpreted as referring to high acuity of hearing. Second, the current study used data from a cross-sectional general population survey and therefore could only provide an indication of association

between variables rather than inferring any causality or temporality between psychopathology and the validity indicators. Third, the current study estimated factor mixture models by constraining all parameters except for latent means as equivalent across latent classes. Prior studies have indicated that these restrictive models rarely fit real data and additional models are required to freely estimate additional parameters however the alternative recommended models failed to converge in the current sample and could not generate usable estimates (Clark et al., 2013). Additional independent samples are required to replicate the findings demonstrated here as well as examine the possibility of less restrictive factor mixture models.

In conclusion, among a large population-based sample of Australian young people (14-17 years), psychopathology might be best conceptualised and organized using a hierarchical structure of latent factors that differ in their specificity. These higher order latent factors can adequately and more efficiently account for a majority of the multiple pairwise associations evident between indicators of psychopathology and additional validators including poor sleep, self-harm, suicidality, risky sexual behaviour, and low self-esteem. However, there were significant and relatively large direct associations identified predominately between the validators and factors at the first-order level (i.e. with internalizing and externalizing). These findings provide further evidence for the utility of estimating multiple levels of the psychopathology hierarchy simultaneously as well as identifying the possibility of a hierarchy of clinical risk factors and/or validators that have differential effects at different levels of psychopathology. This is of particular importance among adolescent populations where early intervention and prevention programs offer the greatest potential to reduce the burden of psychopathology.

REFERENCES

- Afzali, M. H., Sunderland, M., Carragher, N., & Conrod, P. (2017). The Structure of Psychopathology in Early Adolescence: Study of a Canadian Sample. *The Canadian Journal of Psychiatry*, 63(4), 223–230. <https://doi.org/10.1177/0706743717737032>
- Albott, C. S., Forbes, M. K., & Anker, J. J. (2018). Association of Childhood Adversity With Differential Susceptibility of Transdiagnostic Psychopathology to Environmental Stress in Adulthood. *JAMA Network Open*, 1(7), e185354. <https://doi.org/10.1001/jamanetworkopen.2018.5354>
- Baglioni, C., Nanovska, S., Regen, W., Spiegelhalder, K., Feige, B., Nissen, C., ... Riemann, D. (2016). Sleep and mental disorders: A meta-analysis of polysomnographic research. *Psychological Bulletin*, 142(9), 969–990. <https://doi.org/10.1037/bul0000053>
- Borschmann, R., Stark, P., Prakash, C., & Sawyer, S. M. (2018). Risk profile of young people admitted to hospital for suicidal behaviour in Melbourne, Australia. *Journal of Paediatrics and Child Health*, 54(11), 1213–1220. <https://doi.org/10.1111/jpc.13938>
- Carragher, N., Teesson, M., Sunderland, M., Newton, N. C., Krueger, R. F., Conrod, P. J., ... Slade, T. (2016). The structure of adolescent psychopathology: A symptom-level analysis. *Psychological Medicine*, 46(5). <https://doi.org/10.1017/S0033291715002470>
- Caspi, A., Houts, R. M., Belsky, D. W., Goldman-Mellor, S. J., Harrington, H., Israel, S., ... Moffitt, T. E. (2014). The p Factor: One General Psychopathology Factor in the Structure of Psychiatric Disorders? *Clinical Psychological Science : A Journal of the Association for Psychological Science*, 2(2), 119–137. <https://doi.org/10.1177/2167702613497473>
- Caspi, A., & Moffitt, T. E. (2018). All for One and One for All: Mental Disorders in One

Dimension. *American Journal of Psychiatry*, 175(9), 831–844.

<https://doi.org/10.1176/appi.ajp.2018.17121383>

Castellanos-Ryan, N., Brière, F. N., O’Leary-Barrett, M., Banaschewski, T., Bokde, A.,

Bromberg, U., ... IMAGEN Consortium. (2016). The structure of psychopathology in adolescence and its common personality and cognitive correlates. *Journal of Abnormal Psychology*, 125(8), 1039–1052. <https://doi.org/10.1037/abn0000193>

Clark, S. L., Muthén, B., Kaprio, J., D’Onofrio, B. M., Viken, R., & Rose, R. J. (2013).

Models and Strategies for Factor Mixture Analysis: An Example Concerning the Structure Underlying Psychological Disorders. *Structural Equation Modeling: A Multidisciplinary Journal*, 20(4), 681–703.

<https://doi.org/10.1080/10705511.2013.824786>

Conway, C. C., Forbes, M. K., Forbush, K. T., Fried, E. I., Hallquist, M. N., Kotov, R., ...

Eaton, N. R. (2019). A Hierarchical Taxonomy of Psychopathology Can Transform Mental Health Research. *Perspectives on Psychological Science*, 14(3), 419–436.

<https://doi.org/10.1177/1745691618810696>

Daraganova, G. (2017). Self-harm and suicidal behaviour of young people aged 14-15 years

old. In *LSAC Annual Statistical Report 2016* (pp. 119–144). Melbourne: Australian Institute of Family Studies.

Erskine, H. E., Moffitt, T. E., Copeland, W. E., Costello, E. J., Ferrari, A. J., Patton, G., ...

Scott, J. G. (2015). A heavy burden on young minds: The global burden of mental and substance use disorders in children and youth. *Psychological Medicine*, 45(7), 1561–1563. <https://doi.org/10.1017/S0033291714002888>

Forbes, M. K., Rapee, R. M., & Krueger, R. F. (2019). Opportunities for the prevention of

mental disorders by reducing general psychopathology in early childhood. *Behaviour*

- Research and Therapy*, 119, 103411. <https://doi.org/10.1016/j.brat.2019.103411>
- Greene, A. L., Eaton, N. R., Li, K., Forbes, M. K., Krueger, R. F., Markon, K. E., ... Kotov, R. (2019). Are fit indices used to test psychopathology structure biased? A simulation study. *Journal of Abnormal Psychology*. <https://doi.org/10.1037/abn0000434>
- Hafekost, J., Boterhoven de Haan, K., Lawrence, D., Sawyer, M. G., & Zubrick, S. R. (2017). *Validation of the Adolescent self-esteem questionnaire: technical report*. Perth, Western Australia.
- Hafekost, J., Johnson, S., Lawrence, D., Sawyer, M., Ainley, J., Mihalopoulos, C., & Zubrick, S. R. (2016). Introducing ‘Young Minds Matter.’ *Australian Economic Review*, 49(4), 503–514. <https://doi.org/10.1111/1467-8462.12163>
- Hafekost, J., Lawrence, D., Boterhoven de Haan, K., Johnson, S. E., Saw, S., Buckingham, W. J., ... Zubrick, S. R. (2016). Methodology of Young Minds Matter: The second Australian Child and Adolescent Survey of Mental Health and Wellbeing. *Australian & New Zealand Journal of Psychiatry*, 50(9), 866–875. <https://doi.org/10.1177/0004867415622270>
- Haltigan, J. D., Aitken, M., Skilling, T., Henderson, J., Hawke, L., Battaglia, M., ... Andrade, B. F. (2018). “P” and “DP:” Examining Symptom-Level Bifactor Models of Psychopathology and Dysregulation in Clinically Referred Children and Adolescents. *Journal of the American Academy of Child & Adolescent Psychiatry*, 57(6), 384–396. <https://doi.org/10.1016/J.JAAC.2018.03.010>
- Hirshkowitz, M., Whiton, K., Albert, S. M., Alessi, C., Bruni, O., DonCarlos, L., ... Adams Hillard, P. J. (2015). National Sleep Foundation’s sleep time duration recommendations: methodology and results summary. *Sleep Health*, 1(1), 40–43. <https://doi.org/10.1016/j.sleh.2014.12.010>

- Hu, L., & Bentler, P. M. (1999). Cutoff criteria for fit indexes in covariance structure analysis: conventional criteria versus new alternatives. *Structural Equation Modeling*, 6(1), 1–55.
- Keyes, K. M., Eaton, N. R., Krueger, R. F., McLaughlin, K. A., Wall, M. M., Grant, B. F., & Hasin, D. (2012). Childhood maltreatment and the structure of common psychiatric disorders. *British Journal of Psychiatry*, 200, 107–115.
- Kotov, R., Waszczuk, M. A., Krueger, R. F., Forbes, M. K., Watson, D., Clark, L. A., ... Zimmerman, M. (2017). The hierarchical taxonomy of psychopathology (HiTOP): A dimensional alternative to traditional nosologies. *Journal of Abnormal Psychology*, 126(4), 454–477. <https://doi.org/10.1037/abn0000258>
- Laceulle, O. M., Vollebergh, W. A. M., & Ormel, J. (2015). The Structure of Psychopathology in Adolescence. *Clinical Psychological Science*, 3(6), 850–860. <https://doi.org/10.1177/2167702614560750>
- Lubke, G. H., & Muthén, B. (2005). Investigating population heterogeneity with factor mixture models. *Psychological Methods*, 10(1), 21–39. <https://doi.org/10.1037/1082-989X.10.1.21>
- Lubman, D. I., Allen, N. B., Rogers, N., Cementon, E., & Bonomo, Y. (2007). The impact of co-occurring mood and anxiety disorders among substance-abusing youth. *Journal of Affective Disorders*, 103(1–3), 105–112. <https://doi.org/10.1016/j.jad.2007.01.011>
- Mars, B., Heron, J., Crane, C., Hawton, K., Lewis, G., Macleod, J., ... Gunnell, D. (2014). Clinical and social outcomes of adolescent self harm: population based birth cohort study. *BMJ (Clinical Research Ed.)*, 349, g5954. <https://doi.org/10.1136/bmj.g5954>
- Masselink, M., Van Roekel, E., & Oldehinkel, A. J. (2018). Self-esteem in Early

- Adolescence as Predictor of Depressive Symptoms in Late Adolescence and Early Adulthood: The Mediating Role of Motivational and Social Factors. *Journal of Youth and Adolescence*, 47(5), 932–946. <https://doi.org/10.1007/s10964-017-0727-z>
- Masyn, K. E., Henderson, C. E., & Greenbaum, P. E. (2010). Exploring the Latent Structures of Psychological Constructs in Social Development Using the Dimensional-Categorical Spectrum. *Social Development (Oxford, England)*, 19(3), 470–493. <https://doi.org/10.1111/j.1467-9507.2009.00573.x>
- McElroy, E., Belsky, J., Carragher, N., Fearon, P., & Patalay, P. (2018). Developmental stability of general and specific factors of psychopathology from early childhood to adolescence: dynamic mutualism or p-differentiation? *Journal of Child Psychology and Psychiatry*, 59(6), 667–675. <https://doi.org/10.1111/jcpp.12849>
- Muthen, L. K., & Muthen, B. O. (2015). *Mplus Users' Guide*. Los Angeles, CA.: Muthen & Muthen.
- Nylund, K. L., Asparouhov, T., & Muthén, B. O. (2007). Deciding on the Number of Classes in Latent Class Analysis and Growth Mixture Modeling: A Monte Carlo Simulation Study. *Structural Equation Modeling: A Multidisciplinary Journal*, 14(4), 535–569. <https://doi.org/10.1080/10705510701575396>
- Paus, T., Keshavan, M., & Giedd, J. N. (2008). Why do many psychiatric disorders emerge during adolescence? *Nature Reviews Neuroscience*, 9(12), 947–957. <https://doi.org/10.1038/nrn2513>
- Prendergast, L. E., Toumbourou, J. W., McMorris, B. J., & Catalano, R. F. (2019). Outcomes of Early Adolescent Sexual Behavior in Australia: Longitudinal Findings in Young Adulthood. *Journal of Adolescent Health*, 64(4), 516–522. <https://doi.org/10.1016/J.JADOHEALTH.2018.10.006>

- Raftery, A. E. (1995). Bayesian model selection in social research. *Sociological Methodology*, 25, 111–163.
- Rodriguez, A., Reise, S. P., & Haviland, M. G. (2016). Evaluating Bifactor Models : Calculating and Interpreting Statistical Indices. *Psychological Methods*, 21(2), 137–150. <https://doi.org/10.1037/met0000045>
- Shaffer, D., Fisher, P., Lucas, C., Hilsenroth, M. J., & Segal, D. L. (2004). The diagnostic interview schedule for children (DISC). In *Comprehensive handbook of psychological assessment* (pp. 256–270).
- Shaffer, D., Fisher, P., Lucas, C. P., Dulcan, M. K., & Schwab-Stone, M. E. (2000). NIMH Diagnostic Interview Schedule for Children Version IV (NIMH DISC- IV): Description, differences from previous versions, and reliability of some common diagnoses. *Journal of the American Academy of Child and Adolescent Psychiatry*, 39(1), 28–38. <https://doi.org/10.1097/00004583-200001000-00014>
- Slade, T., McEvoy, P. M., Chapman, C., Grove, R., & Teesson, M. (2015). Onset and temporal sequencing of lifetime anxiety, mood and substance use disorders in the general population. *Epidemiology and Psychiatric Sciences*, 24(1), 45–53. <https://doi.org/10.1017/S2045796013000577>
- Starr, L. R., Conway, C. C., Hammen, C. L., & Brennan, P. A. (2014). Transdiagnostic and disorder-specific models of intergenerational transmission of internalizing pathology. *Psychological Medicine*, 44(1), 161–172. <https://doi.org/10.1017/S003329171300055X>
- Sunderland, M., Carragher, N., Chapman, C., Mills, K., Teesson, M., Lockwood, E., ... Slade, T. (2016). The shared and specific relationships between exposure to potentially traumatic events and transdiagnostic dimensions of psychopathology. *Journal of Anxiety Disorders*, 38, 102–109. <https://doi.org/10.1016/j.janxdis.2016.02.001>

- Sunderland, M., & Slade, T. (2014). The relationship between internalizing psychopathology and suicidality, treatment seeking, and disability in the Australian population. *Journal of Affective Disorders, 171*, 6–12. <https://doi.org/10.1016/j.jad.2014.09.012>
- Tackett, J. L. (2006). Evaluating models of the personality–psychopathology relationship in children and adolescents. *Clinical Psychology Review, 26*(5), 584–599. <https://doi.org/10.1016/J.CPR.2006.04.003>
- Teesson, M., Slade, T., & Mills, K. (2009). Comorbidity in Australia: Findings of the 2007 National Survey of Mental Health and Wellbeing. *Australian and New Zealand Journal of Psychiatry, 43*(7), 606–614. <https://doi.org/10.1080/00048670902970908>
- Vandenberg, R. J., & Lance, C. E. (2000). A Review and Synthesis of the Measurement Invariance Literature: Suggestions, Practices, and Recommendations for Organizational Research. *Organizational Research Methods, 3*(1), 4–70. <https://doi.org/10.1177/109442810031002>
- Waldman, I. D., Poore, H. E., van Hulle, C., Rathouz, P. J., & Lahey, B. B. (2016). External validity of a hierarchical dimensional model of child and adolescent psychopathology: Tests using confirmatory factor analyses and multivariate behavior genetic analyses. *Journal of Abnormal Psychology, 125*(8), 1053–1066. <https://doi.org/10.1037/abn0000183>
- Wright, A. G. C., Krueger, R. F., Hobbs, M. J., Markon, K. E., Eaton, N. R., & Slade, T. (2013). The structure of psychopathology: Toward an expanded quantitative empirical model. *Journal of Abnormal Psychology, 122*(1), 281–294.

Table 1: Sample descriptive statistics

		n	%	SE
Clinical/lifestyle factors	Poor sleep	451	21.4	1.05
	Self-harm	214	10.0	0.70
	Suicidal ideation	190	9.1	0.69
	Suicide attempt	65	3.0	0.40
	Multiple sexual partners	116	5.0	0.51
	Sex without condoms	255	10.4	0.73
	Low self esteem	79	3.9	0.44
Psychopathology	Daily smoker	181	7.8	0.69
	Binge drinker	367	15.4	0.85
	Cannabis use	330	14.0	0.86
	Other illegal drug use	90	4.0	0.45
	Separation anxiety disorder	94	4.4	0.49
	Social anxiety disorder	70	3.6	0.47
	Generalized anxiety disorder	74	3.5	0.43
	Major depressive disorder	136	6.2	0.59
	Attention deficit hyperactivity disorder	107	5.6	0.60
	Conduct disorder	36	1.9	0.40
	Paranoia	294	15.1	0.90
	Special messages	77	3.7	0.45
	Mind reading	115	6.1	0.61
	Visual hallucinations	225	12.1	0.84

Auditory hallucinations	308	15.9	0.91
-------------------------	-----	------	------

Table 2: Tetrachoric correlation matrix for indicators of psychopathology

	SAD	SA	GAD	MDD	ADHD	CD	BINGE	TOB	CAN	DRG	PSY1	PSY2	PSY3	PSY4	PSY5
SAD	1.000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
SA	0.588	1.000	-	-	-	-	-	-	-	-	-	-	-	-	-
GAD	0.684	0.665	1.000	-	-	-	-	-	-	-	-	-	-	-	-
MDD	0.619	0.675	0.718	1.000	-	-	-	-	-	-	-	-	-	-	-
ADHD	0.357	0.201	0.364	0.274	1.000	-	-	-	-	-	-	-	-	-	-
CD	0.182	0.283	0.009	<i>0.430</i>	<i>0.539</i>	1.000	-	-	-	-	-	-	-	-	-
Binge	0.033	0.056	0.110	0.177	0.120	0.416	1.000	-	-	-	-	-	-	-	-
Tob	0.219	0.239	0.174	<i>0.339</i>	0.268	0.698	0.675	1.000	-	-	-	-	-	-	-
Can	0.057	0.130	0.111	0.292	0.131	0.554	0.697	0.797	1.000	-	-	-	-	-	-
Drg	0.107	0.051	0.167	0.277	0.136	0.303	0.602	0.721	0.736	1.000	-	-	-	-	-
Psy1	0.263	0.191	0.277	0.281	0.106	0.156	0.196	0.276	0.282	<i>0.359</i>	1.000	-	-	-	-
Psy2	0.113	0.156	0.189	0.114	0.127	0.250	0.144	0.174	0.299	0.139	0.486	1.000	-	-	-
Psy3	0.112	0.131	0.144	0.244	0.038	0.299	0.047	0.165	0.185	0.223	0.560	0.612	1.000	-	-

Psy4	0.193	0.184	<i>0.328</i>	0.194	0.083	0.184	0.170	<i>0.334</i>	0.261	<i>0.375</i>	0.600	0.642	0.565	1.000	-
Psy5	0.108	0.129	<i>0.395</i>	0.269	0.074	0.235	0.121	<i>0.328</i>	<i>0.321</i>	<i>0.386</i>	0.592	0.518	0.496	0.794	1.000

Notes: SAD=social anxiety disorder, SA=separation anxiety, GAD=generalised anxiety disorder, MDD= major depressive disorder,

ADHD=attention deficit hyperactivity disorder, CD=conduct disorder, BINGE= binge drinking, TOB=daily smoking, CAN=cannabis use,

DRG=other illegal drug use, PSY1=paranoia, PSY2=special messages, PSY3=mind reading, PSY4=visual hallucinations, PSY5=auditory

hallucinations. Bold indicate correlations across disorders from conceptually related domains. Italics indicate moderate to large correlations

(>0.3) across disorders from different domains

Table 3: Fit statistics for CFA, LCA, and FMMs.

CFA	LL	k	BIC	saBIC	LMR	Entropy	CFI*	TLI*	RMSEA*
One factor	-7122.06	30	14472.18	14376.87	-	-	0.739	0.695	0.071
Three correlated factors	-6677.50	33	13605.87	13501.03	-	-	0.984	0.981	0.018
Bifactor	-6656.04	45	13654.11	13511.21	-	-	0.988	0.983	0.017
Higher order (1 + 3)	-6677.50	33	13605.87	13501.03	-	-	0.984	0.981	0.018
LCA									
2 class	-7145.94	31	14527.56	14429.07	<0.001	0.798	-	-	-
3 class	-6903.15	47	14163.61	14014.29	<0.001	0.858	-	-	-
4 class	-6759.38	63	13997.72	13797.56	0.014	0.897	-	-	-
5 class	-6682.79	79	13966.16	13715.18	0.022	0.880	-	-	-
6 class	-6645.01	95	14012.25	13710.43	0.106	0.883	-	-	-
FMM									
FMM - 3 factor 2 class	-7145.94	31	14527.56	14429.07	<0.001	0.798	-	-	-
FMM - 3 factor 3 class	-6917.51	35	14101.11	13989.91	<0.001	0.859	-	-	-
FMM - 3 factor 4 class	-6784.60	39	13865.69	13741.78	0.056	0.897	-	-	-

FMM - 3 factor 5 class	-6718.23	43	13763.36	13626.75	0.007	0.880	-	-	-
FMM - 3 factor 6 class	-6690.71	47	13738.73	13589.40	0.235	0.881	-	-	-
FMM - 3 factor 7 class	-6671.87	51	13731.46	13569.43	0.033	0.880	-	-	-
FMM - 3 factor 8 class	-6661.68	55	13741.49	13566.76	0.529	0.827	-	-	-

Notes: * estimated using WLSMV in Mplus otherwise all models were estimated using MLR. All FMM models were estimated with variances and covariances fixed to 0 and factor loadings and thresholds invariant across classes.

Figure 1: Path diagram of example model examining total, direct and indirect effects between poor sleep and the structural model of psychopathology.

Figure 2: Path diagram of the best fitting factor model. SA=separation anxiety, SAD=social anxiety disorder, GAD=generalised anxiety disorder, MDD= major depressive disorder, ADHD=attention deficit hyperactivity disorder, BINGE= binge drinking, TOB=daily smoking, CAN=cannabis use, DRG=other illegal drug use, CD=conduct disorder, PSY1=paranoia, PSY2=unusual beliefs, PSY3=mind reading, PSY4=visual hallucinations, PSY5=auditory hallucinations, INT=internalizing, EXT=externalizing, PSY=psychotic like experiences, p=general psychopathology. All loadings are standardized and significant at $p < 0.05$.

Figure 3: Best fitting seven class 3 factor FMM predicted probabilities. SA=separation anxiety, SAD=social anxiety disorder, GAD=generalised anxiety disorder, MDD= major depressive disorder, ADHD=attention deficit hyperactivity disorder, Binge= binge drinking, TOB=daily smoking, CAN=cannabis use, DRG=other illegal drug use, CD=conduct disorder, PSY1=paranoia, PSY2=unusual beliefs, PSY3=mind reading, PSY4=visual hallucinations, PSY5=auditory hallucinations.

Figure 4: Ratio of absolute standardized direct effects and absolute standardized indirect effects for each indicator of psychopathology on key validity indicators in the higher order model. SA=separation anxiety, SAD=social anxiety disorder, GAD=generalised anxiety disorder, MDD= major depressive disorder, ADHD=attention deficit hyperactivity disorder, Binge= binge drinking, TOB=daily smoking, CAN=cannabis use, DRG=other illegal drug use, CD=conduct disorder, PSY1=paranoia, PSY2=unusual beliefs, PSY3=mind reading, PSY4=visual hallucinations, PSY5=auditory hallucinations, INT = internalizing, EXT = externalizing, PSY=psychotic like experiences.