GROUPS 4 HEALTH versus Cognitive Behaviour Therapy in young people with depression and loneliness: A randomized, Phase 3, non-inferiority trial with 12-month follow-up

Tegan Cruwys PhD ^{1*}	tegan.cruwys@anu.edu.au
Catherine Haslam PhD ²	c.haslam@uq.edu.au
Joanne A. Rathbone BPsychHons ²	j.rathbone@uq.edu.au
Elyse Williams PhD ²	e.williams@uq.edu.au
S. Alexander Haslam PhD ²	a.haslam@uq.edu.au
Zoe C. Walter PhD ²	z.walter@uq.edu.au

1. Research School of Psychology, The Australian National University, Canberra, ACT, Australia
2. School of Psychology, The University of Queensland, Brisbane, QLD, Australia

* Correspondence to: Dr Tegan Cruwys, Research School of Psychology, 39 Science Rd, The Australian National University, Canberra ACT 2601 AUSTRALIA
tegan.cruwys@anu.edu.au

Accepted for publication in *British Journal of Psychiatry* on 17 August 2021.

Citation:

Cruwys, T., Haslam, C., Rathbone, J. A., Williams, E., Haslam, S. A. & Walter, Z. C. (2021). GROUPS 4 HEALTH versus Cognitive Behaviour Therapy in young people with depression and loneliness: A randomized, Phase 3, non-inferiority trial with 12-month follow-up. *British Journal of Psychiatry*.

Abstract

Background. Depression treatments are typically less effective for young people than for adults. However, extant treatments rarely target loneliness, which is a key risk factor in the onset, maintenance, and development of depression.

Aims. This study evaluated the efficacy of a novel loneliness intervention, *GROUPS 4 HEALTH* (G4H), relative to best-practice treatment *Cognitive Behaviour Therapy* (CBT) in reducing loneliness and depression over a 12 month period.

Methods. The study was a Phase 3, randomised non-inferiority trial comparing G4H to dose-controlled group CBT. Participants were 174 young people aged 15-25 years experiencing loneliness and clinically significant symptoms of depression, who were not in receipt of adjunct treatment. Participants were recruited from mental health services in Southeast Queensland, Australia. Randomisation was conducted using computer software. Follow-up assessments and statistical analyses were blind to allocation. Both interventions consisted of 5 x 75 min group-based psychotherapy sessions. The primary outcomes were depression and loneliness, with a non-inferiority margin of 2.20 for depression. The trial was prospectively registered on the Australian New Zealand Clinical Trial Registry ([ACTRN12618000440224](https://www.anzctr.org.au/Trial/Registration/TrialRegistration.aspx?ACTRN12618000440224)) on 27 March 2018 and is closed to new participants.

Results. 174 participants were enrolled between 24 April 2018 and 25 May 2019, with 84 in the G4H condition and 90 in the CBT condition. All randomised participants were included in the intention-to-treat analyses (N=174). The pre-post effect sizes for depression were $d=-.71_{G4H}$ and $d=-.91_{CBT}$. For loneliness, they were $d=-1.07_{G4H}$ and $d=-0.89_{CBT}$. At 12-month follow-up, the absolute difference between groups in depression was 1.176 (95%CI: -1.94, 4.29), and in loneliness was -0.679 (95%CI: -.1.43, .07). No adverse effects were observed.

Conclusions. G4H was non-inferior to CBT for depression and showed a slight advantage over CBT for loneliness that emerged after treatment completion.

Keywords

loneliness; group psychotherapy; mood disorders; depression; mental health; adolescence; youth; social isolation; social identity

GROUPS 4 HEALTH versus Cognitive Behaviour Therapy in young people with depression and loneliness: A randomized, Phase 3, non-inferiority trial with 12-month follow-up

Depression is the leading global cause of disability among young people aged 15-25 years.¹ A recent population study of over 600,000 United States residents found that depression prevalence was not only greatest among those aged 12-25 years, but had also risen over the last ten years among this group.² Ineffective management of depression can set up a young person for a lifetime of recurrent episodes and associated functional impairments, including academic underperformance and failure to attain adult milestones (e.g., career progression, independent living).³

Existing treatments for depression in young people are effective, but suffer from a number of drawbacks. First, psychopharmacological options that constitute frontline depression treatment among adults have lower efficacy and compliance among young people.⁴ Studies have suggested that young people and their parents prefer non-drug treatments for depression.⁵ Unfortunately, psychotherapeutic treatment efficacy is more modest in young people than in adults. One meta-analysis found that effect sizes of cognitive behaviour therapy (CBT) for depression were 30% smaller for children and/or adolescents than for adults.⁶ Another meta-analysis included samples with a mean age of less than 18 years and found that CBT for depression was effective in this population but had a smaller effect size ($g=.35$) than among adults and was no better than other forms of psychotherapy.⁷

Loneliness as a target of treatment

There is a need for innovation in depression treatment, particularly among young people, with an emphasis on strategies that not only improve symptoms, but also reduce the risk of relapse. In seeking to address this need, a growing evidence base has pointed to the role of social and contextual factors in the development and maintenance of depression. In particular, loneliness has been identified as a particularly strong and robust risk factor in the onset, maintenance and relapse of depression.⁸ Loneliness is related to other constructs

such as social isolation, but distinctly captures the *subjective* absence of meaningful connections to others. Loneliness has probable causal links to mental illness and causes significant distress and impairment in its own right. Although most research evaluating loneliness interventions has been conducted with older people⁹, loneliness is more common among young people.¹⁰

Previous trials of interventions targeting loneliness and its downstream mental health consequences have had mixed success, with a 2011 meta-analysis of loneliness interventions describing their benefit as “modest”.⁹ This meta-analysis included 50 studies, and among the 20 randomized trials the average effect size was small ($d=-0.19$). Another review found that loneliness interventions conducted in group contexts tended to be more effective.¹¹

In the years since these reviews, GROUPS 4 HEALTH (G4H) has emerged as a promising intervention for loneliness, with concomitant benefits for depression and general wellbeing.^{12,13} G4H is a manualised group therapy program founded on the social identity theoretical framework. It seeks to increase group-based belonging using a combination of educational, psychotherapeutic, and behavioural strategies. In a Phase 2 randomized controlled trial (RCT) published in 2019 with 120 participants reporting loneliness and mental illness, G4H outperformed treatment-as-usual in reducing loneliness ($d=-1.04$) at four-month follow-up, and performed comparably to treatment-as-usual in reducing depression ($d=-.63$).¹³ Two relevant meta-analyses were also published in 2019. The first found a negative association ($r_z=-.15$) between social identity and depression in 76 (mostly correlational) studies.¹⁴ The second evaluated the efficacy of interventions that build social identity, including G4H.¹⁵ Among the 27 studies identified, 8 included depression as an outcome variable, with an average effect size of $g=-.58$. This emerging evidence base suggests that social identity-based interventions, and G4H in particular, show promise in reducing loneliness and depression.

What is not known, however, is how these beneficial effects of G4H on depression compare to best-practice evidence-based treatment for young people: CBT. For G4H to be widely implemented in practice, it needs not only to perform strongly on loneliness relative to treatment-as-usual, but also to show benefits for depression that are not inferior to gold-standard treatment. It is also crucial that the relative benefits of these interventions are assessed over both the short and longer term, because depression tends to follow a relapsing-remitting pattern.

The current study

The overarching aim of this study was to compare G4H to CBT on primary outcomes of loneliness and depression across a one-year follow-up period. We conducted a Phase 3 RCT in which young people were recruited from mental health services and assigned to receive either G4H or a dose-controlled, manualized format of group-based CBT. Our hypotheses were that:

H1: At program completion, G4H would be superior to CBT in reducing loneliness. H1 would be considered supported if there were evidence of significant improvements in loneliness in the G4H condition (H1a), and a significant condition by time interaction, such that improvements in loneliness over time were greater for the G4H condition than in the CBT condition (H1b).

H2: At program completion, G4H would be as effective as CBT in reducing symptoms of depression (i.e., be non-inferior). H2 would be considered supported if there were evidence of significant improvements in depression in both conditions, with a non-significant condition by time interaction. The non-inferiority margin was a DASS-21 depression change score difference of 2.20 (see protocol¹⁶ for details of calculation).

H3: The benefits of G4H for loneliness and depression would be sustained at 12-month follow-up. H3 would be considered supported if there were no significant increase in

depression and loneliness scores between 6-month and 12-month follow-up in the G4H condition.

Method

Study Design

The study was a randomized, Phase 3, non-inferiority trial with 12-month follow-up. The trial profile is provided in Figure 1. SPIRIT and CONSORT Guidelines were followed. The trial was conducted at two mental health services in South-East Queensland, Australia: a community-based youth mental health service (*headspace*) and a university-based psychology clinic (*University of Queensland*). A total of 26 therapy groups were completed (13 of each type). The trial recruited 15-25 year olds because this corresponds to the population mean age of depression onset and peak prevalence.³

Written informed consent was obtained from all participants. For young people aged under 16 who wished to participate, parental consent was also sought in accordance with the organizational policy of the healthcare services where the group was taking place. The authors assert that all procedures complied with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008. All procedures were approved by the Brisbane Metro South HREC/18/QPAH/54, ratified by the University of Queensland #2018000420 and were prospectively registered on the Australian New Zealand Clinical Trial Registry ([ACTRN12618000440224](https://www.anzctr.org.au/Trial/Registration/TrialReview.aspx?id=12618000440224)) on 27 March 2018. The published study protocol is [available](#).¹⁶

Participants

Participants were 174 young people seeking mental healthcare. Table 1 shows baseline characteristics of the sample. The sample ranged in age from 15-25 years with an average of 19 (SD = 2.01); 159 participants were female (74%) and 55 were male (26%).

Participants were invited to participate in the trial if they were subjectively experiencing low mood and loneliness. Inclusion criteria were: (1) aged 15-25 years, (2) English speaking, at

a sufficient level to understand and engage with group discussions, with the presence of at least one of the following: (3) depressive symptoms of at least mild clinical impairment on the PHQ-9 (≥ 5)¹⁷, or (4) a mental health condition diagnosed by a health professional (e.g., major depressive disorder), or (5) elevated loneliness on the UCLA-20 (≥ 40 ; approximately 1 SD above the mean for 16 year olds¹⁸).

Exclusion criteria were: (1) receiving other concurrent evidence-based treatment for depression (psychopharmacological or psychological), (2) severe suicide ideation, (3) severe psychological symptoms that would interfere with ability to participate (e.g., psychotic episode, substance dependence/intoxication), and (4) severe neurological condition or learning impairment that would limit capacity to engage with the intervention. Participants were monitored by group facilitators for changes that might affect their eligibility (e.g., commencing adjunct treatment, suicidality).

Recruitment used a variety of strategies. Staff at each service were provided with referral information and directed clients to the trial, particularly where they would otherwise be placed on a treatment waitlist. Postcards and posters were displayed at each service, and used to advertise the trial within the community and on university campuses to increase sample representativeness of the intended population, rather than just those presenting to healthcare services.⁷

To improve retention and data quality, participants were offered incentives for completing measures at program completion (\$AUD15 voucher), six months (\$AUD50) and 12 months (\$AUD50). Those who were enrolled in undergraduate psychology at the university where the trial was taking place were offered course credit.

Randomization and masking

In order to reduce expectation effects, the *noninferiority* focus of the trial was emphasized to participants (e.g., “we are investigating whether both treatments are equally effective”). After providing informed consent, participants were assigned to a numbered therapy group. Once

a minimum number of participants had been recruited to commence a group (5 people), the group (and all participants within it) were randomly allocated to G4H or CBT. Randomization was conducted via a computer-generated random number in a 1:1 ratio.

Trial managers conducted group assignment and were not blind. Participants were blind to allocation until after they had confirmed their attendance. All other researchers were blind to allocation. Assessment at follow-up timepoints was conducted via online questionnaire which was identical for all participants, to reduce experimenter bias. Invitations to complete the assessments were identical across conditions and were sent via email and SMS. Statistical analysis was conducted by the lead author without knowledge of allocation (i.e., the meaning of the condition codes [1=G4H; 2=CBT] was only provided after hypothesis testing was complete). All authors had access to the data and project files via a shared university drive. To ensure blinding was maintained, the non-blind trial database was password protected with access only possible for trial managers.

Procedures

Group facilitators were provisional psychologists completing a professional psychology graduate program. In addition to their standard training in supportive counselling, risk assessment, and group therapy, all facilitators were provided with focused training to run the manualized program that they were delivering (G4H or CBT), and 1 hr/week supervision by a clinical psychologist with expertise in G4H or CBT. Adherence to the manualised program was maintained via direct observation, weekly supervision, recorded focus groups with participants and facilitators after program completion, and a validated anonymous questionnaire completed by facilitators after each session.^{12,13}

GROUPS 4 HEALTH. G4H¹² is a psychotherapeutic program that seeks to reduce loneliness by building group-based belonging. It is delivered in groups of 5-8 people and comprises 5 sessions of approximately 75 mins each. Session content is informed by social identity theorising. Participants first learn about the importance of social connectedness and group-based belonging for health (Session 1). They are then guided to develop a visual *social*

identity map to gauge current group belonging (Session 2), before developing goals and plans for engaging with existing groups (Session 3) and for joining new groups (Session 4). These are delivered weekly, before participants meet one month later to review their progress and goals (Session 5).

Cognitive Behaviour Therapy. The comparison condition was dose-controlled manualised group-based CBT for depression: The Blues Program.¹⁹ Minor manual adaptations were made to match the timing and amount of face-to-face contact to G4H; the 6 x 60 minute format was adapted to a 5 x 75 min format. A meta-analysis found that The Blues Program was at least as effective as CBT of longer duration, and better in reducing depression symptoms among high-risk young people than several active comparator interventions.²⁰

Follow-ups. Assessments were completed at eight timepoints: baseline (T0), after each of the five intervention sessions (weekly for 4 weeks [T1-T4], and then 1 month later at program completion [T5]) then at 6 months (T6) and 12 months (T7). All randomised participants were invited to complete all timepoints, including those who discontinued (see below).

Outcomes

Primary outcomes. The primary outcomes were *depression symptom severity*, measured with the depression subscale of the DASS-21²¹, and *loneliness*, measured with the UCLA-4.²²

Secondary outcomes. Additional outcome and process variables were measured at only four timepoints (T0, T5, T6, and T7). Secondary outcomes were social anxiety, general practitioner visits, wellbeing, life satisfaction, self-esteem, and subjective physical health. Process measures were multiple group membership, group compatibility, mental illness identity, group cohesion, therapy group identification, and service identification. Full details of these measures are provided in the protocol¹⁶; analyses are not reported here due to space constraints.

Assessment of safety and adverse events. Facilitators monitored participant wellbeing in session and via assessment data. The protocol allowed facilitators to provide brief ad-hoc supportive counselling as needed without compromising participant eligibility. Facilitators reported all concerns about participants to their supervisor for case management. Any severe decline in mental health would render participants ineligible for the trial, leading to referral for more intensive mental health treatment, and be recorded as an adverse event. No adverse events were recorded.

Statistical Analysis

Full justification of our approach to analysis is provided in the protocol.¹⁶ Two mixed-model repeated-measures (MMRM) models were used to test hypotheses: one to predict loneliness (Figure 2) and one to predict depression (Figure 3). Each model had a nested structure which included fixed effects for timepoint (Level 1), condition, and their interaction, random intercepts for participant (Level 2), and random intercepts for therapy group (Level 3).

Follow-up planned contrasts using estimated marginal means were calculated as appropriate, with correction for multiple comparisons using the multivariate t distribution. Analyses were conducted in R (v.3.4.0), supported by packages *lme4* and *emmeans*.

Timepoint was treated as a categorical variable, with patterns of interaction assessed not only for linear but also non-linear patterns (up to the 6th degree). This was because the time between follow-ups in the experimental design was non-linear and the duration of treatment was T0-T5 only. The analyses were therefore sensitive to non-linear symptom change (e.g., if a treatment led to large initial improvements and symptoms subsequently stabilised).

Missing data were managed using Full Information Likelihood Maximization, which is readily implemented in MRMM to honour the intention-to-treat principle.²³ This approach utilises all available data for all participants. Overall, retention was high throughout the trial. Among the full (intention-to-treat) sample, the median and modal number of timepoints completed in both groups was 8 (i.e., complete data). Similarly, the median and modal number of therapy

sessions attended was 5 (i.e., full attendance) in both groups. 173 participants had at least two timepoints of data, and 169 participants completed at least one of the three follow-up timepoints (T5, T6, T7). T5 had the lowest sample size with 155 respondents. An *a priori* power analysis estimated that 67 people per condition would be needed to have 0.80 power to detect the hypothesised effect with an alpha level of .05. All timepoints met this criterion.

Per-protocol analyses were also conducted. These excluded participants who did not meet eligibility criteria throughout the trial period, were not randomly assigned, did not complete the intervention according to the protocol, or who were missing data at either baseline (T0) or follow-up (T5, T6, and T7). This led to the exclusion of 22 participants who did not attend at least three of the five sessions ($n=17$), who commenced adjunct treatment ($n=3$), or did not complete any of the follow-up timepoints ($n=2$). The likelihood of discontinuation did not differ between the G4H and CBT groups, $\chi^2(1)=2.38, p=.123$.

Results

Hypothesis testing

Full details of analyses are provided in the Supplementary Appendix. H1a predicted significant improvement in loneliness in the G4H condition. Contrasts were specified to evaluate this within the G4H condition from baseline to program completion, $t(1086)=-7.96, p<.0001, d=-.91$, to 6-month follow-up, $t(1085)=-7.85, p<.0001, d=-.89$, and to 12-month follow-up, $t(1084)=-9.12, p<.0001, d=-1.07$. Each of these analyses were significant, supporting H1a.

H1b predicted that improvements in loneliness would be greater in the G4H condition than the CBT condition. To assess this, the interaction term between time and condition was examined. Although the linear interaction was not significant, $t(1083)=0.51, p=.613$, the *quadratic* interaction was significant, $t(1090)=2.12, p=.034$ (supporting H1b). This indicated a non-linear trajectory of recovery that differed between the two groups. Underpinning this significant interaction, simple comparisons indicated that although loneliness decreased steeply between T0 and T2 in both the CBT and G4H conditions ($t_{\text{CBT}}(1077)=6.61, p<.001$;

$t_{G4H}(1085)=4.02, p=.002$), loneliness then stabilised in the CBT condition between T2 and T7, but continued to improve in the G4H condition, ($t_{CBT}(1078)=1.94, p=.524$; $t_{G4H}(1086)=4.86, p<.001$). The absolute difference between groups in loneliness at T7 was -0.679 (95%CI: -.1.43, .07), $t(67.4)=-1.81, p=.075$.¹

H2 predicted that G4H would be non-inferior to CBT in improving depression symptom severity, conceptualized as a change score difference that was both non-significant and less than 2.20 at all follow-up timepoints. Supporting H2, all seven planned contrasts between T0 and each other timepoint were non-significant ($ps>.143$), and at no timepoint did the difference exceed 2.2 (the largest was -2.12 for the T0-T7 comparison). The absolute difference between groups in depression at the 12-month follow-up was 1.176 (95%CI: -1.94, 4.29), $t(61.7)=1.18, p=.453$.

H3 predicted that the benefits of G4H for loneliness and depression would be sustained over time. Supporting H3, planned contrasts indicated no significant change in loneliness between program completion and 6-month follow-up, $t(1080)=-.17, p>.999$, or between 6- and 12-month follow-up, $t(1079)=-1.24, p=.912$. Similarly, for depression, there was no significant change in the G4H condition between program completion and 6-month follow-up, $t(1085)=.34, p>.999$, and limited evidence of change between 6- and 12-month follow-up: $t(1079)=-2.84, p=.087$.

Per-protocol analyses

All hypothesis testing was repeated using only the subsample ($n=152$) who completed the program in accordance with the trial protocol. All of the findings were replicated and each hypothesis supported (see full details in the Supplementary Appendix).

1. There are no formal guidelines for clinically meaningful change on the UCLA-4. However, previous research has treated a change of 2 points on the UCLA-20 (3.3%) as clinically significant and this would correspond to a change of 0.4 points on the UCLA-4. By this benchmark, the difference of 0.68 between G4H and CBT groups (0.69 in the per-protocol analysis) at T7 would be considered clinically meaningful.

Discussion

This non-inferiority RCT compared GROUPS 4 HEALTH (G4H) to dose-controlled group cognitive behaviour therapy (CBT) across a 12-month follow-up. Loneliness improved significantly in both conditions, with G4H showing a slight advantage at 12-month follow-up. Depression improved comparably in both conditions. The reason for the different patterns in loneliness and depression outcomes in the two groups is not clear. However, on theoretical grounds, this could tentatively be attributed to the explicit focus on social group connection in G4H (but not CBT) and on depression in CBT (but not G4H).

Meta-analytic evidence shows that depression treatments are typically less effective for young people than for adults.^{4,6} However, few extant treatments target loneliness, which is a key risk factor in the onset, maintenance, and development of depression.⁸ Interventions that do target loneliness have limited efficacy and focus almost exclusively on older people.⁹ Promising interventions that address this lacuna focus on building group-based belonging in accordance with a social identity theoretical framework.¹² Indeed, recent meta-analysis showed that social identity-based interventions of this form had effect sizes for depression that rivalled those of established treatments.¹⁵

Extending this line of research, the present Phase 3 trial is the first to evaluate a social identity intervention against another evidence-based treatment on loneliness and depression. This represented a conservative test of G4H, given that its efficacy has been attributed, at least in part, to the group-based nature of the intervention (which the CBT condition also had) and that it targets depression symptoms only indirectly (while the CBT condition targeted these symptoms directly). This trial found that both G4H ($d=-.71$) and CBT ($d=-.91$) were effective in reducing depression symptoms. At no timepoint were there significant differences between conditions on depression, and all numeric differences were smaller than the non-inferiority margin of 2.20. The effect sizes, particularly for CBT, were greater than the average reported in previous meta-analyses.^{6,7} This may be partly attributable to the controlled setting of the trial, including the manualised nature of the

treatments, monitoring of facilitator adherence, and the group delivery format (as explained further below).

The trial yielded several other novel findings. First, previous treatment efforts for loneliness have had mixed success.⁹ This is the strongest evidence to date for the benefit of G4H, with effect sizes much greater than the average of extant interventions ($d=-1.07$ vs. -0.19). The effect size in the present trial for loneliness in the G4H condition was almost identical to that found in the Phase 2 trial,²¹ further increasing our confidence in its efficacy. Furthermore, few loneliness intervention studies provide evidence of sustained benefit over time. Here, we found evidence that benefits for loneliness were sustained a year later. This is consistent with both our predictions, and a key goal of G4H which is to develop skills to build and maintain group-based belonging, including during life transitions and times of stress.¹²

Second, this trial provides the first evidence of the efficacy of depression-focused CBT in reducing loneliness ($d=-0.89$). Although these benefits for loneliness were not as large or sustained as G4H, they were larger than average effect sizes for other loneliness-focused interventions. Importantly, though, this trial utilized a group-based format of CBT. Therefore, these benefits may be attributable to the group environment as much as to the content of the intervention itself. Indeed, previous studies have indicated that loneliness interventions are most effective when delivered in a group format.¹¹ Furthermore, CBT is more effective for depression when people feel a sense of belonging with their therapy group.²⁴ These results thus speak to the value of *group-based delivery* of psychotherapy, not only to the value of CBT.

This trial had many strengths: (1) it had a well-powered design and high rate of retention, (2) it included an active dose-controlled comparison group, (3) it recruited from real-world health services across public and private sectors, and (4) it assessed outcomes over a long follow-up period against reliable and valid measures. However, like all studies, it also had limitations. In particular, it was not possible to blind participants or facilitators to their

assignment once the groups had commenced. In order to reduce demand characteristics related to this, the *noninferiority* focus of the trial was emphasized at all times.

The COVID-19 pandemic has raised the profile of loneliness, not only in terms of its prevalence and its effects on mental health, but also the limited evidence-base on which we can intervene to address it.²⁵ Having established the efficacy of G4H, future research needs to also evaluate its effectiveness, particularly when delivered by health professionals with minimal researcher oversight. It will also be important to establish its scalability in large implementation trials and adaptability for remote delivery. Acceptability and feasibility of the intervention in a wide variety of healthcare contexts is yet to be established. Although attrition from this trial did not differ significantly between groups, further investigation of the rate and reasons for treatment drop-out in G4H is warranted. The group-based delivery format of G4H, and its suitability for delivery by trainees, mean that the intervention is likely to be a cost-effective alternative to treatment-as-usual. However, this too will be important to establish quantitatively in a formal cost-effectiveness evaluation.

This study has implications for clinical practice, patient care, and policy. Its findings suggest that a clinical focus on loneliness is warranted in treatment of depressed young people. Importantly, this also aligns with the lived experience of young people, who often identify a social trigger for the onset of their first depressive episode and typically prefer an active, skills-based approach to treatment.⁵ However, as the evidence presented here illustrates, loneliness-oriented interventions are likely to be most effective when they draw upon the established science of social relationships embodied by the social identity framework. Indeed, this framework holds promise for the improvement of clinical interventions more broadly – an opportunity that is yet to be fully realised. Crucially, the two interventions compared here both achieved large effect sizes for depression and loneliness despite both being brief, group-based, and manualised; making them more resource-efficient than treatment-as-usual. Depression typically has a remitting-recurring course and causes substantial impairment in functioning for young people. The efficacy of group-based G4H

and CBT suggests that policymakers should prioritise efforts to make them accessible for young people.

Declaration of Interest. The authors TC, CH and AH are the developers of the GROUPS 4 HEALTH program, and The University of Queensland holds its associated Intellectual Property. These authors have an academic, rather than commercial, interest in the program, with training offered at cost and additional funds directed to ongoing program development and evaluation. The other authors have no interests to declare.

Funding Statement. This research was funded by Australian Rotary Health and the National Health and Medical Research Council (#1173270).

Acknowledgements. We are grateful for the funding and support provided for this trial by Australian Rotary Health and the National Health and Medical Research Council (#1173270). The trial would not have been possible without the generous support of our group facilitators, the staff at *headspace* and the University of Queensland clinic, and all the young people involved in the trial.

Sarah V. Bentley, Genevieve A. Dingle, and Jolanda Jetten co-authored GROUPS 4 HEALTH and were essential to its development. Victoria Gore-Jones and Francis Dark assisted in the early stages of the trial development and implementation.

Author Contribution. TC led the grant, oversaw the trial, did the statistical analyses and wrote the first draft of the report. CH provided conceptual input at all stages of the grant and project, and led the design of the intervention and the training of facilitators. JR and EW were the trial managers. AH provided conceptual input and led grants that supported program development. ZW provided expertise on protocol development and analyses. All authors provided critical feedback on the report.

Data Availability. The data that supports the findings of this study will be made available online upon publication on the [Open Science Framework](#).

Table 1. Baseline characteristics of the intention-to-treat population^a

	Groups 4 Health (N=84)	Cognitive Behaviour Therapy (N=90)
Gender	59 (70.2%) female 25 (29.8%) male	72 (80.0%) female 18 (20.0%) male
Age	<i>M (SD)</i> 18.86 (2.06) range 15-25 years	19.02 (1.87) range 15-25 years
Education	7 (8.3%) Year 10 or less 63 (75.0%) Year 12 9 (10.7%) certificate or diploma 5 (6.0%) Bachelor degree	2 (2.2%) Year 10 or less 69 (76.7%) Year 12 8 (8.9%) certificate or diploma 10 (11.1%) Bachelor degree 1 (1.1%) Graduate degree
Nationality	55 (65.5%) Australian 15 (17.9%) Chinese 14 (16.7%) Other	48 (53.3%) Australian 22 (24.4%) Chinese 20 (22.2%) Other
Ethnicity	44 (52.4%) White 33 (39.3%) Asian 3 (3.6%) Indigenous ^b 2 (2.4%) Black 1 (1.2%) Middle Eastern 1 (1.2%) Not disclosed	35 (38.9%) White 50 (55.6%) Asian 3 (3.3%) Indigenous ^b 1 (1.1%) Black 1 (1.1%) Middle Eastern
<i>Inclusion criteria</i>		
Formal Mental Health Diagnosis ^c	23 (31.1%) Yes 17 (77.3%) Mood disorder 16 (72.7%) Anxiety disorder 6 (27.3%) Other 51 (68.9%) No	25 (30.5%) Yes 19 (79.2%) Mood disorder 15 (60.0%) Anxiety disorder 3 (12.5%) Other 57 (69.5%) No
Depression score (PHQ-9)	14.01 (5.21) 98.8% ≥5 (mild clinical range) 83.3% ≥10 (moderate clinical range) 44% ≥15 (severe clinical range)	14.08 (5.14) 100% ≥5 (mild clinical range) 78.9% ≥10 (moderate clinical range) 42.2% ≥15 (severe clinical range)
Loneliness (UCLA-20)	51.95 (9.94) 97.6% ≥33 (population mean ¹⁸) 84.5% >42 (population mean + 1 SD) 50% >51 (population mean + 2 SD)	53.60 (9.90) 98.9% ≥33 (population mean ¹⁸) 87.8% >42 (population mean + 1 SD) 57.8% >51 (population mean + 2 SD)
<i>Primary outcomes</i>		
Depression score (DASS-21)	19.76 (8.54)	20.80 (9.52)
Loneliness (UCLA-4)	10.68 (2.07)	11.07 (2.21)

Note.

- T-tests (for continuous variables) and chi-square tests (for categorical variables) indicated no baseline differences between conditions were statistically significant ($ps > .135$).
- This category included people identifying as Aboriginal Australian, Torres Strait Islander, and/or Maori.
- People were prompted to skip this question if they did not wish to disclose, and more than 10% of the sample did so. Percentages provided here are a proportion of those who responded to this question, which likely underestimate the true proportion of the participants with a formal diagnosis.

Figure 1. Trial profile.

Note. All 174 randomised participants were invited to participate in the follow-up assessments, and so 17 of the 22 discontinued participants have outcome data available for inclusion in the intention-to-treat analysis.

Figure 2. Change in loneliness over time.

Figure 3. Change in depression symptom severity over time.

References

- 1 Ferrari AJ, Charlson FJ, Norman RE, *et al.* Burden of depressive disorders by country, sex, age, and year: Findings from the Global Burden of Disease Study 2010. *PLoS Med* 2013; **10**. DOI:10.1371/journal.pmed.1001547.
- 2 Weinberger AH, Gbedemah M, Martinez AM, Nash D, Galea S. Trends in depression prevalence in the USA from 2005 to 2015 : widening disparities in vulnerable groups. 2018.
- 3 Peters AT, A M, Jacobs RH, *et al.* Trajectories of functioning into emerging adulthood following treatment for adolescent depression. *J Adolesc Heal* 2016; **58**: 253–9.
- 4 Cipriani A, Zhou X, Del Giovane C, *et al.* Comparative efficacy and tolerability of antidepressants for major depressive disorder in children and adolescents: a network meta-analysis. *Lancet* 2016; **388**: 881–90.
- 5 Jaycox LH, Asarnow JR, Sherbourne CD, Rea MM, LaBorde AP, Wells KB. Adolescent primary care patients' preferences for depression treatment. *Adm Policy Ment Heal Ment Heal Serv Res* 2006; **33**: 198–207.
- 6 García-Escalera J, Chorot P, Valiente RM, Reales JM, Sandín B. Efficacy of transdiagnostic cognitive-behavioral therapy for anxiety and depression in adults, children and adolescents: A meta-analysis. *Rev Psicopatol y Psicol Clin* 2016; **21**: 147–75.
- 7 Weisz JR, McCarty CA, Valeri SM. Effects of psychotherapy for depression in children and adolescents: A meta-analysis. *Psychol Bull* 2006; **132**: 132–49.
- 8 Erzen E, Çikrikci Ö. The effect of loneliness on depression: A meta-analysis. *Int J Soc Psychiatry* 2018; **64**: 427–35.
- 9 Masi CM, Chen HY, Hawkey LC, Cacioppo JT. A meta-analysis of interventions to reduce loneliness. *Personal Soc Psychol Rev* 2011; **15**: 219–66.
- 10 Mance P. Is Australia experiencing an epidemic of loneliness? Evidence from 16 waves of the Housing, Income and Labour Dynamics in Australia Survey. *Working Paper. Canberra, 2018*.

- 11 Cattan M, White M, Bond J, Learmouth A. Preventing social isolation and loneliness among older people: a systematic review of health promotion interventions. *Ageing Soc* 2005; **25**: 41–67.
- 12 Haslam C, Cruwys T, Haslam SA, Dingle G, Chang MX-L. Groups 4 Health: Evidence that a social-identity intervention that builds and strengthens social group membership improves mental health. *J Affect Disord* 2016; **194**: 188–95.
- 13 Haslam C, Cruwys T, Chang MXL, *et al.* GROUPS 4 HEALTH reduces loneliness and social anxiety in adults with psychological distress: Findings from a randomized controlled trial. *J Consult Clin Psychol* 2019; **87**: 787–801.
- 14 Postmes T, Wichmann LJ, van Valkengoed AM, van der Hoef H. Social identification and depression: A meta-analysis. *Eur J Soc Psychol* 2019; **49**: 110–26.
- 15 Steffens NK, Larue CJ, Haslam C, *et al.* Social identification-building interventions to improve health : a systematic review and meta- analysis. *Health Psychol Rev* 2019; **0**: 1–28.
- 16 Cruwys T, Haslam C, Walter ZC, Rathbone J, Williams E. The connecting adolescents to reduce relapse (CARR) trial: Study protocol for a randomized controlled trial comparing the efficacy of Groups 4 Health and cognitive behaviour therapy in young people. *BMC Public Health* 2019; **19**: 1–10.
- 17 Spitzer RL, Kroenke K, Williams JBW, and the PHQPCS Group. Validation and utility of a self-report version of PRIME-MD. *JAMA* 1999; **282**: 1737–44.
- 18 Shevlin M, Murphy S, Murphy J. The Latent Structure of Loneliness: Testing competing factor models of the UCLA loneliness scale in a large adolescent sample. *Assessment* 2015; **22**: 208–15.
- 19 Stice E, Burton E, Bearman SK, Rohde P. Randomized trial of a brief depression prevention program. *Behav Res Ther* 2007; **45**: 863–76.
- 20 Stice E, Shaw H, Bohon C, Marti CN, Rohde P. A meta-analytic review of depression prevention programs for children and adolescents: Factors that predict magnitude of intervention effects. *J Consult Clin Psychol* 2009; **77**: 486–503.

- 21 Henry JD, Crawford JR. The short-form version of the Depression Anxiety Stress Scales (DASS-21): construct validity and normative data in a large non-clinical sample. *Br J Clin Psychol* 2005; **44**: 227–39.
- 22 Hays RD, DiMatteo MR. A Short-Form Measure of Loneliness. *J Pers Assess* 1987; **51**: 69–81.
- 23 Molenberghs G, Thijs H, Jansen I, *et al.* Analyzing incomplete longitudinal clinical trial data. *Biostatistics* 2004; **5**: 445–64.
- 24 Cruwys T, Haslam SA, Dingle GA, *et al.* Feeling connected again: Interventions that increase social identification reduce depression symptoms in community and clinical settings. *J Affect Disord* 2014; **159**: 139–46.
- 25 Jetten J, Reicher SD, Haslam SA, Cruwys T. Together apart: The psychology of COVID-19. Los Angeles: SAGE Publications, 2020
DOI:10.12968/prps.2015.sup172.2.