Aerobic exercise for cognition in mild cognitive impairment

A Bayesian hierarchical re-analysis of Zheng et al 2016 BJSM

John R. Best*

05 June 2020

Introduction

Hierarchical modeling is a promising approach to understanding how an intervention—*e.g.*, exercise training—impacts multiple, related outcomes—*e.g.*, multiple measures of cognitive performance. Such models can be applied to individual-level data within the context of a single exercise trial, but can also be applied to study-level data within the context of a meta-analysis of a set of relevant trials. Rather than treat each outcome (*e.g.*, delayed recall) as distinct and therefore submitted to separate meta-analytic models, all outcomes are included together in a single model. Specific outcomes are considered random, drawn from a population of possible outcomes. The result is the estimation of the overall population meta-analytic effect (*e.g.*, the effect of exercise on ‘cognition’) and estimation of outcome-specific random effects (*e.g.*, the effect of exercise on delayed recall). Critically, in estimation of outcome-specific random effects, there is partial pooling of information across outcomes. This is advantageous for many reasons, including minimizing over-fitting of data and handling imbalance across outcomes, as occurs when some outcomes are sampled more frequently than others. These less-sampled outcomes *borrow* information from the more-sampled outcomes during the estimation process.

Additionally, a Bayesian-based approach to data analysis and reporting offers advantages over the more commonly-used frequentist approach to meta-analysis. Rather than conclude whether the meta-analytic effect can be distinguished from zero according to some acceptable level of error, Bayesian analysis focuses on the range of parameters θ that would be compatible with the set of meta-analyzed studies in light of the prior belief about the range of these parameters. In turn, one address more interesting questions such as what is the probability that there is any sign of efficacy on cognition— $P(\theta > 0)$ —or even better, what is the probability that there is efficacy that exceeds some agreed-upon meaningful minimum, perhaps 1/4 of a standard deviation— $P(\theta > .25)$?

Combining the two to conduct a Bayesian hierarchical meta-analysis could be beneficial to researchers conducting meta-analysis. It could be especially valuable in addressing a common dilemma in which researchers use a frequentist approach, set an arbitrary α error rate of 0.05, and then observe a meta-analytic effect with a p value that slightly exceeds this threshold. This can lead to a contradictory stance by the meta-analyst such that they state that the effect was not statistically significant but also state that the evidence appears to support efficacy of the treatment. Further, this approach can lead to black and white statements such that treatment affects cognitive domain X but not domain Y, as if these are wholly separate outcomes.

*Simon Fraser University and the University of British Columbia, john[underscore]best@sfu.ca

Data

As an exercise, I re-analyzed the studies meta-analyzed by Zheng and colleagues published in the *British Journal of Sports Medicine* (Zheng et al. 2016). In this meta-analysis the authors focus on older persons (60 years and older) who meet a broadly-defined criteria of mild cognitive impairment but who were free of dementia or a diagnosed neurological or neurodegenerative condition (Alzheimer’s or Parkinson’s disease).

For the current analyses, I used the 11 studies that Zheng et al identified. Prior to analyses, I examined the original studies, and noted that the data from one study (Barnes et al. 2013) appeared to be incorrectly analyzed. Specifically, raw baseline values on the outcome measures (see Table 2 of this paper) appeared to be added to *standardized* change scores (see Table 3) to estimate post-test performance. For the current analyses, I used the standardized within-group change scores reported in Table 3 of Barnes et al. with an assumed within-group change variance of 1. Additional changes in comparison to Zheng et al. was maintaining separation of men and women from the study by van Uffelen and colleagues (Uffelen et al. 2008), and the inclusion of additional cognitive outcomes that were reported by the two studies from Lam and colleagues (Lam et al. 2012, 2015), from Nagamatsu and colleagues (Nagamatsu et al. 2013) and the study by Scherder et al (Scherder et al. 2005). Finally, for standardized effect sizes, I used Hedge’s g because it corrects for small-sample bias more so than Cohen’s d , which was used by Zheng and colleagues.

The first 20 rows of data are shown below in **Table 1**. Data are in long format to capture nesting of effect sizes under individual studies (“Author”) and cognitive domains. A complete spreadsheet of data used for the current analyses can be found **here**.

Table 1: Table 1. Slice of data used

Author	Domain	hedge_g	se
Barnes_2013	Attention and Speed	-0.18	0.25
Barnes_2013	Attention and Speed	0.04	0.25
Barnes_2013	Attention and Speed	-0.19	0.25
Barnes_2013	Attention and Speed	-0.04	0.25
Barnes_2013	Attention and Speed	-0.24	0.25
Barnes_2013	Attention and Speed	0.04	0.25
Barnes_2013	Executive Function	-0.05	0.25
Barnes_2013	Executive Function	-0.04	0.25
Barnes_2013	Executive Function	0.13	0.25
Barnes_2013	Executive Function	-0.13	0.25
Barnes_2013	Delayed Memory	0.01	0.25
Barnes_2013	Immediate Memory	-0.19	0.25
Hu_2014	Delayed Memory	0.41	0.14
Hu_2014	Immediate Memory	0.64	0.15
Lam_2012	Attention and Speed	0.20	0.13
Lam_2012	Executive Function	0.36	0.13
Lam_2012	Executive Function	0.26	0.13
Lam_2012	Global Cognition	0.42	0.13
Lam_2012	Global Cognition	0.36	0.13
Lam_2012	Delayed Memory	0.39	0.13

Analysis and Results

Bayesian hierarchical models were conducted in R version 4.0.0 using the package `brms` with the following specifications:

$$\begin{aligned}d_i &\sim \text{Normal}(\theta_i, \sigma_i = se_i) \\ \theta_i &\sim \text{Normal}(\mu, \tau) \\ \mu &\sim \text{Normal}(0, 1) \\ \tau &\sim \text{HalfCauchy}(0, 1)\end{aligned}$$

Thus, the model estimated the overall population-level meta-analytic effect of aerobic training versus control (μ) as well as domain- and study-specific deviations from population mean (τ). Somewhat informative priors were given for the population level effect and variance estimates. For example, the normal prior centered at 0 and variance of 1 would suggest that overall *Hedge's g* of 1 (a fairly large effect) would be plausible though less likely than effects closer to zero.

In R code:

```
zheng_md1 <- brm(data=zheng_data, family=gaussian,
  hedge_g | se(se) ~ 1 + (1|Domain) + (1|Author),
  prior = c(prior(normal(0,1), class = Intercept),
    prior(cauchy(0,1), class = sd)
  ),
  iter = 5000, warmup = 1000, chains = 4, cores = 1,
  control = list(adapt_delta = 0.99, max_treedepth = 15),
  seed = 12)
```

The model converged without issue, with $\hat{R}$ values of 1.00 for all parameter estimates. Examination of trace plots showed good mixing of the 4 chains.

The posterior distribution for the meta-analytic average effect on cognition had a mean of 0.23 with a 95% credible interval from 0 to 0.46. This is in-line with the meta-analytic effects reported for the memory domains by Zheng and colleagues.

Figure 1 provides a posterior predictive check, showing the observed distribution of effect sizes versus 50 simulations from the posterior predictive distribution. The simulated distribution of effect sizes matches well the observed; there is greater mass given to negative effect sizes (i.e., suggesting a harmful effect of aerobic exercise) that is due to the neutral normal prior centered as zero.

The domain-specific and overall distributions of posterior draws of the effect of aerobic exercise on cognition are shown in **Figure 2** below. Effects appear to vary modestly by domain with global cognition having the greatest proportion of mass in favour of a positive effect of aerobic training and attention and speed having the smaller proportion of mass in favour of a positive effect.

Various questions can be asked of the posterior distributions. For example, **Table 2** below summarizes the probability of any efficacy— $P(\theta > 0)$ —and of modest efficacy— $P(\theta > .25)$ —for the pooled effect and for the domain-specific effects. Assuming the prior distribution $\text{Normal}(0, 1)$ for the pooled effect and a $\text{Cauchy}(0, 1)$ for the variance parameters, and the validity of the data, there appears to be a bit less than even odds that aerobic exercise has *at least* a modest effect of cognition. The odds appear noticeably higher when global measures of cognition are used (e.g., MMSE, MoCA, ADAS-Cog) and less for measures of attention and speed. With respect to any sign of efficacy (i.e., the effect size on average is greater than zero), the odds are pretty good in favour of a positive effect.

Of course, these conclusions are only as good as the data analyzed. If there is systematic bias—publication bias, questionable research practices (including selective reporting), design bias (e.g., inability to blind participants to condition)—that inflates the effects in favour of aerobic exercise, then Bayesian-based conclusions

Figure 1: Distribution of observed compared to 50 simulated effect sizes

Figure 2: Meta-analytic effects of aerobic training across cognitive domains

will be undermined just as frequentist ones would be. Employment of skeptical priors would somewhat counteract that effect but it is still an important caution.

Table 2: Table 2. Probability of efficacy by domain

Domain	Any efficacy	Modest efficacy
Global.Cognition	1.00	0.86
Delayed.Memory	0.98	0.45
Immediate.Memory	0.98	0.44
Pooled Effect	0.98	0.41
Executive.Function	0.97	0.28
Working.Memory	0.95	0.27
Attention.and.Speed	0.90	0.17

Variation across studies is shown below along with the pooled effect in **Figure 3**. In comparison to cognitive domain, variation across studies appears larger. The large variation in the width of the distributions of posterior draws reflects the fact that some studies report very few effect sizes, *e.g.*, (Nascimento et al. 2015)

Figure 3: Meta-analytic effects of aerobic training across studies

A more direct comparison of variation across studies and across domains is shown below in **Figure 4**. There appears to be more study-related variation than domain-related variation. This could reflect any number of things, including differences in aerobic training protocol, nuanced differences in sample characteristics, to name two.

Figure 4: Comparison of variation across authors and across domains

Conclusions

What to conclude from this re-analysis of Zheng et al. (Zheng et al. 2016)? The conclusions are similar to those Zheng and colleagues in that the aerobic exercise appears to have a positive effect on cognition (broadly defined), according to the studies under study. The benefit of the current Bayesian hierarchical analysis is that more specific conclusions can be made in contrast to simple rejection of a null hypothesis that the effect is zero. There is some variation by cognitive domain—perhaps global cognitive measures and memory-based measures are especially sensitive—though this should be considered cautiously.

There is still uncertainty as to whether aerobic exercise has at least modest efficacy on cognition—here defined as $d > .25$. And the issue of no study being able to double-blind should be remembered. Though the fact that aerobic exercise has known primary benefits to cardiorespiratory health should be noted in determination in whether to recommend aerobic exercise to an individual with mild cognitive impairment.

References

- Barnes, D. E., W. Santos-Modesitt, G. Poelke, A. F. Kramer, C. Castro, L. E. Middleton, and K. Yaffe. 2013. "The Mental Activity and eXercise (Max) Trial: A Randomized Controlled Trial to Enhance Cognitive Function in Older Adults." Journal Article. *JAMA Internal Medicine*, 1–8. <https://doi.org/10.1001/jamainternmed.2013.189>.
- Lam, L. C., W. C. Chan, T. Leung, A. W. Fung, and E. M. Leung. 2015. "Would Older Adults with Mild Cognitive Impairment Adhere to and Benefit from a Structured Lifestyle Activity Intervention to Enhance Cognition?: A Cluster Randomized Controlled Trial." Journal Article. *PLoS One* 10 (3): e0118173. <https://doi.org/10.1371/journal.pone.0118173>.
- Lam, L. C., R. C. Chau, B. M. Wong, A. W. Fung, C. W. Tam, G. T. Leung, T. C. Kwok, T. Y. Leung, S. P. Ng, and W. M. Chan. 2012. "A 1-Year Randomized Controlled Trial Comparing Mind Body Exercise (Tai Chi) with Stretching and Toning Exercise on Cognitive Function in Older Chinese Adults at Risk of Cognitive Decline." Journal Article. *J Am Med Dir Assoc* 13 (6): 568 e15–20. <https://doi.org/10.1016/j.jamda.2012.03.008>.
- Nagamatsu, L. S., A. Chan, J. C. Davis, B. L. Beattie, P. Graf, M. W. Voss, D. Sharma, and T. Liu-Ambrose. 2013. "Physical Activity Improves Verbal and Spatial Memory in Older Adults with Probable Mild Cognitive Impairment: A 6-Month Randomized Controlled Trial." Journal Article. *J Aging Res* 2013: 861893. <https://doi.org/10.1155/2013/861893>.
- Nascimento, C. M., J. R. Pereira, L. Pires de Andrade, M. Garuffi, C. Ayan, D. S. Kerr, L. L. Talib, M. R. Cominetti, and F. Stella. 2015. "Physical Exercise Improves Peripheral Bdnf Levels and Cognitive Functions in Mild Cognitive Impairment Elderly with Different Bdnf Val66met Genotypes." Journal Article. *J Alzheimers Dis* 43 (1): 81–91. <https://doi.org/10.3233/JAD-140576>.
- Scherder, E. J., J. Van Paasschen, J. B. Deijen, S. Van Der Knokke, J. F. Orlebeke, I. Burgers, P. P. Devriese, D. F. Swaab, and J. A. Sergeant. 2005. "Physical Activity and Executive Functions in the Elderly with Mild Cognitive Impairment." Journal Article. *Aging Ment Health* 9 (3): 272–80. <https://doi.org/10.1080/13607860500089930>.
- Uffelen, J. G. van, M. J. Chinapaw, W. van Mechelen, and M. Hopman-Rock. 2008. "Walking or Vitamin B for Cognition in Older Adults with Mild Cognitive Impairment? A Randomised Controlled Trial." Journal Article. *Br J Sports Med* 42 (5): 344–51. <https://doi.org/10.1136/bjsm.2007.044735>.
- Zheng, G., R. Xia, W. Zhou, J. Tao, and L. Chen. 2016. "Aerobic Exercise Ameliorates Cognitive Function in Older Adults with Mild Cognitive Impairment: A Systematic Review and Meta-Analysis of Randomised Controlled Trials." Journal Article. *British Journal of Sports Medicine*. <https://doi.org/10.1136/bjsports2015-095699>.