Interoception modulates the self-prioritization effect

Tatsuru Honda^{1*}, Takashi Nakao^{2*}

¹Graduate School of Education, Hiroshima University, Hiroshima, Japan.

²Graduate School of Humanities and Social Sciences, Hiroshima University, Hiroshima, Japan.

*** Correspondence:**

Tatsuru Honda

tatsuruhnd@gmail.com

Keywords: self-prioritization effect, interoception, interoceptive accuracy, cardiac cycle, self-relevance

Abstract

Self-relevant information is processed faster and more accurately than non-self-relevant information. Such a bias is developed even for newly associated information with the self, also known as the self-prioritization effect (SPE). Interoception, which refers to the overall processing of information from inside the body, is crucial for self-relevant processing; however, little is known about its role in SPE. In this study, we investigated the relationship between the magnitude of SPE and interoceptive accuracy (IAc), defined as an individual's ability to accurately perceive one's own interoceptive state. Additionally, to explore the causal relationship, we measured SPE by presenting self- or non-self-relevant stimuli based on the participant's cardiac cycle in the shape-label matching task. We demonstrated that IAc was negatively correlated with the magnitude of SPE in terms of discrimination of the relevance of the stimuli. In addition, a correlation was observed only when the stimuli were presented during cardiac systole. Furthermore, IAc was negatively correlated with the processing of self-relevant stimuli but not with non-self-relevant stimuli. Our findings suggest that individuals with high IAc were less able to discriminate whether an external neutral stimulus was self-relevant when the stimulus was presented at systole. Our results may reflect the tendency to recognize the self-relevance of stimuli based on interoception in individuals with high IAc. Since the present study used geometric shapes, which are not easily recognized as stimuli that can induce changes in the interoception, individuals with high IAc assigned less self-relevance to the stimuli, resulting in weaker SPE. From this perspective, we further discussed the conditions that lead to stronger SPE in individuals with high IAc, in contrast to the present study results.

1 Introduction

The self has attracted the interest of researchers in psychology for a long time. In the field of experimental psychology, researchers have attempted to understand the self from a functional perspective by quantifying the self-bias effect (Sui and Humphreys, 2015; Sui and Gu, 2017). For example, it is well known that people can be aware of their own name being called, even in situations where they are not paying attention—known as the cocktail party effect (Moray, 1959). Information related to the self, such as one's own name, attracts more attention than non-self-relevant stimuli. Biases have been demonstrated in a variety of external stimuli and cognitive processes (Cunningham and Turk, 2017; Sui and Humphreys, 2015).

38 Recently, a novel paradigm has demonstrated that biases emerge even for neutral stimuli newly
39 associated with one's self. In the shape-label matching paradigm (Sui et al., 2012), participants were
40 first asked to learn the associations between geometric shapes and personal labels (e.g., circles
41 represent self, triangles represent best friends, squares represent strangers). Following the learning
42 phase, the participants judged whether the presented shape-label pairings matched the previously
43 learned associations. The results showed that their responses to self-relevant stimuli (i.e., circle –
44 self) were faster and more accurate than non-self-relevant stimuli. This is known as the self-
45 prioritization effect (SPE). This paradigm demonstrated that individuals rapidly develop biases, even
46 for newly associated stimuli with the self. In addition, although self-relevance and intimacy cannot be
47 separated when using familiar stimuli such as one's own name or face, which have been widely used
48 in conventional studies, the shape-label matching paradigm allowed us to examine self-bias without
49 the influence of familiarity (Kim and Florack, 2021). SPE has been observed in the shape-label
50 matching paradigm for various types of stimuli (Payne et al., 2017; Sel et al., 2019; Sui et al., 2012)
51 and other sensory modalities (Frings and Wentura, 2014; Schäfer et al., 2016).

52 Although SPE is observed when an external stimulus is associated with the current self, no such
53 preferential processing is observed when associated with the past or future self (Golubickis et al.,
54 2017; Kim and Florack, 2021). Golubickis et al. (2017) examined whether SPE could emerge when
55 associating a geometric shape with a temporally distant self (i.e., current self vs future self) in the
56 shape-label matching paradigm. The results showed that SPE did not emerge in the stimuli associated
57 with the temporally distant self, but only with the current self. These results can be explained by the
58 fact that we sometimes treat our temporally distant selves as strangers (Pronin et al., 2008; Pronin
59 and Ross, 2006). Given that such “self-becomes-other” effects are caused by the lack of information
60 on internal states (Pronin et al., 2008; Pronin and Ross, 2006), that information may play a crucial
61 role in the emergence of SPE.

62 Interoception is one of the internal states of the self. It refers to the overall process of how the
63 nervous system senses, interprets, and integrates information arising from within the body, providing
64 a moment-by-moment representation of internal bodily states with or without awareness (Berntson
65 and Khalsa, 2021). Interoception supports homeostatic control, allows allostatic adaptation, and
66 drives behavior through feelings such as hunger and thirst (Tsakiris and Critchley, 2016). Several
67 theories have proposed that interoception is the foundation of the self (Craig, 2009; Seth, 2013), and
68 previous empirical findings support these proposals. For instance, heartbeat evoked responses as an
69 index of the temporal dynamics of brain activity in relation to cardiac activity (Baransukas et al.,
70 2021) covaried with the activity of the default mode network node during self-relevant thinking
71 (Babo-rebelo et al., 2016). Qin et al. (2020) conducted a meta-analysis of neuroimaging studies and
72 concluded that interoceptive information is necessary for processing self-relevant information.
73 Ambrosini et al. (2019) reported that interoceptive information speeds up recognizing one's own face.
74 Therefore, the findings of interoception and SPE suggest that SPE may be observed when the
75 external stimuli are associated with the current self, driven by interoception.

76 However, we could also assume an opposite relationship between SPE and interoception. Studies of
77 body ownership have reported that interoceptive accuracy (IAC), an individual's ability to accurately
78 perceive their own interoceptive state (Garfinkel et al., 2015), modulates the degree of self-
79 association with external bodily stimuli (Schauder et al., 2015; Tsakiris et al., 2011). It was
80 demonstrated that individuals with high IAC tend to experience a less rubber hand illusion, that is,
81 they have difficulty recognizing external objects (i.e., rubber hand) as their own by synchronized
82 visuo-tactile stimulation. Similarly, individuals with high IAC show less susceptibility to enfacement
83 illusion, that is, recognizing others' faces as their own by seeing another person's face being stroked

in synchrony with their own face (Tajadura-Jiménez and Tsakiris, 2014). These findings indicate that individuals with high IAc are less likely to recognize external stimuli as being associated with the self. Although the typical shape-label matching paradigm investigates the association of the self with geometric shapes rather than the body, the findings of body-ownership studies allow us to predict that individuals with high IAc show weaker SPE.

Thus, there are two possibilities for the relationship between SPE and interoception: interoception either promotes or inhibits SPE. However, to the best of our knowledge, the relationship between SPE and interoception has not yet been investigated.

The present study aimed to elucidate the relationship between SPE and interoception. As mentioned above, a positive association between SPE and interoception is expected from studies of the shape-label matching paradigm and interoception. In the shape-label matching paradigm, it has been reported that stimuli associated with a temporally distant self can dampen the emergence of SPE and do not facilitate information processing in the current self (Golubickis et al., 2017; Kim and Florack, 2021). This may be because a self that is temporally distant from its current self lacks access to and availability of information about the internal state, that is, interoceptive information (Pronin and Ross, 2006). Individuals with high IAc, who can perceive their interoceptive information accurately, are expected to show a stronger SPE.

On the other hand, studies on the relationship between IAc and body ownership predict a negative association between SPE and IAc. Individuals with high IAc have difficulty recognizing external bodily stimuli with the self through synchronized visuo-tactile stimulation (Schauder et al., 2015; Tsakiris et al., 2011; Tajadura-Jiménez and Tsakiris, 2014). Therefore, assuming that the same relationship applies to the association between the self and external geometric shapes in the shape-label matching paradigm, it is expected that individuals with high IAc will show a weaker SPE in the shape-label matching paradigm.

Additionally, to get closer to an understanding of the causal relationship between IAc and SPE, we investigated whether the perception of interoceptive information affects SPE. Cardiac interoceptive information (i.e., strength and timing of individual heartbeats) is transmitted to the brain by the firing of arterial baroreceptors during systole in the cardiac cycle (Critchley and Garfinkel, 2018). Therefore, in studies examining whether interoceptive information affects cognitive processes, the stimuli that trigger the cognitive process were presented in coincidence with either the systolic or diastolic phases of the cardiac cycle (e.g., Garfinkel et al., 2014; Gray et al., 2012). These studies have reported that presenting stimuli based on the timing of the cardiac cycle modulates cognitive processing, including self-face recognition (Ambrosini et al., 2019). Moreover, several previous studies have reported that individuals with high IAc are more affected by stimuli presented at systole (Sel et al., 2016; von Mohr et al., 2021). Based on these findings, if the perception of interoceptive information affects SPE, the following results are expected: individuals with high IAc can perceive interoceptive information sent to the brain during systole, thus affecting SPE. In contrast, individuals with low IAc cannot perceive interoceptive information even at systole, and SPE is not affected. As a result, a correlation between IAc and SPE is expected when the stimuli are presented coincident with systole. However, since there is no interoceptive information to perceive during diastole, whether IAc is high or low, it is expected that there will be no correlation between IAc and SPE at diastole. Thus, if the perception of interoceptive information affects SPE, the correlation between IAc and SPE would be observed only when the stimuli are presented at systole and not at diastole.

In summary, we hypothesized that (1) IAC is positively or negatively correlated with SPE, and (2) the correlation will be observed only under conditions where the stimuli are presented at systole in the cardiac cycle.

2 Materials and Methods

2.1 Participants

Thirty-eight Japanese university students participated in the study. Two participants were excluded due to hardware failure or low-quality of recorded ECG data. Thus, a total of 36 participants (18 women, aged 20.6 ± 1.9 years) were included in the final analyses. The participants had normal or corrected-to-normal visual acuity.

The protocol was reviewed and approved by the Ethical Committee of the Graduate School of Education, Hiroshima University. All participants signed an informed consent form and were instructed that the experiment could be discontinued at any time. The participants received 2,000 yen as compensation after finishing the experiment.

2.2 Materials and Apparatus

The experiment was programmed, and all data were recorded using MATLAB 2019b software (Mathworks). Stimuli were presented using a 24-inch monitor (1920×1080 at 60 Hz) placed at a distance of approximately 60 cm from the participant. ECG was continuously recorded to measure the heart activity of the participants. Three Ag/AgCl electrodes were placed in lead II configuration: two electrodes were positioned underneath the left and right collarbone and the other on the lower left rib of each participant. The ECG signal was recorded at 400 Hz with a Biopac MP160 and a BN-RSPEC amplifier (Biopac System, USA) using MATLAB 2019b software.

2.3 Procedure

All participants were tested individually in a shielded room. The participants were instructed about the experimental procedure and signed a consent form. After placement of the ECG electrodes, the participants were instructed to sit in a comfortable chair. Subsequently, the participants engaged in two tasks: one was a shape-label matching task with presentation timing manipulated based on the cardiac cycles, and the other was a heartbeat counting task (Schandry, 1981).

2.3.1 Perceptual matching task

We conducted a shape-label matching task to quantify individual SPEs. This task consisted of two phases: a learning phase and a matching phase. In the learning phase, participants were expected to learn three associations between geometrical shapes and social labels (e.g., circle-self, triangle-best friend, and square-stranger) in 60 seconds. The shapes were described linguistically and were not presented as images during this phase.

After the learning phase, a matching phase was conducted (Figure 1A). Participants were required to judge whether the presented shape-label pairs were matched based on the associations learned during the learning phase. Additionally, to examine the relationship between SPE and interoception, the stimuli were presented either during systole or diastole. Corresponding to the previous studies (Garfinkel et al., 2014; Gray et al., 2012), in this experiment, stimuli were presented at the T-wave peaks in the Systole, and at 250 ms post-T-wave peaks in the Diastole conditions respectively (Figure 1B). The T-wave peaks were detected in real time when the amplitude of the ECG signal exceeded a

predefined threshold. The threshold was determined by the experimenter based on the resting state of the heart for 15 s before beginning the experimental task.

In the matching phase, each trial began with the presentation of the fixation cross, the duration of which was determined by the detection of the T-wave peaks. The mean time of the fixation cross was 1323.66 ms ($SD=106.65$ ms). Subsequently, a shape and a label were presented above and below the fixation cross for 100 ms, respectively, followed by a blank screen. The presentation of the shape-label pair was time-locked to either the T-wave peaks (Systole condition) or delayed 250 ms from the T-wave peaks (Diastole condition). Within the blank screen, participants were required to judge whether the presented shape-label pairing was matched by pressing one of the two buttons (J key or F key on the keyboard) as quickly and accurately as possible, based on the previously learned associations. The corresponding of the J key or the F key to the matching or non-matching response was counterbalanced across the participants. Feedback was given (“CORRECT,” “INCORRECT,” and “LATE”) for 500 ms according to the participant's response at the end of each trial. If the participant did not respond within 1000 ms, it was defined as “LATE.” Participants conducted 30 trials for each of the 12 conditions (self-matching at systole, self-nonmatching at systole, friend-matching at systole, friend-nonmatching at systole, stranger-matching at systole, stranger-nonmatching at systole, and similar combinations at diastole). There were five blocks of 72 trials each. In each block, six trials for each of the 12 conditions were conducted in a pseudo-randomized order. All stimuli were colored black and presented on a gray background.

2.3.2 Heartbeat counting task

Individual differences in IAc were quantified using a heartbeat counting task. Participants were asked to silently count only their perceived heartbeats without taking their pulse (Desmedt et al., 2018). Each trial began with a warning tone at 800 Hz, and approximately 3 s later, a 1000 Hz tone was given indicating the start of the counting interval, and participants began counting their heartbeats. Finally, a 1000 Hz tone was sounded to indicate the end of the counting interval, and the participants reported their perceived number of heartbeats. There were six trials with variable time windows of 25, 30, 35, 40, 45, and 50 s for the counting intervals. The trials were presented in a randomized order. The participants received no information about the length of each interval.

2.4 Data Analysis

2.4.1 Shape-label matching task

We assessed reaction time (RT) and sensitivity (d-prime) as indices of SPE. Only correct responses in matching trials were included in the RT analysis. The signal detection approach was used to calculate d-prime. In the case where hit rates and false alarm rates are 1.0 or 0, the corresponding z-scores are infinite and d-prime cannot be calculated. Therefore, we corrected all hit rates and false alarm rates by adding 0.5 to the number of hits and false alarms and dividing by $N + 1$, where N is the number of “match” or “mismatch” trials (Snodgrass and Corwin, 1988). Following this, we also assessed the magnitude of SPE (i.e., the magnitude to which self-relevant stimuli are processed faster or more accurately than other-relevant stimuli) for each individual for both RT and d-prime. The magnitude of SPE on RT was calculated by the difference between the Stranger and Self conditions, divided by the sum of the same two conditions [that is, $(\text{Stranger} - \text{Self}) / (\text{Self} + \text{Stranger})$]. The magnitude of SPE on d-prime was calculated by the difference between the Self and Stranger conditions, divided by the sum of the same two conditions [that is, $(\text{Self} - \text{Stranger}) / (\text{Self} + \text{Stranger})$]. Higher values for both indicated that self-relevant stimuli were processed more preferentially than other-relevant stimuli. The reason for including a Friend condition in the shape-

label matching task, although it was not used to assess the magnitude of SPE, was to make it sufficiently challenging to avoid ceiling effects (Verplanken and Sui, 2019).

2.4.2 Heartbeat counting task

IAC was assessed according to the following formula: $1/6\Sigma[1-(\text{ECG recorded heartbeats} - \text{participant counted heartbeats}) / \text{ECG recorded heartbeats}]$ (Schandry, 1981). IAC scores range from 0 to 1, with higher scores indicating better IAC.

3 Results

3.1 Validation of stimuli onset times

First, in the shape-label matching task, we examined whether the stimuli were accurately presented during systole or diastole. This is because the timing of stimuli presentation was based on the participants' T-wave peaks, but the detection criteria of T-wave peaks were set by the experimenter for each participant to detect T-wave peaks in real time, which may have differed from the actual T-wave peaks. Therefore, T-wave peaks (actual T-wave peaks) were determined a posteriori from the recorded ECG waveforms of each participant during the shape-label matching task, and the temporal difference from the T-wave peaks set during the task (established T-wave peaks) was calculated (actual - established). The results showed that the temporal difference was 18.32 ms ($SD=7.28$ ms), indicating that the T-wave peaks were detected almost accurately during the experimental task. The stimuli in the Systole and Diastole conditions were presented at T-wave peaks - 11.92 ms ($SD=7.55$ ms) and T-wave peaks + 243.72 ms ($SD=7.11$ ms), respectively, which were close to the target timing. In addition, we confirmed that all stimuli were within the R-R interval at which the T-wave peaks were detected as a criterion. These results indicated that the timing of the stimuli presentation was appropriate in the shape-label matching task.

3.2 SPE in the shape-label matching task

Table 1 summarizes the mean RT and d-prime of each condition. To confirm the emergence of SPE in the shape-label matching task, we conducted a 3 (Label) $\times$ 2 (Cardiac cycle) repeated-measures ANOVA for both RTs and d-prime. For RT, the result showed a significant main effect of the Label ($F(2, 70)=18.01, p<0.01, \eta_p^2=0.34$), shorter RT in the Self condition than either the Friend or Stranger conditions (both $p<0.01$), and shorter RT in the Friend condition than in the Stranger condition ($p<0.01$). We did not find any significant difference in the Cardiac cycle factor ($F(1, 35)=0.81, p=0.38, \eta_p^2=0.02$) and interaction effects ($F(2, 70)=1.18, p=0.31, \eta_p^2=0.03$). For d-prime, the result showed a significant main effect of the Label ($F(2, 70)=4.59, p<0.05, \eta_p^2=0.12$), with larger d-prime in the Self condition than the Stranger condition ($p<0.05$), and with no difference between the Self and Friend or Friend and Stranger conditions. No significant difference was found in the Cardiac cycle factor ($F(1, 35)=3.25, p=0.08, \eta_p^2=0.09$) and interaction effects ($F(2, 70)=0.52, p=0.60, \eta_p^2=0.02$). These results indicate that SPE emerged in the shape-label matching task.

3.3 Correlation Analyses

Correlation analyses were conducted to explore the relationship between the magnitude of SPE on both RT and d-prime and IAC, respectively (Hypothesis 1). The results showed that there was a significant negative correlation between the magnitude of SPE on d-prime and IAC ($\rho=-0.36, p<0.05, 95\%CI=-0.61, 0.03$; Figure 2A), but no significant correlation between the magnitude of SPE on RT and IAC ($\rho=-0.04, p=0.82, 95\%CI=-0.36, 0.29$).

Similarly, we examined the relationship separately in the Systole and Diastole conditions (Hypothesis 2). For the Systole condition, the magnitude of SPE on d-prime significantly negatively correlated with IAc ($\rho=-0.34$, $p<0.05$, 95%CI=-0.60, 0.01; Figure 2B), but not with RT ($\rho=-0.01$, $p=0.94$, 95%CI=-0.34, 0.32). In the Diastole condition, there was no significant correlation between the magnitude of SPE and IAc (RT: $\rho=-0.08$, $p=0.63$, 95%CI=-0.40, 0.25; d-prime: $\rho=-0.27$, $p=0.11$, 95%CI=-.55, .06; Figure 2C).

Although the results showed that the magnitude of SPE on d-prime in the Systole condition correlates with IAc, it is unclear how IAc is related to the sensitivity of processing self- and other-relevant information. To better understand this, we explored the relationship between d-prime and IAc in the Self and Stranger conditions, respectively. The results showed that there was a significant negative correlation between d-prime in the Self-Systole condition and IAc ($\rho=-0.38$, $p<0.05$, 95%CI=-0.63, -0.06; Figure 2D), but there was no significant correlation between d-prime in the Stranger-Systole condition and IAc ($\rho=0.15$, $p=0.38$, 95%CI=-0.19, 0.46; Figure 2E).

3.4 Effects of the cardiac cycle without considering IAc

Although this was not our main interest, we conducted paired *t*-tests on the magnitude of SPE on RT and d-prime, respectively, to examine the effects of the cardiac cycle. The results showed that there were no significant differences between the Systole and Diastole conditions for both RT ($t(35)=0.74$, $p=0.47$, $d=0.12$, 95% CI=-0.21, 0.45) and d-prime ($t(35)=.91$, $p=0.37$, $d=0.15$, 95% CI=-0.18, 0.48). There was no evidence that the cardiac cycle independently affected the magnitude of SPE.

4 Discussion

The present study investigated the relationship between the magnitude of SPE measured using the shape-label matching task (Sui et al., 2012) with the presented stimuli based on the participant's cardiac cycle and IAc measured using the heartbeat counting task (Schandry, 1981). We proposed two possibilities for interoception, namely, that it promotes or inhibits SPE, and hypothesized that (1) IAc is positively or negatively correlated with SPE, and (2) the correlation is observed only under conditions where the stimuli are presented at systole in the cardiac cycle where the interoceptive information is transmitted to the brain. The results showed a significant negative correlation between the magnitude of SPE on d-prime and IAc. Moreover, a significant correlation was observed only in the Systole condition but not in the Diastole condition. Additionally, IAc was significantly correlated with d-prime in the Self-Systole condition but not in the Stranger-Systole condition. Our results are the first to indicate a relationship between SPE and interoception. More specifically, individuals with high IAc were found to be less able to discriminate whether an external stimulus (i.e., a geometric shape) was self-relevant when the stimulus was presented coincident with the timing of interoceptive information being transmitted to the brain.

Why were individuals with high IAc less able to discriminate self-relevance when presented at systole? This may reflect the tendency to recognize the self-relevance of stimuli based on interoception in individuals with high IAc. More specifically, they may tend to recognize the stimuli that can induce changes in interoception as self-relevant stimuli, whereas they may recognize stimuli that are difficult to attribute as stimuli that can induce changes in interoception as non-self-relevant stimuli. Since the present study used geometric shapes that are not easily recognized as stimuli, which can induce changes in interoception, individuals with high IAc assigned less self-relevance to the stimuli, resulting in weaker SPE.

295 This idea is an application of the findings on body ownership and interoception (e.g., Suzuki et al.,
 296 2013; Tsakiris et al., 2011). Studies of body ownership have reported that individuals with high IAc
 297 are less likely to experience a sense of body ownership by synchronized visuo-tactile stimulation;
 298 that is, they have difficulty recognizing external fake bodily stimuli as relevant to their own
 299 (Schauder et al, 2015; Tsakiris et al., 2011; Tajadura-Jiménez and Tsakiris, 2014). This result has
 300 been interpreted to indicate that individuals with high IAc preferentially process interoceptive
 301 information from the actual body rather than exteroceptive information (i.e., visual and tactile
 302 information), thus making them less likely to experience a sense of body ownership (Tsakiris et al.,
 303 2011). In contrast, in situations where the fake hand is more likely to be perceived as inducing
 304 changes in one's own interoception, individuals with high IAc are more likely to experience a sense
 305 of body ownership. Suzuki et al. (2013) demonstrated that individuals with high IAc experienced a
 306 stronger sense of body ownership by presenting synchronized visuo-tactile stimuli to a hand that has
 307 a close visual similarity to the participant's real hand projected in augmented reality. They also
 308 reported that there was a trend for individuals with high IAc to experience a stronger sense of body
 309 ownership for the virtual hand, which flashed in synchronization with the participants' heartbeats
 310 without presenting any tactile stimuli. Therefore, body ownership studies suggest that interoceptive
 311 information serves as an anchor of body perception for individuals with high IAc, and that they are
 312 more likely to recognize self-relevance for external stimuli that could be related to their own
 313 interoception.

314 As described before, the current finding that high IAc individuals were less able to discriminate self-
 315 relevance when it was presented at systole may indicate a recognition style of self-relevance in
 316 individuals with high IAc. The present study used geometric shapes, which are neutral stimuli and
 317 seem less likely to be recognized as inducing changes in interoception even if they are presented at
 318 systole in which interoceptive information is transmitted to the brain. Therefore, it is possible that
 319 individuals with high IAc, who recognize self-relevance based on interoception, may have had
 320 difficulty recognizing geometric shapes as self-relevant, resulting in a smaller magnitude of SPE.

321 Conversely, individuals with high IAc may show stronger SPE (i.e., be able to discriminate self-
 322 relatedness when it is presented at systole) if stimuli are more likely to be recognized as inducing
 323 changes in their own interoception, such as emotional stimuli (Garfinkel et al., 2014). In the present
 324 study, we also expected that IAc would be positively correlated with the magnitude of SPE. Although
 325 this expectation was not supported, it is possible that a positive correlation could be found if
 326 emotional stimuli were used. However, since this prediction was derived based on existing body
 327 ownership studies described above, further study of SPE is needed.

328 Despite our insights, we should consider the following limitations. The first is the validity of the
 329 heartbeat counting task (Schandry, 1981) used to measure IAc. Although the task is commonly used
 330 in interoceptive studies because it is non-invasive and easy to perform (e.g., Ambrosini et al., 2019;
 331 Tsakiris et al., 2011), it has been argued that the performance of the task is affected by non-
 332 interoceptive factors (Ainley et al., 2020; Corneille et al., 2020; Ring et al., 2015; Zamariola et al.,
 333 2018; Zimprich et al., 2020). To exclude the effect of knowledge about heartbeats on performance
 334 (Ring et al., 2015), we referred to Desmedt et al. (2018) and instructed the participants to count only
 335 their perceived heartbeats, but response bias may still have affected it (Corneille et al., 2020).

336 The second limitation is the causal relationship between IAc and SPE. To get closer to the
 337 elucidation of the causal relationship, we presented stimuli based on the participant's cardiac cycle in
 338 the shape-label matching task. However, the present study assumed but did not confirm that
 339 individuals with high IAc would perceive interoception at systole in the cardiac cycle because

interoceptive information is transmitted to the brain at the time (Garfinkel et al., 2014; Gray et al., 2012). In addition, the relationship between IAc and SPE shown in this study is related to individual differences; thus, it is unclear whether changing within-individual IAc affects the processing of self-relevant information. Given that IAc could be improved by non-invasive vagus nerve stimulation (Villani et al., 2019) and physical activity (Quadt et al., 2021), further research would be able to investigate a more direct causality by using these methods.

Finally, we did not observe any relationship between SPE, RT, and IAc. Both d-prime and RT have been used as indices of SPE in several previous studies, and the high robustness of the emergence of SPE has been confirmed (e.g., Sui et al., 2012). Although the present study also observed SPE both in RT and d-prime, the association with IAc was observed only in d-prime and not in RT. The reason for this one-sided result remains unclear. Nevertheless, our results suggest that IAc is significant for exploring the differences in background processes reflected in RT and d-prime in SPE.

5 Conclusion

In the present study, we demonstrated that (1) IAc is negatively correlated with SPE, and (2) the correlation was observed only when the self- and other-relevant stimuli were presented at the cardiac systole phase. In addition, IAc was negatively correlated with the processing of self-relevant but not other-relevant stimuli. Thus, our results showed that individuals with high IAc were less able to discriminate whether an external neutral stimulus (i.e., a geometric shape) was self-relevant when the stimulus was presented simultaneously as interoceptive information was transmitted to the brain. The results provide implications for how individuals with high IAc recognize the self-relevance of external stimuli: individuals with high IAc tend to recognize the self-relevance of stimuli based on interoception. More specifically, they may tend to recognize the stimuli that are difficult to attribute as stimuli that can induce changes in interoception as non-self-relevant stimuli, whereas they may recognize stimuli that can induce changes in interoception as self-relevant stimuli. Our findings have important implications for understanding how external stimuli are recognized as self-relevant.

6 Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

7 Author Contributions

TH and TN contributed to the design of the study; TH conducted the study, analyzed the data, and wrote the first draft; both authors contributed to the revision of the manuscript.

8 Funding

This research was supported by the Center of Innovation Program of the Japan Science and Technology Agency. (JST) JPMJCE1311.

9 Data Availability Statement

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

10 Acknowledgments

I would like to thank Editage (<https://www.editage.com/>) for proofreading my English.

11 References

- Ainley, V., Tsakiris, M., Pollatos, O., Schulz, A., and Herbert, B. M. (2020). Comment on “Zamariola et al. (2018), Interoceptive accuracy scores are problematic: Evidence from simple bivariate correlations”—The empirical data base, the conceptual reasoning and the analysis behind this statement are misconceived and do not support the authors’ conclusions. *Biol. Psychol.*, 152, 107870. <https://doi.org/10.1016/j.biopsycho.2020.107870>
- Ambrosini, E., Finotti, G., Azevedo, R. T., Tsakiris, M., and Ferri, F. (2019). Seeing myself through my heart: Cortical processing of a single heartbeat speeds up self-face recognition. *Biol. Psychol.*, 144, 64–73. <https://doi.org/10.1016/j.biopsycho.2019.03.006>
- Babo-Rebelo, M., Richter, C. G., and Tallon-Baudry, C. (2016). Neural responses to heartbeats in the default network encode the self in spontaneous thoughts. *J. Neurosci.*, 36(30), 7829–7840. <https://doi.org/10.1523/JNEUROSCI.0262-16.2016>
- Baranauskas, M., Grabauskaitė, A., Griškova-Bulanova, I., Lataitytė-Šimkevičienė, B., and Stanikūnas, R. (2021). Heartbeat evoked potentials (HEP) capture brain activity affecting subsequent heartbeat. *Biomed. Signal Process Control*, 68, 102731. <https://doi.org/10.1016/j.bspc.2021.102731>
- Berntson, G. G., and Khalsa, S. S. (2021). Neural circuits of interoception. *Trends Neurosci.*, 44, 17–28. <https://doi.org/10.1016/j.tins.2020.09.011>
- Corneille, O., Desmedt, O., Zamariola, G., Luminet, O., and Maurage, P. (2020). A heartfelt response to Zimprich et al. (2020), and Ainley et al. (2020)’s commentaries: Acknowledging issues with the HCT would benefit interoception research. *Biol. Psychol.*, 152, 107869. <https://doi.org/10.1016/j.biopsycho.2020.107869>
- Craig, A. D. (2009). How do you feel — now? The anterior insula and human awareness. *Nat. Rev. Neurosci.*, 10, 59–70. <https://doi.org/10.1038/nrn2555>
- Critchley, H. D., and Garfinkel, S. N. (2018). The influence of physiological signals on cognition. *Curr. Opin. Behav. Sci.*, 19, 13–18. <https://doi.org/10.1016/j.cobeha.2017.08.014>
- Cunningham, S. J., and Turk, D. J. (2017). Editorial: A review of self-processing biases in cognition. *Q. J. Exp. Psychol. (Hove)*, 70(6), 987–995. <https://doi.org/10.1080/17470218.2016.1276609>
- Desmedt, O., Luminet, O., and Corneille, O. (2018). The heartbeat counting task largely involves non-interoceptive processes: Evidence from both the original and an adapted counting task. *Biol. Psychol.*, 138, 185–188. <https://doi.org/10.1016/j.biopsycho.2018.09.004>
- Frings, C., and Wentura, D. (2014). Self-prioritization processes in action and perception. *J. Exp. Psychol. Hum. Percept. Perform.*, 40. <https://doi.org/10.1037/a0037376>
- Garfinkel, S. N., Minati, L., Gray, M. A., Seth, A. K., Dolan, R. J., and Critchley, H. D. (2014). Fear from the heart: Sensitivity to fear stimuli depends on individual heartbeats. *J. Neurosci.*, 34(19), 6573–6582. <https://doi.org/10.1523/JNEUROSCI.3507-13.2014>
- Garfinkel, S. N., Seth, A. K., Barrett, A. B., Suzuki, K., & Critchley, H. D. (2015). Knowing your own heart: Distinguishing interoceptive accuracy from interoceptive awareness. *Biol. Psychol.*, 104, 65–74. <https://doi.org/10.1016/j.biopsycho.2014.11.004>
- Golubickis, M., Falben, J. K., Sahraie, A., Visokomogilski, A., Cunningham, W. A., Sui, J., and Macrae, C. N. (2017). Self-prioritization and perceptual matching: The effects of temporal construal. *Mem. Cogn.*, 45(7), 1223–1239. <https://doi.org/10.3758/s13421-017-0722-3>
- Gray, M. A., Beacher, F. D., Minati, L., Nagai, Y., Kemp, A. H., Harrison, N. A., and Critchley, H. D. (2012). Emotional appraisal is influenced by cardiac afferent information. *Emotion*,

- 12(1), 180–191. <https://doi.org/10.1037/a0025083>
- Kim, H., and Florack, A. (2021). Immediate self-information is prioritized over expanded self-information across temporal, social, spatial, and probability domains. *Q. J. Exp. Psychol. (Hove)*, 74(9), 1615–1630. <https://doi.org/10.1177/17470218211004208>
- Moray, N. (1959). Attention in dichotic listening: Affective cues and the influence of instructions. *Q. J. Exp. Psychol. (Hove)*, 11(1), 56–60. <https://doi.org/10.1080/17470215908416289>
- Payne, S., Tsakiris, M., and Maister, L. (2017). Can the self become another? Investigating the effects of self-association with a new facial identity. *Q. J. Exp. Psychol. (Hove)*, 70(6), 1085–1097. <https://doi.org/10.1080/17470218.2015.1137329>
- Pronin, E., Olivola, C. Y., and Kennedy, K. A. (2008). Doing Unto Future Selves As You Would Do Unto Others: Psychological Distance and Decision Making. *Pers. Soc. Psychol. Bull.*, 34(2), 224–236. <https://doi.org/10.1177/0146167207310023>
- Pronin, E., and Ross, L. (2006). Temporal differences in trait self-ascription: When the self is seen as an other. *J. Pers. Soc. Psychol.*, 90(2), 197–209. <https://doi.org/10.1037/0022-3514.90.2.197>
- Qin, P., Wang, M., and Northoff, G. (2020). Linking bodily, environmental and mental states in the self—A three-level model based on a meta-analysis. *Neurosci. Biobehav. Rev.*, 115, 77–95. <https://doi.org/10.1016/j.neubiorev.2020.05.004>
- Quadt, L., Garfinkel, S. N., Mulcahy, J. S., Larsson, D. E., Silva, M., Jones, A.-M., Strauss, C., and Critchley, H. D. (2021). Interoceptive training to target anxiety in autistic adults (ADIE): A single-center, superiority randomized controlled trial. *EClinicalMedicine*, 39, 101042. <https://doi.org/10.1016/j.eclinm.2021.101042>
- Ring, C., Brener, J., Knapp, K., and Mailloux, J. (2015). Effects of heartbeat feedback on beliefs about heart rate and heartbeat counting: A cautionary tale about interoceptive awareness. *Biol. Psychol.*, 104, 193–198. <https://doi.org/10.1016/j.biopsycho.2014.12.010>
- Schäfer, S., Wesslein, A.-K., Spence, C., Wentura, D., and Frings, C. (2016). Self-prioritization in vision, audition, and touch. *Exp. Brain Res.*, 234(8), 2141–2150. <https://doi.org/10.1007/s00221-016-4616-6>
- Schandry, R. (1981). Heart beat perception and emotional experience. *Psychophysiology*, 18(4), 483–488. <https://doi.org/10.1111/j.1469-8986.1981.tb02486.x>
- Schauder, K. B., Mash, L. E., Bryant, L. K., and Cascio, C. J. (2015). Interoceptive ability and body awareness in autism spectrum disorder. *J. Exp. Child Psychol.*, 131, 193–200. <https://doi.org/10.1016/j.jecp.2014.11.002>
- Sel, A., Azevedo, R. T., and Tsakiris, M. (2017). Heartfelt self: cardio-visual integration affects self-face recognition and interoceptive cortical processing. *Cereb. Cortex*, 27(11), 5144–5155. <https://doi.org/10.1093/cercor/bhw296>
- Sel, A., Sui, J., Shepherd, J., and Humphreys, G. (2019). Self-association and attentional processing regarding perceptually salient items. *Rev. Phil. Psych.*, 10(4), 735–746. <https://doi.org/10.1007/s13164-018-0430-3>
- Seth, A. K. (2013). Interoceptive inference, emotion, and the embodied self. *Trends Cogn. Sci.*, 17(11), 565–573. <https://doi.org/10.1016/j.tics.2013.09.007>
- Snodgrass, J. G., and Corwin, J. (1988). Pragmatics of Measuring Recognition Memory: Applications to Dementia and Amnesia. *J. Exp. Psychol. Gen.*, 117(1), 34–50. <https://doi.org/10.1037/0096-3445.117.1.34>
- Sui, J., and Gu, X. (2017). Self as object: Emerging trends in self research. *Trends Neurosci.*, 40(11), 643–653. <https://doi.org/10.1016/j.tins.2017.09.002>
- Sui, J., He, X., and Humphreys, G. W. (2012). Perceptual effects of social salience: Evidence from self-prioritization effects on perceptual matching. *J. Exp. Psychol. Hum. Percept.*

- Perform., 38(5), 1105–1117. <https://doi.org/10.1037/a0029792>
- Sui, J., and Humphreys, G. W. (2015). Super-size me: Self biases increase to larger stimuli. *Psychon. Bull. Rev.*, 22(2), 550–558. <https://doi.org/10.3758/s13423-014-0690-6>
- Suzuki, K., Garfinkel, S. N., Critchley, H. D., and Seth, A. K. (2013). Multisensory integration across exteroceptive and interoceptive domains modulates self-experience in the rubber-hand illusion. *Neuropsychologia*, 51(13), 2909–2917. <https://doi.org/10.1016/j.neuropsychologia.2013.08.014>
- Tajadura-Jiménez Tsakiris, M., and Critchley, H. (2016). Interoception beyond homeostasis: Affect, cognition and mental health. *Philos. Trans. R. Soc. Lond. B: Biol. Sci.*, 371(1708), 20160002. <https://doi.org/10.1098/rstb.2016.0002>
- Tsakiris, M., Jiménez, A. T., and Costantini, M. (2011). Just a heartbeat away from one's body: Interoceptive sensitivity predicts malleability of body-representations. *Proc. R. Soc. B: Biol. Sci.*, 278(1717), 2470–2476. <https://doi.org/10.1098/rspb.2010.2547>
- Verplanken, B., and Sui, J. (2019). Habit and identity: Behavioral, cognitive, affective, and motivational facets of an integrated self. *Front. Psychol.*, 10. <https://doi.org/10.3389/fpsyg.2019.01504>
- Villani, V., Tsakiris, M., and Azevedo, R. T. (2019). Transcutaneous vagus nerve stimulation improves interoceptive accuracy. *Neuropsychologia*, 134, 107201. <https://doi.org/10.1016/j.neuropsychologia.2019.107201>
- von Mohr, M., Finotti, G., Villani, V., and Tsakiris, M. (2021). Taking the pulse of social cognition: Cardiac afferent activity and interoceptive accuracy modulate emotional egocentricity bias. *Cortex*, 145, 327–340. <https://doi.org/10.1016/j.cortex.2021.10.004>
- Zamariola, G., Maurage, P., Luminet, O., and Corneille, O. (2018). Interoceptive accuracy scores from the heartbeat counting task are problematic: Evidence from simple bivariate correlations. *Biol. Psychol.*, 137, 12–17. <https://doi.org/10.1016/j.biopsycho.2018.06.006>
- Zimprich, D., Nusser, L., and Pollatos, O. (2020). Are interoceptive accuracy scores from the heartbeat counting task problematic? A comment on Zamariola et al. (2018). *Biol. Psychol.*, 152, 107868. <https://doi.org/10.1016/j.biopsycho.2020.107868>

12 Figure legends

Figure 1. The procedures of the matching phase and the timings of stimulus presentation in the shape-label matching task.

(A) The participants judged whether the presented shape-label pair was matched based on the associations learned during the learning phase. (B) The shape-label pairs were presented in coincidence with either systole (T-wave peak) or diastole (T-wave peak + 250 ms) in the participant's cardiac cycle.

Figure 2. Results of correlation analyses

Correlations between IAc and the magnitude of SPE on d-prime in the both Systole, and Diastole (A), Systole (B) and Diastole (C) conditions. Correlations between IAc and d-prime in the Self-Systole (D) and Stranger-Systole (E) conditions.

Table 1. Mean reaction time (RT) and d-prime as a function of shape category and cardiac cycle.

Cardiac cycle	Shape category	Mean RT (ms)		d-prime	
Both (Systole and Diastole)	Self	561.13	(66.30)	3.12	(0.80)
	Friend	596.08	(64.45)	2.83	(0.82)
	Stranger	624.58	(62.88)	2.64	(0.67)
Systole	Self	559.71	(64.97)	2.99	(0.69)
	Friend	599.71	(65.46)	2.70	(0.80)
	Stranger	626.19	(66.34)	2.53	(0.65)
Diastole	Self	562.48	(70.19)	3.02	(0.83)
	Friend	591.95	(66.92)	2.88	(0.92)
	Stranger	623.14	(63.72)	2.68	(0.77)

Note. Standard deviations (SDs) are shown in parentheses.