A Simple Diffusion Model for Psychometric Analyses

Udo Boehm^a, Maarten Marsman^a, Han L. J. van der Maas^a, Gunter Maris^b

Author Note

^aDepartment of Psychology, University of Amsterdam, Amsterdam, The Netherlands, email Udo Boehm: u.bohm@uva.nl; email Maarten Marsman: M.Marsman@uva.nl; email Han L. J. van der Maas: h.l.j.vandermaas@uva.nl

^bACT, Iowa City, USA, email: gunter.maris@act.org

The authors declare no competing financial interests. This research was supported by a Netherlands Organisation for Scientific Research (NWO) creative industry grant (314-99-107) to GM and a Veni grant (451-17-017) to MM.

Correspondence concerning this article should be addressed to Udo Boehm, Department of Psychology, University of Amsterdam, Nieuwe Prinsengracht 129B, 1018 WS Amsterdam, The Netherlands, Email: u.bohm@uva.nl

All programming code for the simulations is available at <https://osf.io/fh4wp/>

Abstract

The emergence of computer-based assessments has made response times, in addition to response accuracies, available as a source of information about test-takers' latent abilities. The predominant approach to jointly account for response times and accuracies are statistical models. Substantive approaches such as the diffusion model, on the other hand, have been slow to gain traction due to their unwieldy functional form. In the present work we show how a single simplifying assumption yields a highly tractable diffusion model. This simple diffusion model is straightforward to analyse using Gibbs sampling and can be readily extended with a latent regression framework. We demonstrate the superior computational efficiency of our model compared to the standard diffusion model in a simulation study and showcase the theoretical merit of our model in an example application.

keywords: Diffusion model, response time data, cognitive psychometrics

A Simple Diffusion Model for Psychometric Analyses

Introduction

The emergence of computer-based assessments has introduced new opportunities to record additional sources of information – other than response accuracy – that can be used to infer the latent abilities of test-takers. One additional source of information that has received much attention in the psychometric literature is response time (RT), and several approaches have been developed that use both RT and response accuracy to assess the latent trait level of test-takers. These approaches broadly fall into one of three categories (Tuerlinckx, Molenaar, & van der Maas, 2016). The collateral information approach treats RTs as an additional source of information that allows the item response parameters to be estimated more efficiently (e.g., Van Breukelen, 2005; Van der Linden, 2006; Van der Linden, 2007). The normative approach views RTs as the result of respondents complying with a particular scoring rule (e.g., Maris & van der Maas, 2012; Van Rijn & Ali, 2017). The process approach attempts to model the cognitive processes underlying the observed RTs (e.g., Ranger & Kuhn, 2018; Rouder, Province, Morey, Gomez, & Heathcote, 2015; Tuerlinckx & De Boeck, 2005; Van der Maas, Molenaar, Maris, Kievit, & Borsboom, 2011).

Although all three approaches have the common goal to model the RT and accuracy distribution in terms of latent traits, they differ markedly in the way that RTs are related to the latent variables. The most popular instantiation of the collateral information approach, Van der Linden's (2007) Hierarchical Model framework, specifies separate measurement models for RTs, in terms of latent speed parameters, and for accuracies, in terms of latent ability parameters. The relationship between RTs and accuracies is established via a second level of modelling where population models specify the joint distribution of the latent speed and ability factors. This approach is computationally very efficient and affords the user a high degree of flexibility in choosing the measurement models. However, the separation of speed and ability factors in the measurement models has resulted in some debates in the literature (e.g., Van der Linden, 2009; Van der Maas et al., 2011).

Models from the normative approach and the cognitive model approach, on the other hand, typically directly relate latent ability parameters to RT and accuracy. Maris and van der Maas's (2012) Signed Residual-Time Model, for instance, provides a normative model for settings in which the Signed Residual Time scoring rule is used. This model proceeds from the assumption that the SRT score is a sufficient statistic for a person's latent ability, which leads to a highly tractable exponential family model that has the standard two-parameter logistic model as its item response function. However, one disadvantage of this model is that applications are limited to niche-settings where the SRT scoring rule is used (Klinkenberg, Straatemeier, & Van Der Maas, 2011). The Diffusion Model framework (DM, Tuerlinckx & De Boeck, 2005; Van der Maas et al., 2011) was developed in mathematical psychology as a process model in which different model parameters represent different cognitive process involved in answering an item (Ratcliff, 1978). Besides this attractive process interpretation of the model parameters, the DM can be applied in a wide variety of settings as it is based on a more general scoring rule that only supposes that decision makers minimise their response time for a given level of accuracy. Moreover, Tuerlinckx and De Boeck (2005) point out that, for an appropriate latent-trait parameterisation, the DM's item response function reduces to the standard two-parameter logistic model. The DM thus combines desirable features of the HM and the SRTM framework.

Unfortunately, current models that are formulated in the DM framework are computationally intractable, at least for psychometric applications, which typically involve hundreds or several thousand test-takers responding to dozens of test questions. The reason for this intractability is that the DM parameters appear in a series expression that has no analytic solution. As a result, the series needs to be approximated repeatedly, for each test-taker-by-question combination, and each iteration of the estimation procedure. Hence, whilst in the HM framework it is straightforward to build a tractable model, including complete hierarchical modelling of the psychometric data, applications of the DM framework are currently limited to relatively constrained settings.

The objective of the current paper is to show how a single simplifying assumption results in a fully tractable latent trait DM for RTs and accuracy. One appealing feature of our simplifi-

cation is that the kernel of the DM is then proportional to a normal density in each of its parameters. When combined with conjugate prior distributions in a Bayesian framework, we show that the joint posterior distribution is easily sampled from using the Gibbs sampler as each of the full-conditional posterior distributions are easy-to-sample-from (truncated) normal densities. In sum, this paper introduces a fully tractable version of the DM that can be feasibly applied to large-scale psychometric problems.

The Diffusion Model

The DM was originally developed in mathematical psychology as a process model for memory retrieval (Ratcliff, 1978). Over the years the model has been applied to numerous other domains of cognitive psychology, the most prominent of which is decision-making (for a review, see Ratcliff & McKoon, 2008). The DM describes the process by which a test-taker chooses between two response options, A and B, as the noisy accumulation of information. In its basic form, the DM includes five model parameters, t_0 , v , s , a , and z (see Figure 1) that account for the joint distribution of response time $t \in \mathbb{R}^+$ and choice $x \in \{0, 1\}$, where $x = 1$ corresponds to choosing option A and $x = 0$ corresponds to choosing option B. The model decomposes the response time as $t = t_0 + t_d$. The constant t_0 is the non-decision time, which accounts for the time spent on cognitive processes not related to the choice process, such as reading the item and carrying out a motor response. The distribution of the decision time t_d is determined by the remaining four model parameters. The DM represents the choice process as a repeated sampling process where the test-taker samples and integrates noisy information about the two response options from a normal distribution with mean given by the drift rate parameter v and standard deviation given by the diffusion coefficient s . Each response option is associated with a decision threshold, one located at 0 and the other located at a , and a is referred to as the boundary separation. The sampling process starts at the point z , located between the two decision thresholds, and continues until a decision threshold is reached. The point where the accumulated evidence first crosses a decision threshold determines the decision time t_d , and the test-taker commits to the corresponding response option.

Figure 1: Diffusion Model (DM) and its parameters. Distributions of decision times for option A and option B are shown above and below the corresponding decision threshold.

The joint density of t and x is:

$$\begin{aligned} \mathbb{P}(t, x) = & \frac{\pi s^2}{a^2} \exp \left([ax - z] \frac{v}{s^2} - (t - t_0) \frac{v^2}{2s^2} \right) \\ & \times \sum_{k=1}^{\infty} \left[k \sin \left(\frac{\pi k [ax - 2xz + z]}{a} \right) \exp \left(-\frac{\pi^2 s^2 k^2}{2a^2} (t - t_0) \right) \right]. \end{aligned} \quad (1)$$

As can be seen, this density is only identified up to a multiplicative constant; multiplying v , s , a , and z by a constant would yield the same density. The standard way to identify the model is to set the diffusion coefficient to a fixed value, typically $s = 1$. Substantively, this identification choice means that the decision process is equally noisy across test-takers and items, which is an appropriate assumption in experimental applications but might be less appropriate in a psychometric context. We will revisit this issue when we present our simplified diffusion model.

The effect of the non-decision time parameter t_0 is to shift the decision time distribution. The remaining parameters are most easily interpreted in terms of the speed and accuracy of responding on an ability test. Therefore, assume that option A is the correct response to an item on an ability test. The drift rate v is typically interpreted as the speed of information processing. It determines how much information each sample conveys about the correct choice; a higher drift rate therefore leads to faster and more accurate choices. The diffusion coefficient s determines the volatility of the information. A higher diffusion coefficient means that the accumulated informa-

tion is more likely to jump to one of the decision thresholds, which results in faster choices. At the same time, due to the increased volatility, a higher diffusion coefficient also implies a higher probability of an incorrect choice. The boundary separation a is typically interpreted as the response caution with which a person processes an item. It determines the amount of information required for a choice; a larger boundary separation leads to slower but more accurate choices as more information needs to be accumulated. Finally, the starting point z reflects a bias in the choice process towards one response option. A starting point that is closer to the upper threshold means that less information is required to choose option A, thus increasing the probability of choosing option A.

From the joint density (1), expressions can be derived for the mean response time (Cox & Miller, 1970; Laming, 1973):

$$\mathbb{E}(T) = \frac{a}{2v} \frac{1 - \exp\left(-\frac{va}{s^2}\right)}{1 + \exp\left(-\frac{va}{s^2}\right)}, \quad (2)$$

and the probability of choosing option A:

$$\mathbb{P}(X = 1) = \frac{\exp\left(-\frac{2zv}{s^2}\right) - 1}{\exp\left(-\frac{2av}{s^2}\right) - 1}. \quad (3)$$

The latter expression suggests a close link between the DM and the logistic item response model. As pointed out by Tuerlinckx and De Boeck (2005), the choice probability of the two-parameter logistic model can be obtained from the choice probability of the DM by making two simplifying assumptions. Firstly, if the diffusion coefficient is fixed to one and the decision process is assumed to be unbiased, that is, $z = a/2$, Equation (3) simplifies to $\exp(va) / (\exp(va) + 1)$. Secondly, the drift rate for person p answering item i can be decomposed as $v_{ip} = \theta_p + \beta_i$, where θ_p is the person ability parameter and β_i is the item easiness parameter. This yields the expression for the choice probability in the 2-PLM, $\exp(a(\theta_p + \beta_i)) / (\exp(a(\theta_p + \beta_i)) + 1)$, where a plays the role of the item discrimination parameter. Substantively, this parameterisation suggests that the speed and accuracy of information processing is determined by the interaction between a person's ability and the item easiness, with higher ability and higher easiness resulting

in faster and more accurate responding. Moreover, the boundary separation reflects the response caution elicited by an item, with more discriminating items resulting in slower and more accurate responding.

Simplifying the Diffusion Model

The main obstacle to a computationally efficient implementation of the DM is the dependence of the series expression in the density (1) on the model parameters a , z , and s . The goal of psychometric applications is typically to estimate model parameters for a large number of persons answering a comparatively small number of items. If, for example, the boundary separation parameter a is assumed to differ between persons, estimation of this parameter will require the evaluation of the likelihood function (1) once for each person at each step of the estimation algorithm. As a appears in the series expression in the likelihood function, each evaluation will require approximating this series. Even the most efficient approximation methods require computing the series to at least 4 or 5 terms at each iteration (Navarro & Fuss, 2009). The computational costs increase exponentially as further DM parameters depend on item or person effects.

To resolve the computational inefficiency, we first introduce the reparameterisation $c = s/a$. This yields for the diffusion density:

$$\mathbb{P}(t, x) = \pi c^2 \exp\left([ax - z] \frac{v}{s^2} - t \frac{v^2}{2s^2}\right) \times \sum_{k=1}^{\infty} \left[k \sin\left(\frac{\pi k[ax - 2xz + z]}{a}\right) \exp\left(-\frac{\pi^2 c^2 k^2}{2} t\right) \right].$$

If we assume that c is constant, say $c = 2$, the dependence of the exp function on the parameters s and a in the series expression is removed. Substantively, such a constant ratio of boundary separation to the diffusion coefficient means that test-takers will accumulate more information on items for which information is more volatile.

Furthermore, the assumption $z = a/2$ introduced by Tuerlinckx and De Boeck (2005) also removes the dependence of the sin function on the parameters a and z in the series expression and thus yields a fully tractable model:

$$\mathbb{P}(t, x) = \pi c^2 \exp\left([x - 1/2] \frac{v}{cs} - t \frac{v^2}{2s^2}\right) \times \sum_{k=1}^{\infty} \left[k \sin\left(\frac{\pi k}{2}\right) \exp\left(-\frac{\pi^2 c^2 k^2}{2} t\right) \right].$$

Substantively, a starting point that is equidistant from the two decision thresholds means that test-takers are not biased toward one response option.

This latter simplifying assumption is shared with several other simplifications of the DM (e.g. Tuerlinckx & De Boeck, 2005; Van der Maas et al., 2011; Wagenmakers, Van der Maas, & Grasman, 2007) and seems justified on conceptual grounds in psychometric applications. There is no reason to assume that test-takers a priori prefer one response option over another because the assignment of the correct response to a response option is typically balanced across items. Moreover, an a priori preference for one response option would vary randomly across test-takers and thus only be identifiable at the level of individual test-takers.

The first simplifying assumption that the ratio of the diffusion coefficient and boundary separation is constant might appear less obvious. Firstly, the diffusion coefficient is treated as a model parameter that needs to be estimated here whilst in traditional applications it is set to a fixed value. A fixed level of volatility in the diffusion process is clearly an appropriate assumption in mathematical psychology, where experiments consist of the repeated presentation of homogeneous stimuli and researchers have full control over all statistical properties of the stimuli. In the popular Random Dot Motion task, for example, stimuli consist of a cloud of pseudo-randomly moving dots that is constructed to elicit the same level of visual noise across experimental trials (Britten, Shadlen, Newsome, & Movshon, 1992). Researchers can control the difficulty of the experimental task by increasing or decreasing the proportion of consistently moving dots independently of the level of visual noise. In psychometric applications, however, different statistical aspects of the items are more difficult to control. Although test items might have the same visual appearance, it is not clear whether the difficulty of an item can be manipulated independently of the level of visual or physiological noise elicited by the item. Therefore, it seems more appropriate to allow for variation in the volatility of the diffusion process across items and treat the diffusion coefficient as a model parameter that needs to be estimated.

Secondly, a constant ratio between boundary separation and the diffusion coefficient implies that test-takers adjust to changes in the volatility of information across items by chang-

ing their response caution. This assumption was recently debated in mathematical psychology where it has been suggested that decision-makers adjust their boundary separation in response to changes in the signal-to-noise ratio (e.g., Deneve, 2012; Ditterich, 2006; Shadlen & Kiani, 2013). There is some evidence to suggest that decision-makers are capable of such adjustments in the context of perceptual decision-making, even within a single decision (e.g., Drugowitsch, Moreno-Bote, Churchland, Shadlen, & Pouget, 2012; Hanks, Mazurek, Kiani, Hopp, & Shadlen, 2011; Thura, Beauregard-Racine, Fradet, & Cisek, 2012). How well this assumption generalises to psychometric applications will need to be tested in future applications of our model.

Together, our simplifying assumptions yield a model with two parameters, v and s . It is natural to assume that the drift rate v depends on the item difficulty and the test-taker's ability, as suggested by Tuerlinckx and De Boeck (2005). That is, for person p , $p = 1, \dots, P$, answering items i , $i = 1, \dots, I$, we assume additive effects of the person's ability θ_p , and the item easiness β_i , on drift rate:

$$v_{ip} = \beta_i + \theta_p.$$

Moreover, following our discussion above, the diffusion coefficient should act as the item discrimination parameter α_i :

$$s_{ip} = \alpha_i.$$

The resulting likelihood function for our simple diffusion model (sDM) is:

$$\begin{aligned} \mathbb{P}(t_{ip}, x_{ip} \mid \beta_i, \theta_p, \alpha_i) = & \pi c^2 \exp \left([x_{ip} - 1/2] \frac{\beta_i + \theta_p}{c \alpha_i} - t_{ip} \frac{(\beta_i + \theta_p)^2}{2 \alpha_i^2} \right) \\ & \sum_{k=1}^{\infty} \left[k \sin \left(\frac{\pi k}{2} \right) \exp \left(- \frac{\pi^2 c^2 k^2}{2} t_{ip} \right) \right] \end{aligned} \quad (4)$$

and dropping all constants not related to β_i , θ_p , or α_i and letting $\alpha_i^{-1} = \tilde{\alpha}_i$:

$$\propto \exp \left([x_{ip} - 1/2] \tilde{\alpha}_i \frac{(\beta_i + \theta_p)}{c} - t_{ip} \tilde{\alpha}_i^2 \frac{(\beta_i + \theta_p)^2}{2} \right).$$

Note that this model is only identified up to a multiplicative constant, which we will address below when we introduce random effects.

The form of the kernel of our sDM suggests that a Bayesian analysis of our model might proceed by choosing a multivariate normal distribution as joint prior for the model parameters β_i , θ_p and $\tilde{\alpha}_i$, and an inverse Wishart prior for the variance-covariance matrix, which would yield a joint multivariate normal posterior distribution for all parameters. However, the interpretation of $\tilde{\alpha}_i$ in terms of the diffusion coefficient means that only positive values are admissible for this parameter, and the prior distribution would therefore need to be truncated at zero. The approach we will take here is to assign independent (truncated) normal priors to the model parameters and set up a Gibbs sampler to analyse our model, which allows us to retain the cognitive process interpretation of the model parameters.

The probability of a correct response in our sDM is given by:

$$\mathbb{P}(X = 1) = \frac{\exp(\tilde{\alpha}_i/c(\theta_p + \beta_i))}{1 + \exp(\tilde{\alpha}_i/c(\theta_p + \beta_i))}. \quad (5)$$

This is the choice probability in the 2-PLM, where $\tilde{\alpha}_i/c$ plays the role of the item discrimination parameter, θ_p is the person ability, and β_i is the item easiness parameter. Hence, in the absence of RT data our sDM reduces to the 2-PLM.

Implicit in our development of the sDM are two further simplifications. Firstly, we ignore the non-decision component of the response time and assume that the response time equals the decision time. Although the role of the non-decision component might be negligible in most psychometric applications, researchers might nevertheless be interested in estimating non-decision time. We will show below how our model can be extended to allow for explicit estimation of a non-decision component. Secondly, we assume that the ratio of boundary separation and diffusion coefficient c is a known constant. However, in practical applications c itself might be a variable of interest that researchers would like to estimate from their data. Below we will suggest how this problem might be addressed.

Random Effects

The likelihood function of our sDM, Equation (4), has a simple form that makes it amenable to extension. The terms of the series expression do not depend on any of the model parameters

and can therefore be dropped in analyses of the likelihood function, which leaves a simple exponential family model. Moreover, the item and person effects on drift rate, β_i and θ_p , and the item effects on the diffusion coefficient, $\tilde{\alpha}_i$, only appear in the first and second powers. Hence, the full conditional distributions introduced below take the form of a normal kernel, which allows for simple conjugate analysis and easy sampling-based estimation.

One problem that is regularly of interest in psychometrics is the modelling of unobserved heterogeneity. We show here how our sDM can be extended to accommodate a random effects structure. Extensions to other statistical problems, such as missing data, inclusion of manifest predictor variables, or the addition of autoregressive components are straightforward.

To model heterogeneity in the data we assume the following population-level distributions:

$$\beta_i \sim \mathcal{N}(\mu_I, \sigma_I^2), \quad \theta_p \sim \mathcal{N}(\mu_P, \sigma_P^2), \quad \tilde{\alpha}_i \sim \mathcal{N}(\mu_S, \gamma_S^2)[0, \infty).$$

Here $\mathcal{N}(\mu, \sigma^2)$ denotes the normal distribution with mean μ and variance σ^2 . The expression $[0, \infty)$ indicates the support of the truncated distribution. The population-level parameters can be assigned prior distributions in a similar manner:

$$\mu_P \sim \mathcal{N}(v_{P0}, \tau_{P0}^2), \quad \sigma_P^2 \sim \text{invG}(a_{P0}, b_{P0}), \quad \mu_I \sim \mathcal{N}(v_{I0}, \tau_{I0}^2), \quad \sigma_I^2 \sim \text{invG}(a_{I0}, b_{I0}).$$

Here $\text{invG}(a, b)$ denotes the inverse gamma distribution with shape parameter a and scale parameter b . As μ_S and γ_S^2 will usually not be the target of statistical inference, these parameters can simply fixed to specific values.

The full-conditional conjugate posterior distributions for the person and item effects on drift rate v_{ip} for N_P persons answering N_I items are:

$$\beta_i \sim \mathcal{N}(\hat{\beta}_{i,N_P}, \hat{\sigma}_{i,N_P}^2), \quad \theta_p \sim \mathcal{N}(\hat{\theta}_{p,N_I}, \hat{\sigma}_{p,N_I}^2),$$

where

$$\begin{aligned} \hat{\sigma}_{i,N_P}^2 &= \left(\sum_{p=1}^{N_P} \frac{t_{ip}}{\alpha_i^2} + \frac{1}{\sigma_I^2} \right)^{-1}, \quad \hat{\beta}_{i,N_P} = \left(\sum_{p=1}^{N_P} \frac{(x_{ip}-1/2)}{c\alpha_i} - \sum_{p=1}^{N_P} \frac{t_{ip}\theta_p}{\alpha_i^2} + \frac{\mu_I}{\sigma_I^2} \right) \hat{\sigma}_{i,N_P}^2 \\ \hat{\sigma}_{p,N_I}^2 &= \left(\sum_{i=1}^{N_I} \frac{t_{ip}}{\alpha_i^2} + \frac{1}{\sigma_P^2} \right)^{-1}, \quad \hat{\theta}_{p,N_I} = \left(\sum_{i=1}^{N_I} \frac{(x_{ip}-1/2)}{c\alpha_i} - \sum_{i=1}^{N_I} \frac{t_{ip}\beta_i}{\alpha_i^2} + \frac{\mu_P}{\sigma_P^2} \right) \hat{\sigma}_{p,N_I}^2. \end{aligned}$$

Similarly, the full-conditional conjugate posterior distribution for item effects on the reciprocal diffusion coefficient s_{ip}^{-1} is:

$$\tilde{\alpha}_i \sim \mathcal{N}(\hat{\alpha}_{i,N_p}, \hat{\gamma}_{i,N_p}^2)[0, \infty),$$

where

$$\hat{\gamma}_{i,N_p}^2 = \gamma_S^2 \left(\sum_{p=1}^{N_p} t_{ip} (\beta_i + \theta_p)^2 \gamma_S^2 + 1 \right)^{-1}, \quad \hat{\alpha}_{i,N_p} = \left(\mu_S / \gamma_S^2 + \sum_{p=1}^{N_p} (x_{ip} - 1/2) (\beta_i + \theta_p) / c \right) \hat{\gamma}_{i,N_p}^2.$$

The full-conditional posterior distributions for the population-level parameters are:

$$\begin{aligned} \mu_P &\sim \mathcal{N}(v_{P,N_p}, \tau_{P,N_p}^2), & \text{where } v_{P,N_p} &= \frac{\tau_P^2 \sum_{p=1}^{N_p} \theta_p + \sigma_P^2 v_{P,0}}{\tau_{P,N_p}^2 + \sigma_P^2}, & \tau_{P,N_p}^2 &= \frac{\tau_P^2 \sigma_P^2}{\tau_{P,N_p}^2 + \sigma_P^2} \\ \sigma_P^2 &\sim \text{invG}(a_{P,N_p}, b_{P,N_p}), & \text{where } a_{P,N_p} &= a_{P,0} + N_p/2, & b_{P,N_p} &= \sum_{p=1}^{N_p} (\theta_p - \mu_P)^2/2 + b_{P,0} \\ \mu_I &\sim \mathcal{N}(v_{I,N_I}, \tau_{I,N_I}^2), & \text{where } v_{I,N_I} &= \frac{\tau_I^2 \sum_{i=1}^{N_I} \beta_i + \sigma_I^2 v_{I,0}}{\tau_{I,N_I}^2 + \sigma_I^2}, & \tau_{I,N_I}^2 &= \frac{\tau_I^2 \sigma_I^2}{\tau_{I,N_I}^2 + \sigma_I^2} \\ \sigma_I^2 &\sim \text{invG}(a_{I,N_I}, b_{I,N_I}), & \text{where } a_{I,N_I} &= a_{I,0} + N_I/2, & b_{I,N_I} &= \sum_{i=1}^{N_I} (\beta_i - \mu_I)^2/2 + b_{I,0} \end{aligned}$$

Let $\vec{t} = [t_{11}, \dots, t_{IP}]$ be the vector of response times and let $\vec{x} = [x_{11}, \dots, x_{IP}]$ be the vector of accuracies. Moreover, denote by Θ the set of all item, person, and population-level parameters, then:

$$\begin{aligned} \mathbb{P}(\vec{t}, \vec{x} \mid \Theta) \mathbb{P}(\Theta) &\propto \exp \left(\sum_{i=1}^{N_I} \sum_{p=1}^{N_p} \left(\tilde{\alpha}_i [x_{ip} - 1/2] \frac{\beta_i + \theta_p}{c} - t_{ip} \tilde{\alpha}_i^2 \frac{(\beta_i + \theta_p)^2}{2} \right) \right) \\ &\cdot \exp \left(- \left(\sum_{i=1}^{N_I} \frac{(\beta_i - \mu_I)^2}{2\sigma_I^2} + \sum_{p=1}^{N_p} \frac{(\theta_p - \mu_P)^2}{2\sigma_P^2} + \sum_{i=1}^{N_I} \frac{(\tilde{\alpha}_i - \mu_S)^2}{2\gamma_S^2} \right) \right) \\ &\cdot \exp \left(- \left(\frac{(\mu_I - \mu_{I0})^2}{2\tau_{I0}^2} + \frac{(\mu_P - \mu_{P0})^2}{2\tau_{P0}^2} + \frac{b_I}{\sigma_I^2} + \frac{b_P}{\sigma_P^2} \right) \right) \\ &\cdot \sigma_I^{-2(a_I+1+N_I/2)} \sigma_P^{-2(a_P+1+N_P/2)}. \end{aligned}$$

This random effects extension of our model is not identified, which we address by fixing three model parameters:

$$\mu_I = 0, \quad \sigma_I^2 = 1, \quad \sigma_P^2 = 1.$$

The first constraint addresses the trade-off between the β_i and θ_p , where adding a constant to all β_i and subtracting the same constant from all θ_p yields the same value for the likelihood. By

requiring $\mu_I = 0$ we fix the location of the item effects on drift rate. The second constraint addresses the indeterminate scale of the item and person effects, where multiplying all β_i and θ_p by a constant is compensated for by multiplying the corresponding variance terms by the same constant. The third constraint addresses the trade-off between the drift rate v_{ip} and the diffusion coefficient α_i , which is again only determined up to a multiplicative constant. In addition to these technical constraints, we propose fixing $\mu_S = 1$ and $\gamma_S^2 = 1/2$ as these parameters are not usually the target of statistical inference.

MCMC Sampling

The analysis of our model relies on MCMC sampling from the posterior distribution. The full conditional posterior distributions for all parameters are conjugate to the prior distributions and have standard functional forms. We can therefore easily obtain MCMC samples using the Gibbs sampler (Gelfand & Smith, 1990), which is implemented by alternately sampling from the full conditional posterior distributions.

Simulation Study

We tested the sampling behaviour of our simplified DM in a simulation study. We used the RTdists (Singmann et al., 2016) R package to generate RT and choice data for 120 persons answering 70 items. The generating population-level parameters were set to $\mu_I = 0$, $\sigma_I^2 = 1$, $\mu_P = 1$, $\sigma_P^2 = 1$, $\mu_S = 1$, and $\gamma_\alpha^2 = 1/2$. We truncated the generating distribution for $\tilde{\alpha}_i$ at $1/2$ instead of 0 to avoid unrealistically large diffusion coefficients in the simulated data. We furthermore set the ratio of diffusion coefficient and boundary separation to $c = 2$. As the DM was developed for experimental applications in mathematical psychology, where RTs usually do not exceed 2.5s, we multiplied the generated RTs by 10 to cover a wider range of values. Panel A of Figure 2 shows the joint distribution of mean accuracies and mean RTs for the simulated items. As can be seen, both variables spanned a realistic range of values. Panel B of Figure 2 shows the marginal distribution of RTs. All RTs fell in the range below 10s, which is comparable to the range seen in performance data from an ability test. Moreover, the distribution exhibits the skew

Figure 2: Simulation study with 120 persons answering 70 items. (A) Accuracy and mean RT for simulated items. (B) Marginal RT distribution over items and persons. (C) Estimated and true values for the item (β_i) and person (θ_p) effects on drift rate, and for the item effects on the reciprocal diffusion coefficient ($\tilde{\alpha}_i$). (D) Example MCMC chains for item and person effects after a burn-in period of 2000 samples.

and long tail that is characteristic of RT data.

We implemented our random effects model using custom-written R code (available at <https://osf.io/fh4wp/>) in combination with the algorithm for sampling from truncated normal distributions implemented in the truncnorm package (Mersmann, Trautmann, Steuer, & Bornkamp, 2018). For each model parameter we obtained 3 chains with 22000 MCMC samples. We discarded the first 2000 samples as burn-in samples. Panel C of Figure 2 shows the posterior mean estimates of the person and item effects θ_p , β_i , and $\tilde{\alpha}_i$. As can be seen, all parameters could be recovered accurately. Panel D of Figure 2 shows three example chains for the person and item effects. All three parameters, β_2 , θ_2 and $\tilde{\alpha}_2$ showed good sampling behaviour with relatively small autocorrelations.

Table 1

Convergence results for simulated data.

Parameter	$\hat{R}$		
μ_P	1.014		
	Quantiles		
	0.25	0.5	0.75
β_i	1.004	1.007	1.010
θ_p	1.009	1.010	1.010
$\tilde{\alpha}_i$	<1.001	1.001	1.001

Note. Quantiles are computed across persons or items.

Table 1 shows a summary of the potential scale reduction factor $\hat{R}$ (Gelman & Rubin, 1992). As can be seen, $\hat{R}$ was approximately 1.01 for the population parameter μ_p . Similarly, $\hat{R}$ did not exceed 1.01 for all chains for all parameters β_i , θ_p and $\tilde{\alpha}_i$. Taken together, the results of our simulation study show that person and item effects on drift rate and item effects on the diffusion coefficient can be estimated with good efficiency and generating values can be recovered with high accuracy.

To gauge the efficiency of our sDM relative to the classical DM, we compared the computing time of the two models when applied to small subsets of a simulated data set. Specifically, we fitted the sDM and DM model to RT and accuracy data from 50 test-takers answering 10, 20, 30, 40, 50, 75, 100, 150, or 200 items. The data were generated with the same parameter values as in the parameter recovery simulation above. To fit the DM, we used HDM module (Vandekerckhove, Tuerlinckx, & Lee, 2011; Wabersich & Vanderkerckhove, 2014) available in Stan (Carpenter et al., 2017) and specified the model as follows:

$$\begin{aligned}
\mu_a &\sim \mathcal{N}(1, 1/2)[0,], & \sigma_a^2 &\sim \text{invG}(1, 1), & a_p &\sim \mathcal{N}(\mu_a, \sigma_a^2)[0,] \\
\mu_v &\sim \mathcal{N}(1, 2), & \sigma_v^2 &\sim \text{invG}(1, 1), & v_p &\sim \mathcal{N}(\mu_v, \sigma_v^2) \\
(\vec{t}_p, \vec{x}_p) &\sim \text{wiener}(a_p, v_p, t_0, z).
\end{aligned}$$

Here, $\vec{r}_p$ is the vector of RTs for test-taker p , $\vec{x}_p$ is the vector of the corresponding response accuracies, and a_p and v_p are the boundary separation and drift rate for test-taker p test-taker. $\mathcal{N}(\mu, \sigma^2)[0, \infty)$ denotes the normal distribution with mean μ and variance σ^2 , truncated at 0, and $wiener(a, v, t_0, z)$ is the density of DM given in Equation 1. We set the parameters $t_0 = 0$ and $z = 1/2$ for all test-takers. We ignored item effects on drift rate and boundary separation in this model specification as including such effects would make it practically impossible to obtain posterior samples for the DM density. Our modelling approach also aligns with the standard approach used in applications of the DM to comparable data sets (Wagenmakers, Ratcliff, Gomez, & McKoon, 2008; Yap, Balota, Sibley, & Ratcliff, 2012; Yap, Sibley, Balota, Ratcliff, & Rueckl, 2015).

Figure 3 shows the CPU time require to obtain 6000 MCMC samples, of which 1000 are burn in samples, for the DM (black line) and the sDM (grey line). Axes are on the logarithmic scale with base 2. As can be seen, the computing time for both models increases approximately linearly with increasing numbers of items. However, on the natural scale, the rate of increasing is 700 times as high for the classical DM as for the sDM.

Application

We assessed the performance of our model under realistic conditions by applying it to a psychometric data set. The Amsterdam Chess Test (ACT; Van der Maas & Wagenmakers, 2005) was developed as a quicker and more efficient way to gauge a chess player's skill level than the official ELO rating, which is based on the player's performance over several tournaments. The ACT comprises two parallel versions of the test, A and B, each of which consists of 40 items that vary in difficulty. The items are subdivided into a set of 20 tactical items, a set of 10 positional items, and a set of 10 endgame items. Van der Maas and Wagenmakers validated both versions of the test in a sample of 259 Dutch chess players. Here we apply our sDM model to the RT and accuracy data for the A version of the test. We excluded the data of three test-takers who did not complete all items.

We fitted our model, obtaining three chains of 12,000 samples for each model parameter, of which we discarded the first 2,000 samples as burn in. As before, we set the ratio of diffusion

Figure 3: Computing time for sDM and DM. Dots indicate the CPU time required to obtain 6000 MCMC samples for 50 test-takers and the indicated number of items. Axes are logarithmic with base 2.

coefficient and boundary separation to $c = 2$. Panel A of Figure 4 shows the relationship between the ELO rating and mean RT, mean accuracy, and estimated person effect on drift rate for the 256 test-takers. As can be seen, there was a trend for mean RT to be lower for players with higher ELO ratings, and there was a clear increase in mean accuracy with higher ELO ratings. Similarly, the posterior mean of the person effects θ_p clearly increased with higher ELO ratings. The cognitive process interpretation of this latter result is that higher skilled players have a higher drift rate, that is, these players process chess-specific information more quickly than less skilled players.

Panels B and C of Figure 4 show the relationship between the posterior means for the β_i and α_i , and the behavioural results for the test items. As can be seen in panel A, the posterior estimates of β tended to increase with mean RT and the posterior estimates of $\tilde{\alpha}$ showed a near perfect, slightly nonlinear relationship with mean RT. A polynomial regression model of degree three with the logarithm of the mean RT as dependent variable and β_i and $\tilde{\alpha}_i$ as predictors yielded a multiple correlation coefficient of $R^2 = .85$ and $R^2 = .99$, respectively. Panel C shows a similar pattern for the mean accuracy; mean accuracy had a near perfect, slightly nonlinear relationship

Table 2

Convergence results for ACT data.

Parameter	$\hat{R}$		
μ_P	1.0192		
	Quantiles		
	0.25	0.5	0.75
β_i	1.0002	1.0003	1.0005
θ_p	1.0010	1.0013	1.0017
α_i	<1.0001	1.0002	1.0002

Note. Quantiles are computed across persons or items.

with the posterior estimates of β and mean accuracy tended to increase for larger values of $\tilde{\alpha}$. A polynomial regression model of degree three with the arc sine of the mean accuracy as dependent variable and β_i and α_i as predictors yielded a correlation coefficient of $R^2 = .99$ and $R^2 = .77$, respectively. The cognitive process interpretation of these results is that differences in mean RT between items were largely due to differences in the volatility of the accumulated information, whilst differences in mean accuracy between items were due to differences in drift rate.

Panel C of Figure 4 shows three exemplary MCMC chains from our model fit. All chains show overall good mixing behaviour. There are some slow drifts visible in the chain for $\tilde{\alpha}_1$. Nevertheless, the values of the potential scale reduction factor, shown in Table 2, indicate excellent convergence, despite the large number of model parameters and the relatively small number of MCMC samples.

Extensions

Our development of the sDM so far has focused on a Bayesian implementation of the core model. In this section we will discuss the scope for extension of our core model.

Figure 4: Results of application of our sDM to validation data for the Amsterdam Chess Test (ACT; Van der Maas & Wagenmakers, 2005). The sample included data from 256 persons answering 40 items. (A) Relationship between mean RT and ELO rating (left plot; correlation is between ELO rating and logarithm of RT), between mean accuracy and ELO rating (middle plot; correlation is between ELO rating and arc sine of accuracy), and between posterior mean of θ and ELO rating (right plot). (B) Relationship between mean RT and posterior mean of β (left plot) and between mean RT and posterior mean of $\tilde{\alpha}$ (right plot). (B) Relationship between mean accuracy and posterior mean of β (left plot) and between mean accuracy and posterior mean of $\tilde{\alpha}$ (right plot). (C) Example MCMC chains for item and person effects after a burn-in period of 2000 samples.

Non-Decision Time

The full DM given in Equation (1) decomposes RT into a decision time component and a non-decision time component. The effect of the non-decision component is to shift the leading edge of the marginal RT distribution away from 0. In our development of the sDM we assumed the non-decision time to be 0. This assumption seems justified on conceptual grounds. The decision time component accounts for the time-intensive cognitive processes related to the decision problem at hand whereas the non-decision time component accounts for quicker cognitive processes not related to the decision problem, such as the time taken to read a question. In psychometric applications response times are typically in the range of several seconds and the contribution non-decision component is often negligible. Nevertheless, if researchers are interested in separating decision and non-decision components of RTs, this can be accomplished in several ways. Firstly, researchers could determine the shortest response time for each test-taker and subtract that response time from all other response times, thus shifting the leading edge of the marginal RT distribution for each test-taker to 0. Secondly, researchers could estimate non-decision time directly. To this end, non-decision time can be assigned a truncated normal prior distribution. This yields a truncated normal distribution for the full-conditional posterior distribution, which can easily be sampled from. Hence, estimation of the non-decision component is straightforward within the Gibbs sampling framework we used for the remaining model parameters. For other choices of the prior distribution, such as the uniform distribution often used in applications in mathematical psychology (e.g., Turner, van Maanen, & Forstmann, 2015), the full-conditional posterior distribution is no longer conjugate to the prior distribution. In this case posterior sampling can be implemented using the Adaptive Rejection Sampling (ARS) algorithm (Gilks & Wild, 1992).

Estimating the ratio of diffusion coefficient and boundary separation

Throughout the development of our core model, we assumed a fixed ratio $c = 2$ of the diffusion coefficient to boundary separation. Although the assumption of a fixed ratio is the key to our simplification of the diffusion likelihood in Equation (4), the value of this constant itself

might be of interest in some applications. Researchers might, for instance, be interested in testing whether the ratio is higher in one subpopulation of test-takers than in another, that is, whether one subpopulation of test-takers has a higher response caution when answering items than another. Questions of this kind can be addressed by specifying two versions of the sDM with different values for c . Both models can then be applied to the data of both subpopulations and standard methods for model comparison, such as AIC (Akaike, 1973), BIC (Schwarz, 1978), or Bayes factors (Jeffreys, 1961; Kass & Raftery, 1995), can be used to assess the relative fit of both models in each subpopulation.

Regression Models for Person and Item Effects

In psychometric applications, and in the context of educational testing in particular, researchers are often interested in the relationship between test-takers' latent traits and external covariates such as age, education, or other test scores. This is typically accomplished by regressing the latent variables on the covariates of interest. We will illustrate how such a regression model can be built into our sDM with a simple example where one latent variable is regressed on a set of manifest covariates. A more complete discussion of latent regression models can be found in Fox and Glas (2001) and Fox (2006).

Due to the functional form of our sDM, the addition of a regression extension is straightforward. The random effects structure for the latent variable of interest can be replaced by a regression equation. For an appropriately chosen prior distribution on the regression coefficients and residual variance, this yields conjugate normal-inverse gamma posterior distributions for the regression coefficients and residual variance, and leaves the normal-normal conjugacy for the latent variable itself intact.

Assume, for instance, that we are interested in regressing the person effect on drift rate, θ_p , on a set of covariates. We specify the regression model as:

$$\theta_p = \lambda_0 + \lambda_1 Y_{p,1} + \dots + \lambda_K Y_{p,K} + \varepsilon_i, \quad \varepsilon_i \sim \mathcal{N}(0, \sigma_{res}^2), \quad (6)$$

where $Y_{p,1}, \dots, Y_{p,K}$ are the manifest covariate values for person p , $\lambda_0, \dots, \lambda_K$ are the regression

weights, and ε_i is a normally distributed error term with mean 0 and variance σ_{res}^2 . If we assign the non-informative prior distribution $[\vec{\lambda}, \sigma_{res}^2] \propto 1/\sigma_{res}^2$ and replace the random effects term for θ_p by the regression model, the kernel of our model that involves θ_p becomes:

$$\begin{aligned} \mathbb{P}(\vec{t}, \vec{x} \mid \vec{\theta}, \vec{\lambda}, \sigma_{res}^2) \mathbb{P}(\vec{\lambda}) \mathbb{P}(\sigma_{res}^2) &\propto \exp \left(- \sum_{p=1}^{N_p} \theta_p^2 \left(\sum_{i=1}^{N_I} \frac{t_{ip} \tilde{\alpha}_i^2}{2} + \frac{1}{2\sigma_{res}^2} \right) \right) \\ &\cdot \exp \left(2 \sum_{p=1}^{N_p} \theta_p \left(\sum_{i=1}^{N_I} \tilde{\alpha}_i \frac{[x_{ip} - 1/2]}{2c} - \sum_{i=1}^{N_I} t_{ip} \tilde{\alpha}_i^2 \frac{\beta_i}{2} + \frac{(\lambda_0 + \lambda_1 Y_1 + \dots + \lambda_K Y_K)}{2\sigma_{res}^2} \right) \right) \\ &\cdot (\sigma_{res}^2)^{-(N_p/2+1)}. \end{aligned}$$

Here, $\vec{\theta}$ denotes the $N_p \times 1$ vector of person effects on drift rate, θ_p , and $\vec{\lambda}$ denotes the $K \times 1$ vector of regression weights. Note that replacing the expression for the random effect on θ_p by the regression model replaces the variance term σ_p^2 by the residual variance σ_{res}^2 . As before, this extended model is only identified up to a multiplicative constant, which we address by fixing $\sigma_{res}^2 = 1$ (instead of the constraint $\sigma_p^2 = 1$). From the expression above it can easily be seen that the full conjugate posterior distributions for $\vec{\lambda}$, σ_{res}^2 , and θ_p have the form indicated above (for details on the derivation, see chapter 14 in Gelman et al., 2013):

$$\vec{\lambda} \sim \mathcal{N}(\hat{\lambda}, V_{\lambda} \sigma_{res}^2), \quad \sigma_{res}^2 \sim \text{invG}(\frac{n}{2}, \frac{n-K}{2} \hat{\sigma}_{res}^2), \quad \theta_p \sim \mathcal{N}(\hat{\theta}_{p, N_I}, \hat{\sigma}_{p, N_I}^2),$$

where

$$\begin{aligned} \hat{\lambda} &= (Y^T Y)^{-1} X \theta, & V_{\lambda} &= (Y^T Y)^{-1}, & \hat{\sigma}_{res}^2 &= 1/(n-K)(\theta - Y\lambda)^T(\theta - Y\lambda) \\ \hat{\sigma}_{p, N_I}^2 &= \left(\sum_{i=1}^{N_I} \frac{t_{ip}}{\alpha_i^2} + \frac{1}{\sigma_{res}^2} \right)^{-1}, & \hat{\theta}_{p, N_I} &= \left(\sum_{i=1}^{N_I} \frac{(x_{ip}-1/2)}{c\alpha_i} - \sum_{i=1}^{N_I} \frac{t_{ip}\beta_i}{\alpha_i^2} + \frac{\lambda_0 + \lambda_1 Y_{p,1} + \dots + \lambda_K Y_{p,K}}{\sigma_{res}^2} \right) \hat{\sigma}_{p, N_I}^2. \end{aligned}$$

Here, Y is the $N_p \times (K+1)$ design matrix with all entries of the first column equal to 1 and the remaining columns the K predictors $Y_1, \dots, Y_K$. Hence, MCMC samples for latent regression sDM can be easily obtained using Gibbs-sampling.

As an illustrative example, we applied the regression extension for our sDM to the ACT data. We used the test-takers' age (Age), ELO-rating (ELO), and whether or not they had prior experience using a computer (CompExp) as predictors. We excluded data from 47 test-takers from this analysis as these persons either failed to respond to some test items, did not indicate

whether they had computer experience, or had no ELO-rating. We z-standardised all predictor variables before entering them into the model so that the estimated regression weights λ can be directly compared across predictors. To obtain MCMC-samples for the regression weights and sDM model parameters, we implemented a Gibbs-sampler using custom-written R code in combination with algorithms for sampling from truncated normal (truncnorm R package; Mersmann et al., 2018), multivariate normal (MASS R package; Ripley et al., 2018), and inverse gamma distributions (invgamma R package; Kahle, 2017). For each model parameter we generated three MCMC chains with 12,000 samples, of which we discarded the first 2,000 samples as burn in. To identify the model, we set $\sigma_{res}^2 = 1$ and used the settings $\mu_I = 0$, $\sigma_I^2 = 1$, and $c = 2$ as before.

The top row of Figure 5 shows the relationship between the three predictors and the estimates of θ_p obtained in our application of the sDM above. The leftmost plot suggests a negative relationship between drift rate and age, which means that older test-takers should tend to process chess-related items more slowly than younger test takers. The middle plot suggests a positive relationship between ELO-rating and drift rate, which means that more highly skilled players process chess-related information more quickly. Finally, in the rightmost plot there is no clear relationship between computer experience and θ_p , which means that test-takers' processing speed is independent of their experience using computers. The plots in the bottom row of Figure 5 show the distribution of posterior samples for the three regression weights, λ_{Age} , λ_{ELO} , and $\lambda_{CompExp}$. As can be seen in the leftmost plot, a large proportion of the posterior mass for λ_{Age} lay left of 0 and the posterior mean was -0.11, which confirms that there was a slight negative relationship between drift rate and age. The middle plot shows that most posterior mass for λ_{ELO} lay right of 0 and the posterior mean 0.24 indicates a comparatively strong positive relationship between drift rate and ELO-rating. Finally, the rightmost plot shows that the posterior mass for $\lambda_{CompExp}$ was symmetrically distributed around 0 and the posterior mean -0.01 confirms that there was no relationship between computer experience and drift rate.

Figure 5: Regression of the person effects on drift rate, θ , on different predictors from the Amsterdam Chess Test sample (ACT; Van der Maas & Wagenmakers, 2005). The top row shows the relationship between the estimated drift rate effects from the model without latent regression component, and test-takers' age (left panel), ELO rating (middle panel), and computer experience (right panel; data points are jittered for improved visibility). The bottom row shows the distribution of posterior samples for the standardised regression weights for the three predictors. Dashed grey lines indicate a regression weight of 0.

Discussion

In the present paper we developed a fully tractable latent trait diffusion model. The key to achieving tractability was the observation that if the ratio of boundary separation to the diffusion coefficient is constant, the DM likelihood is proportional to a normal density for each model parameter. This allowed us to build a Bayesian implementation of the DM with a random effects structure for items and persons. Our model is straightforward to analyse using Gibbs sampling and is easy to extend. Moreover, a simulation study demonstrated the superior computational efficiency compared to a standard implementation of the DM.

Our sDM combines the efficiency of a psychometric model with the favourable properties of a cognitive model. As the DM originated from mathematical psychology, there are well-established interpretations of the model parameters in terms of cognitive processes (e.g., Forstmann, Ratcliff, & Wagenmakers, 2016; Ratcliff & McKoon, 2008). Moreover, the model's functional form establishes a direct link between latent traits on the one hand, and RT and response accuracy on the other. At the same time, the DM's specialised development for experimental applications means that there are some limitations that might need to be addressed to make the model applicable in a wider psychometric context. Firstly, the DM was developed to account for RTs in the range below two or three seconds (Ratcliff & McKoon, 2008) whereas RTs in psychometric applications are typically in the range of several seconds. One way to address this problem is to scale observed RTs to the appropriate range. Nevertheless, the good performance of our model in the application example suggest that even without such rescaling, the sDM might provide an appropriate account of psychometric RT data. Secondly, the DM was developed for two-alternative forced choice settings, where each decision threshold corresponds to one response option and the sign of the drift rate parameter indicates which response option is favoured by the accumulating evidence. In a multiple-choice setting, there is no such clear correspondence between the response options, and the two decision thresholds and the sign of the drift rate parameter. If there is only one correct option and the incorrect options can be assumed to be equally attractive, they might be subsumed under one threshold and the correct response might be associated with the

other threshold (Van der Maas et al., 2011). However, in a setting with multiple correct answers, the DM will generally not be applicable (e.g., Ratcliff, Smith, Brown, & McKoon, 2016).

The development of latent trait models for RT data always involves a trade-off between statistical convenience and substantive scope, as highlighted by our discussion of the strengths and limitations of our sDM, and the question arises how this trade-off should be made. At one end of the spectrum are highly tractable statistical models that are void of any conceptual commitments. However, the validity of a psychometric test hinges on the psychological processes that connect the item presentation and item response. It therefore seems paramount to construct psychological theories of item response (Van der Maas et al., 2011). At the other of the spectrum are models from mathematical psychology that are conceptually rich but cannot be applied to the large data sets typically found in psychometric settings. The standard approach to making cognitive models applicable in psychometric settings is to systematically remove conceptual commitments that are embodied in the model's functional form until the resulting model is sufficiently tractable for psychometric applications. In recent years, several authors have suggested such simplified cognitive models for psychometric applications. In the remainder of this section we will discuss how the trade-off between statistical convenience and substantive scope in our sDM compares to the trade-offs made in these models.

Our development of the sDM here is one of several attempts at making the DM applicable in psychometric contexts. As such, our model shares many of the advantages of earlier developments but at the same time overcomes some of the challenges other models face. Vandekerckhove, Verheyen, and Tuerlinckx (2010) developed a hierarchical Bayesian implementation of the four-parameter DM. Their hierarchical diffusion model (HDM) affords users a high level of flexibility in implementing random effects and regression components, as does our sDM. Moreover, as HDM estimates the four core parameters of the DM it provides a substantive account for many standard phenomena in experimental applications, such as speed-accuracy trade-offs and response biases. However, as shown in our simulations, this broad scope comes at the cost of considerable computational inefficiency, which limits the applicability of the model in psychome-

tric settings.

Wagenmakers et al. (2007) presented a simplified version of the DM, the EZ model, which assumes the starting point of the diffusion process to be equidistant from the two decision thresholds. This version of the DM has a similar substantive scope as our sDM because it explains RT and accuracy in terms of three substantively meaningful model parameters, non-decision time, boundary separation, and drift rate. Moreover, estimation of the model parameters is highly efficient as it relies on closed-form expressions for the model parameters in terms of the probability of a correct response and the mean and variance of correct RTs. At the same time, this reliance on the moments of the marginal likelihood functions for RT and accuracy severely limits the model's flexibility. The three model parameters can only be estimated from either the marginal RT distribution for each item or the marginal RT distribution for each test-taker, which precludes the estimation of separate person and item effects (see also Ratcliff, 2008). Moreover, the lack of a likelihood function means that the model cannot be implemented hierarchically or extended in any meaningful sense.

Another class of cognitive models that have been adapted for psychometric applications are ballistic accumulator models. These models replace the stochastic information accumulation process in the DM by a ballistic process and allow information to accumulate independently for each choice option, which yields more tractable models (e.g., Brown & Heathcote, 2008). One particular instantiation of a ballistic accumulator model, the log-normal race model (LNR; Heathcote & Love, 2012), has recently been adapted to account for RTs in attitudinal and personality tests with multiple response categories (Ranger & Kuhn, 2018). Moreover, Rouder et al.'s (2015) hierarchical Bayesian implementation of the LNR can account for RT data in ability tests with multiple response options. Their model can furthermore be extended to include autocorrelative and latent regression components, and is straightforward to analyse using Gibbs sampling. However, although the LNR model parameters have a cognitive process interpretation similar to the DM parameters, the LNR model lacks the key distinction between boundary separation and the rate of information accumulation. Thus, whilst the LNR model shares the statistical convenience

of our sDM and can readily be applied in settings with more than two choice options, the lack of a differentiation between boundary separation and the rate of information accumulation represents a considerable limitation of the substantive scope of the LNR.

Taken together, our sDM combines a high level of computational efficiency and statistical flexibility with a conceptually meaningful interpretation of its model parameters. It thus outperforms earlier attempts to make the DM applicable in psychometric contexts. Within the larger class of accumulator models of decision-making, only the LNR model provides a similar level of statistical convenience but has a more limited substantive scope.

References

- Akaike, H. (1973). Information theory and an extension of the maximum likelihood principle. In B. N. Petrov & S. Caski (Eds.), *Proceedings of the second international symposium on information theory*. (pp. 267–281). Akademiai Kiado, Budapest, Hungary.
- Britten, K. H., Shadlen, M. N., Newsome, W. T., & Movshon, A. J. (1992). The analysis of visual motion: A comparison of neuronal and psychophysical performance. *Journal of Neuroscience*, 12(12), 4745–4765. doi:10.1523/JNEUROSCI.12-12-04745.1992
- Brown, S. D., & Heathcote, A. (2008). The simplest complete model of choice response time: linear ballistic accumulation. *Cognitive Psychology*, 57(3), 153–178. doi:10.1016/j.cogpsych.2007.12.002
- Carpenter, B., Gelman, A., Hoffman, M., Lee, D., Goodrich, B., Betancourt, M., ... Riddell, A. (2017). Stan: A probabilistic programming language. *Journal of Statistical Software*, 76(1). doi:10.18637/jss.v076.i01
- Cox, D. R., & Miller, H. D. (1970). *The theory of stochastic processes*. London, England: Chapman & Hall/CRC Press.
- Deneve, S. (2012). Making decisions with unknown sensory reliability. *Frontiers in Neuroscience*, 6. doi:10.3389/fnins.2012.00075
- Ditterich, J. (2006). Evidence for time-variant decision making. *The European Journal of Neuroscience*, 24(12), 3628–3641.
- Drugowitsch, J., Moreno-Bote, R., Churchland, A. K., Shadlen, M. N., & Pouget, A. (2012). The cost of accumulating evidence in perceptual decision making. *Journal of Neuroscience*, 32(11), 3612–3628.
- Forstmann, B. U., Ratcliff, R., & Wagenmakers, E. (2016). Sequential sampling models in cognitive neuroscience: Advantages, applications, and extensions. *Annual Review of Psychology*, 67, 641–666. doi:10.1146/annurev-psych-122414-033645
- Fox, J.-P. (2006). Fixed Effects IRT Model. *Behaviormetrika*, 33(1), 1–16.

- Fox, J.-P., & Glas, C. A. W. (2001). Bayesian estimation of a multilevel IRT model using Gibbs sampling. *Psychometrika*, 66(2), 271–288.
- Gelfand, A. E., & Smith, A. F. (1990). Sampling-based approaches to calculating marginal densities. *Journal of the American Statistical Association*, 85(410), 398–409. doi:10.1080/01621459.1990.10476213
- Gelman, A., Carlin, J. B., Stern, H. S., Dunson, D. B., Vehtari, A., & Rubin, D. B. (2013). *Bayesian Data Analysis* (3rd ed.). London: Chapman and Hall/ CRC.
- Gelman, A., & Rubin, D. B. (1992). Inference from iterative simulation using multiple sequences. *Statistical Science*, 7(4), 457–511. doi:10.1214/ss/1177011136
- Gilks, W. R., & Wild, P. (1992). Adaptive Rejection Sampling for Gibbs Sampling. *Journal of the Royal Statistical Society. Series C (Applied Statistics)*, 41(2), 337–348.
- Hanks, T. D., Mazurek, M. E., Kiani, R., Hopp, E., & Shadlen, M. N. (2011). Elapsed decision time affects the weighting of prior probability in a perceptual decision task. *Journal of Neuroscience*, 31(17), 6339–63352.
- Heathcote, A., & Love, J. (2012). Linear deterministic accumulator models of simple choice. *Frontiers in Psychology*, 3. doi:http://doi.org/10.3389/fpsyg.2012.00292
- Jeffreys, H. (1961). *Theory of probability*. Oxford: Oxford University Press.
- Kahle, D. (2017). invgamma: The inverse Gamma distribution (R package version 1.1) [Computer software]. Retrieved from <https://cran.r-project.org/web/packages/invgamma/index.html>
- Kass, R., & Raftery, A. (1995). Bayes factors. *Journal of the American Statistical Association*, 90, 773–795.
- Klinkenberg, S., Straatemeier, M., & Van Der Maas, H. L. (2011). Computer adaptive practice of Maths ability using a new item response model for on the fly ability and difficulty estimation. *Computers and Education*, 57(2), 1813–1824. doi:10.1016/j.compedu.2011.02.003
- Laming, D. R. J. (1973). *Mathematical psychology*. London, England: Academic Press.

- Maris, G., & van der Maas, H. (2012). Speed-Accuracy Response Models: Scoring Rules based on Response Time and Accuracy. *Psychometrika*, 77(4), 615–633. doi:10.1007/s11336-012-9288-y
- Mersmann, O., Trautmann, H., Steuer, D., & Bornkamp, B. (2018). truncnorm: Truncated normal distribution (R package version 1.0-8) [Computer software]. Retrieved from <https://cran.r-project.org/web/packages/truncnorm/index.html>
- Navarro, D. J., & Fuss, I. G. (2009). Fast and accurate calculations for cumulative first-passage time distributions in Wiener diffusion models. *Journal of Mathematical Psychology*, 53, 222–230. doi:10.1016/j.jmp.2012.09.002
- Ranger, J., & Kuhn, J. (2018). Modeling Responses and Response Times in Rating Scales With the Linear Ballistic Accumulator. *Methodology*, 14(3), 119–132. doi:10.1027/1614-2241/a000152
- Ratcliff, R. (1978). A theory of memory retrieval. *Psychological Review*, 85(2), 59–108.
- Ratcliff, R. (2008). The EZ-diffusion method: Too EZ? *Psychonomic Bulletin & Review*, 15(6), 1218–1228. doi:10.3758/PBR.15.6.1218
- Ratcliff, R., & McKoon, G. (2008). The diffusion decision model: theory and data for two-choice decision tasks. *Neural Computation*, 20(4), 873–922. doi:10.1162/neco.2008.12-06-420
- Ratcliff, R., Smith, P. L., Brown, S. D., & McKoon, G. (2016). Diffusion Decision Model: Current Issues and History. *Trends in Cognitive Sciences*, 20(4), 260–281. doi:10.1016/j.tics.2016.01.007
- Ripley, B., Venables, B., Bates, D. M., Hornik, K., Gebhardt, A., & Firth, D. (2018). MASS: Support functions and datasets for venables and Ripley's MASS(R package version 7.3-51.1) [Computer software]. Retrieved from <https://cran.r-project.org/web/packages/MASS/index.html>
- Rouder, J. N., Province, J. M., Morey, R. D., Gomez, P., & Heathcote, A. (2015). The Lognormal Race: A Cognitive-Process Model of Choice and Latency with Desirable Psychometric Properties. *Psychometrika*, 80(2), 491–513. doi:10.1007/s11336-013-9396-3

- Schwarz, G. (1978). Estimating the dimension of a model. *The Annals of Statistics*, 6(2), 461–464.
- Shadlen, M. N., & Kiani, R. (2013). Decision making as a window on cognition. *Neuron*, 80(3), 791–806.
- Singmann, H., Scott, B., Gretton, M., Heathcote, A., Voss, A., Voss, J., & Terry, A. (2016). rt-dists: Response time distributions (R package version 0.6-6) [Computer software]. Retrieved from <https://cran.r-project.org/web/packages/rtdists/index.html>
- Thura, D., Beauregard-Racine, J., Fradet, C.-W., & Cisek, P. (2012). Decision making by urgency gating: Theory and experimental support. *Journal of Neurophysiology*, 108(11), 2912–2930.
- Tuerlinckx, F., Molenaar, D., & van der Maas, H. (2016). Diffusion-based item response modeling. In W. van der Linden (Ed.), *Handbook of modern item response theory* (Vol. 1, pp. 283–300). Chapman and Hall/CRC Press.
- Tuerlinckx, F., & De Boeck, P. (2005). Two interpretations of the discrimination parameter. *Psychometrika*, 70(4), 629–650. doi:10.1007/s11336-000-0810-3
- Turner, B. M., van Maanen, L., & Forstmann, B. U. (2015). Informing cognitive abstractions through neuroimaging: The neural drift diffusion model. *Psychological Review*, 122(2), 312–336. doi:10.1037/a0038894
- Van Breukelen, G. J. P. (2005). Psychometric modeling of response speed and accuracy with mixed and conditional regression. *Psychometrika*, 70(2), 359–376. doi:10.1007/s11336-003-1078-0
- Van der Linden, W. J. (2006). A Lognormal Model for Response Times on Test Items. *Journal of Educational and Behavioral Statistics*, 31(2), 181–204. doi:10.3102/10769986031002181
- Van der Linden, W. J. (2007). A Hierarchical Framework For Modeling Speed and Accuracy on Test Items. *Psychometrika*, 72(3), 287–308.
- Van der Linden, W. J. (2009). Conceptual Issues in Response-Time Modeling. *Journal of Educational Measurement*, 46(3), 247–272.

- Van der Maas, H. L. J., Molenaar, D., Maris, G., Kievit, R. A., & Borsboom, D. (2011). Cognitive Psychology Meets Psychometric Theory: On the Relation Between Process Models for Decision Making and Latent Variable Models for Individual Differences. *Psychological Review*, 118(2), 339–356. doi:10.1037/a0022749
- Van der Maas, H. L. J., & Wagenmakers, E.-J. (2005). A psychometric analysis of chess expertise. *American Journal of Psychology*, 118(1), 29–60.
- Van Rijn, P. W., & Ali, U. S. (2017). A comparison of item response models for accuracy and speed of item responses with applications to adaptive testing. *British Journal of Mathematical and Statistical Psychology*, 70(2), 317–345. doi:10.1111/bmsp.12101
- Vandekerckhove, J., Tuerlinckx, F., & Lee, M. D. (2011). Hierarchical Diffusion Models for Two-Choice Response Times. *Psychological Methods*, 16(1), 44–62. doi:10.1037/a0021765
- Vandekerckhove, J., Verheyen, S., & Tuerlinckx, F. (2010). A crossed random effects diffusion model for speeded semantic categorization decisions. *Acta Psychologica*, 133(3), 269–282. doi:10.1016/j.actpsy.2009.10.009
- Wabersich, D., & Vanderkerckhove, J. (2014). Extending JAGS: A tutorial on adding custom distributions to JAGS (with a diffusion model example). *Behavior Research Methods*, 46, 15–28. doi:10.1016/j.cognition.2008.05.007
- Wagenmakers, E. J., Van der Maas, H. L. J., & Grasman, R. P. P. P. (2007). An EZ-diffusion model for response time and accuracy. *Psychonomic Bulletin & Review*, 14(1), 3–22. doi:10.3758/BF03194023
- Wagenmakers, E.-J., Ratcliff, R., Gomez, P., & McKoon, G. (2008). A diffusion model account of criterion shifts in the lexical decision task. *Journal of Memory and Language*, 58, 140–159. doi:10.1016/j.jml.2007.04.006
- Yap, M. J., Balota, D. A., Sibley, D. E., & Ratcliff, R. (2012). Individual differences in visual word recognition: Insights from the English lexicon project. *Journal of Experimental Psychology: Human Perception and Performance*, 38(1), 53–79. doi:10.1037/a0024177

Yap, M. J., Sibley, D. E., Balota, D. a., Ratcliff, R., & Rueckl, J. (2015). Responding to nonwords in the lexical decision task: Insights from the English Lexicon Project. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, 41(3), 597–613. doi:10.1037/xlm0000064