High Vagal Tone and Rapid Extinction Learning as Potential Transdiagnostic Protective Factors

Following Childhood Violence Exposure

Eli S. Susman^{1,*}

David G. Weissman PhD^{1,*}

Margaret A. Sheridan PhD²

Katie A. McLaughlin PhD¹

¹Department of Psychology, Harvard University, Cambridge, MA

²Department of Psychology and Neuroscience, University of North Carolina at Chapel Hill,
Chapel Hill, NC

*These authors contributed equally to the work.

Corresponding Author:

Eli S. Susman, William James Hall, 33 Kirkland St. Cambridge, MA 02138.

esusman@fas.harvard.edu

Acknowledgements

This research was funded by the National Institute of Mental Health (R01-MH103291 to McLaughlin), an Early Career Research Fellowship from the Jacobs Foundation (McLaughlin), and a OneMind Institute Rising Star Award (McLaughlin). We are tremendously grateful to Debbie Bitran, Andrea Duys, and Azure Reid-Russell for help with participant recruitment and testing. The authors have no conflicts of interest to declare.

Data Availability Statement

The data that support the findings of this study are openly available in the Open Science Framework at <https://doi.org/10.17605/OSF.IO/XUR87> (Susman, Weissman, & McLaughlin, 2020).

Abstract

Childhood exposure to violence is strongly associated with psychopathology. High resting respiratory sinus arrhythmia (RSA) is associated with lower levels of psychopathology in children exposed to violence. High RSA may help to protect against psychopathology by facilitating fear extinction learning, allowing more flexible autonomic responses to learned threat and safety cues. In the present study, 165 youth (79 female, ages 9-17; 86 exposed to violence) completed assessments of violence exposure, RSA, and psychopathology, and a fear extinction learning task; 134 participants returned and completed psychopathology assessments two years later. Resting RSA moderated the longitudinal association of violence exposure with PTSD symptoms and externalizing psychopathology, such that the association was weaker among youths with higher RSA. Higher skin conductance responses (SCR) during extinction learning to the threat cue (CS+) was associated with higher internalizing symptoms at follow-up and greater SCR to the safety cue (CS-) was associated with higher PTSD, internalizing, and externalizing symptoms, as well as the p-factor, controlling for baseline symptoms. Findings suggest that higher RSA may protect against emergence of psychopathology among children exposed to violence. Moreover, difficulty extinguishing learned threat responses and elevated autonomic responses to safety cues may be associated with risk for future psychopathology.

Keywords: psychopathology, adolescent, respiratory sinus arrhythmia, aversive learning, childhood trauma

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

Childhood violence exposure—such as physical abuse, sexual abuse, or witnessing domestic violence—is one of the most pressing public health crises in the world (Kolk, 2017), and explains a substantial portion of mental health problems in the population (Green et al., 2010a; Kessler et al., 2010; McLaughlin et al., 2012). Violence exposure is associated with increased risk for many forms of psychopathology across the internalizing and externalizing spectrum (Cicchetti & Toth, 2005; Green et al., 2010; Kessler et al., 2010; McGrath et al., 2017; McLaughlin et al., 2012; Schaefer et al., 2018), as well as with the transdiagnostic general psychopathology factor (i.e., p-factor) (Caspi et al., 2014; Schaefer et al., 2018; Weissman, Guyer et al. 2019). Identifying protective factors that buffer against the onset of psychopathology following childhood exposure to violence is critical to developing more targeted and effective interventions (Duffy et al., 2019).

High vagal tone may be one such protective factor. Vagal tone reflects parasympathetic nervous system (PNS) regulation of heart rate through the vagus nerve (Berntson et al., 1997; Porges, 2007). The vagus nerve acts as a brake on the firing of the sinoatrial node, the heart's primary pacemaker, following activation of vagal motor neurons in the nucleus ambiguus and the dorsal motor nucleus (Allen et al., 2007; Porges, 1995, 2007; Thayer et al., 2009). Respiratory sinus arrhythmia (RSA) is a noninvasive measure that reflects vagal tone as variability in heart rate within the frequency band associated with respiration (Allen et al., 2007; Berntson et al., 1993). High vagal tone, indicated by higher levels of RSA at rest, has been shown to moderate the associations of exposure to stress, adversity and violence with psychopathology, such that the link between adversity and psychopathology is weaker or absent in children with high vagal tone (El-Sheikh et al., 2001; Katz & Gottman, 1997; Katz & Gottman, 1995; McLaughlin, Alves, et al., 2014; McLaughlin et al., 2015; Porges, 2007). The specific mechanisms through which high RSA

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

may help to protect against psychopathology following childhood exposure to violence and adversity are unknown. However, one such mechanism could be the way in which vagal tone influences associative learning processes involved in fear extinction.

Extinction learning arises when new associations are formed that compete with those acquired through prior aversive learning (Bouton, 1993; Bouton, 2004; Milad & Quirk, 2012). Aversive learning, or fear learning, occurs when a previously neutral stimulus becomes associated with threat due to repeated pairings with an aversive unconditioned stimulus (US), such as a shock or loud noise. Extinction learning occurs when the conditioned stimulus (CS+), that had previously been paired with the aversive US, is repeatedly presented without the US. During fear extinction learning, repeated presentation of the CS+ in the absence of the US creates a new memory trace associating the CS+ with safety that competes with the original conditioned fear memory (Bouton, 1993; Bouton, 2004; Milad & Quirk, 2012).

Some studies suggest that vagal activity may moderate fear extinction. For example, vagal nerve stimulation has been shown to reduce behavioral indicators of fear (e.g. freezing) responses to the CS+ in animal models (Peña et al., 2013). Alternatively, vagal tone may index central nervous system mechanisms that underlie both vagal activity and fear extinction. Vagal tone has been hypothesized to reflect greater tonic inhibition of the amygdala by the medial prefrontal cortex (mPFC) (Thayer et al., 2012), and mPFC-amygdala coupling has been shown to play an integral role in extinction learning (Giustino & Maren, 2015; Kim et al., 2011; Phelps et al., 2004). Because the inhibitory influence of the mPFC on the amygdala can contribute to both higher vagal tone and the extinction of learned fear responses, higher vagal tone may index a greater capacity for extinguishing learned fear associations (Beauchaine, 2015; Jennings et al., 2016; Mather & Thayer, 2018; Porges, 2007; Sakaki et al., 2016; Thayer et al., 2012).

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

The neural circuits involved in fear extinction learning undergo considerable maturation in middle childhood and adolescence (Pattwell et al., 2012). In older adolescents and adults, extinction learning occurs more quickly and recruitment of the mPFC occurs to a greater degree than for children, who recruit the amygdala more strongly for threat and safety learning than adolescents and adults (Lau et al., 2011; Morriss et al., 2019). This developmental shift may be mediated by increasing structural integrity of the uncinate fasciculus—a white matter tract connecting the mPFC and amygdala (Morriss et al., 2019). Lower structural integrity of the uncinate fasciculus in adolescence is associated both with childhood maltreatment and increased internalizing symptoms following exposure to stressful life events (Hanson et al., 2015). Likewise, altered amygdala-vmPFC functional connectivity during fear extinction learning is associated with increased risk for anxiety disorders (Gold et al., 2016). Moreover, altered extinction learning is associated with elevated risk for numerous forms of internalizing psychopathology in adults (Craske et al., 2011; Garcia & Zoellner, 2017; Gazendam et al., 2013; Grillon & Morgan III, 1999; Guthrie & Bryant, 2006; Lissek et al., 2005; Norrholm et al., 2011; Staples-Bradley et al., 2018), as well as children (Britton et al., 2013; Craske et al., 2008; Dvir et al., 2019; Shechner et al., 2015).

Given that smaller physiological responses to the CS+ during extinction learning is associated with decreased risk for PTSD (Blechert et al., 2007; Norrholm et al., 2011; Orr et al., 2000; Peri et al., 2000; Zuj, Palmer, Hsu et al., 2016; Zuj, Palmer, Lommen et al., 2016), and other forms of psychopathology (Craske et al., 2008; McGuire et al., 2016; Michael et al., 2007), higher vagal tone may protect against transdiagnostic psychopathology through associated reductions in fear responses during extinction learning. Indeed, higher resting RSA moderated the association between violence exposure and PTSD avoidance symptoms in children and

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

adolescents in a prior study, such that violence exposure was only associated with elevated symptoms among youths with low resting RSA (Jenness et al., 2019). Higher resting RSA also moderated the association between violence exposure and skin conductance responses (SCR) during early fear extinction, such that, among violence-exposed youths, higher RSA was associated with lower SCR responses to the CS+ during early extinction learning, indicating greater extinction of learned fear. Finally, this pattern of lower SCR during early fear extinction mediated the association between higher RSA and lower PTSD symptoms among youth exposed to violence, suggesting that the protective influence of high vagal tone against PTSD symptoms among youths exposed to violence may operate, at least partially, through enhanced extinction learning (Jenness et al., 2019).

The present study seeks to replicate and extend these findings in a large, longitudinal sample of children and adolescents with high levels of exposure to violence. We evaluate the extent to which vagal tone and fear extinction might promote resilience to trauma-related psychopathology. We first attempt to replicate the findings of Jenness et al. (2019) by conducting analyses in relation to PTSD, a form of psychopathology that is common among children exposed to violence and associated with alterations in extinction learning mechanisms (Milad & Quirk, 2012). We further examined whether vagal tone and fear extinction act as transdiagnostic mechanisms of resilience by evaluating their associations with internalizing, externalizing, and a measure of general psychopathology (i.e. p-factor). In preregistered analyses (<https://osf.io/xur87/>). We examine several hypotheses. First, we expected that violence exposure would be positively associated with PTSD, internalizing problems, externalizing problems, and p-factor. Second, we predicted that resting RSA would moderate the association between violence exposure and PTSD, internalizing problems, and p-factor, such that violence exposure

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

would be associated with greater symptoms of psychopathology among youth with low resting RSA, but not among those with high RSA. Third, we hypothesized that larger skin conductance responses (SCR) to the CS+ during early extinction learning would be associated with higher PTSD symptoms, internalizing problems, or p-factor. Fourth, we predicted that resting RSA would moderate the association between violence exposure and SCR to the CS+ during early extinction learning, such that violence exposure would be associated with higher SCR reactivity to the CS+, among youth with low resting RSA, but not among those with high resting RSA. Finally, we predicted indirect effects of violence exposure on PTSD symptoms, internalizing problems, and p-factor via higher SCR reactivity to the CS+ during early extinction, conditional on resting RSA, such that higher SCR reactivity to the CS+ would mediate the association between violence exposure and symptoms of transdiagnostic psychopathology among youths with low resting RSA, but not among youths with high resting RSA. In all analyses of psychopathology, we first examined associations cross-sectionally, and then longitudinally in predicting symptom changes over a two-year follow-up period.

Methods

Participants

Participants were children and adolescents between the ages of 8 and 17 living in the Seattle area. A total of 262 youth aged 8-16 years were enrolled into the study. Youth and caregivers were recruited for participation at schools, after-school and prevention programs, adoption programs, food banks, shelters, parenting programs, medical clinics, and the general community in Seattle, WA between January 2015 and June 2017. Recruitment efforts were targeted at recruiting a sample with variation in exposure to maltreatment-related trauma. To do so, we recruited from neighborhoods with high levels of violent crime, clinics that served a predominantly low-SES catchment area, and agencies that work with families who have been victims of violence (e.g., domestic violence shelters, programs for parents mandated to receive intervention by Child Protective Services). Exposure to violence and other inclusion and exclusion criteria were assessed during the first study visit. Inclusion criteria for the violence-exposed group included exposure to physical or sexual abuse or direct witnessing of domestic violence. Children in the control group were matched to children in the violence-exposed group on age, sex, and handedness; inclusion criteria required an absence of exposure to maltreatment or other forms of significant interpersonal violence. Exclusion criteria included $IQ < 80$, presence of pervasive developmental disorder, active psychotic symptoms or mania, active substance abuse, and presence of safety concerns. Of the 262 children enrolled in the first study visit, three were excluded from all analysis due to low IQ ($n=1$), presence of pervasive developmental disorder ($n=1$), and presence of psychotic symptoms and drug abuse ($n=1$).

The data on RSA and fear learning were acquired during a second study visit, on a smaller sample of children and adolescents invited to participate in a more involved

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

neuroimaging and physiological assessment. A total of 169 participants participated in the resting electrocardiogram and fear learning tasks at baseline. Equipment malfunctions resulted in loss of all physiological data from four participants. Four participants declined to complete the fear extinction task, and 5 participants discontinued the task before the end of extinction phase. Eleven non-responders during the conditioning phase were dropped from all SCR analyses, as fear extinction learning cannot occur if fear conditioning has not first been established. The full analytic sample for this study consisted of 165 subjects (86 violence-exposed) at the baseline assessment.

A longitudinal follow-up assessment was conducted approximately two years following the baseline assessments ($M = 21.96$ months, $SD = 7.88$ months) to assess symptoms of psychopathology. A total of 134 subjects (67 violence-exposed) of the 165 participants with usable data on RSA and fear learning at baseline (79.3%) participated in this follow-up. Of the 31 children in the analytic sample that dropped out, 19 were maltreated (22% attrition), and 12 were not (15% attrition) ($\chi^2 = 0.87, p = .350$).

All procedures were approved by the Institutional Review Board at the University of Washington. Written informed consent was obtained from legal guardians; children provided written assent. Maltreatment not previously reported to the relevant authorities was reported to Child Protective Services using standard clinical procedures.

Fear Learning Task

We utilized a fear learning task validated for children and adolescents that included three phases: preconditioning, conditioning, and extinction (McLaughlin et al., 2016; Shechner et al., 2015). During the pre-acquisition phase, participants viewed the CS+ and CS- (blue and yellow bells, counterbalanced across participants, 4 trials each) without the US (an aversive 96 dB alarm

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

noise). The acquisition phase consisted of 20 trials of the CS+ and CS– (10 of each). The CS+ co-terminated with the US in 80% of acquisition trials. During the extinction phase, the CS+ and CS– (8 trials each) were presented without the US. Intervals between trials ranged between 8 to 12 s (mean = 10 s).

Skin conductance responses (SCR) provided the primary metric of fear conditioning. Consistent with Jenness et al. (2019), we focused specifically on responses to the CS+ during the early fear extinction phase of the task, as fear responses tend to return to baseline in the latter half of the extinction task (McLaughlin et al., 2016; Shechner et al., 2015). Moreover, the early extinction phase has been the main focus of studies on extinction learning, and is typically where participants exhibit the greatest variability in fear response intensity during the extinction phase (Milad et al., 2009; Milad & Quirk, 2002).

Measures

Violence Exposure.

As described in more detail previously (Weissman, Jenness, et al., 2019), participants exposed to physical abuse, sexual abuse, or domestic violence were categorized as violence-exposed. Otherwise, they were classified as controls. Children were classified as experiencing physical or sexual abuse if abuse was endorsed by the child on the Childhood Experiences of Care and Abuse (CECA) interview (Bifulco et al., 1994), PTSD-RI trauma screen, or above the validated Child Trauma Questionnaire (CTQ) threshold (Bernstein et al., 1997; Walker et al., 1999), or by parent on the Juvenile Victimization Questionnaire (JVQ) (Finkelhor et al., 2005), or PTSD-RI trauma screen (Steinberg et al., 2013). Exposure to domestic violence reported on the Violence Exposure Scale for Children-Revised (VEX-R) (Fox & Leavitt, 1995) or PTSD-RI

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

trauma screen was determined based on child report only. Consistency of reporting across these measures is summarized in supplemental materials table S2.

Vagal Tone.

Vagal tone was assessed using noninvasive Electrocardiogram (ECG) recordings, obtained with a Biopac ECG amplifier (Goleta, CA) according to accepted guidelines (Fox & Leavitt, 1995). Participants were instructed to sit quietly without moving for 5 minutes. We used a modified Lead II configuration, placing electrodes on the right clavicle, left lower torso, and right leg (ground). ECG data, sampled at 1.0 kHz, were collected using Biopac MP150 hardware and Acknowledge software. Trained raters, blinded to violence exposure of participants, visually inspected automatic R-peak detection of the ECG data using Mindware Heart Rate Variability (HRV) Software (Mindware Technologies, Gahannah, OH). Visual inspection focused on detecting ectopic beats and confirming accurate detection of R-peaks in the ECG waveform. From IBI time series, we calculated RSA in 1-minute bins. To calculate RSA, the HRV module detrended the data using a first order polynomial to remove the mean and any linear trends, cosine tapered the data, submitted it to Fast Fourier Transformation, and took the natural log integral of high frequency power in the appropriate respiratory frequency band based on participant age (0.24–0.40 Hz for 8-12 years of age, and 0.18–0.40 Hz for 13-16 years of age), (Shader et al., 2018).

Skin Conductance Response.

Skin conductance response (SCR) served as the primary measure of fear extinction learning. For each participant, an experimenter filled two silver/silver chloride electrodes with sodium chloride gel. The tips (fingerprint side) of the participant's non-dominant index and middle finger were exfoliated and wiped with rubbing alcohol, where an experimenter then

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

fastened the two electrodes. We acquired SCR using a Biopac galvanic skin response module (Goleta, CA), which collected skin conductance response at a sampling rate of 250 Hz throughout the fear conditioning task. Using AcqKnowledge 4.0 software (Biopac Systems, Goleta, CA), we calculated SCRs as the difference in amplitude between 1 second pre-CS baseline and peak response to CS+ in the 1–4 seconds following onset of the stimulus in accordance with standard practice (Dawson et al., 2017). The aforementioned 1-4 second response window is a standard procedure used to avoid collecting SCR unrelated to the CS+, and to capture only peak skin conductance response to CS+ (Dawson et al., 2017). In order to ensure that SCR changes during the conditioning phase did not result from the US, the response window terminates before onset of the US. During the conditioning phase of the task, responses were only counted if SCR exceeded 0.02 microsiemens (μs) averaged across trials. Eleven participants were non-responders during the conditioning phase of the task based on this threshold. Eight were from the violence-exposed group, and three were from the control group. These non-responders were removed from all SCR-related analyses as extinction learning cannot occur if fear conditioning has not first been established. All SCRs from the extinction phase were included in analyses, regardless of their magnitude. Because the residuals from our models using raw SCR amplitudes were skewed (all $S_s > 2.01$), we square root transformed the raw SCR amplitudes in order to reduce skew, which is a common approach in developmental fear conditioning studies (Abend et al., 2020; Gamwell et al., 2015; Jovanovic et al., 2014; Kreutzmann et al., 2021; McLaughlin et al., 2016; Morriss et al., 2019; Shechner et al., 2015; Waters & Pine, 2016).

Psychopathology Symptoms

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

Post-traumatic stress disorder (PTSD) symptoms. The PTSD-RI is a widely used assessment of PTSD symptom severity with strong convergent validity and internal consistency for the overall measure and among subscales (Jenness et al., 2019; Steinberg et al., 2013). The PTSD-RI measures re-experiencing, avoidance/numbing, and hyper-arousal, which are PTSD symptoms specifically related to childhood exposure to violence. The PTSD-RI has sound psychometric properties (Steinberg et al., 2013) and had excellent internal consistency in our sample ($\alpha = .94$). The higher of the parent and youth-reported PTSD symptom severity was used in our analyses, consistent with the standard “or” rule for determining the presence of psychopathology symptoms in children (Kessler et al., 2012; Merikangas et al., 2010). PTSD scores at both baseline and follow-up fall within assumptions of normality ($S=0.702$ and $S=1.212$ respectively), and more importantly the residuals of PTSD scores for every model in which PTSD is used fall within assumptions of normality ($0.191 < Ss < 0.590$) (Hair, 2009; Lumley et al., 2002).

Internalizing and Externalizing Psychopathology. Internalizing symptoms were assessed based on the PTSD-RI, the Children’s Depression Inventory 2 (CDI), and the Screen for Child Anxiety Related Emotional Disorders (SCARED). The CDI is one of the most widely used assessments for depression in the literature, with good test–retest reliability, internal consistency, and construct validity (Rivera et al., 2005; Sitarenios & Kovacs, 1999), particularly over repeated administrations (Finch Jr et al., 1987). The SCARED is a broadly used, validated instrument with good internal consistency, discriminant validity, and test-retest reliability (particularly between depressive and anxiety disorders) used to screen children for anxiety disorders (Birmaher et al., 1997; Birmaher et al., 1999; Hale et al., 2005).

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

Externalizing symptoms were assessed with the externalizing problems subscale of the CBCL, which is comprised of subscales assessing aggressive behaviors, rule breaking Behaviors, and attention problems. The CBCL is one of the most widely used validated measures of youth emotional and behavioral problems in both clinical and research settings, and demonstrates strong reliability, as well as convergent and discriminative validity (Achenbach et al., 2003; Ebesutani et al., 2010; Nakamura et al., 2009). The higher of the parent and youth-reported externalizing symptom severity was used in our analyses.

Transdiagnostic Psychopathology. As described in previous publications (Weissman, Bitran, et al., 2019; Weissman, Jenness, et al., 2019) we performed confirmatory factor analysis (CFA) to test a correlated-factors model specifying Internalizing and Externalizing latent factors. All CFA analyses were performed in MPlus version 8.1 (Muthén & Muthén, 2012). Given that our observed indicator variables were slightly skewed and kurtotic, we used the robust maximum likelihood estimator (MLR), which employs a sandwich estimator to arrive at standard errors robust to non-normality of observations. MLR performs well in modest sample sizes with skewed data, as in the present study (Li, 2016). We assessed the relative fit of each model using the Akaike Information Criterion (AIC), Bayesian Information Criterion (BIC) and the Sample Adjusted BIC. Fit indices for the correlated-factors model were: AIC = 6056.31, BIC = 6249.00, Sample adjusted BIC = 6077.79. Standardized factor loadings for the Internalizing (CDI, SCARED, PTSD) latent factor ranged from 0.65 to 0.73, all p 's < .001. Standardized factor loadings for the Externalizing (Aggressive Behaviors, Rule Breaking Behaviors, Attention Problem) latent factor ranged from 0.76 to 0.87, all p 's < .001. Fit indices for the correlated factors model at follow-up were: AIC = 4507.81, BIC = 4682.09, Sample Adjusted BIC = 4514.19. Standardized factor loadings for the Internalizing Factor ranged from 0.48 to 0.95, all

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

p 's $< .001$. Standardized factor loadings for the Externalizing Factor ranged from 0.76 to 0.84, all p 's $< .001$.

The general psychopathology factor (or p-factor) was estimated in a bi-factor model specifying both a General Psychopathology latent factor ("p") and residual internalizing and externalizing factors. Fit indices for the bi-factor model were: AIC = 6018.23, BIC = 6228.77, Sample adjusted BIC = 6041.71. Standardized factor loadings for the latent p-factor (CDI, SCARED, PTSD, Aggressive Behaviors, Rule Breaking Behaviors, Attention Problems) ranged from 0.44 to 0.80, all p 's $< .001$. Fit indices for the bi-factor model at the longitudinal follow-up were: AIC = 4481.00, BIC = 4665.15, Sample Adjusted BIC = 4487.74. Standardized factor loadings for the latent P factor ranged from 0.46 to 0.81, all p 's $< .001$. For more details, see supplemental materials (Table S1). As assessed by relative fit indices and factor loadings, both the correlated factor model and the bi-factor model fit the data similarly well at the baseline assessment, with a relatively better fit for the bi-factor model.

Data Analysis

Analyses were preregistered, and all data and analysis code are publicly available (osf.io/6hvuy). We deviated from the pre-registration in two instances. First, we incorporated externalizing problems as an additional psychopathology outcome based on a reviewer request. Second, we evaluated whether associations between SCR during early extinction learning and psychopathology were specific to the CS+ or reflect a more general hyperresponsivity, even to previously learned safety cues, by incorporating analyses of SCR to the CS- during extinction learning.

Analyses were conducted in R Version 3.6.2. All analyses controlled for age, sex, and family income-to-needs ratio, the latter of which differed for youths with and without exposure

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

to trauma. All analyses with psychopathology at follow-up as the outcome controlled for the corresponding measure of psychopathology at baseline.

To evaluate whether vagal tone moderated the association between violence exposure and psychopathology, we estimated linear regression models with violence exposure, RSA, and their interaction in predicting each of the relevant psychopathology outcomes. To examine whether vagal tone moderated the association between violence exposure and SCR to the CS+ during early extinction, we estimated linear regression models with violence exposure, RSA, and their interaction in predicting SCR to the CS+ during early extinction as an outcome. Because SCR was positively skewed, even after square root transformation, asymmetrical bootstrapped confidence intervals were estimated using the “boot” package in R (Canty & Ripley, 2019; Davison & Hinkley, 1997) based on 10,000 bootstrapped samples.

To evaluate whether higher SCR to the CS+ during early extinction was associated with psychopathology, we estimated linear regression models with early extinction SCR to the CS+ in predicting psychopathology outcomes as baseline and follow-up, controlling for violence exposure. To determine whether associations between SCR during early fear extinction learning and psychopathology were specific to the CS+, we also conducted analyses with SCR to the CS– as described above. Specifically, we estimated linear regression models with early extinction SCR to the CS– in predicting psychopathology symptoms at baseline and follow-up, controlling for violence exposure. We examined responses to the threat and safety cues separately because these reflect different psychological processes mediated by distinct neural networks (Fullana et al., 2016). Examining the CS+ at the beginning of extinction provides an index of the degree of responding to the threat cue when – if extinction has occurred – it is no longer associated with

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

that threat. This response is largely independent from responses to the safety cue that has never been associated with threat.

Missing income-to-needs ratio data for 12 participants were handled with multiple imputation, based on 100 imputations using the “mice” package (Buuren & Groothuis-Oudshoorn, 2011).

Results

Descriptive statistics

Demographic information and descriptive statistics for all primary variables for both the violence-exposed and control group can be found in Table 1. Further details on regression analyses can be found in Tables S6-S9 in the supplementary materials.

Violence Exposure and Psychopathology

We first investigated the association of violence exposure with baseline psychopathology. Violence exposure was positively associated with PTSD symptoms, ($\beta = .67, p < .001$), internalizing problems ($\beta = .53, p < 0.001$), externalizing problems ($\beta = 0.57, p < .001$), and the p-factor ($\beta = 0.62, p < .001$) at baseline. When accounting for symptoms at baseline, we did not find any association between violence exposure and PTSD symptoms ($\beta = .08, p = 0.468$), externalizing problems ($\beta = -.14, p = 0.194$), or the p-factor ($\beta = -.10, p = 0.301$) at two-year follow-up. However, we did find an association between violence exposure and fewer internalizing problems at two-year follow-up, when accounting for the higher symptoms at baseline among trauma-exposed youth ($\beta = -.20, p = 0.048$).

RSA, Violence Exposure, and Psychopathology

Resting RSA did not moderate the association between violence exposure and baseline symptoms of PTSD ($\beta = -0.11, p = .170$), internalizing problems ($\beta = -.09, p = .298$),

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

externalizing problems ($\beta = .04$, $p = .639$), or the p-factor ($\beta = .00$, $p = .955$). However, resting RSA did moderate the association between violence exposure and PTSD symptoms ($\beta = -0.20$, $p = .010$), and externalizing problems at two-year follow-up ($\beta = -0.20$, $p = .039$) (Figure 1). RSA did not moderate the association of violence exposure with either internalizing symptoms ($\beta = -0.09$, $p = .340$) or p-factor ($\beta = -.15$, $p = .078$) at two-year follow-up.

RSA, Violence Exposure, and Fear Extinction

Violence exposure was not associated with SCR during the early extinction phase ($B = -.01$, 95% CI = $-.037, .014$) and this association was not moderated by RSA ($B = -.01$, 95% CI = $-.028, .022$). Thus, we did not proceed with our pre-registered analyses examining extinction learning as a mediator of the link between violence exposure and PTSD symptoms, conditional on RSA. This model did, however, reveal a significant negative association between age and SCR to the CS+ during early extinction ($B = -.01$, 95% CI = $-.011, -.001$), indicating that fear extinction learning was greater as age increased. To evaluate the specificity of this association to the learned threat cue, we repeated this analysis with SCR to the CS- as the outcome. There was also a negative association between age and SCR to the CS- during early extinction ($B = -.01$, 95% CI = $-.013, -.001$), suggesting that physiological reactivity to the safety cue during early extinction also decreased with age.

Fear Extinction and Psychopathology

Higher SCR to the CS+ during early extinction was associated with higher internalizing symptoms ($\beta = 0.14$, $p = .042$), but not PTSD ($\beta = .08$, $p = .180$), externalizing problems ($\beta = 0.13$, $p = .068$) or p-factor scores ($\beta = .12$, $p = .053$) at two-year follow-up (Figure 2). In contrast, higher SCR to the CS- (i.e., safety cue) during early extinction was associated with higher PTSD symptoms ($\beta = .11$, $p = .0496$), internalizing symptoms ($\beta = .16$, $p = .013$), externalizing

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

problems ($\beta = .17$, $p = .016$), and p-factor ($\beta = .18$, $p = .004$) at the two-year follow-up (Figure 2).

Discussion

This longitudinal study investigated the extent to which vagal tone and autonomic responses during fear extinction learning act as transdiagnostic mechanisms of resilience to psychopathology following childhood exposure to violence. Vagal tone moderated the longitudinal association of violence exposure with PTSD symptoms and externalizing problems, such that the association of violence exposure with psychopathology was lower among youths with high vagal tone than among those with low vagal tone. Greater skin conductance responses to threat cues during fear extinction was associated with higher internalizing symptoms two-years later, whereas elevated skin conductance responses to safety cue during fear extinction was associated with higher symptoms of PTSD, internalizing, and externalizing psychopathology as well as p-factor scores two-years later. However, the protective influence of high vagal tone on the association between violence exposure and psychopathology was not explained by fear extinction learning in this sample.

Vagal tone moderated the longitudinal associations of violence exposure with both PTSD symptoms and externalizing problems, such that violence exposure was associated with worsening symptoms of psychopathology among youth with low RSA, but not among youth with high RSA. These findings replicate findings from a small study examining PTSD symptoms (Jenness et al., 2019) and extends prior work focused on internalizing and externalizing symptoms (El-Sheikh et al., 2001; Jenness et al., 2019; McLaughlin, Alves, et al., 2014; McLaughlin et al., 2015; Porges, 2007), by demonstrating that the association of violence exposure with PTSD symptoms is lower among youth with high RSA than among those with low

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

RSA. These findings align with the Polyvagal Theory, which proposes that high vagal tone enables contextually appropriate autonomic nervous system functioning that promotes greater resilience following exposure to stress and adversity (Porges, 2007; Sullivan et al., 2018). High vagal tone may reflect greater tonic inhibition of the amygdala by the medial prefrontal cortex (mPFC), which plays a critical role in effective regulation of negative emotion (Banks et al., 2007; Thayer et al., 2012). A greater capacity for emotion regulation might be one mechanism through which high vagal tone protects against the emergence of psychopathology following exposure to violence (Edwards & Pinna, 2020; Porges et al., 1994; Weissman, Bitran, et al., 2019).

Higher sympathetic nervous system (SNS) reactivity to both threat and safety cues during extinction learning was associated with subsequent increases in transdiagnostic psychopathology. Specifically, higher SCR to learned threat cues during early extinction learning predicted higher internalizing symptoms two years later. Elevated physiological reactivity to threat cues may reflect persistent fear to cues that were previously associated with threat that contributes to increased risk for internalizing problems in youth. Indeed, prior work suggests that heightened SNS responses to threat cues during early extinction learning is associated with increased risk for anxiety disorders in children (Craske et al., 2008). The present study extends these findings by demonstrating that higher SCR to learned threat cues during early extinction learning is associated with a broad range of internalizing problems in a longitudinal design. However, this association was not specific to learned threat cues. Higher SCR to learned safety cues during early extinction learning was associated with higher symptoms of PTSD, internalizing problems, externalizing problems, and general psychopathology two years later. Poor threat–safety discrimination during fear conditioning has been associated with externalizing psychopathology

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

in children and adolescents in cross-sectional studies (Fairchild et al., 2008; McLaughlin et al., 2016). To our knowledge, the present study is the first to link heightened SNS responses to the threat and safety cues during extinction learning with internalizing, externalizing, and general psychopathology (p-factor) in youth. This pattern of increased reactivity following fear learning, to both learned threat and safety cues is consistent with the recently proposed “better safe than sorry” information processing strategy (Bergh et al., 2020). This strategy in the context of aversive learning is thought to reflect oversimplified perception of stimulus features (e.g. recognizing the bell’s shape but not its color) that may lead one to expect that situations containing features of learned threat cues (e.g. the bell’s shape) are dangerous, even if these situations also contain features that designate safety (e.g. the bell’s color) (Bergh et al., 2020). This “better safe than sorry” information processing strategy may result in chronic deviations of expectations from reality that lead to greater levels of psychopathology (Bergh et al., 2020). In situations indicative of safety, failure to differentiate between threat and safety cues may result in contextually inappropriate fear responses that contribute to increased risk for multiple forms of psychopathology in youth, ranging from anxiety and internalizing problems to reactive aggression. While these findings related to the safety cue provide important context about the specificity of associations between responses to the conditioned threat cue and psychopathology, analyses related to the CS– were not preregistered, and thus should be considered preliminary until replicated in future studies.

We also found a negative association between age and SNS responses to both threat and safety cues. This suggests that the neural systems involved in extinguishing learned threat responses are continuing to develop across childhood and adolescence. Indeed, SCR during extinction learning tends to be higher in children than adults (Britton et al., 2013). Thus,

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

heightened reactivity to threat and safety cues among youth with psychopathology may reflect altered maturation of processes involved with fear extinction learning, including perceptual memory and the regulation of SNS activity (Bergh et al., 2020; Britton et al., 2013). Indeed, developmental change in fear extinction learning may be most pronounced during childhood and adolescence (Pattwell et al., 2012), suggesting this may be a period in which the mechanisms underlying threat and safety learning are particularly susceptible to environmental influences. These findings underscore the need for further research examining the influence of threat and safety learning on psychopathology across development.

Neither vagal tone, violence exposure, nor their interaction were associated with fear extinction learning. Thus, we failed to replicate the finding that enhanced fear extinction learning may be a mechanism through which vagal tone protects against psychopathology in children exposed to trauma (Jenness et al., 2019). We also failed to replicate the prior finding that vagal tone moderates the association between violence exposure and psychopathology cross-sectionally. This could be because past cross-sectional studies examining such associations, on average, used samples comprised largely of adolescents (Jenness et al., 2019; McLaughlin et al., 2014, 2015). Developmental differences in vagal tone might arise as a result of increasing inhibition of the amygdala by the mPFC with age, a pattern consistent with a number of studies on functional connectivity of the mPFC and amygdala (Gee et al., 2013; Gentzler et al., 2012; Weissman, Guyer et al., 2018, 2019), albeit in different types of emotion processing tasks than the aversive learning task utilized here. The link between vagal tone and enhanced extinction learning in violence-exposed youth may therefore depend on the maturation of this brain circuit. Thus, it is possible that the protective influence of vagal tone manifests differently across development as the neural circuits that underlie threat and safety learning mature.

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

Our findings underscore the need for future research investigating whether interventions that target vagal tone might also mitigate the effects of exposure to violence in youth. For example, mindfulness-based interventions, which have been shown to increase vagal tone in response to laboratory stressors, have also been shown to reduce symptoms of psychopathology among adults exposed to interpersonal violence (Ditto et al., 2006; Gallegos et al., 2015; Kelly & Garland, 2016). However, we are unaware of work demonstrating the effectiveness of interventions on psychopathology that strengthen vagal tone in youth. Thus, future research should explore, for example, whether interventions that are personalized to this developmental period and target threat and safety learning in a more scaffolded manner (e.g. mindfulness-based interventions) enhance the effectiveness of exposure-based treatments (Björkstrand et al., 2019; Gard et al., 2014; McGuire & Storch, 2019; Pattwell et al., 2012; Sullivan et al., 2018).

Limitations

The current study has several limitations. First, differences in aversive learning have been observed in children exposed to trauma and thus differences in extinction learning may also partially reflect differences in the patterns of learning of conditioned fear (Fairchild et al., 2008; McLaughlin et al., 2016). Furthermore, socioeconomic status is confounded with exposure to violence in our sample. While we include family income as a covariate in our analyses, future studies that more carefully match on socioeconomic status may extend this work to further disentangle the unique influence of this environmental factor on psychopathology. Finally, given that exposure to violence in our sample predates measures of vagal tone, we are unable to establish the extent to which individual differences in vagal tone may be a protective factor that exists before exposure to trauma or may reflect differential trajectories of physiological adaptation or calibration following trauma (Giudice et al., 2011). Future research using

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

genetically controlled and/or multi-wave longitudinal designs are necessary to further understand how development of vagal tone influences psychopathology among children exposed to adversity. Finally, the bifactor model has been criticized because fit indices tend to be biased in its favor relative to other model types (e.g. Greene et al., 2019). This is an important point for research aimed at uncovering the latent structure of psychopathology. However, for the purposes of individual differences analyses, as in the current report, the rank ordering of participants on general psychology measures is highly stable regardless of the statistical approach used to derive these scores (Clark et al., 2021; Schaefer et al., 2018).

Conclusion

We examined the role of fear extinction learning and high vagal tone as potential protective factors for psychopathology among youths exposed to violence. We found that the longitudinal association of violence exposure with PTSD symptoms and externalizing problems was weaker among children with high vagal tone than among those with low vagal tone. These associations were not explained, however, by differences in extinction learning. We also provide novel evidence that the magnitude of SNS responses to learned threat and safety cues decrease with age, suggesting developmental improvements in extinction learning, and that greater SNS reactivity to both threat and safety cues may contribute to transdiagnostic psychopathology over time. Altered maturation of processes involved in threat and safety learning may increase risk for psychopathology as children grow older and require more refined information processing strategies and contextually appropriate autonomic nervous system functioning in order to adapt to new environments and function with greater independence. Taken together, these findings suggest that interventions that target increasing vagal tone, while also decreasing reactivity to learned threat and safety cues in a developmentally appropriate manner, may have potential to

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

enhance the effectiveness of current evidence-based treatments and prevention efforts for psychopathology in youth.

References

- Abend, R., Gold, A. L., Britton, J. C., Michalska, K. J., Shechner, T., Sachs, J. F., ... Pine, D. S. (2020). Anticipatory Threat Responding: Associations With Anxiety, Development, and Brain Structure. *Biological Psychiatry*, 87(10), 916–925.
<https://doi.org/10.1016/j.biopsych.2019.11.006>
- Allen, J. J. B., Chambers, A. S., & Towers, D. N. (2007). The many metrics of cardiac chronotropy: A pragmatic primer and a brief comparison of metrics. *Biological Psychology*, 74(2), 243–262. <https://doi.org/10.1016/j.biopsycho.2006.08.005>
- Banks, S. J., Eddy, K. T., Angstadt, M., Nathan, P. J., & Phan, K. L. (2007). Amygdala–frontal connectivity during emotion regulation. *Social Cognitive and Affective Neuroscience*, 2(4), 303–312. <https://doi.org/10.1093/scan/nsm029>
- Beauchaine, T. P. (2015). Respiratory Sinus Arrhythmia: A Transdiagnostic Biomarker of Emotion Dysregulation and Psychopathology. *Current Opinion in Psychology*, 3, 43–47.
<https://doi.org/10.1016/j.copsyc.2015.01.017>
- Bergh, O. V. den, Brosschot, J., Critchley, H., Thayer, J. F., & Ottaviani, C. (2020). Better Safe Than Sorry: A Common Signature of General Vulnerability for Psychopathology: *Perspectives on Psychological Science*. (Sage CA: Los Angeles, CA).
<https://doi.org/10.1177/1745691620950690>
- Bernstein, D. P., Ahluvalia, T., Pogge, D., & Handelsman, L. (1997). Validity of the Childhood Trauma Questionnaire in an adolescent psychiatric population. *Journal of the American Academy of Child & Adolescent Psychiatry*, 36(3), 340–348.
<https://doi.org/10.1097/00004583-199703000-00012>

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

Berntson, G. G., Bigger, J. T., Eckberg, D. L., Grossman, P., Kaufmann, P. G., Malik, M., ...

van der Molen, M. W. (1997). Heart rate variability: Origins, methods, and interpretive caveats. *Psychophysiology*, 34(6), 623–648. <https://doi.org/10.1111/j.1469-8986.1997.tb02140.x>

Berntson, G. G., Cacioppo, J. T., & Quigley, K. S. (1993). Respiratory sinus arrhythmia:

Autonomic origins, physiological mechanisms, and psychophysiological implications.

Psychophysiology, 30(2), 183–196. <https://doi.org/10.1111/j.1469-8986.1993.tb01731.x>

Bifulco, A., Brown, G. W., & Harris, T. O. (1994). Childhood Experience of Care and Abuse

(CECA): A retrospective interview measure. *Journal of Child Psychology and*

Psychiatry, and Allied Disciplines, 35(8), 1419–1435. <https://doi.org/10.1111/j.1469-7610.1994.tb01284.x>

Birmaher, BORIS, Brent, D. A., Chiappetta, L., Bridge, J., Monga, S., & Baugher, M. (1999).

Psychometric Properties of the Screen for Child Anxiety Related Emotional Disorders

(SCARED): A Replication Study. *Journal of the American Academy of Child &*

Adolescent Psychiatry, 38(10), 1230–1236. [https://doi.org/10.1097/00004583-](https://doi.org/10.1097/00004583-199910000-00011)

199910000-00011

Birmaher, Boris, Khetarpal, S., Brent, D., Cully, M., Balach, L., Kaufman, J., & Neer, S. M.

(1997). The Screen for Child Anxiety Related Emotional Disorders (SCARED): Scale

Construction and Psychometric Characteristics. *Journal of the American Academy of*

Child & Adolescent Psychiatry, 36(4), 545–553. [https://doi.org/10.1097/00004583-](https://doi.org/10.1097/00004583-199704000-00018)

199704000-00018

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

- Björkstrand, J., Schiller, D., Li, J., Davidson, P., Rosén, J., Mårtensson, J., & Kirk, U. (2019). The effect of mindfulness training on extinction retention. *Scientific Reports*, 9(1), 19896. <https://doi.org/10.1038/s41598-019-56167-7>
- Blechert, J., Michael, T., Vriends, N., Margraf, J., & Wilhelm, F. H. (2007). Fear conditioning in posttraumatic stress disorder: Evidence for delayed extinction of autonomic, experiential, and behavioural responses. *Behaviour Research and Therapy*, 45(9), 2019–2033. <https://doi.org/10.1016/j.brat.2007.02.012>
- Bouton, M. E. (1993). Context, time, and memory retrieval in the interference paradigms of Pavlovian learning. *Psychological Bulletin*, 114(1), 80–99. <https://doi.org/10.1037/0033-2909.114.1.80>
- Bouton, Mark E. (2004). Context and behavioral processes in extinction. *Learning & Memory (Cold Spring Harbor, N.Y.)*, 11(5), 485–494. <https://doi.org/10.1101/lm.78804>
- Britton, J. C., Grillon, C., Lissek, S., Norcross, M. A., Szuhany, K. L., Chen, G., ... Pine, D. S. (2013). Response to Learned Threat: An fMRI Study in Adolescent and Adult Anxiety. *American Journal of Psychiatry*, 170(10), 1195–1204. <https://doi.org/10.1176/appi.ajp.2013.12050651>
- Buuren, S. van, & Groothuis-Oudshoorn, K. (2011). mice: Multivariate Imputation by Chained Equations in R. *Journal of Statistical Software*, 45(3), 1–67.
- Canty, A., & Ripley, B. D. (2019). *boot: Bootstrap R (S-Plus) Functions*.
- Caspi, A., Houts, R. M., Belsky, D. W., Goldman-Mellor, S. J., Harrington, H., Israel, S., ... Moffitt, T. E. (2014). The p factor: One general psychopathology factor in the structure of psychiatric disorders? *Clinical Psychological Science*. <https://doi.org/10.1177/2167702613497473>

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

- Caspi, A., & Moffitt, T. E. (2018). All for one and one for all: Mental disorders in one dimension. *American Journal of Psychiatry*, appi.ajp.2018.17121383-
appi.ajp.2018.17121383. <https://doi.org/10.1176/appi.ajp.2018.17121383>
- Cicchetti, D., & Toth, S. L. (2005). Child Maltreatment. *Annual Review of Clinical Psychology*, 1(1), 409–438. <https://doi.org/10.1146/annurev.clinpsy.1.102803.144029>
- Clark, D. A., Hicks, B. M., Angstadt, M., Rutherford, S., Taxali, A., Hyde, L., Weigard, A. S., Heitzeg, M. M., & Sripada, C. (2021). The General Factor of Psychopathology in the Adolescent Brain Cognitive Development (ABCD) Study: A Comparison of Alternative Modeling Approaches. *Clinical Psychological Science*, 9(2), 169–182.
<https://doi.org/10.1177/2167702620959317>
- Craske, M. G., Rauch, S. L., Ursano, R., Prenoveau, J., Pine, D. S., & Zinbarg, R. E. (2011). What Is an Anxiety Disorder? *FOCUS*, 9(3), 369–388.
<https://doi.org/10.1176/foc.9.3.foc369>
- Craske, M. G., Waters, A. M., Lindsey Bergman, R., Naliboff, B., Lipp, O. V., Negoro, H., & Ornitz, E. M. (2008). Is aversive learning a marker of risk for anxiety disorders in children? *Behaviour Research and Therapy*, 46(8), 954–967.
<https://doi.org/10.1016/j.brat.2008.04.011>
- Davison, A. C., & Hinkley, D. V. (1997). *Bootstrap Methods and Their Applications*. Cambridge: Cambridge University Press. Retrieved from
<http://statwww.epfl.ch/davison/BMA/>
- Dawson, M. E., Schell, A. M., & Filion, D. L. (2017). The electrodermal system. In *Cambridge Handbooks in Psychology. Handbook of psychophysiology*, 4th ed (pp. 217–243). New York, NY, US: Cambridge University Press.

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

Ditto, B., Eclache, M., & Goldman, N. (2006). Short-term autonomic and cardiovascular effects of mindfulness body scan meditation. *Annals of Behavioral Medicine: A Publication of the Society of Behavioral Medicine*, 32(3), 227–234.

https://doi.org/10.1207/s15324796abm3203_9

Duffy, M. E., Twenge, J. M., & Joiner, T. E. (2019). Trends in Mood and Anxiety Symptoms and Suicide-Related Outcomes Among U.S. Undergraduates, 2007–2018: Evidence From Two National Surveys. *Journal of Adolescent Health*, 65(5), 590–598.

<https://doi.org/10.1016/j.jadohealth.2019.04.033>

Dvir, M., Horovitz, O., Aderka, I. M., & Shechner, T. (2019). Fear conditioning and extinction in anxious and non-anxious youth: A meta-analysis. *Behaviour Research and Therapy*, 120, 103431. <https://doi.org/10.1016/j.brat.2019.103431>

Edwards, D. J., & Pinna, T. (2020). A Systematic Review of Associations Between Interoception, Vagal Tone, and Emotional Regulation: Potential Applications for Mental Health, Wellbeing, Psychological Flexibility, and Chronic Conditions. *Frontiers in Psychology*, 11. <https://doi.org/10.3389/fpsyg.2020.01792>

El-Sheikh, M., Harger, J., & Whitson, S. M. (2001). Exposure to interparental conflict and children's adjustment and physical health: The moderating role of vagal tone. *Child Development*, 72(6), 1617–1636. <https://doi.org/10.1111/1467-8624.00369>

Fairchild, G., Van Goozen, S. H., Stollery, S. J., & Goodyer, I. M. (2008). Fear conditioning and affective modulation of the startle reflex in male adolescents with early-onset or adolescence-onset conduct disorder and healthy control subjects. *Biological Psychiatry*, 63(3), 279–285. <https://doi.org/10.1016/j.biopsych.2007.06.019>

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

- Finkelhor, D., Hamby, S. L., Ormrod, R., & Turner, H. (2005). The Juvenile Victimization Questionnaire: Reliability, validity, and national norms. *Child Abuse & Neglect*, 29(4), 383–412. <https://doi.org/10.1016/j.chiabu.2004.11.001>
- Fox, N. A., & Leavitt, L. A. (1995). *The Violence Exposure Scale for Children-Revised (VEX-R)*. College Park, MD: University of Maryland.
- Fullana, M. A., Harrison, B. J., Soriano-Mas, C., Vervliet, B., Cardoner, N., Àvila-Parcet, A., & Radua, J. (2016). Neural signatures of human fear conditioning: An updated and extended meta-analysis of fMRI studies. *Molecular Psychiatry*, 21(4), 500–508. <https://doi.org/10.1038/mp.2015.88>
- Gallegos, A. M., Lytle, M. C., Moynihan, J. A., & Talbot, N. L. (2015). Mindfulness-Based Stress Reduction to Enhance Psychological Functioning and Improve Inflammatory Biomarkers in Trauma-Exposed Women: A Pilot Study. *Psychological Trauma : Theory, Research, Practice and Policy*, 7(6), 525–532. <https://doi.org/10.1037/tra0000053>
- Gamwell, K., Nylocks, M., Cross, D., Bradley, B., Norrholm, S. D., & Jovanovic, T. (2015). Fear Conditioned Responses and PTSD Symptoms in Children: Sex Differences in Fear-Related Symptoms. *Developmental Psychobiology*, 57(7), 799–808. <https://doi.org/10.1002/dev.21313>
- Garcia, N. M., & Zoellner, L. A. (2017). Fear generalisation in individuals with high neuroticism: Increasing predictability is not necessarily better. *Cognition and Emotion*, 31(8), 1647–1662. <https://doi.org/10.1080/02699931.2016.1259160>
- Gard, T., Noggle, J. J., Park, C. L., Vago, D. R., & Wilson, A. (2014). Potential self-regulatory mechanisms of yoga for psychological health. *Frontiers in Human Neuroscience*, 8. <https://doi.org/10.3389/fnhum.2014.00770>

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

- Gazendam, F. J., Kamphuis, J. H., & Kindt, M. (2013). Deficient safety learning characterizes high trait anxious individuals. *Biological Psychology*, 92(2), 342–352.
<https://doi.org/10.1016/j.biopsycho.2012.11.006>
- Gee, D. G., Humphreys, K. L., Flannery, J., Goff, B., Telzer, E. H., Shapiro, M., ... Tottenham, N. (2013). A Developmental Shift from Positive to Negative Connectivity in Human Amygdala–Prefrontal Circuitry. *The Journal of Neuroscience*, 33(10), 4584–4593.
<https://doi.org/10.1523/JNEUROSCI.3446-12.2013>
- Gentzler, A. L., Rottenberg, J., Kovacs, M., George, C. J., & Morey, J. (2012). Atypical development of resting respiratory sinus arrhythmia in children at high risk for depression. *Developmental Psychobiology*, 54(5), 556–567.
<https://doi.org/10.1002/dev.20614>
- Giudice, M. D., Ellis, B. J., & Shirtcliff, E. A. (2011). The Adaptive Calibration Model of stress responsivity. *Neuroscience and Biobehavioral Reviews*, 35(7), 1562–1592.
<https://doi.org/10.1016/j.neubiorev.2010.11.007>
- Giustino, T. F., & Maren, S. (2015). The Role of the Medial Prefrontal Cortex in the Conditioning and Extinction of Fear. *Frontiers in Behavioral Neuroscience*, 9.
<https://doi.org/10.3389/fnbeh.2015.00298>
- Gold, A. L., Shechner, T., Farber, M. J., Spiro, C. N., Leibenluft, E., Pine, D. S., & Britton, J. C. (2016). Amygdala-Cortical Connectivity: Associations with Anxiety, Development, and Threat. *Depression and Anxiety*, 33(10), 917–926. <https://doi.org/10.1002/da.22470>
- Green, J. G., McLaughlin, K. A., Berglund, P. A., Gruber, M. J., Sampson, N. A., Zaslavsky, A. M., & Kessler, R. C. (2010a). Childhood adversities and adult psychiatric disorders in the national comorbidity survey replication I: Associations with first onset of DSM-IV

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

- disorders. *Archives of General Psychiatry*, 67(2), 113–123.
<https://doi.org/10.1001/archgenpsychiatry.2009.186>
- Green, J. G., McLaughlin, K. A., Berglund, P. A., Gruber, M. J., Sampson, N. A., Zaslavsky, A. M., & Kessler, R. C. (2010b). Childhood adversities and adult psychiatric disorders in the national comorbidity survey replication I: Associations with first onset of DSM-IV disorders. *Archives of General Psychiatry*, 67(2), 113–123.
<https://doi.org/10.1001/archgenpsychiatry.2009.186>
- Greene, A. L., Eaton, N. R., Li, K., Forbes, M. K., Krueger, R. F., Markon, K. E., ... Kotov, R. (2019). Are fit indices used to test psychopathology structure biased? A simulation study. *Journal of Abnormal Psychology*, 128(7), 740–764. <https://doi.org/10.1037/abn0000434>
- Grillon, C., & Morgan III, C. A. (1999). Fear-potentiated startle conditioning to explicit and contextual cues in Gulf War veterans with posttraumatic stress disorder. *Journal of Abnormal Psychology*, 108(1), 134–142. <https://doi.org/10.1037/0021-843X.108.1.134>
- Guthrie, R. M., & Bryant, R. A. (2006). Extinction learning before trauma and subsequent posttraumatic stress. *Psychosomatic Medicine*, 68(2), 307–311.
<https://doi.org/10.1097/01.psy.0000208629.67653.cc>
- Hale, W. W., Raaijmakers, Q., Muris, P., & Meeus, W. (2005). Psychometric Properties of the Screen for Child Anxiety Related Emotional Disorders (SCARED) in the General Adolescent Population. *Journal of the American Academy of Child & Adolescent Psychiatry*, 44(3), 283–290. <https://doi.org/10.1097/00004583-200503000-00013>
- Hanson, J. L., Knodt, A. R., Brigidi, B. D., & Hariri, A. R. (2015). Lower structural integrity of the uncinate fasciculus is associated with a history of child maltreatment and future

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

- psychological vulnerability to stress. *Development and Psychopathology*, 27(4 0 2), 1611–1619. <https://doi.org/10.1017/S0954579415000978>
- Jenness, J. L., Miller, A. B., Rosen, M. L., & McLaughlin, K. A. (2019). Extinction Learning as a Potential Mechanism Linking High Vagal Tone with Lower PTSD Symptoms among Abused Youth. *Journal of Abnormal Child Psychology*, 47(4), 659–670. <https://doi.org/10.1007/s10802-018-0464-0>
- Jennings, J. R., Sheu, L. K., Kuan, D. C.-H., Manuck, S. B., & Gianaros, P. J. (2016). Resting state connectivity of the medial prefrontal cortex covaries with individual differences in high-frequency heart rate variability. *Psychophysiology*, 53(4), 444–454. <https://doi.org/10.1111/psyp.12586>
- Jovanovic, T., Nylocks, K. M., Gamwell, K. L., Smith, A., Davis, T. A., Norrholm, S. D., & Bradley, B. (2014). Development of Fear Acquisition and Extinction in Children: Effects of Age and Anxiety. *Neurobiology of Learning and Memory*, 113, 135–142. <https://doi.org/10.1016/j.nlm.2013.10.016>
- Jr, A. J. F., Saylor, C. F., Edwards, G. L., & McIntosh, J. A. (1987). Children's Depression Inventory: Reliability Over Repeated Administrations. *Journal of Clinical Child Psychology*, 16(4), 339–341. https://doi.org/10.1207/s15374424jccp1604_7
- Katz, L. F., & Gottman, J. M. (1997). Buffering children from marital conflict and dissolution. *Journal of Clinical Child Psychology*, 26(2), 157–171. https://doi.org/10.1207/s15374424jccp2602_4
- Katz, Lynn Fainsilber, & Gottman, J. M. (1995). Vagal tone protects children from marital conflict. *Development and Psychopathology*, 7(1), 83–92. <https://doi.org/10.1017/S0954579400006350>

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

Kelly, A., & Garland, E. L. (2016). Trauma-Informed Mindfulness-Based Stress Reduction for Female Survivors of Interpersonal Violence: Results From a Stage I RCT. *Journal of Clinical Psychology*, 72(4), 311–328. <https://doi.org/10.1002/jclp.22273>

Kessler, R. C., Avenevoli, S., Costello, E. J., Georgiades, K., Green, J. G., Gruber, M. J., ... Merikangas, K. R. (2012). Prevalence, persistence, and sociodemographic correlates of DSM-IV disorders in the National Comorbidity Survey Replication Adolescent Supplement. *Archives of General Psychiatry*, 69(4), 372–380. <https://doi.org/10.1001/archgenpsychiatry.2011.160>

Kessler, R. C., McLaughlin, K. A., Green, J. G., Gruber, M. J., Sampson, N. A., Zaslavsky, A. M., ... Williams, D. R. (2010). Childhood adversities and adult psychopathology in the WHO World Mental Health Surveys. *The British Journal of Psychiatry: The Journal of Mental Science*, 197(5), 378–385. <https://doi.org/10.1192/bjp.bp.110.080499>

Kessler, R. C., McLaughlin, K. A., Green, J. G., Gruber, M. J., Sampson, N. A., Zaslavsky, A. M., ... Williams, D. R. (2010). Childhood adversities and adult psychopathology in the WHO World Mental Health Surveys. *British Journal of Psychiatry*, 197(05), 378–385. <https://doi.org/10.1192/bjp.bp.110.080499>

Kim, M. J., Loucks, R. A., Palmer, A. L., Brown, A. C., Solomon, K. M., Marchante, A. N., & Whalen, P. J. (2011). The structural and functional connectivity of the amygdala: From normal emotion to pathological anxiety. *Behavioural Brain Research*, 223(2), 403–410. <https://doi.org/10.1016/j.bbr.2011.04.025>

Kolk, B. A. van der. (2017). Developmental Trauma Disorder: Toward a rational diagnosis for children with complex trauma histories. *Psychiatric Annals*, 35(5), 401–408. <https://doi.org/10.3928/00485713-20050501-06>

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

Kreutzmann, J. C., Marin, M.-F., Fendt, M., Milad, M. R., Ressler, K., & Jovanovic, T. (2021).

Unconditioned response to an aversive stimulus as predictor of response to conditioned fear and safety: A cross-species study. *Behavioural Brain Research*, 402, 113105.

<https://doi.org/10.1016/j.bbr.2020.113105>

Lau, J. Y., Britton, J. C., Nelson, E. E., Angold, A., Ernst, M., Goldwin, M., ... Pine, D. S.

(2011). Distinct neural signatures of threat learning in adolescents and adults.

Proceedings of the National Academy of Sciences, 108(11), 4500–4505.

<https://doi.org/10.1073/pnas.1005494108>

Li, C.-H. (2016). Confirmatory factor analysis with ordinal data: Comparing robust maximum

likelihood and diagonally weighted least squares. *Behavior Research Methods*, 48(3),

936–949. <https://doi.org/10.3758/s13428-015-0619-7>

Lissek, S., Powers, A. S., McClure, E. B., Phelps, E. A., Woldehawariat, G., Grillon, C., & Pine,

D. S. (2005). Classical fear conditioning in the anxiety disorders: A meta-analysis.

Behaviour Research and Therapy, 43(11), 1391–1424.

<https://doi.org/10.1016/j.brat.2004.10.007>

Mather, M., & Thayer, J. (2018). How heart rate variability affects emotion regulation brain

networks. *Current Opinion in Behavioral Sciences*, 19, 98–104.

<https://doi.org/10.1016/j.cobeha.2017.12.017>

McGrath, J. J., Saha, S., Lim, C. C. W., Aguilar-Gaxiola, S., Alonso, J., Andrade, L. H., ...

Wojtyniak, B. (2017). Trauma and psychotic experiences: Transnational data from the

World Mental Health survey. *British Journal of Psychiatry*.

<https://doi.org/10.1192/bjp.bp.117.205955>

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

McGuire, J. F., Orr, S. P., Essoe, J. K.-Y., McCracken, J. T., Storch, E. A., & Piacentini, J.

(2016). Extinction learning in childhood anxiety disorders, obsessive compulsive disorder and posttraumatic stress disorder: Implications for treatment. *Expert Review of Neurotherapeutics*, 16(10), 1155–1174. <https://doi.org/10.1080/14737175.2016.1199276>

McGuire, J. F., & Storch, E. A. (2019). An Inhibitory Learning Approach to Cognitive-

Behavioral Therapy for Children and Adolescents. *Cognitive and Behavioral Practice*, 26(1), 214–224. <https://doi.org/10.1016/j.cbpra.2017.12.003>

McLaughlin, K. A., Alves, S., & Sheridan, M. A. (2014). Vagal regulation and internalizing

psychopathology among adolescents exposed to childhood adversity. *Developmental Psychobiology*, 56(5), 1036–1051. <https://doi.org/10.1002/dev.21187>

McLaughlin, K. A., Green, J. G., Gruber, M. J., Sampson, N. A., Zaslavsky, A. M., & Kessler,

R. C. (2012a). Childhood adversities and first onset of psychiatric disorders in a national sample of adolescents. *Archives of General Psychiatry*, 69(11), 1151–1160.

<https://doi.org/10.1001/archgenpsychiatry.2011.2277>

McLaughlin, K. A., Green, J. G., Gruber, M. J., Sampson, N. A., Zaslavsky, A. M., & Kessler,

R. C. (2012b). Childhood adversities and first onset of psychiatric disorders in a national sample of US adolescents. *Archives of General Psychiatry*.

<https://doi.org/10.1001/archgenpsychiatry.2011.2277>

McLaughlin, K. A., Rith-Najarian, L., Dirks, M. A., & Sheridan, M. A. (2015). Low vagal tone

magnifies the association between psychosocial stress exposure and internalizing psychopathology in adolescents. *Journal of Clinical Child and Adolescent Psychology*:

The Official Journal for the Society of Clinical Child and Adolescent Psychology,

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

American Psychological Association, Division 53, 44(2), 314–328.

<https://doi.org/10.1080/15374416.2013.843464>

McLaughlin, K. A., Sheridan, M. A., Gold, A. L., Duys, A., Lambert, H. K., Peverill, M., ...

Pine, D. S. (2016). Maltreatment Exposure, Brain Structure, and Fear Conditioning in Children and Adolescents. *Neuropsychopharmacology: Official Publication of the American College of Neuropsychopharmacology, 41(8), 1956–1964.*

<https://doi.org/10.1038/npp.2015.365>

Merikangas, K. R., He, J.-P., Burstein, M., Swanson, S. A., Avenevoli, S., Cui, L., ... Swendsen,

J. (2010). Lifetime prevalence of mental disorders in U.S. adolescents: Results from the National Comorbidity Survey Replication--Adolescent Supplement (NCS-A). *Journal of the American Academy of Child and Adolescent Psychiatry, 49(10), 980–989.*

<https://doi.org/10.1016/j.jaac.2010.05.017>

Michael, T., Blechert, J., Vriends, N., Margraf, J., & Wilhelm, F. H. (2007). Fear conditioning in panic disorder: Enhanced resistance to extinction. *Journal of Abnormal Psychology, 116(3), 612–617.*

<https://doi.org/10.1037/0021-843X.116.3.612>

Milad, M. R., Pitman, R. K., Ellis, C. B., Gold, A. L., Shin, L. M., Lasko, N. B., ... Rauch, S. L.

(2009). Neurobiological Basis of Failure to Recall Extinction Memory in Posttraumatic Stress Disorder. *Biological Psychiatry, 66(12), 1075–1082.*

<https://doi.org/10.1016/j.biopsych.2009.06.026>

Milad, M. R., & Quirk, G. J. (2002). Neurons in medial prefrontal cortex signal memory for fear extinction. *Nature, 420(6911), 70–74.* <https://doi.org/10.1038/nature01138>

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

Milad, M. R., & Quirk, G. J. (2012). Fear extinction as a model for translational neuroscience:

Ten years of progress. *Annual Review of Psychology*, 63, 129–151.

<https://doi.org/10.1146/annurev.psych.121208.131631>

Morriss, J., Christakou, A., & van Reekum, C. M. (2019). Multimodal evidence for delayed

threat extinction learning in adolescence and young adulthood. *Scientific Reports*, 9(1),

7748. <https://doi.org/10.1038/s41598-019-44150-1>

Muthén, B., & Muthén, B. O. (2012). *Statistical analysis with latent variables*. New York:

Wiley.

Norrholm, S. D., Jovanovic, T., Olin, I. W., Sands, L. A., Karapanou, I., Bradley, B., & Ressler,

K. J. (2011). Fear Extinction in Traumatized Civilians with Posttraumatic Stress

Disorder: Relation to Symptom Severity. *Biological Psychiatry*, 69(6), 556–563.

<https://doi.org/10.1016/j.biopsych.2010.09.013>

Orr, S., Metzger, L., Lasko, N., Macklin, M., Peri, T., & Roger, P. (2000). De novo conditioning

in trauma-exposed individuals with and without posttraumatic stress disorder. *Journal of*

Abnormal Psychology, 109, 290–298. <https://doi.org/10.1037/0021-843X.109.2.290>

Pattwell, S. S., Duhoux, S., Hartley, C. A., Johnson, D. C., Jing, D., Elliott, M. D., ... Lee, F. S.

(2012). Altered fear learning across development in both mouse and human. *Proceedings*

of the National Academy of Sciences, 109(40), 16318–16323.

<https://doi.org/10.1073/pnas.1206834109>

Peña, D. F., Engineer, N. D., & McIntyre, C. K. (2013). Rapid remission of conditioned fear

expression with extinction training paired with vagus nerve stimulation. *Biological*

Psychiatry, 73(11), 1071–1077. <https://doi.org/10.1016/j.biopsych.2012.10.021>

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

- Peri, T., Ben-Shakhar, G., Orr, S. P., & Shalev, A. Y. (2000). Psychophysiological assessment of aversive conditioning in posttraumatic stress disorder. *Biological Psychiatry*, 47(6), 512–519. [https://doi.org/10.1016/S0006-3223\(99\)00144-4](https://doi.org/10.1016/S0006-3223(99)00144-4)
- Phelps, E. A., Delgado, M. R., Nearing, K. I., & LeDoux, J. E. (2004). Extinction Learning in Humans: Role of the Amygdala and vmPFC. *Neuron*, 43(6), 897–905. <https://doi.org/10.1016/j.neuron.2004.08.042>
- Porges, S. W., Doussard-Roosevelt, J. A., & Maiti, A. K. (1994). Vagal tone and the physiological regulation of emotion. *Monographs of the Society for Research in Child Development*, 59(2–3), 167–186.
- Porges, Stephen W. (1995). Orienting in a defensive world: Mammalian modifications of our evolutionary heritage. A Polyvagal Theory. *Psychophysiology*, 32(4), 301–318. <https://doi.org/10.1111/j.1469-8986.1995.tb01213.x>
- Porges, Stephen W. (2007). The polyvagal perspective. *Biological Psychology*, 74(2), 116–143. <https://doi.org/10.1016/j.biopsycho.2006.06.009>
- Rivera, C. L., Bernal, G., & Rosselló, J. (n.d.). *The Children Depression Inventory (CDI) and the Beck Depression Inventory (BDI): Their validity as screening measures for major depression in a group of Puerto Rican adolescents*. 5, 15.
- Sakaki, M., Yoo, H. J., Nga, L., Lee, T.-H., Thayer, J. F., & Mather, M. (2016). Heart rate variability is associated with amygdala functional connectivity with MPFC across younger and older adults. *NeuroImage*, 139, 44–52. <https://doi.org/10.1016/j.neuroimage.2016.05.076>
- Schaefer, J. D., Moffitt, T. E., Arseneault, L., Danese, A., Fisher, H. L., Houts, R., ... Caspi, A. (2018). Adolescent Victimization and Early-Adult Psychopathology: Approaching Causal

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

Inference Using a Longitudinal Twin Study to Rule Out Noncausal Explanations.

Clinical Psychological Science, 6(3), 352–371.

<https://doi.org/10.1177/2167702617741381>

Shader, T. M., Gatzke-Kopp, L. M., Crowell, S. E., Jamila Reid, M., Thayer, J. F., Vasey, M.

W., ... Beauchaine, T. P. (2018). Quantifying respiratory sinus arrhythmia: Effects of misspecifying breathing frequencies across development. *Development and*

Psychopathology, 30(1), 351–366. <https://doi.org/10.1017/S0954579417000669>

Shechner, T., Britton, J. C., Ronkin, E. G., Jarcho, J. M., Mash, J. A., Michalska, K. J., ... Pine,

D. S. (2015). Fear Conditioning and Extinction in Anxious and Nonanxious Youth and Adults: Examining a Novel Developmentally Appropriate Fear-Conditioning Task.

Depression and Anxiety, 32(4), 277–288. <https://doi.org/10.1002/da.22318>

Sherwood(Chair), A., Allen, M. T., Fahrenberg, J., Kelsey, R. M., Lovallo, W. R., & Doornen,

L. J. P. van. (1990). Methodological Guidelines for Impedance Cardiography.

Psychophysiology, 27(1), 1–23. <https://doi.org/10.1111/j.1469-8986.1990.tb02171.x>

Sitarenios, G., & Kovacs, M. (1999). Use of the Children's Depression Inventory. In *The use of psychological testing for treatment planning and outcomes assessment, 2nd ed* (pp. 267–298). Mahwah, NJ, US: Lawrence Erlbaum Associates Publishers.

Snyder, H. R., & Hankin, B. L. (2017). All Models Are Wrong, but the p Factor Model Is

Useful: Reply to Widiger and Oltmanns (2017) and Bonifay, Lane, and Reise (2017).

Clinical Psychological Science, 5(1), 187–189.

<https://doi.org/10.1177/2167702616659389>

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

- Staples-Bradley, L. K., Treanor, M., & Craske, M. G. (2018). Discrimination between safe and unsafe stimuli mediates the relationship between trait anxiety and return of fear. *Cognition & Emotion*, 32(1), 167–173. <https://doi.org/10.1080/02699931.2016.1265485>
- Steinberg, A. M., Brymer, M. J., Kim, S., Briggs, E. C., Ippen, C. G., Ostrowski, S. A., ... Pynoos, R. S. (2013). Psychometric Properties of the UCLA PTSD Reaction Index: Part I. *Journal of Traumatic Stress*, 26(1), 1–9. <https://doi.org/10.1002/jts.21780>
- Sullivan, M. B., Erb, M., Schmalzl, L., Moonaz, S., Noggle Taylor, J., & Porges, S. W. (2018). Yoga Therapy and Polyvagal Theory: The Convergence of Traditional Wisdom and Contemporary Neuroscience for Self-Regulation and Resilience. *Frontiers in Human Neuroscience*, 12. <https://doi.org/10.3389/fnhum.2018.00067>
- Susman, E., Weissman, D., & McLaughlin, K. A. (2020). *Vagal Tone and Extinction Learning as Potential Transdiagnostic Protective Factors Following Childhood Violence Exposure*. <https://doi.org/10.17605/OSF.IO/XUR87>
- Thayer, J. F., Ahs, F., Fredrikson, M., Sollers, J. J., & Wager, T. D. (2012). A meta-analysis of heart rate variability and neuroimaging studies: Implications for heart rate variability as a marker of stress and health. *Neuroscience and Biobehavioral Reviews*, 36(2), 747–756. <https://doi.org/10.1016/j.neubiorev.2011.11.009>
- Thayer, J. F., Hansen, A. L., Saus-Rose, E., & Johnsen, B. H. (2009). Heart Rate Variability, Prefrontal Neural Function, and Cognitive Performance: The Neurovisceral Integration Perspective on Self-regulation, Adaptation, and Health. *Annals of Behavioral Medicine*, 37(2), 141–153. <https://doi.org/10.1007/s12160-009-9101-z>
- Walker, E. A., Unutzer, J., Rutter, C., Gelfand, A., Saunders, K., VonKorff, M., ... Katon, W. (1999). Costs of health care use by women HMO members with a history of childhood

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

abuse and neglect. *Archives of General Psychiatry*, 56(7), 609–613.

<https://doi.org/10.1001/archpsyc.56.7.609>

Waters, A. M., & Pine, D. S. (2016). Evaluating differences in Pavlovian fear acquisition and extinction as predictors of outcome from cognitive behavioural therapy for anxious children. *Journal of Child Psychology and Psychiatry*, 57(7), 869–876.

<https://doi.org/10.1111/jcpp.12522>

Weissman, D. G., Bitran, D., Miller, A. B., Schaefer, J. D., Sheridan, M. A., & McLaughlin, K.

A. (2019). Difficulties with emotion regulation as a transdiagnostic mechanism linking child maltreatment with the emergence of psychopathology. *Development and Psychopathology*, 31(3), 899–915. <https://doi.org/10.1017/S0954579419000348>

Weissman, D. G., Guyer, A. E., Ferrer, E., Robins, R. W., & Hastings, P. D. (2018).

Adolescents' brain-autonomic coupling during emotion processing. *NeuroImage*, 183, 818–827. <https://doi.org/10.1016/j.neuroimage.2018.08.069>

Weissman, D. G., Guyer, A. E., Ferrer, E., Robins, R. W., & Hastings, P. D. (2019). Tuning of brain–autonomic coupling by prior threat exposure: Implications for internalizing problems in Mexican-origin adolescents. *Development and Psychopathology*, 31(3), 1127–1141. <https://doi.org/10.1017/S0954579419000646>

Weissman, D. G., Jenness, J. L., Colich, N. L., Miller, A. B., Sambrook, K. A., Sheridan, M. A., & McLaughlin, K. A. (2019). Altered Neural Processing of Threat-Related Information in Children and Adolescents Exposed to Violence: A Transdiagnostic Mechanism Contributing to the Emergence of Psychopathology. *Journal of the American Academy of Child & Adolescent Psychiatry*, 0(0). <https://doi.org/10.1016/j.jaac.2019.08.471>

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

Zuj, D. V., Palmer, M. A., Hsu, C.-M. K., Nicholson, E. L., Cushing, P. J., Gray, K. E., &

Felmingham, K. L. (2016). Impaired Fear Extinction Associated with Ptsd Increases with Hours-Since-Waking. *Depression and Anxiety*, 33(3), 203–210.

<https://doi.org/10.1002/da.22463>

Zuj, D. V., Palmer, M. A., Lommen, M. J. J., & Felmingham, K. L. (2016). The centrality of fear extinction in linking risk factors to PTSD: A narrative review. *Neuroscience &*

Biobehavioral Reviews, 69, 15–35. <https://doi.org/10.1016/j.neubiorev.2016.07.014>

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

Table 1 Descriptive statistics by violence exposure.

	<u>Exposed to Violence (n = 86)</u>		<u>Controls (n = 79)</u>			
	%	N	%	N	χ^2	P-value
<u>Sex</u>					1.08	.300
Female	52.3	45	43	34		
Male	47.7	41	57	45		
	M	(SD)	M	(SD)	t-Value	P-value
Income to Needs	2.22	2.28	5.39	2.26	8.64	<.001
Age	12.71	2.74	12.61	2.58	−0.22	.825
PTSD (Baseline)	29.15	15.55	3.95	8.81	−12.94	<.001
PTSD (Follow-up)	21.52	14.94	1.90	6.51	−9.80	<.001
Internalizing (Baseline)	0.54	0.75	−0.62	0.69	−10.36	<.001
Internalizing (Follow-up)	0.36	1.03	−0.30	0.76	−4.219	<.001
P-factor (Baseline)	0.53	0.67	−0.67	0.69	−11.28	<.001
P-Factor (Follow-up)	0.41	0.91	−0.39	0.71	−5.69	<.001
Resting RSA	5.91	1.10	5.83	1.18	−0.49	.622
Early Extinction CS+	1.04	0.07	1.06	0.08	1.99	.049
Early Extinction CS−	1.06	0.08	1.06	0.06	0.25	.803

Note: RSA = respiratory sinus arrhythmia; CS + = conditioned stimulus paired with aversive noise;

PTSD = posttraumatic stress disorder.

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

Figure 1 A) The association between baseline respiratory sinus arrhythmia (RSA) and PTSD symptoms at two-year follow-up in violence-exposed, and unexposed youth. B) The association between baseline RSA and externalizing problems at two-year follow-up, in violence-exposed, and unexposed youth.

VAGAL TONE EXTINCTION & PSYCHOPATHOLOGY

Figure 2 A) Higher SCR to the CS+ during early extinction learning did not predict higher PTSD symptoms at follow-up, controlling for baseline ($B = 16.84$, $\beta = .08$, $SE = 12.48$, $t = 1.35$, $p = .180$). B) Higher SCR to the CS+ during early extinction learning predicted higher internalizing problems at follow-up, controlling for baseline ($B = 1.86$, $\beta = 0.14$, $SE = 0.13$, $t = 2.06$, $p = .042$). C) Higher early extinction SCR to the CS+ marginally predicted higher externalizing problems at follow-up, controlling for baseline ($B = 1.78$, $\beta = 0.13$, $SE = 0.97$, $t = 1.85$, $p = .068$). D) Higher early extinction SCR to the CS+ marginally predicted higher p-factor at follow-up, controlling for baseline ($B = 1.58$, $\beta = .12$, $SE = 0.81$, $t = 1.94$, $p = .053$). E) Higher early extinction SCR to the CS- predicted PTSD symptoms at follow-up, controlling for baseline ($B = 24.20$, $\beta = .11$, $SE = 12.19$, $t = 1.98$, $p = .0496$). F) Higher early extinction SCR to the CS- predicted higher internalizing-factor at follow-up, controlling for baseline ($B = 2.23$, $\beta = .16$, $SE = 0.88$, $t = 2.54$, $p = .013$). G) Higher early extinction SCR to the CS- predicted higher externalizing problems at follow-up, controlling for baseline ($B = 2.29$, $\beta = .17$, $SE = 0.94$, $t = 2.44$, $p = .016$). H) Higher early extinction SCR to the CS- predicted higher p-factor at follow-up, controlling for baseline ($B = 2.31$, $\beta = .18$, $SE = 0.78$, $t = 2.96$, $p = .004$).