Increasing Anticipatory Pleasure in Major Depression through Enhancing Episodic Future Thinking: A Randomized Single-Case Series Trial

Hallford, D. J.^{1,3}, Sharma, M. K.², & Austin, D. W.¹

¹ School of Psychology, Deakin University, 1 Gheringhap Street, Geelong, Victoria 3220, Australia

² Department of Clinical Psychology, National Institute of Mental Health and Neurosciences, Hosur Road, Bangalore, Karnataka, India.

³ Corresponding Author, School of Psychology, Deakin University, 221 Burwood Highway, Burwood, Victoria 3125, Melbourne, Australia, Phone: + 61 3 9251 7777, Email: david.hallford@deakin.edu.au

Conflict of Interest: None to declare

Acknowledgements

The authors would like to acknowledge all staff and consumers at the National Institution of Mental Health and Neuroscience for their contributions to this study. D. J. Hallford was funded by an Australian Academy of Sciences Australia-India Fellowship and a Deakin Health Postdoctoral Research Fellowship

Keywords: major depression; anticipatory pleasure; anhedonia; episodic future thinking

Abstract

Depressed individuals have difficulty anticipating pleasure, which can impact motivation and functioning. One factor in this may be impairments in their episodic future thinking (EFT). This study examined whether enhancing EFT through increasing detail/vividness and mental imagery would increase anticipatory pleasure among individuals with Major Depressive Disorder. A randomized start-point, single case series design was used. Depressed outpatients ($N = 7$) completed surveys through the day over two weeks to nominate upcoming positive events and rate them on EFT detail/vividness, mental imagery, and anticipatory pleasure. At a randomized start-point, activities to enhance the detail/vividness and mental imagery for these upcoming events were introduced. Significant increases in detail and imagery were observed when EFT activities were introduced, which correlated with increases in how pleasurable it was thought the activities would be and how pleasurable it was thinking about them. Enhancing EFT may be a mechanism to increase anticipatory pleasure in depression. Implications for treatment are discussed.

Keywords: major depression; anticipatory pleasure; anhedonia; episodic future thinking

Anhedonia refers to difficulty feeling pleasure as a response to experiences that might otherwise elicit a pleasurable, hedonic response, and is a core diagnostic feature of depressive disorders (American Psychiatric Association, 2013; World Health Organization, 1992). Some evidence-based psychological therapies, such as cognitive therapy (Beck, 2011), behavioural activation (Hopko, Lejuez, Ruggiero, & Eifert, 2003) and acceptance and commitment therapy (Hayes, Strosahl, & Wilson, 2011), put a particular emphasis on increasing rewarding and meaningful activities as a way of increasing hedonic responses and addressing anhedonia. Recently, anhedonia has been increasingly conceptualised as a deficit in the ability to not only experience pleasure from behaviour that would usually be rewarding, but also as a deficit in *anticipating* the experience of pleasure (Gard et al., 2006; Gard, Kring, Gard, Horan, & Green, 2007; Treadway & Zald, 2011). Indeed, evidence has accumulated that indicates large, significant deficits in self-reported levels of anticipatory pleasure (*removed for blinding*; Wu et al., 2017; Yang et al., 2014, 2017), and abnormalities in the neural correlates of reward anticipation amongst those who are clinically depressed (Stringaris et al., 2018).

Anticipatory pleasure can be conceptualized as a maintaining factor in clinical depression, given it is a proximal influence in the motivation and intention to engage in rewarding and goal-directed behavior (Engel et al., 2013; Sherdell et al., 2012). Low anticipatory pleasure is predictive of low motivation for reward (Foussias & Remington, 2008; Treadway & Zald, 2011), low reward responsiveness (Foussias & Remington, 2008), and generally poorer outcomes on psychosocial functioning in mental illness (Buck & Lysaker, 2013; Foussias et al., 2011). Therefore, where one anticipates little pleasure from future experiences, intrinsic motivation for these experiences and intention to engage in them may also be low, positively-reinforcing behaviours are less likely to be engaged in, and this process may over time pervasively disrupt psychosocial functioning. Difficulty anticipating positive events in conjunction with a reduced experience of reward may then give rise to other factors that maintain dysfunction, such as generalized hopelessness about having positive

experiences and outcomes for oneself in the future (MacLeod et al., 2005; O'Connor, Connery, & Cheyne, 2000).

One possible reason for low anticipatory pleasure amongst depressed individuals may be the impairments they have in mentally simulating possible future events in specific and detailed ways. Activating and maintaining contextually detailed cognitive representations of experience are thought to be crucial for the anticipation of positive affect and reward (Kring & Caponigro, 2010). However, studies have shown that when depressed individuals engage in episodic future thinking (EFT), that is, thinking of specific, spatially and temporally-located future events they might experience, compared to healthy controls they provide more general descriptions of categories of events or events that extend over long periods (*removed for blinding*). Further, when specific instances of future experiences are generated, they typically have less contextual detail (King et al., 2011). Kahnemann and Tversky (1982) theorized that individuals use a simulation heuristic when engaging in EFT, whereby mental representations that are more realistically represented in mind and easier to generate are perceived as more plausible and likely to occur. This might also then give rise to a stronger experience of anticipatory emotion. Accordingly, where depressed individuals lack the ability to cognize detailed representations of experiences that might be rewarding in their future, they may also be less likely to anticipate pleasure from the thought of that experience. Following from this, if the specificity and detail of EFT for positive events could be enhanced, then corresponding changes in anticipatory pleasure might also be expected.

Several studies to date have assessed the emotional effects of engaging in EFT for positive events in samples with depressive symptoms. Higher ratings of vividness for positive imagery have been shown to correlate with lower depressive symptoms in dysphoric individuals (Holmes, Lang, Moulds, & Steele, 2008), and imagining positive events using mental imagery shown to produce a stronger intensity of positive effect relative to verbal conditions (Holmes, Coughtrey, & Connor, 2008, Holmes, Lang, & Shah, 2009). In a sample of dysphoric individuals, simulations of future events prompted by cues of abstract experiences (e.g., “people will admire you”, “you will be able

to cope easily with pressure”) were shown to increase the self-reported vividness of the experiences, and the perceived likelihood that they would occur (Boland, Riggs, & Anderson, 2018). As part of a trial of the effects of positive mental imagery on depressive symptoms, Renner, Pictet, Holmes, and Blackwell (2017) showed that higher vividness ratings of mental simulations of pre-scripted scenarios with positive outcomes were related to increased behavioural activation. Although not focused on depression, a pilot study of anticipatory skills training with five individuals with schizophrenia incorporated sensory experiences of pleasurable events and examination of the possible rewards of behaviours in order to increase anticipatory pleasure (Favrod, Giuliani, Ernst, & Bonsack, 2010). The results showed change over time on a trait measure of anticipatory pleasure, although the possible mediating mechanism of increased detail in mental representations of positive events was not assessed.

A common feature of the depression-focused studies discussed above is that they did not assess anticipatory pleasure. This is crucial in terms of assessing outcomes of simulating future experiences on the expectations of a hedonic response from these experiences, rather than using variables as a proxy for this that are only modestly correlated, such as positive affect (Gard et al., 2006), generalized optimism (Bryant, 2003), and happiness (Bryant, 2003; Saperia et al., 2018). As Renner et al. (2017) suggest, studies that assess and promote EFT for specific, personally-relevant events are needed to match the actual content and affective outcomes of the cognitions with desired behavioural outcomes. The ecological validity of examining EFT for events in daily life then, rather than abstractions or events that are not personally-relevant, seems important if effects are intended to be observed for experiences that actually occur in daily life. This approach may also increase the feasibility of translating such findings for use in established treatments that focus on increasing meaningful and rewarding behaviours. Renner and colleagues (2019) have recently shown that engaging in motivational imagery activities for planned behaviours can increase anticipatory pleasure in a non-clinical population (Renner, Murphy, Ji, Manly, & Holmes, 2019). However, it has not been tested whether or not such effects would also be found in those with depression.

The Current Study

As outlined above, anticipatory pleasure deficits are linked with poorer motivation and psychosocial functioning, and represent a target for intervention in clinical depression. Given that depressed individuals have difficulties in effectively mentally-representing positive future events, one mechanism through which to increase anticipatory pleasure for upcoming events may be to enhance their episodic future thinking. The aim of this study was to examine whether providing EFT tasks to depressed individuals would increase the level of contextual detail in their mental simulations of upcoming positive events, and increase their anticipatory pleasure for these events. These activities were designed to encourage characteristics of EFT that are known to be related to increased affective response, such as using mental imagery (Pictet, Coughtrey, Mathews, Holmes, 2011), using a first-person perspective (Holmes et al., 2008), and being temporally close (D'Argembeau & Van Der Linden, 2004). It was hypothesized that when engaging in enhanced EFT activities of upcoming positive events, relative to a control phase, depressed individuals would report higher levels of detail/vividness and mental imagery for these events, and higher levels of anticipatory pleasure. The current study was designed to focus on increasing the knowledge of mechanisms with which to improve treatments, and not on treatment *per se*. However, given the opportunity, we also measured relevant variables of mood, depressive symptoms, trait anticipatory pleasure, and behavioural activation. We anticipated that if there were changes observed on these variables that they would be in the direction of improved mood, lower symptoms, and increased anticipatory pleasure and behavioural activation.

Methods

Design

The study used a replicated, randomized start-point single case series A-B design to test the effects of enhanced EFT on anticipatory pleasure across repeated measurements. The study was pre-registered prior to commencing recruitment and data collection (Australian New Zealand Clinical Trials Registry: 12618000257268). There were two phases in the study: Control (completion of

measures) and enhanced EFT (EFT activities + completion of measures). Randomized start-point designs involve randomizing participants to one of a pre-determined number of points where they switch from the control phase to the enhanced EFT phase. The number of possible points in which a participant can switch between phases is restricted so as to avoid having too few measurement points.

In this study, measures were taken five times per day (8:30am, 11:30am, 2pm, 4:30pm, and 7pm) over the course of 14 days. The control phase lasted for a minimum of 15 data-points (three days). There were then 25 subsequent data-points (the 16th data-point at the start of the fourth day to the 40th data-point at the end of the eighth day) that participants could be switched to the enhanced EFT phase. The enhanced EFT phase would then last for a minimum of six days or more dependent on the start-point participants were randomized to, until the last data-point (70th data-point at the end of the two weeks). The primary outcome was anticipatory pleasure. The study was not designed as an intervention for depressive symptoms, however, we did include secondary outcomes of state mood (assessed in the daily surveys), depressive symptoms, behavioural activation (both for the specific upcoming events, and in general), and trait anticipatory pleasure. The latter three were assessed at baseline, when participants switched between the control and enhanced EFT phases, and at the end of the two weeks. As a manipulation check, we assessed self-reported detail/vividness and use of mental imagery in relation to the upcoming events in both phases.

Participants

The inclusion criteria were: aged between 18 and 65 years, a current Major Depressive Disorder (single or recurrent) as the primary complaint, and as diagnosed according to the International Classification of Diseases - 10th edition (World Health Organization), residing in Bengaluru, India, and fluency in the English language. Exclusion criteria were: psychotic depressive features, psychotic disorders, substance use disorder, neurodevelopmental or neurodegenerative disorders, obsessive-compulsive disorder, borderline personality disorder, and any concurrent psychosocial or psychological treatment. Antidepressant medication was not an

exclusion criterion, but there was a requirement of being stable in type and dose for at least four weeks prior to commencing the study. All participants taking antidepressant medication remained stable in type and dose throughout the study period.

A total of 10 participants were recruited for the study, but three were excluded prior to analyses. One participant advised at completion of the study that they had consistently violated the protocol by answering questions in the daily surveys at random, one participant completed only 22.5% of data-points, and the remaining participant reported depressive symptoms in the normal range at baseline indicating they no longer met study criteria. The characteristics of the sample used in analyses are shown in Table 1. As indicated, the sample was wholly male, with Major Depressive Disorder of mostly mild to moderate severity for at least three months duration (median of 10 months). Most had previous episodes and were on medication, only two had co-morbid mental illnesses.

Materials

The daily surveys were sent to the participants' phone via email, in which a link to the survey was provided, and EFT activities in the second phase. In each of the daily surveys, participants were asked to choose an activity that they were most looking forward to or could possibly do over the next two hours that they would find enjoyable or would give them a sense of achievement. Participants were told that if they nominated something they were currently engaged in and would be doing until the next data-point then they might try to think about an aspect of that experience that might be positive in some way. They then selected from a number of pre-set categories of events consistent with Wu et al. (2017): *work/school/study, TV/internet/games/media, conversation/socializing, errand/chores, a hobby (not physical activity), physical activity/exercise, eating/drinking, or other activity not listed above* (and asked to specify what the activity was within this category). Some examples of each activity were provided to prompt responses, for example, for the hobby category the following were listed: reading a book or magazine, doing a puzzle, playing an instrument, or listening to music.

To assess the variables of interest in the daily surveys a series of single-item measures were used, a common approach in studies that repeatedly assess episodic thinking throughout the day (Barsics, Van Der Linden, & D'Argembeau, 2016; Beaty, Seli, & Schacter, 2018). This was also in consideration of the large number of surveys, to reduce burden and increase response rate. All responses were made using a 0 – 100 scale with a sliding bar. Mood was assessed at the start of each daily survey with the items, “*Please rate how sad you are right now*”, (0 = *not sad at all*, 100 = *extremely sad*) and “*Please rate how happy you are right now*” (0 = *not at all happy*, 100 = *extremely happy*).

The mental detail/vividness of the upcoming activity was assessed using the item “*How vivid and detailed was your thought of the activity?*”, and mental imagery was assessed with the item, “*How much did you find yourself thinking in pictures/mental images about the activity?*” (both 0 = *not at all*, 100 = *very much so*). Participants were then asked to indicate whether they completed the activity that was chosen in the previous survey (*Yes/No*).

State anticipatory pleasure was then assessed using two items; one referring to how enjoyable the upcoming event would be, “*How pleasant/enjoyable do you think it will be to do this activity?*”, and the other referring to how enjoyable it was to think about doing the activity, “*How pleasurable/enjoyable is it to just think about doing this activity?*”, (both 0 = *not at all*, 100 = *very much so*).

Depression, Anxiety, and Stress Scale (DASS; Lovibond & Lovibond, 1995). The 7-item self-report depression subscale from the DASS 21-item short-form was used to assess for depressive symptoms. The short-form of the DASS depression subscale assess the core features of depression (e.g., *low mood, loss of interest, self-worth, and motivation*) and has good psychometric properties (Antony, Bieling, Cox, Enns, & Swinson, 1998). The items were summed, with higher scores indicating a higher severity of depressive symptoms. Internal reliability was found to be acceptable in the current study (Cronbach’s α across time-points = .82)

Temporal Experience of Pleasure Scale (TEPS; Gard et al., 2006). The 10-item anticipatory pleasure subscale of the self-report TEPS was used to assess for trait levels of anticipatory pleasure. The TEPS uses a combination of concrete (e.g., “*When ordering something off the menu, I imagine how good it will taste*”) and abstract items (e.g., “*When something exciting is coming up in my life, I really look forward to it*”) to assess the general ability to experience anticipatory pleasure. Responses were given on a 6-point scale (1 = *Very false for me*, 6 = *Very true for me*), and items were summed so that higher scores indicated a stronger tendency to anticipate pleasure. Internal reliability was found to be acceptable in the current sample (Cronbach’s α across time-points = .71)

Behavioral Activation for Depression Scale (BADS; Kanter, Mulick, Busch, Berlin & Martell, 2006). The BADS is a 25-item, self-report measure of behavioral activation. The 25 items assess goal-directed activity and completion of scheduled activities, avoidance of aversive states and rumination, and impairment in work/school and social domains. The items were summed, with higher scores indicating less avoidance and more behavioral engagement. Internal reliability was found to be acceptable (Cronbach’s α across time-points = .81)

Although there were initial plans to administer the Episodic Future Thinking-Test (*blinded for review*) this was abandoned due to time restrictions for assessment with the first two participants, and it was subsequently deemed to be too burdensome.

Enhanced Episodic Future Thinking Activities. The EFT activities were designed to increase the detail with which participants thought of upcoming events, and evoke a related emotional response. To do this, participants were asked to use a first-person perspective when thinking about the activity, and provide written accounts of thoughts, feelings, contextual details, and sequences of actions that would occur in the forthcoming activity. They were first given the instructions: “*Now we would like you think very intensely about this enjoyable activity that you chose. Think strongly about it and would it would be like to do. Imagine it happening from your own eyes/perspective. Really think strongly about this possible positive activity and use pictures in your*

mind, like a movie. Imagine what it would feel like to do this activity". They then provided answers to a number of open-ended question designed to help them generate episodic detail about the possible event: "*Where will you be?*", "*What will happen during this positive experience?*", "*Describe some positive things that you might think*", "*Describe the positive feelings that you will have during this activity*", and "*What will be the best part of it?*" They were then asked to choose which positive emotions they might feel during this activity from a list of adjectives from the positive affect subscale on the Positive and Negative Affect Schedule (Watson, Clark, & Tellegan, 1988); "*happy, excited, strong, enthusiastic, proud, inspired, determined, attentive, interested, and hopeful*", with the words "*calm*" and "*amused*" added to increase the range of responses.

Procedure

Prior to commencing the study, ethics approval was obtained from the university research human ethics committee and the *blinded for review* ethics committee. Potential participants were identified through the intake procedures of the adult outpatient department of *blinded for review*. Individuals who fit criteria for participation were identified by clinicians (psychiatrists, clinical psychologists, and psychiatric social workers) and permission was asked to be followed up by the research team. The diagnosis of Major Depressive Disorder and other eligibility criteria were confirmed with the referring clinician prior to this follow-up. The participant was then contacted via phone, or in person at their next scheduled appointment at the outpatient department. They were provided with a plain language consent form, and provided signed consent. Participants then completed baseline measures using a laptop computer, and were guided through how to complete the daily surveys of the control phase by a researcher. They were then randomized to a start-point for the enhanced EFT phase using an online randomizer (www.randomizer.org), and the daily surveys commenced the following morning at 8:30am. They attended the outpatient department again at the time and date of the enhanced EFT phase start-point to complete another set of measures, and were guided through how to complete the EFT activities that would be added to the daily surveys. At the end of the two weeks they again met with a researcher to complete the final set

of measures and to debrief on their experience. The researchers contacted each participant shortly after the commencement of each phase, to encourage completion of the daily surveys. The participants were reimbursed with 500 Indian Rupees upon each visit to the outpatient department.

Data Analysis

Descriptive statistics and Spearman rank correlation coefficients were generated using SPSS 24.0. Baseline Corrected Tau (BCT) was calculated for each participant on the study variables (Tarlow, 2017), which provides a non-parametric inferential test for differences in scores between the phases. BCT analyses can correct for trends in the control phase by first detecting whether a trend is occurring then statistically removing this trend so it equals or approaches zero before data is analyzed. This is particularly relevant where it is possible that an initial control phase might itself produce changes in scores on some variable. It is noted then that BCT analyses may produce effects that are in the opposite direction to mean changes between phases where significant trends in an initial phase are found. This procedure also provides an effect size estimate that is based on rank correlations, and therefore is less biased by outliers than the standardized mean difference. Suggested interpretations for effect sizes of the related Tau-U statistic are as follows: below .20 a small effect, .20 - .60 moderate, .60 - .80 large, and above .80 very large (Vannest & Ninci, 2015). It is also noted that the Tau-U statistic is not always bounded between -1 and +1, whereas the BCT used here is. Therefore, these effect size interpretations may underestimate the effect size of BCT statistics. A combined *p*-value for all participants was calculated using meta-analysis with Edgington's additive method (Edgington & Onghena, 2007) in the WINPEPI software (Abramson, 2011).

Inferential tests are not reported for depressive symptoms, trait anticipatory pleasure, and behavioural activation, given the small sample and inadequate power to detect changes, however, standardized mean differences are reported as a description of mean change and direction.

An average of 32.2% of data was missing for each of the participants (range 20.75 – 45.65%), which was unsurprising given the large amount of data-points, and repetitive nature of the

daily activities. This missing data was replaced using expectation maximization, a single imputation technique which has been shown to preserve inferential integrity in single case designs that are missing significant amounts of data (Smith, Borckardt, & Nash, 2012).

Results

The frequency of endorsement of the upcoming activity categories, in descending order, were: work/school/study (23.6%), TV/internet/games/media (21.6%), other activity not listed above (18.4%), conversation/socializing (12.1%), eating/drinking (11%), errand/chores (7.9%), hobby (not physical activity; 2.7%), physical activity/exercise (2.7%). To assess whether there were differences on the anticipatory pleasure items between the different activity categories, a MANOVA was conducted, using phase as a covariate to control for scores differences between control and enhanced EFT phases. The results indicated a significant multivariate effect for activity category, $F(14, 708) = 3.53, p < .001$, and univariate effects for how enjoyable participants thought the upcoming activity would be, $F(7, 345) = 5.72, p < .001$, and how enjoyable it was to just think about doing, $F(7, 345) = 5.26, p < .001$. Post-hoc tests were conducted to assess differences between the categories, using Games-Howell corrections in the context of unequal sample variances and sizes. The results showed differences in categories were consistent for the two anticipatory pleasure items: TV/internet/games/media and conversation/socializing evoked significantly more anticipatory pleasure than work/school/study, eating/drinking, and other activity not listed (all $p < .05$). No other differences were found.

Mechanisms of Change/Manipulation Check

The results from the BCT analyses indicated significant increases in detail/vividness from control phase to enhanced EFT phases for five participants, with one showing no significant change (P6) and one showing a significant decrease after baseline trend correction (P1). The meta-analysis, with combined p -values, indicated an overall statistically significant increase in detail/vividness (BCT = .271, $p = .046$). For mental imagery, the results indicated significant increases in mental imagery for five participants, with the remaining showing no significant change (P6 and P7). The

meta-analysis with combined p -values indicated an overall statistically significant increase in detail/vividness ($BCT = .336, p = .001$). In summary, the majority of participants increased in detail and mental imagery, and meta-analyses showed evidence for overall effects.

The association between detail/vividness and mental imagery and anticipatory pleasure was then assessed. Using the data prior to missing values replacement, strong, positive correlations between higher detail/vividness and higher anticipatory pleasure were observed for expected pleasure, $r(355) = .68, p < .001$, and pleasure thinking about the activities, $r(361) = .80, p < .001$, and for the use of mental imagery and expected pleasure, $r(361) = .68, p < .001$, and pleasure thinking about the activities, $r(360) = .80, p < .001$. Detail/vividness was also found to correlate highly with the use of mental imagery, $r(354) = .88, p < .001$. All correlations remained the same after controlling for phase. This indicated that higher detail/vividness and mental imagery were both associated with stronger anticipatory pleasure.

Multiple regressions were then conducted to explore whether detailed/vividness and mental imagery independently predicted the anticipatory pleasure variables, while controlling for phase. The model for expected pleasure for the event was significant, $F(2, 350) = 184.5, p < .001$, indicating the variables predicted 54% of variance in expected pleasure. Phase was a significant predictor ($\beta = .20, p < .001$), and both detail/vividness ($\beta = .32, p < .001$) and mental imagery ($\beta = .39, p < .001$) explained independent variance. The model for pleasure thinking about the upcoming event was also significant, $F(2, 350) = 386.5, p < .001$, and indicated the variables predicted 69% of variance in pleasure from thinking about the upcoming event. Again, phase was a significant predictor ($\beta = .06, p = .023$), and detail/vividness ($\beta = .40, p < .001$) and mental imagery ($\beta = .44, p < .001$) both explained unique variance. This indicated that higher detail/vividness and mental imagery independently predicted stronger anticipatory pleasure.

Primary Outcome: Anticipatory Pleasure

Table 2 shows the descriptive statistics for the primary outcomes, and Figure 1 and 2 show graphical representations of scores over time. As indicated, all participants increased in their mean

scores on how enjoyable they thought it would be to do the upcoming activities. The BCT analyses showed three participants had significant increases between the phases, with the remaining showing no significant change. The meta-analysis with combined p -values indicated an overall statistically significant increase in how enjoyable the participants thought it would be to do the upcoming activities ($BCT = .243, p < .001$). For how enjoyable it was to think about doing the upcoming activities, the results indicated significant increases for all participants. As expected, the meta-analysis with combined p -values indicated an overall statistically significant increase on how enjoyable it was to think about doing the upcoming activities ($BCT = .298, p < .001$). In summary, the participants reported increases in anticipatory pleasure, although this was more consistent for how enjoyable it was to think about doing the upcoming activities relative to how enjoyable it was thought they would be.

Secondary Outcomes

Turning to analyses of the frequency of engaging in the nominated behaviours, it is noted that P3 was excluded as all of their responses were “yes” and therefore there was no variance to estimate missing data with (see Supplementary Table 2 for all participants statistics). The results showed significant decreases in frequency of engaging in the nominated behaviours for two participants (P6 and P7), and no other significant changes amongst the participants. The meta-analysis combining effect sizes and p -values indicated that overall participants did not collectively change between the phases ($BCT = .027, p = .555$).

There was inconsistency across participants regarding mean changes in sadness and happiness between the phases. The results for sadness indicated significant decreases for four participants, and a significant increase for one (P5). The meta-analysis indicated a statistically significant decrease in sadness (sadness: $BCT = -.108, p < .001$). For happiness, there were significant decreases for two participants (P5 and P7) and no other significant changes. The meta-analysis indicated no overall statistically significant change in happiness (sadness: $BCT = .100, p = 1.00$).

Changes in mean scores (see Table 3) were not apparent from baseline to the end of the enhanced EFT phase for the measures of depressive symptoms ($d = .07$) and trait temporal experience of pleasure ($d = -.15$), but some mean change was observed for behavioural activation ($d = .42$). To examine the associations between these three measures, Spearman rank correlations were conducted. The average correlation across time-points between the DASS and TEPS-A was $r = -.44$, the average correlation between DASS and BADS was $r = -.58$, and the average correlation between TEPS-A and BAD was $r = .56$.

Discussion

This study aimed to investigate whether enhancing episodic future thinking for upcoming positive events in daily life would increase the anticipatory pleasure related to these events in individuals with Major Depressive Disorder. Taken collectively, the results showed that the majority of participants' self-reported detail/vividness and use of mental imagery for upcoming events increased when the EFT activities were introduced, and they reported anticipating more pleasure when thinking about upcoming events, relative to the control phase. Meta-analytic analyses of single-case results supported these findings. Further, the reports of detail/vividness and mental imagery were strongly and independently associated with higher scores on both anticipatory pleasure items. This suggests that enhancing EFT was a mechanism for increasing anticipatory pleasure, and that detail/vividness and mental imagery were distinct characteristics of EFT that, when increased in conjunction, promoted higher expectation and experience of pleasure in this sample.

Depressed individuals demonstrate significantly less specificity and detail when simulating future events (*removed for blinding*). Contextual detail/vividness in the simulation of future events is important in determining how plausible and realistic these events are judged to be (Boland et al., 2018; Kahneman & Tversky, 1982), and mental imagery (Holmes et al., 2008, 2009; Holmes, Mathews, Dalgleish, & Mackintosh, 2006) can help boost the positive feelings that they evoke. Therefore, a paucity of detail, low salience of incentives, and less evoked positive emotion

(Whitton, Treadway, & Pizzagalli, 2015) may explain why depressed individuals also report less anticipation of pleasure for upcoming events (Wu et al., 2017), and decreased responsiveness to the notion of reward (Henriques & Davidson, 2000; McFarland & Klein, 2009). This study provides evidence that anticipatory pleasure can be increased among depressed individuals by enhancing the detail/vividness and mental imagery that they use to imagine these future events.

As noted, mean changes in responses were seen in the hypothesised direction on the EFT and anticipatory pleasure variables for the majority of participants. One participant, however, reported a decrease in enjoyment thinking about the events (P6). It is unclear why this happened, although it is notable that this same participant did not increase on detail and mental imagery, instead reporting decreases, albeit not statistically significant. Therefore, these changes are at least consistent with the overall associations between detail, mental imagery, and anticipatory pleasure variables, and theory that anticipatory pleasure will increase as a product of changes in these EFT variables. It is possible that this individual found it difficult to engage in the activities to enhance their EFT, which may have lead to feeling demoralised about upcoming positive events over time, although this is of course speculative. It was notable that the participant did not reduce on the expectation of enjoyment though. Findings among previous studies have been inconsistent in regards to the the strength of association between expecting enjoyment and experiencing enjoyment in thinking about a future event, with some showing strong correlations (Barsics et al., 2016), and others weak (Baumgartner, Pieters, & Bagozzi, 2008). The variables strongly correlated in this study, and both changed across the sample to a similar degree. Therefore, it is reasonable to believe that, overall, participants experienced pleasure at the time of thinking about events that they perceived would be more pleasurable, or at least that these constructs significantly overlapped in how participants conceptualized them.

No changes were observed on the trait measure of anticipatory pleasure, although this was not a primary outcome and, given the brevity of the study, unlikely to be affected. Strong correlations were found between anticipatory pleasure, behavioural activation, and depressive

symptoms, supporting previous findings of associations between anticipating pleasure and behavioural functioning in clinical samples (Buck & Lysaker, 2013; Foussias et al., 2011). It would be useful to include these measures in future studies of depression treatments to track if changes are associated with one another over time. To our knowledge, only one study has assessed trait anticipatory pleasures as an outcome of an intervention, reporting evidence for increases in a small group of schizophrenia patients after a course of structured psychological treatment (Favrod et al., 2010).

The current study was designed to test a plausible mechanism to increase anticipatory pleasure in a clinically-depressed sample. Unsurprisingly then, simply asking depressed individuals to repeatedly engage in EFT activities for positive events that could occur in their daily life did not significantly impact on depressive symptoms. No overall significant change was observed on engagement in the behaviours that were nominated in the EFT activities either. We note that on debriefing from the study, almost all participants reported that although they had not always engaged in the nominated behaviour (e.g., because of practical reasons, plans changing due to circumstances, changing their mind) they felt that it encouraged them to engage in other activities they might not have otherwise. The evidence for an increase on the behavioural activation measure is consistent with this, although this was not a primary outcome of the study and these findings are considered to be preliminary in nature.

There are a number of considerations in regards to the methodology used in this study. Firstly, participants commented that the data-points were too frequent across the day, which made it difficult for them to continue to generate a great variety of activities that they would enjoy. This suggests there are boundary conditions in terms of the level of creativity and diversity in positive activities that individuals will choose in daily life. Reducing the number of data-points in future studies would help rectify this to some degree. All participants indicated that at times they tended to nominate events that they were already engaged, especially if they lasted over several data-points. Although focusing on aspects of tasks they were already engaged in was consistent with the

protocol, this involves reappraisal of a current activity rather than generation of a separate, novel experience. It would be interesting to investigate whether the effects of enhanced EFT activities are different dependent on whether the nominated events are already underway. Different types of EFT activities might be needed to guide individuals to modify their current expectations of their short-term future. It also raises the issue of temporal distance. In this study we focused on daily experiences, but it would be interesting to examine whether EFT activities for more distal events or goals would also have effects on anticipatory pleasure, and goal-directed behaviour.

Although expectations of pleasure are related to decision-making about engaging in behaviours (Mellers, Schwartz & Ritov, 1999), so are other factors that were not assessed in the current study. Perceived control over future events may be a key factor, as an individual's view of how capable they are in enacting a behaviour or experiencing an event is related to their attitudes towards it, and their intention (Armitage & Conner, 2001). Anticipated enjoyment from an event may also be partially contingent on the importance that it has in the context of someone's personal goals or values. Assessing the personal significance of upcoming events would help establish if this moderates any effects. Measuring the strength of different types of feelings that people anticipate they might experience would also be useful. This might be particularly useful in differentiating between activities that produce pleasure due to their hedonistic nature (e.g., eating something delicious, enjoying conversation with another), or positive emotion in the form of pride or achievement, which might be more closely related to self-efficacy (e.g., finishing a set task at work, completing a chore at home). Some instrumental, goal-directed behaviours provide positive emotional reward when their end-point is reached, rather than through the process of reaching an end-point. The category of work/school/study was one of the most frequently chosen, and some of these activities might generally be perceived as producing a sense of achievement rather than being a purely hedonistic pursuit, such as consuming entertainment. Given that engaging in EFT activities appears to be related to self-efficacy, and perceived ability to cope (Brown, MacLeod, Tata, & Goddard, 2002), this indicates the need for ordinal measurement of anticipated feelings of

achievement, pride, accomplished, or the like. Activities geared towards instrumental tasks may benefit from a more nuanced approach to EFT than presented here for breaking down the steps in achieving an end-point.

A further consideration is whether activating mental representations of past positive events in conjunction with generating future events might further enhance anticipatory pleasure.

Reactivating mental representations of experiences that are similar in nature to those that are or could be upcoming might help prime episodic thinking to incorporate these details, increasing the plausibility of an event, and the anticipation of it as being enjoyable. Indeed, engaging in episodic thinking for past events does have effects on detail in episodic future thinking (*removed for blinding*; Madore & Schacter, 2014; Madore & Schacter, 2016), and in one clinical study with schizophrenia patients, the prediction of pleasure from future events was comparable to healthy controls only when first recalling and describing memories (Painter & Kring, 2016). This priming of similar past experiences may therefore be another mechanism through which to enhance EFT and increase anticipatory pleasure.

These findings may have implications for intervention. Behavioural activation treatments currently have good evidence for reducing depressive symptoms (Ekers et al., 2014), and could be potentially enhanced by the addition of EFT activities to promote rich mental representations of positive events that evoke anticipatory pleasure prior to engaging in them. The mobile phone-based approach used in this study was feasible to implement, and may be useful to promote motivation and, ultimately, positively-reinforcing behaviours. In contrast to the current study, this might be particularly effective where the events are pre-planned and factors such as the possible barriers to engaging in it, and the personal significance, have been methodically discussed by the therapist and patient. Interventions using positive mental imagery have already been utilized in the form of cognitive-bias modification to treat depressive symptoms, and reduce anhedonia (Blackwell et al., 2015). Such interventions could also be brought to bear on the daily events in depressed people's lives as a way of increasing their anticipated pleasure for proximal events. In the current study,

future events were simulated only once each. Repeatedly simulating specific experiences that could occur in the future has been shown to increase their emotional valence and plausibility (Szpunar & Schacter, 2013), but conflicting results have been found for dysphoric individuals (Boland et al., 2018). If repeated EFT is tested as a method of further enhancing anticipated pleasure, it may require fewer events to be nominated to avoid burnout or fatigue in relation to thinking about the experiences.

Another possibility is the development of programs to increase generalized specificity and detail in episodic future thinking. Memory specificity training, whereby depressed individuals are taught to recall specific and detailed memories to redress deficits in this ability, has been shown over a series of clinical trials to be effective in reducing depressive symptoms (Eigenhuis, Seldenrijk, van Schaik, Raes, & van Oppen, 2017; Neshat-Doost et al., 2012; Werner-Seidler et al., 2018). Training in specificity and detail in thinking about future events might also be beneficial in helping depressed individuals develop a generalized tendency to simulate positive events using characteristics discussed here that increase the anticipated pleasure. Including physiological measurements of changes in the reward systems of the brain may be another method of assessing the impact of training for EFT. For example, the use of electroencephalography to assess for changes in event-related potentials in areas associated with reward expectation and responsiveness, such as in the parietal lobes, (P2 and P3 areas; Flores, Münte, & Doñamayor, 2015; Novak & Foti, 2015), and increases in asymmetry in frontal lobe activity, favoring the left frontal lobe, which would indicate higher approach motivation (Nelson, Kessel, Klein, & Shankman, 2018; Pizzagalli, Sherwood, Henriques, & Davidson, 2005).

In conclusion, the current study provides some support for the theory that the anticipation and experience of pleasure for upcoming positive events can be increased in those with clinical depression increasing the detail/vividness and use of mental imagery when using episodic future thinking to simulate these events.

References

- Abramson, J. H. (2011). WINPEPI updated: computer programs for epidemiologists, and their teaching potential. *Epidemiologic Perspectives & Innovations*, 8(1), 1. doi:10.1186/1742-5573-8-1
- American Psychiatric Association. (2013). Diagnostic and statistical manual of mental disorders (5th ed.). Arlington, VA: American Psychiatric Publishing.
- Antony, M. M., Bieling, P. J., Cox, B. J., Enns, M. W., & Swinson, R. P. (1998). Psychometric properties of the 42-item and 21-item versions of the Depression Anxiety Stress Scales in clinical groups and a community sample. *Psychological Assessment*, 10(2), 176-181. doi:10.1037/1040-3590.10.2.176
- Armitage, C. J., & Conner, M. (2001). Efficacy of the theory of planned behaviour: A meta-analytic review. *British Journal of Social Psychology*, 40(4), 471-499.
doi:10.1348/014466601164939
- Barsics, C., Van der Linden, M., & D'Argembeau, A. (2016). Frequency, characteristics, and perceived functions of emotional future thinking in daily life. *The Quarterly Journal of Experimental Psychology*, 69(2), 217-233. doi:10.1080/17470218.2015.1051560
- Baumgartner, H., Pieters, R. & Bagozzi, R. P. (2008). Future-oriented emotions: Conceptualization and behavioral effects. *European Journal of Social Psychology*, 38 (4), 685–696.
doi:10.1002/ejsp.467
- Beaty, R. E., Seli, P., & Schacter, D. L. (2018). Thinking about the past and future in daily life: an experience sampling study of individual differences in mental time travel. *Psychological Research*, 1-12. doi:10.1007/s00426-018-1075-7
- Beck, J. S. (2011). *Cognitive behavior therapy: Basics and beyond*. New York, NY: Guilford press.
- Boland, J., Riggs, K. J., & Anderson, R. J. (2018). A brighter future: The effect of positive episodic simulation on future predictions in non-depressed, moderately dysphoric & highly dysphoric individuals. *Behaviour Research and Therapy*, 100, 7-16. doi:10.1016/j.brat.2017.10.010

- Brown, G.P., MacLeod, A.K., Tata, P., & Goddard, L. (2002). Worry and the simulation of future outcomes. *Anxiety, Stress, and Coping*, 15, 1–17. doi:10.1080/10615800290007254
- Bryant, F. (2003). Savoring Beliefs Inventory (SBI): A scale for measuring beliefs about savouring. *Journal of Mental Health*, 12(2), 175-196. doi:10.1080/0963823031000103489
- Buck, B., & Lysaker, P. H. (2013). Consummatory and anticipatory anhedonia in schizophrenia: stability, and associations with emotional distress and social function over six months. *Psychiatry Research*, 205(1-2), 30-35. doi:10.1016/j.psychres.2012.09.008
- Bulté, I., & Onghena, P. (2013). The single-case data analysis package: Analysing single-case experiments with R software. *Journal of Modern Applied Statistical Methods*, 12(2), 450-478. doi:10.22237/jmasm/1383280020
- D'Argembeau, A., & Van der Linden, M. (2004). Phenomenal characteristics associated with projecting oneself back into the past and forward into the future: Influence of valence and temporal distance. *Consciousness and Cognition*, 13(4), 844-858. doi:10.1016/j.concog.2004.07.007
- Eigenhuis, E., Seldenrijk, A., van Schaik, A., Raes, F., & van Oppen, P. (2017). Feasibility and effectiveness of memory specificity training in depressed outpatients: A pilot study. *Clinical Psychology & Psychotherapy*, 24(1), 269-277. doi:10.1002/cpp.1995
- Ekers, D., Webster, L., Van Straten, A., Cuijpers, P., Richards, D., & Gilbody, S. (2014). Behavioural activation for depression; an update of meta-analysis of effectiveness and subgroup analysis. *PloS One*, 9(6), e100100.
- Engel, M., Fritzsche, A., & Lincoln, T. M. (2013). Anticipatory pleasure and approach motivation in schizophrenia-like negative symptoms. *Psychiatry Research*, 210(2), 422-426. doi:10.1016/j.psychres.2013.07.025
- Favrod, J., Giuliani, F., Ernst, F., & Bonsack, C. (2010). Anticipatory pleasure skills training: a new intervention to reduce anhedonia in schizophrenia. *Perspectives in Psychiatric Care*, 46(3), 171-181. doi:10.1111/j.1744-6163.2010.00255

- Flores, A., Münte, T. F., & Doñamayor, N. (2015). Event-related EEG responses to anticipation and delivery of monetary and social reward. *Biological Psychology*, *109*, 10-19. doi: 10.1016/j.biopsycho.2015.04.005
- Foussias, G., & Remington, G. (2008). Negative symptoms in schizophrenia: avolition and Occam's razor. *Schizophrenia Bulletin*, *36*(2), 359-369. doi:10.1093/schbul/sbn094
- Foussias, G., Mann, S., Zakzanis, K. K., van Reekum, R., Agid, O., & Remington, G. (2011). Prediction of longitudinal functional outcomes in schizophrenia: the impact of baseline motivational deficits. *Schizophrenia Research*, *132*(1), 24-27. doi:10.1016/j.schres.2011.06.026
- Gard, D. E., Gard, M. G., Kring, A. M., & John, O. P. (2006). Anticipatory and consummatory components of the experience of pleasure: a scale development study. *Journal of Research in Personality*, *40*(6), 1086-1102. doi:10.1016/j.jrp.2005.11.001
- Gard, D. E., Kring, A. M., Gard, M. G., Horan, W. P., & Green, M. F. (2007). Anhedonia in schizophrenia: distinctions between anticipatory and consummatory pleasure. *Schizophrenia Research*, *93*(1-3), 253-260. doi:10.1016/j.schres.2007.03.008
- Hayes, S. C., Strosahl, K. D., & Wilson, K. G. (2011). *Acceptance and commitment therapy: The process and practice of mindful change*. New York, NY: Guilford Press
- Henriques, J. B., & Davidson, R. J. (2000). Decreased responsiveness to reward in depression. *Cognition & Emotion*, *14*(5), 711-724. doi:10.1080/02699930050117684
- Holmes, E. A., Coughtrey, A. E., & Connor, A. (2008). Looking through rose-tinted glasses? Imagery perspective and positive mood. *Emotion*, *8*(6), 875-879. doi: 10.1037/a0013617.
- Holmes, E. A., Lang, T. J., & Shah, M. (2009). Developing interpretation bias modification as a 'cognitive vaccine' for depressed mood: Imagining positive events makes you feel better than thinking about the verbally. *Journal of Abnormal Psychology*, *118*(1), 76-88. doi: 10.1037/a0012590.

- Holmes, E. A., Mathews, A., Dalgleish, T., & Mackintosh, B. (2006). Positive interpretation training: Effects of mental imagery versus verbal training on positive mood. *Behavior Therapy*, 37(3), 237-247. doi:10.1016/j.beth.2006.02.002
- Hopko, D. R., Lejuez, C. W., Ruggiero, K. J., & Eifert, G. H. (2003). Contemporary behavioral activation treatments for depression: Procedures, principles, and progress. *Clinical Psychology Review*, 23(5), 699-717. doi:10.1016/S0272-7358(03)00070-9
- Jing, H. G., Madore, K. P., & Schacter, D. L. (2016). Worrying about the future: An episodic specificity induction impacts problem solving, reappraisal, and well-being. *Journal of Experimental Psychology: General*, 145(4), 402-218. doi: 10.1037/xge0000142
- Kahneman, D. (1982). The simulation heuristic. In D. Kahneman, P. Slovic, & A. Tversky (Eds.), *Judgment under uncertainty: Heuristics and biases* (pp. 201-208). Great Britain; Cambridge University Press
- Kanter, J. W., Mulick, P. S., Busch, A. M., Berlin, K. S., & Martell, C. R. (2006). The Behavioral Activation for Depression Scale (BADs): Psychometric properties and factor structure. *Journal of Psychopathology and Behavioral Assessment*, 29, 191–202. doi: 10.1007/s10862-006-9038-5.
- King, M. J., Williams, L. A., MacDougall, A. G., Ferris, S., Smith, J. R., Ziolkowski, N., & McKinnon, M. C. (2011). Patients with bipolar disorder show a selective deficit in the episodic simulation of future events. *Consciousness and Cognition*, 20(4), 1801-1807. doi:10.1016/j.concog.2011.05.005
- Kring, A. M., & Caponigro, J. M. (2010). Emotion in schizophrenia: where feeling meets thinking. *Current Directions in Psychological Science*, 19(4), 255-259. doi:0.1177/0963721410377599
- Lovibond, P. F., & Lovibond, S. H. (1995). The structure of negative emotional states: Comparison of the Depression Anxiety Stress Scales (DASS) with the Beck Depression and Anxiety

Inventories. *Behaviour Research and Therapy*, 33(3), 335-343. doi:10.1016/0005-7967(94)00075-U

McFarland, B. R., & Klein, D. N. (2009). Emotional reactivity in depression: diminished responsiveness to anticipated reward but not to anticipated punishment or to nonreward or avoidance. *Depression and Anxiety*, 26(2), 117-122. doi:10.1002/da.20513

MacLeod, A. K., Tata, P., Tyrer, P., Schmidt, U., Davidson, K., & Thompson, S. (2005). Hopelessness and positive and negative future thinking in parasuicide. *British Journal of Clinical Psychology*, 44(4), 495-504. doi:10.1348/014466505X35704

Madore, K. P., & Schacter, D. L. (2014). An episodic specificity induction enhances means-end problem solving in young and older adults. *Psychology and Aging*, 29(4), 913-924. doi:10.1037/a0038209

Madore, K. P., & Schacter, D. L. (2016). Remembering the past and imagining the future: Selective effects of an episodic specificity induction on detail generation. *The Quarterly Journal of Experimental Psychology*, 69(2), 285-298. doi: 10.1080/17470218.2014.999097

Mellers, B., Schwartz, A., & Ritov, I. (1999). Emotion-based choice. *Journal of Experimental Psychology: General*, 128(3), 332-345. doi:10.1037/0096-3445.128.3.332

Nelson, B. D., Kessel, E. M., Klein, D. N., & Shankman, S. A. (2018). Depression symptom dimensions and asymmetrical frontal cortical activity while anticipating reward. *Psychophysiology*, 55(1), e12892. doi:10.1111/psyp.12892

Neshat-Doost, H. T., Dalgleish, T., Yule, W., Kalantari, M., Ahmadi, S. J., Dyregrov, A., & Jobson, L. (2013). Enhancing autobiographical memory specificity through cognitive training: An intervention for depression translated from basic science. *Clinical Psychological Science*, 1(1), 84-92. doi:10.1177/2167702612454613

Novak, K. D., & Foti, D. (2015). Teasing apart the anticipatory and consummatory processing of monetary incentives: An event-related potential study of reward dynamics. *Psychophysiology*, 52(11), 1470-1482. doi: 10.1111/psyp.12504

- O'Connor, R. C., Connery, H., & Cheyne, W. M. (2000). Hopelessness: The role of depression, future directed thinking and cognitive vulnerability. *Psychology, Health & Medicine*, 5(2), 155-161. doi:10.1080/713690188
- Painter, J. M., & Kring, A. M. (2016). Toward an understanding of anticipatory pleasure deficits in schizophrenia: Memory, prospection, and emotion experience. *Journal of Abnormal Psychology*, 125(3), 442-452. doi:10.1037/abn0000151
- Pictet, A., Coughtrey, A. E., Mathews, A., & Holmes, E. A. (2011). Fishing for happiness: the effects of generating positive imagery on mood and behaviour. *Behaviour Research and Therapy*, 49(12), 885-891. doi:10.1016/j.brat.2011.10.003
- Pizzagalli, D. A., Sherwood, R. J., Henriques, J. B., & Davidson, R. J. (2005). Frontal brain asymmetry and reward responsiveness: a source-localization study. *Psychological Science*, 16(10), 805-813. doi:10.1111/j.1467-9280.2005.01618.x
- Renner, F., Ji, J., Pictet, A., Holmes, E., & Blackwell, S. (2017). Effects of engaging in repeated mental imagery of future positive events on behavioural activation in individuals with major depressive disorder. *Cognitive Therapy And Research*, 41(3), 369-380. doi:10.1007/s10608-016-9776-y
- Renner, F., Murphy, F. C., Ji, J. L., Manly, T., & Holmes, E. A. (2019). Mental imagery as a “motivational amplifier” to promote activities. *Behaviour Research and Therapy*, 114, 51-59. [10.1016/j.brat.2019.02.002](https://doi.org/10.1016/j.brat.2019.02.002)
- Saperia, S., Da Silva, S., Siddiqui, I., McDonald, K., Agid, O., Remington, G., & Foussias, G. (2018). Investigating the predictors of happiness, life satisfaction and success in schizophrenia. *Comprehensive Psychiatry*, 81, 42-47. doi:10.1016/j.comppsy.2017.11.005
- Sherdell, L., Waugh, C. E., & Gotlib, I. H. (2012). Anticipatory pleasure predicts motivation for reward in major depression. *Journal of Abnormal Psychology*, 121(1), 51-60. doi:10.1037/a0024945

- Smith, J. D., Borckardt, J. J., & Nash, M. R. (2012). Inferential precision in single-case time-series data streams: how well does the em procedure perform when missing observations occur in autocorrelated data?. *Behavior Therapy*, 43(3), 679-685. doi:10.1016/j.beth.2011.10.001
- Stringaris, A., O'Callaghan, G., Pine, D. S., Leibenluft, E., Vidal-Ribas, P., & Keren, H. (2018). T90. A conceptual and metanalytic review of reward processing in the pathogenesis of depression. *Biological Psychiatry*, 83(9), S163-S164. doi:10.1016/j.biopsych.2018.02.426
- Szpunar, K. K., & Schacter, D. L. (2013). Get real: Effects of repeated simulation and emotion on the perceived plausibility of future experiences. *Journal of Experimental Psychology: General*, 142(2), 323-327. doi:10.1037/a0028877
- Tarlow, K. R. (2017). An improved rank correlation effect size statistic for single-case designs: Baseline corrected Tau. *Behavior Modification*, 41(4), 427-467. doi:10.1177/0145445516676750
- Treadway, M. T., & Zald, D. H. (2011). Reconsidering anhedonia in depression: lessons from translational neuroscience. *Neuroscience & Biobehavioral Reviews*, 35(3), 537-555. doi:10.1016/j.neubiorev.2010.06.006
- Vannest, K. J., & Ninci, J. (2015). Evaluating intervention effects in single-case research designs. *Journal of Counseling & Development*, 93(4), 403-411. doi:10.1002/jcad.12038
- Watson, D., Clark, L. A., & Tellegan, A. (1988). Development and validation of brief measures of positive and negative affect: The PANAS scales. *Journal of Personality and Social Psychology*, 54(6), 1063–1070. doi:10.1037/0022-3514.54.6.1063
- Werner-Seidler, A., Hitchcock, C., Bevan, A., McKinnon, A., Gillard, J., Dahm, T., ... & Rodrigues, E. (2018). A cluster randomized controlled platform trial comparing group MEMory specificity training (MEST) to group psychoeducation and supportive counselling (PSC) in the treatment of recurrent depression. *Behaviour Research and Therapy*, 105, 1-9. doi:10.1016/j.brat.2018.03.004

- Whitton, A. E., Treadway, M. T., & Pizzagalli, D. A. (2015). Reward processing dysfunction in major depression, bipolar disorder and schizophrenia. *Current Opinion in Psychiatry*, 28(1), 7-12. doi:10.1097/ycp.0000000000000122
- World Health Organization. (1992). *The ICD-10 classification of mental and behavioural disorders: Clinical descriptions and diagnostic guidelines*. Geneva: World Health Organization.
- Wu, H., Mata, J., Furman, D. J., Whitmer, A. J., Gotlib, I. H., & Thompson, R. J. (2017). Anticipatory and consummatory pleasure and displeasure in major depressive disorder: An experience sampling study. *Journal of Abnormal Psychology*, 126(2), 149-159. doi:10.1037/abn0000244
- Yang, X. H., Huang, J., Zhu, C. Y., Wang, Y. F., Cheung, E. F., Chan, R. C., & Xie, G. R. (2014). Motivational deficits in effort-based decision making in individuals with subsyndromal depression, first-episode and remitted depression patients. *Psychiatry Research*, 220(3), 874-882. doi:10.1016/j.psychres.2014.08.056
- Yang, X. H., Tian, K., Wang, D. F., Wang, Y., Cheung, E. F., Xie, G. R., & Chan, R. C. (2017). Anhedonia correlates with abnormal functional connectivity of the superior temporal gyrus and the caudate nucleus in patients with first-episode drug-naïve major depressive disorder. *Journal of Affective Disorders*, 218, 284-290. doi:10.1016/j.jad.2017.04.053

P1

P2

P3

P4

P5

P6

P7

Figure 1. Measurements of anticipatory pleasure (“How pleasant/enjoyable do you think it will be to do this activity?”) across the control and enhanced EFT phase, with anticipatory pleasure ratings on the Y-Axis and data-point over the two weeks on the X-Axis.

P1

P2

P3

P4

P5

Figure 2. Measurements of anticipatory pleasure (“How pleasurable/enjoyable is it to just think about doing this activity?”) across the control and enhanced EFT phase, with anticipatory pleasure ratings on the Y-Axis and data-point over the two weeks on the X-Axis.

Table 1.
Participant Characteristics

ID	Age	Gender	Educational Attainment	Relationship Status	Severity of MDD	Duration of MDD Episode (months)	Previous MDD Episodes	Co-morbidities	Medications (all daily doses)
P1	18	Male	High-school	Single	Severe	13	1	/	Sertraline 100mg
P2	24	Male	Bachelor Degree	Single	Moderate	18	2	/	Fluoxetine 20mg
P3	19	Male	High-school	Single	Mild	3	1	Panic Disorder	/
P4	31	Male	Bachelor Degree	Non-married relationship	Moderate	10	1	/	Sertraline 100mg
P5	22	Male	Bachelor Degree	Single	Mild	3	2	/	Escitalopram 10mg
P6	28	Male	Bachelor Degree	Single	Mild	4	0	/	/
P7	26	Male	High-school	Single	Moderate	48	2	Social Phobia	Fluoxetine 60mg

MDD = Major Depressive Disorder, mg = milligrams

Table 2.

Descriptive Statistics, Effects Sizes and P-values for Anticipatory Pleasure Variables

Participant	Control	Enhanced EFT	Control	Baseline Trend	BCT	BCT <i>SE</i>	BCT <i>p</i> -value
	Mean	Mean	Phase SD	Tau			
<i>How enjoyable the upcoming activity was perceived as being</i>							
P1	26.65	32.33	14.04	.038	.285	0.156	.004
P2	22.50	46.30	13.76	.363**	.043	0.169	.667
P3	49.31	69.46	13.06	-.188	.502	0.146	<.001
P4	54.14	69.92	6.89	.362*	.027	0.169	.792
P5	41.90	64.07	23.34	.050	.516	0.145	<.001
P6	14.30	16.59	9.60	-.125	.162	0.167	.119
P7	45.00	53.78	17.60	.000	.171	0.165	.096
<i>How enjoyable it was thinking about the upcoming activity</i>							
P1	38.35	53.74	9.2	.167	.560	0.140	<.001
P2	42.68	56.16	6.35	.331*	.204	0.167	.041
P3	44.17	57.34	9.82	-.005	.499	0.146	<.001
P4	36.74	54.94	7.86	.266	.661	0.127	<.001

ANTICIPATORY PLEASURE FUTURE THINKING

36

P5	56.32	46.27	10.66	-.373**	.395	0.155	<.001
P6	48.51	50.83	10.43	.348*	-.520	0.144	<.001
P7	45.62	51.51	10.82	-.013	.294	0.158	.003

*Significant baseline trend at the $p < .05$ level, which was corrected for. BCT = Baseline Controlled Tau

Table 3.

Descriptive Statistics for Study Variables

	DASS-D	TEPS-A	BADS
Baseline	14.1 (3.2)	37.1 (10.4)	39.5 (15.9)
Between Phases	14.2 (5.5)	36.8 (9.4)	51.8 (11.1)
Post Enhanced EFT Phase	14.4 (4.5)	35.7 (8.4)	45.8 (13.3)

DASS-D = Depression subscale of the DASS-21, TEPS-A = anticipatory pleasure subscale of the Temporal Experience of Pleasure Scale,

BADS = Behavioural Activation Scale for Depression

Supplementary Table 1.

Descriptive Statistics, Effects Sizes and P-values for Manipulation Check Variables

Participant	Control	Enhanced	Control	Baseline	BCT	BCT <i>SE</i>	BCT <i>p</i> -value
	Mean	EFT Mean	Phase SD	Trend Tau			
<i>Detail/Vividness</i>							
P1	26.65	32.33	14.04	.403*	-.444	0.151	<.001
P2	22.50	46.30	13.76	.148	.395	.156	<.001
P3	49.31	69.46	13.06	.034	.523	.144	<.001
P4	54.14	69.92	6.89	.323	.670	.125	<.001
P5	41.90	64.07	23.34	-.332*	.596	.136	<.001
P6	23.29	16.59	9.60	.003	-.201	.166	.052
P7	45.00	53.78	17.60	-.157	.358	.158	.001
<i>Mental Imagery</i>							
P1	19.35	25.41	12.85	.030	.254	.164	.014
P2	9.12	23.70	11.44	.121	.357	.158	.001
P3	43.62	58.03	7.26	.034	.510	.145	<.001
P4	35.75	55.31	7.46	.262	.670	.126	<.001

ANTICIPATORY PLEASURE FUTURE THINKING

39

P5	50.78	49.54	11.03	-.363*	.552	.141	<.001
P6	54.16	47.69	13.88	.064	-.146	.167	.162
P7	46.09	51.35	10.42	-.072	.159	.167	.125

*Significant baseline trend at the $p < .05$ level, which was corrected for. BCT = Baseline Controlled Tau

Supplementary Table 2.

Descriptive Statistics, Effects Sizes and P-values for Secondary Outcomes

Participant	Control	Enhanced EFT	Control	Baseline	BCT	BCT <i>SE</i>	BCT <i>p</i> -
	Mean	Mean	Phase SD	Trend Tau			value
<i>Behavioural Engagement</i>							
P1	1.65	1.58	0.49	.042	-.121	.168	.293
P2	1.27	1.22	0.41	.145	-.014	.169	.908
P3	/	/	/	/	/	/	/
P4	1.50	1.27	0.44	.007	-.210	.165	.061
P5	1.43	1.06	0.45	.118	-.182	.166	.107
P6	1.18	1.45	0.37	-.187	.381	.156	.001
P7	1.13	1.30	0.32	-.250	.335	.159	.003
<i>Mood- Sad</i>							
P1	43.28	52.15	15.35	-.118	.207	.165	.054
P2	47.55	52.07	10.96	.000	.196	.166	.057
P3	54.13	44.81	11.67	-.240	-.353	.158	.001
P4	43.14	52.55	12.70	.346*	-.511	.145	<.001

ANTICIPATORY PLEASURE FUTURE THINKING

41

P5	46.71	51.95	11.41	.081	.228	.165	.028
P6	55.26	47.08	11.74	-.281	-.314	.160	.002
P7	56.60	47.71	13.67	-.222	-.215	.165	.044
<i>Mood - Happy</i>							
P1	52.29	49.26	13.67	-.025	-.033	.169	.770
P2	52.43	47.96	11.22	.095	-.199	.166	.055
P3	50.77	49.03	12.02	-.256	-.117	.168	.262
P4	48.50	50.56	11.32	.000	.114	.168	.276
P5	56.51	46.16	13.68	-.230	-.271	.163	.014
P6	50.07	49.96	12.73	-.168	.065	.169	.545
P7	54.72	48.37	11.54	-.013	-.263	.168	.011

Note: for behavioural engagement 1 = engaged in nominated upcoming activity, 2 = did not engage. *Significant baseline trend at the $p < .05$ level, which was corrected for. BCT = Baseline Controlled Tau