A Network Perspective on Successful Aging and Resilience in Later Life: A
Protocol Paper

Lotte P. Brinkhof ^{a,b,d*}, K. Richard Ridderinkhof ^{a,d}, Sanne de Wit ^{a,d}, Harm J. Krugers ^{c,d},
Jaap M. J. Murre ^{a,d}

^aDept. of Psychology, Faculty of Behavioural and Social Sciences, University of Amsterdam,
Amsterdam, Netherlands;

^bCentre for Urban Mental Health, University of Amsterdam, Amsterdam, Netherlands;

^cSwammerdam Institute for Life Sciences, Cognitive Neuroscience, Faculty of Science,
University of Amsterdam, Amsterdam, Netherlands;

^dAmsterdam Brain & Cognition (ABC), University of Amsterdam, Amsterdam, Netherlands

*Corresponding Author: E-mail: l.p.brinkhof@uva.nl, Address: Nieuwe Achtergracht 129,
1001 NK Amsterdam, Netherlands.

This paper has not been peer reviewed.

Abstract

Background. Aging inevitably gives rise to many late-life challenges and transitions that impose threats for one's physical and mental health, which can greatly impact our well-being and quality of life. Hence, the key to successful aging may not be the absence of these challenges and transitions, but the ability to demonstrate resilience to them. We recently started a large-scale study ($N > 10.000$) among older adults (55 years or older) in the Netherlands, with the aim to understand resilience as an emergent property that arises through the interplay between various (risk and protective) factors from different domains.

Method. Participants are asked to complete online questionnaires and tests that cover a multitude of relevant factors from multiple domains (e.g., physical, psychological, cognitive, social, environmental). Relationships between those factors will be analysed through network analysis, in which conditional dependencies between factors are depicted in a network of nodes. In this way, critical resilience (protective) and risk factors can be identified.

Discussion. This study offers a unique possibility to obtain insights into the factors that characterize and contribute to resilience in old age and thereby assist in generating hypotheses for follow up work. The contribution of this study in the growing field of resilience research, as well as the limitations, are discussed accordingly.

Keywords: resilience, successful aging, quality of life, well-being, network analysis

Background

As the rapid graying of the world's population continues, a major challenge is to promote successful aging. *Successful aging* is commonly viewed as a multidimensional concept, that encompasses domains of physical, cognitive, psychological and social health (1, 2). However, retaining high and stable levels of functioning across all domains might be an unattainable goal for the majority of people approaching old age. Many age-related changes affect physical and cognitive functions (3–7), leading to increased risks of acquiring chronic diseases (e.g., dementia, osteoarthritis), physical disabilities and mental health problems (e.g., 8–10). Moreover, older adults face particular late-life challenges (e.g., loss of spouse, loneliness) that further impose a threat for their physical and mental health status (11,12). Accordingly, it has been argued that there may be different *successful aging phenotypes*, referring to the variation in the combination of aspects of successful aging in which individuals show more and less favorable trajectories of functioning (13). As such, the absence of disease, illness or disability is not imperative and older adults may also age successfully despite physical or cognitive limitations, as long as these are well controlled and therefore have little influence on well-being and quality of life (14–16). In line with this, we endorse the perspective of the World Health Organization (WHO) and consider successful aging as *the process of developing and maintaining the functional ability and competence that enables well-being and quality of life in older age* (15). This places a stronger emphasis on subjective perceptions and addresses individuals' capacities to thrive and flourish in the face of adversity, something that successful aging may crucially depend on (17–21). *Resilience* in late life can be conceptualized as a protective characteristic that can impact the process of successful aging by maintaining functional ability or competence, well-being, and quality of life despite the presence of internal (e.g., decline, disease) or external (e.g., loss of spouse) challenges or adversity (22–26).

63 In previous studies, high resilience has been positively associated with multiple
64 beneficial outcomes, such as quality of life, well-being, happiness, life satisfaction, mental
65 health, and the subjective experience of successful aging (21,27–33). Furthermore, it has been
66 negatively associated with depression and mortality risk (34–37). High resilience may also
67 support desirable lifestyle habits (38–41).

68 While an increasing number of studies has identified factors that contribute to
69 resilience (e.g., adaptive coping styles, optimism, emotional stability, being physically active,
70 social involvement, green space, working memory capacity; e.g., 20,42–48), little is known
71 about the interplay between various protective and risk factors. We recognize that the field
72 must move beyond capturing resilience in one personality characteristic or trait (49), but
73 rather focus on the evaluation of resilience in terms of potential health outcomes (22,50–52),
74 characterized by multiple risk and protective factors. To take on this challenge, we will adopt
75 a network perspective, which proposes that resilience is an emergent property that arises
76 through interactions among such factors and is captured within several outcome variables that
77 are crucial to (the definition of) successful aging (e.g., quality of life, (mental) well-being).
78 This is in line with the idea that resilience should be understood as an interrelated system of
79 multiple (resilience) factors (e.g., 53–55). We extend this perspective by looking at the
80 interplay with potential risk factors as well (e.g., living alone, history of depression).
81 Altogether, this allows us to infer an individual's response to certain internal or external
82 perturbations. When a certain event or challenge (e.g., loss of friend) interrupts a central
83 factor (e.g., social support), this may ignite a cascade of (aversive) symptoms (e.g., loneliness,
84 depression). If these changes are not sufficiently controlled or managed by other (protective)
85 factors in the network (e.g., coping ability), the network will be dominated by (a critical
86 number of) dysfunctional connections, and resilience to perturbation will be lost. This should
87 negatively impact successful aging, as reflected in low well-being and quality of life.

88 This study sets out to reveal unique, and perhaps unsuspected patterns of associations
89 between demographic, social, environmental, psychological, behavioral, psycho-social and
90 cognitive factors that contribute to seniors' resilience, either positively or negatively. To this
91 end, we will collect data, focusing on the aforementioned domains, of at least 10.000 seniors
92 (55 or older) using an online inventory. Through large-scale network analyses, we aim to
93 explore the factors and interaction that are central to successful aging and underpin resilience
94 in later life. Importantly, unlike other network studies that aim to characterize resilience in
95 terms of stability of the network (e.g., by looking at the number and strength of connections,
96 with sparse networks being considered more resilient; see e.g., 56,57), we specifically focus
97 on the qualitative patterns of interplay between (1) QoL and (mental) well-being as outcomes
98 of resilience and (2) empirically-defined risk and protective factors that are thought to affect
99 resilience in later life. Conceptualizing psychological constructs, such as resilience, as a
100 complex interplay of factors has gained increasing popularity in various research fields (e.g.,
101 53–55,57,58). However, to the best of our knowledge, this is the first time that a network
102 approach is used to model resilience in later life based on a multitude of different factors, in
103 relation to proposed outcomes of resilience.

105 **Methods**

106 *Sample Characteristics*

107 This study aims to recruit 10.000 to 30.000 participants via an online inventory. Participants
108 are eligible if they are 55+ years, living in the Netherlands and have no Alzheimer's Disease,
109 vascular dementia, frontotemporal dementia, Lewy body dementia or other dementia
110 diagnosis. Moreover, insufficient command of Dutch (i.e. having trouble understanding the
111 Dutch language when reading newspapers or brochures), impaired vision or not being able to
112 perform the operations, for example due to severe symptoms expression of a

neurological/neurodegenerative disease such as Parkinson, amyotrophic lateral sclerosis (ALS), multiple sclerosis (MS) or rheumatism or any other physical or mental impairment, required to successfully use a computer or laptop independently (i.e., mouse clicks, pressing keys on the keyboard) are also exclusion criteria.

We strive to include a representative sample that reflects ethnic and urban diversity in the Netherlands, by recruiting participants from both (predominantly) rural areas (e.g., mostly from the least densely populated provinces: Groningen, Friesland, Drenthe and Zeeland) as well as large cities (e.g., Amsterdam, Rotterdam, Utrecht and Den Haag). Six-digit postal codes will be collected to assess the residence of the participants and command of the Dutch language is checked by means of a self-report question during the inventory (see operationalization section; 59)

Operationalization

Online registration. Interested individuals can register online via www.seniorendoenmee.nl, where they will be asked some demographic information, including name, email address, their six-digit postal code, age and gender. In addition, they are asked to declare that they meet the entry requirements (see Sample Characteristics), the registration form has been completed truthfully, the privacy statement has been read and they agree to the storage and processing of the data provided by means of the form in accordance with the privacy statement. Finally, they are asked to indicate whether they are interested to be kept informed about the progress of the study by means of a newsletter.

Participation. Participants receive an email immediately after registration, which includes a link that provides access to an information brochure. This information brochure once again emphasizes the importance of meeting the entry requirements criteria and explains that the

inventory should be completed independently. That is, assisting individuals (e.g., caregiver, family member or technician) may help to explain how the mouse or arrows should be manipulated or facilitate the typing or entering of responses, but are not allowed to provide aid or hints to the participants in order to obtain desirable and accurate results. After that, participants are instructed to explicitly indicate that they meet the entry requirements (for each requirement independently) and to give their informed consent.

Design. The main inventory consists of three parts of questionnaires and (psychological) tests that are delivered online through Qualtrics (questionnaires; www.qualtrics.com), NeuroTask Scripting (tests; www.scripting.neurotask.com) and is regulated by the LOTUS platform. LOTUS is a webtool developed and managed by the University of Amsterdam and is used to guide participants automatically through the different parts of the study. Each part takes approximately 20 to 30 minutes to complete. The first part starts automatically after participants have given their online consent. We have taken precautions to minimize the effects of negative and positive framing by, for example, presenting questionnaires about depression and anxiety prior to a psychological or cognitive test, such that answers on questions later in that package will likely not be influenced by the previous questions anymore. At the end of each part, participants are given the option to 1) start directly with the next part, 2) take a (short) break and continue afterwards or 3) resume the study the next day.

Automatically sent emails are implemented in the design of the study, with the intention to either inform participants that they can start or resume with the inventory or to remind them that they did not finish a certain part of the inventory yet.

162 **Materials**

163 A detailed overview of the materials can be found in the Supplemental A. A list of all the
 164 variables included is shown in Table 1.

165

Domain	Questions/tests
<i>Demographics</i>	Age, gender, socio-economic status (e.g., level of education, employment status), relationship status, nationality, language, religion, neighborhood ^a (e.g., urbanization grade, distance to daily needs, walkability index, COVID-19 reports ^b , temperature ^c).
<i>Mental Wellbeing and Quality of Life</i>	Happiness Scale (60), Warwick-Edinburgh Mental Well-being Scale (61,62), Centre of Epidemiological Studies Short Depression Scale (Andresen et al., 1994), questions about the history of depression, the World Health Organization Quality of Life Instrument-Older Adults Module (64,65), the Hospital Anxiety and Depression Scale – Anxiety (66,67) and questions about the history of anxiety.
<i>Resilience and coping</i>	The Brief Resilience Scale (68,69), and the Positive Appraisal Style and Behavioural Coping Scale (51,70,71).
<i>Stress</i>	The Perceived Stress Scale (72,73), a composite list of major life events (74,75) and COVID-19 related stressors (51).
<i>Personality</i>	The Ten Item Personality Measure (76,77), the General Self-efficacy Scale (78,79), the Rosenberg Self-Esteem Scale (80,81), the Self-Management Ability Scale (82) and the Brief Self-control Scale (83–85).
<i>Social interactions</i>	The Loneliness Scale (86), questions regarding the number of (social) contacts and interactions, questions regarding participation in social activities, the Social Support List (Discrepancies; 122)
<i>Health</i>	The Short-Form Health Survey-36 (88,89), questions regarding physical and neurological diseases.
<i>Lifestyle</i>	Questions regarding subjective lifestyle quality, length and weight, diet, physical activity, electronica use, mental health disorders and

	smoking status, the Pittsburgh Sleep Quality Index (90), the Alcohol Use Disorders Identification Test (91–93)
<i>Living</i>	Questions regarding the living situation and satisfaction with both the living situation and neighbourhood, transport mobility.
<i>Other</i>	The Aging Perceptions Questionnaire (94), Perceived Ageism Questionnaire (self-generated), questions regarding boredom, domestic animals and holidays, the Prospective and Retrospective Memory Questionnaire (95–98), questions regarding the subjective impact of COVID-19) and the Stringency Index ^d .

Table 1. A list of variables included in the study.

^a Six-digit postal code (PC6) of subjects' addresses will be used to link environmental data to other study variables. The data will be retrieved from national registration organisations (for years ranging from 2015 to 2020), such as Statistics Netherlands or the Institute for Public Health and the Environment (RIVM), and will be initiated by the Geoscience and Health Cohort Consortium (GECCO).

^b The number of COVID-19 reports per municipality (over the last 14 days in the Netherlands) will be matched to each participant using the municipality codes of Statistics Netherlands.^{1,2}

^c The average temperature at the day(s) of participation per province will be taken from the KNMI (Royal Netherlands Meteorological Institute).³

^d The stringency Index of the Oxford COVID-19: Government Response Tracker (OxCGRT)⁴ will be used to assess the strictness of 'lockdown style' policies that primarily restrict people's behaviour in the Netherlands at the time of participation

Analyses

Overview. The main analyses can be divided into two parts: (1) the small-scale and (2) the large-scale network analysis.

¹ <https://data.rivm.nl/geonetwork/srv/dut/catalog.search#/metadata/1c0fcd57-1102-4620-9cfa-441e93ea5604?tab=relations>

² <https://www.cbs.nl/nl-nl/onze-diensten/methoden/classificaties/overig/gemeentelijke-indelingen-per-jaar/indeling%20per%20jaar/gemeentelijke-indeling-op-1-januari-2020>

³ <http://projects.knmi.nl/klimatologie/daggegevens/selectie.cgi>

⁴ <https://www.bsg.ox.ac.uk/research/research-projects/coronavirus-government-response-tracker>

- (1) The small-scale network analysis will include 30 factors that are selected based on their theoretical relevance (underlined variables in appendix A, e.g., happiness, mental wellbeing, depression, quality of life, anxiety, positive appraisal style, perceived stress, loneliness). The inclusion of only 30 factors will enable us to estimate a network with a relatively small sample (see ‘Justification Sample Size’ for details), such that findings can be used to assist in generating hypotheses for follow up work and the development of interventions aimed to promote successful aging early in the process.
- (2) The large-scale analysis will include all 83 main factors (numbered variables, listed in appendix A).

Network estimation. The R-packages **mgm** and **bootnet** will be used to estimate pairwise (undirected) Mixed Graphical Models (MGMs) at group level (11,99–101), allowing us to model each relevant factor on its proper domain (i.e., Categorical, Gaussian and Poisson). Lasso regularization will be applied to reduce the inclusion of spurious edges, resulting in a network that is easier to interpret and has a higher specificity (102,103). This penalty will be weighted by a hypertuning parameter (gamma) of 0.25, which will be selected using the conservative Extended Bayesian Information Criterion (EBIC). We will estimate pairwise relations controlling for all other factors in the network (concentration network), hence, edges represent partial correlations or multiple regression coefficients (connection with categorical variables). If there is no edge between two nodes, they are conditionally independent given all other nodes in the network.

The network will be visualized using the R package **qgraph** (104), where nodes represent factors and edges pairwise interactions between two nodes controlled for all other nodes in the network (i.e., partial correlations). This will give us insights into the strength and

value (positive – green; negative – red) of the connections between the different nodes (see Figure 1), with the thickness and saturation of the edges signifying the strength of the relationships

The stability of the edge weights will be assessed by using **non-parametric bootstrapping** (using resampled data with replacement, 1000 samples; 66–68). Here, confidence intervals are constructed around the edge weights, with narrow intervals reflecting more stable edge-weight estimates. A **bootstrapped difference test** will be performed to identify whether the edge weights significantly differ from each other. For each to-be-compared edge pair, the difference between the bootstrapped edge weights is calculated and a confidence interval around this difference score is constructed. Edges are deemed to be significant, if zero is not in the confidence interval.

[INSERT FIGURE 1]

Nodewise predictability. The nodewise predictability (i.e. how well a given node is predicted by all its neighbouring nodes in the network: e.g., proportion explained variance) will also be computed and visualized in the network using the R packages **mgm** and **qgraph** (101,104,108). Predictability provides us with a measure of the extent to which a node is determined by its neighbours (109). For example, we will identify to what extent a certain node (e.g., loneliness) can possibly be changed by intervening on the nodes directly associated with it (e.g., social contacts), but also to what extent nodes are not explained by factors included in the network. When the node-wise predictability is low, there might be other variables, that are not included in the network, that interact with this node.

Identification of communities within network structure. Since interpreting individual interactions in large networks is challenging, we will further analyse the structure of the network by identifying strongly connected subgraphs (or communities) in the network (110). Despite the presence of theoretically defined communities (e.g., mental health, physical health, lifestyle, etc.), we are particularly interested in identifying empirically derived communities and the differential roles of the nodes within and between them. That is, psychological construct with many and strong connections within a certain community could be thought of as the core of a community, whereas factors that belong to multiple communities facilitate the communication between certain domains (see *Local Structure Analysis* for more information on how the differential roles will be established). We will use the CliquePercolation R-package to run the clique percolation algorithm for weighted networks (111–113), and identify k -cliques (i.e., fully connected subgraphs, or communities, with k nodes). This method allows for the identification of nodes that belong to multiple communities (i.e., cross-loading), rather than only one. The edge list of the estimated MGM will be used to optimize k (i.e., the smallest clique size, with $k = 3$ as minimum required by the algorithm) and I (i.e., intensity threshold; how strong the average relation among nodes within a community needs to be in order to be detected as community) using the `cpThreshold` function (for more details see 111,114). The optimal k will be chosen by visually inspecting the community networks and choosing k that corresponds to the community network that does not include a giant component to which most nodes belong, but rather has communities of smaller sizes.

Local Structure Analysis. To examine the local structure of the communities, we will apply a similar approach as described by Blanken and colleagues (114). The edge weight values a certain node has within its community is summed and defined as the **stabilizing index**; and

between its community is summed and defined as **communicating index**. These indices provide specific insights of the factors that play a crucial role at the local level (within and between sub-networks).

We will also perform some additional exploratory analyses, where we will zoom into the subscales of some factors (i.e., QOL, SMA, LS, SF, APQ, PRMQ, see appendix for explanation of these nodes). These analyses will be performed on the community level, where we will estimate separate MGMs for each community and only include those nodes that have been related to the multi-dimensional factor (e.g., QOL) in the primary model. The subscales or dimensions of the multi-dimensional factor itself will be included as separate nodes (e.g., QOL1, QOL2, QOL3, etc.). This approach will reduce the number of nodes included in each model, thereby decreasing the complexity of the estimated models and improving the power of the analysis. This enables us to identify how subscales or dimensions relate to other factors within the same community. Importantly, these analyses will not allow us to say anything about the interplay of the subscales with nodes related to other communities. Although generalization to the primary model may seem compelling, the estimated network structure of each community will change as a consequence of the altered set of variables and will therefore not exactly resemble the structure and corresponding edge weights of nodes within and between communities.

Moderated Network Models. Moderated Mixed Graphical Model (or Moderated Network Models; MNMs) will be estimated to examine differences across networks of subgroups with favourable and less favourable predispositions or outcomes (115). The **mgm** package estimates these MNMs with a nodewise regression approach that allows us to evaluate all moderation effects involving a specific categorical (e.g., living alone) or continuous (e.g., QoL) variable. A positive moderation effect on the relationship between A and B would

indicate that if one increases the value of the moderator, then the pairwise interaction between A and B becomes stronger. In case a moderation effect is found, we will condition on (fix the values of) the moderator. In case of a categorical moderator, the predefined categories are used to fix the values of the moderator. For continuous variables, we will choose the condition values based on the scaled version of the moderator, creating different subgroups. This allows us to distinguish different pairwise MGMs for each conditioned value and subsequently compare the strengths of the edges between the different networks. The stability of the estimates will also be assessed. We will test for moderation effects of our main outcome variables (i.e., mental well-being and quality of life), as well as other categorical and continuous variables (e.g., urbanization grade) that seem relevant based on the network and community structures and outcomes.

Justification sample size

In the current methodological landscape, it is beyond the bounds of possibility to precisely determine the required sample size needed to accurately estimate the network structure. That is, to perform a power analysis for networks, one must have a prior expectation of the weighted (true) network, including the network structure and the strength of the non-zero edges. Since our dataset will include a unique combination of variables, in terms of both the distributions (i.e., combination of categorical and gaussian) and the theoretical level (e.g., social, environmental, psychological, cognitive, etc.), there is no comparable dataset that we can use to estimate the expected network structure. Hence, it is impossible to accurately assess the sensitivity (true-positive rate), specificity (true-negative rate) and correlation between edge weights of the true network and edge weights of the estimated network given certain sample sizes in advance (116,117).

On that account, we aim for large samples, where we will recruit 2000 participants for the small-scale analysis and at least 10.000 observations for the large-scale analysis. Subsequently, we will use post-hoc stability analysis to evaluate the accuracy and stability of the estimated network parameters, as described under ‘Network estimation’.

Data exclusion and missing data

As seniors are sometimes reluctant to discuss the possibility of cognitive impairment and dementia with their general practitioner, some participants may have suspicious cognitive complaints without having a diagnosis. That is, however, beneficial for our sample, as we want to ensure a sample representative of the elderly in society, which includes a large group that show (mild) cognitive impairments or dementia-like forms without having a diagnosis (and perhaps even without subjective complaints). Still, we will exclude data from participants that show considerable deviation from performance and answers of most participants, reducing the possibility that our sample includes participants with severe memory impairments. After visual inspection of the distribution of the variables, highly skewed variables will be subjected to the robust median absolute deviation (MAD) outlier detection mechanism (118), with a threshold of 2.5.

If there is any indication of a change in circumstances during an extended break between sessions that may influence the data, participants will be excluded from analyses as well. Moreover, the MAD will also be used to exclude participants that completed the inventory in an extreme short time.

MGM estimation algorithms cannot handle missing data. For that reason, we will perform imputations in case of Gaussian or Poisson missing variables. Since imputations are not reliable for categorical variables, participants that miss data for those variables will be excluded from all analyses entirely.

Discussion

Research on aging has indicated that successful aging is potentially attainable for many individuals (14,119,120). However, aging inevitably gives rise to certain late life challenges and transitions (e.g., 3,7,81,82), and retaining high and stable levels of functioning across all indicators of successful aging might be a challenge to most people. The key to thriving throughout our later years may therefore not be the absence of these challenges and transitions, but the ability to demonstrate resilience against them. This should allow individuals to maintain functional ability or competence that enables well-being and quality of life.

The current study uses a network approach to understand resilience in later life as an emergent property that arises through associations among a multitude of (protective and risk) factors from different domains. Our analyses allow us to identify the contribution of each factor to specific outcomes that are important to successful aging (i.e., quality of life and well-being), as well as how the risk and resilience factors influence each other. In this manner, all parts of the network act together and determine the overall level of resilience, where some factors enhance resilience and others may contribute to tipping the balance towards vulnerability in the face of adversity. An important assumption of our perspective on resilience is that, while resilience may be determined in part by static traits, predispositions, habits, and/or fixed contextual backgrounds, at least some factors underlying resilience are subject to change (e.g., by promoting through interventions). This in turn suggests that even individuals with (some) unfavorable personality traits, experiences, or backgrounds, may be able to demonstrate resilience, as long as the balance between factors that promote and threat resilience is preserved. This is in line with the suggestion that resilience is possible for most older adults, including those with cognitive impairments (20,123). Finally, the emergent nature of resilience offers opportunities to optimize it, by targeting those features that are

instrumental to reach and maintain resilience at old age. This study may therefore not only assist in generating hypotheses for follow-up work, but also in the development of interventions aimed to promote successful aging.

Obtaining results that can be generalized to the entire older population is of great importance, but there are some limitations in our study that may hamper this. Apart from the individuals with severe physical disabilities and memory impairments that are not eligible to participate (see *Sample Characteristics*), our study may also preclude a specific group of older adults that is not involved with technological communication or does not have the required knowledge and confidence to participate in online research. This may give rise to self-selection bias or non-random attrition (124,125). To minimize the manifestation of this, we created an explanatory video in which we demonstrate how to sign up for the study and which (minimal) steps need to be taken to start. Moreover, participants are automatically directed to a consecutive part of the study when they have finished the previous set of questions or tests. We ensured that the instructions were complete and coherent, and we offer assistance via e-mail 7 days a week to help participants who face any problems. Still, there may be circumstances in which people encounter technical problems due to outdated or suboptimal operating systems that cannot be solved by us.

The representativeness of the Dutch senior's population may also be affected by the required command of the Dutch language. It was estimated that older migrant adults made up 10.2% of the aging population (55 and older) in January 2020 (126). However, participants with a migration background are generally harder to reach, due to language and cultural barriers, as well as lack of interest and mistrust of the research process (127). This may have implications for the generalizability and implementations of the results in terms of ethnic backgrounds. Although we may ultimately not be able to obtain an exact representation of the multi-cultural population in terms of proportions, we strive for enough diversity among

participants to be able to distinguish subgroups with different backgrounds. The same applies to groups of lower socioeconomic status (SES), as we expect this subgroup to be underrepresented as compared to groups of higher SES as well (128).

Finally, any online platform may introduce some uncertainty with respect to the reliability and validity of the results of our measures. Indeed, we have used validated and reliable scales and instruments as much as possible, but participants still have to read the instructions carefully to complete our questions and tests successfully. Some may not pay sufficient attention (e.g., distracted or tired), which could lead to misunderstandings with respect to the actions that must be performed. While a test leader could assist in case of uncertainties in a lab study, receiving additional help requires more effort in online research. In the current study, participants can ask questions about the inventory via e-mail. To minimize the likelihood of questions arising about the cognitive or psychological tests, we integrated practice sessions, in which reminders would pop up while putting the instructions into practice. Nevertheless, it remains important to carefully analyze the results of these tests. Despite the explained disadvantage of a test leader not being present, it may also reduce the likelihood of participants giving socially desirable answers and thereby improve the quality of the data.

The fact that participants can complete the study at home and that there are no geographical boundaries, is a considerable advantage for aging research in general, but is especially convenient with regard to the global COVID-19 pandemic during which this study will be conducted. Nevertheless, the pandemic will unavoidably impact our results. Our study has a strong focus on quality of life and mental well-being, but also includes factors such as social participation and daily living. Those factors are doubtlessly influenced by the pandemic, although the extent of the impact likely differs per individual (129–131). Moreover, the moment of participation may also play a major role, since there is great

variance in the levels of spread and severity of the restrictions over time. An attempt to (at least partially) correct for the latter concern has been made by including the Stringency Index of the OxCGRt, assessing the strictness of lockdown style policies that primarily restrict people's behavior, in our analyses. We expect that higher levels of this index are accompanied by more substantial (objective) impacts on mental health and behavior. The subjective impact of the pandemic on answers given during the study will also be assessed and may further inform us about the consequences for our results and help to interpret the generalizability of our results.

In light of this study, it is important to discuss some critical points regarding resilience research in general. Indeed, it has increasingly been emphasized that resilience can only be understood in the context of responding to significant adversity (132–134). It has been proposed that (one aspect of) resilience pertains to “a good *mental health* outcome following an adverse life event or a period of difficult life circumstances” (132). Accordingly, the need for (prospective) longitudinal studies has been expressed, with multiple measurements, preferably before and after a certain life event. Since our study will examine resilience using a cross-sectional design, it is crucial to emphasize that we do not aim to *measure* resilience. Instead, we plan to *characterize* resilience in terms of relations between factors that influence each other and thereby obtain a picture of the central factors (e.g., health condition, SF-36) that in the event of an interruption (e.g., cerebral haemorrhage or infarction) may result in a cascade of changes within the network, unless certain protective factors (e.g., strong social network, coping ability) may prevent or counteract a transition into a vulnerable phase. As such, we consider age-related challenges, such as decline in physical or cognitive function, loss of partner or lack of personal autonomy, as adversity and aim to explore those factors that could determine whether individuals that experience such challenges are still capable of maintaining good quality of life and well-being. In other words, we mainly focus on the

aspect of resilience that relates to *withstanding of adversity* (preventive), rather than *bouncing back* from it.

This naturally leads to the question as to whether we can identify resilience factors among those who did not experience any age-related adversity (yet). In fact, our analysis approach still allows us to see which (connections between) nodes may be central and, for instance, ignite a chain of symptoms when triggered by some perturbation. Moreover, it also enables us to infer what factors could predict whether someone will or will not succumb to adversity. Suppose we would find a certain pattern of associations between health condition, stress, and depression as shown in Figure 2, where a good health condition is associated with reduced stress, and stress is associated with increased levels of depression. If a strong (adaptive) coping ability would additionally be associated with reduced stress and depression, this could indicate that coping ability is an important *protective* factor that reduces the impact stress can have on depressive symptoms when someone would experience a major health setback, and also tempers depressive symptoms directly. Obviously, one should be cautious in interpreting such results, as causality cannot be inferred from a cross-sectional concentration network. Some associations or chains may even be bi-directional or part of a feedback loop. While positive thinking may, for instance, enhance adaptive coping strategies and therefore lead to higher levels of quality of life, quality of life may also impact the extent to which someone can think positively. Plainly, the distinction between factors that have been claimed to characterize resilience (e.g., positive thinking) or factors that are commonly seen as consequences of high resilience (e.g., good quality of life) can be quite arbitrary. This study will add insights to this interplay and can help us to generate hypotheses for follow-up work.

[INSERT FIGURE 2]

Conclusion

Using a network approach, this study offers a unique possibility to obtain insights into the interactions between factors that characterize and underpin resilience in old age. We acknowledge that successful aging may crucially depend on the ability to thrive in the face of adversity and that the ability to demonstrate resilience may distinguish those who are aging successfully from those who are not.

List of abbreviations

WHO: World Health Organization
ALS: amyotrophic lateral sclerosis
MS: multiple sclerosis
PC6: six-digit postal code
RIVM: the Institute for Public Health and the Environment
GECCO: Geoscience and Health Cohort Consortium
KNMI: Royal Netherlands Meteorological Institute
OxCGRT: Oxford COVID-19 Government Response Tracker
MGM: Mixed Graphical Model
EBIC: Extended Bayesian Information Criterion
CI: confidence interval
CS: correlation stability coefficient
QOL: Quality of Life
SMA: Self-management ability
LS: Loneliness Scale
SF: health condition
APQ: Ageing Perceptions Questionnaire

481 PRMQ: Prospective and Retrospective Memory Questionnaire

482 MNM: Moderated Network Model

483 SES: socioeconomic status

484 MAD: median absolute deviation

485

Declarations

Ethics approval and consent to participate

The study is approved by the local ethics committee of the University of Amsterdam (2020-DP-12556) and will be conducted in accord with relevant laws and institutional guidelines.

Consent for publication

Not applicable

Availability of data and materials

The datasets used and/or analyzed during the current study will be made publicly available after completion of the project, and will until that time be available from the corresponding author on a collaboration basis upon reasonable request.

Competing interests

The authors declare that they have no competing interests

Funding

This research was funded by the Centre for Urban Mental Health, a Research Priority Area at the University of Amsterdam.

Authors' contributions

J.M. and R.R. conceived of the original idea of the network perspective; L.B. conceived of and developed the presented idea and corresponding analytical framework. L.B., J.M., R.R., S.W., and H.K wrote the manuscript, with L.B. taking the lead. All authors provided critical

feedback and helped shape the idea, protocol, and manuscript; all authors approved the manuscript.

Acknowledgements

We thank Karoline Huth and Julian Burger (both University of Amsterdam, Dept. of Methodology) for assistance in selecting the (appropriate) analytical methods, and the anonymous pilot participants for their constructive feedback on the content and design of the study during our pilot-phase.

Authors' information

L.B.: PhD-student

J.M.: expert in neural networks

R.R.: expert in cognitive aging

S.W.: expert in habit-related intervention

H.K.: expert in neurobiology of aging

Footnotes

[1] <https://data.rivm.nl/geonetwork/srv/dut/catalog.search#/metadata/1c0fcd57-1102-4620-9cfa-441e93ea5604?tab=relations>

[2] <https://www.cbs.nl/nl-nl/onze-diensten/methoden/classificaties/overig/gemeentelijke-indelingen-per-jaar/indeling%20per%20jaar/gemeentelijke-indeling-op-1-januari-2020>

[3] <http://projects.knmi.nl/klimatologie/daggegevens/selectie.cgi>

[4] <https://www.bsg.ox.ac.uk/research/research-projects/coronavirus-government-response-tracker>

References

1. Kim SH, Park S. A Meta-Analysis of the Correlates of Successful Aging in Older Adults. *Res Aging* [Internet]. 2017 Jun 1 [cited 2020 Sep 9];39(5):657–77. Available from: <https://journals.sagepub.com/doi/abs/10.1177/0164027516656040>
2. Urtamo A, Jyväkorpi SK, Strandberg TE. Definitions of successful ageing: A brief review of a multidimensional concept [Internet]. Vol. 90, *Acta Biomedica*. Mattioli 1885; 2019 [cited 2020 Sep 9]. p. 359–63. Available from: [/pmc/articles/PMC6776218/?report=abstract](https://pubmed.ncbi.nlm.nih.gov/3276218/)
3. Glisky EL. Changes in Cognitive Function in Human Aging. In: *Brain Aging* [Internet]. CRC Press; 2019 [cited 2020 Aug 5]. p. 3–20. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK3885/>
4. Janssen I, Heymsfield SB, Ross R. Low relative skeletal muscle mass (sarcopenia) in older persons is associated with functional impairment and physical disability. *J Am Geriatr Soc* [Internet]. 2002 May 1 [cited 2020 Oct 20];50(5):889–96. Available from: <https://onlinelibrary.wiley.com/doi/full/10.1046/j.1532-5415.2002.50216.x>
5. Newman AB, Haggerty CL, Goodpaster B, Harris T, Kritchevsky S, Nevitt M, et al. Strength and muscle quality in a well-functioning cohort of older adults: the Health, Aging and Body Composition Study. *J Am Geriatr Soc* [Internet]. 2003 Mar 1 [cited 2020 Oct 20];51(3):323–30. Available from: <https://onlinelibrary.wiley.com/doi/full/10.1046/j.1532-5415.2003.51105.x>
6. Vasconcelos RS, dos Souza SS, Carneiro VLR, dos Alves SC. Strength and ability to implement the activities of daily living in elderly resident in rural areas. *Colomb Med* [Internet]. 2016 [cited 2020 Oct 20];47(3):167–71. Available from: [/pmc/articles/PMC5091276/?report=abstract](https://pubmed.ncbi.nlm.nih.gov/27091276/)
7. Wang DXM, Yao J, Zirek Y, Reijnierse EM, Maier AB. Muscle mass, strength, and

physical performance predicting activities of daily living: a meta-analysis [Internet].
Vol. 11, Journal of Cachexia, Sarcopenia and Muscle. Wiley Blackwell; 2020 [cited
2020 Oct 20]. p. 3–25. Available from:

<https://onlinelibrary.wiley.com/doi/abs/10.1002/jcsm.12502>

8. Jaul E, Barron J. Age-related diseases and clinical and public health implications for the 85 years old and over population. *Front Public Heal* [Internet]. 2017 Dec 11 [cited 2020 Oct 20];5. Available from: [/pmc/articles/PMC5732407/?report=abstract](https://pubmed.ncbi.nlm.nih.gov/32407/)
9. Ohrnberger J, Fichera E, Sutton M. The dynamics of physical and mental health in the older population. *J Econ Ageing* [Internet]. 2017 Jun 1 [cited 2020 Sep 8];9:52–62. Available from: [/pmc/articles/PMC5446314/?report=abstract](https://pubmed.ncbi.nlm.nih.gov/3446314/)
10. Ohrnberger J, Fichera E, Sutton M. The relationship between physical and mental health: a mediation analysis. *Soc Sci Med*. 2017 Dec 1;195:42–9.
11. Fried EI, Bockting C, Arjadi R, Borsboom D, Amshoff M, Cramer AOJ, et al. From loss to loneliness: The relationship between bereavement and depressive symptoms. *J Abnorm Psychol*. 2015 May 1;124(2):256–65.
12. Valtorta NK, Kanaan M, Gilbody S, Ronzi S, Hanratty B. Loneliness and social isolation as risk factors for coronary heart disease and stroke: Systematic review and meta-analysis of longitudinal observational studies. *Heart* [Internet]. 2016 [cited 2020 Oct 7];102(13):1009–16. Available from: <http://dx.doi.org/10.1136/heartjnl-2015-308790>
13. Kok AAL, Aartsen MJ, Deeg DJH, Huisman M, Castle NG. Capturing the diversity of successful aging: An operational definition based on 16-year trajectories of functioning. *Gerontologist* [Internet]. 2017 Apr 1 [cited 2020 Oct 19];57(2):240–51. Available from: <https://academic.oup.com/gerontologist/article/57/2/240/2632001>
14. Strawbridge WJ, Wallhagen MI, Cohen RD. Successful aging and well-being: Self-

rated compared with Rowe and Kahn. *Gerontologist* [Internet]. 2002 Dec 1 [cited 2020 Oct 28];42(6):727–33. Available from:

<https://academic.oup.com/gerontologist/article/42/6/727/671748>

15. WHO. Ageing: healthy ageing and functional ability [Internet]. 2020 [cited 2021 Feb 2]. Available from: <https://www.who.int/westernpacific/news/q-a-detail/ageing-healthy-ageing-and-functional-ability>

16. Young Y, Frick KD, Phelan EA. Can Successful Aging and Chronic Illness Coexist in the Same Individual? A Multidimensional Concept of Successful Aging. Vol. 10, *Journal of the American Medical Directors Association*. Elsevier Inc.; 2009. p. 87–92.

17. Harris PB. Another wrinkle in the debate about successful aging: the undervalued concept of resilience and the lived experience of dementia. *Int J Aging Hum Dev* [Internet]. 2008 [cited 2021 Apr 16];67(1):43–61. Available from:

<https://pubmed.ncbi.nlm.nih.gov/18630190/>

18. Fry PS, Keyes CLM. New frontiers in resilient aging: life-strengths and well-being in late life [Internet]. *New Frontiers in Resilient Aging: Life-Strengths and Well-Being in Late Life*. 2010 [cited 2021 Feb 19]. 1–365 p. Available from:

<https://books.google.com/books?hl=nl&lr=&id=IT47K3GTZvIC&oi=fnd&pg=PR7&ots=t9iBZaZVyO&sig=qwDYcmTFd5O7EpdN3aSPeQovTaE>

19. Greve W, Staudinger UM. Resilience in later adulthood and old age: resources and potentials for successful aging. In: *Developmental Psychopathology: Second Edition* [Internet]. 2006 [cited 2021 Feb 19]. p. 796–840. Available from:

<https://www.researchgate.net/publication/232484142>

20. MacLeod S, Musich S, Hawkins K, Alsgaard K, Wicker ER. The impact of resilience among older adults. *Geriatr Nurs (Minneap)*. 2016 Jul 1;37(4):266–72.

21. Netuveli G, Blane D. Quality of life in older ages [Internet]. Vol. 85, *British Medical*

- 610 Bulletin. Oxford Academic; 2008 [cited 2020 Oct 20]. p. 113–26. Available from:
611 <https://academic.oup.com/bmb/article/85/1/113/291398>
- 612 22. Hildon Z, Montgomery SM, Blane D, Wiggins RD, Netuveli G. Examining resilience
613 of quality of life in the face of health-related and psychosocial adversity at older ages:
614 What is ‘right’ about the way we age? *Gerontologist* [Internet]. 2010 Feb 1 [cited 2020
615 Oct 23];50(1):36–47. Available from:
616 <https://academic.oup.com/gerontologist/article/50/1/36/689060>
- 617 23. Mehta M, Whyte E, Lenze E, Hardy S, Roumani Y, Subashan P, et al. Depressive
618 symptoms in late life: associations with apathy, resilience and disability vary between
619 young-old and old-old. *Int J Geriatr Psychiatry*. 2008 Mar;23(3):238–43.
- 620 24. Ryff CD, Love GD, Essex MJ, Singer B. Resilience in adulthood and later life. In:
621 *Handbook of Aging and Mental Health*. Springer US; 1998. p. 69–96.
- 622 25. Stephens C, Breheny M, Mansvelt J. Healthy ageing from the perspective of older
623 people: A capability approach to resilience. *Psychol Heal* [Internet]. 2015 Jun 3 [cited
624 2021 Feb 19];30(6):715–31. Available from:
625 <https://doi.org/10.1080/08870446.2014.904862>
- 626 26. Hayman KJ, Kerse N, Consedine NS. Resilience in context: the special case of
627 advanced age. *Aging Ment Health* [Internet]. 2017 Jun 3 [cited 2021 Feb 3];21(6):577–
628 85. Available from:
629 <https://www.tandfonline.com/doi/full/10.1080/13607863.2016.1196336>
- 630 27. Bowling A, Iliffe S. Psychological approach to successful ageing predicts future
631 quality of life in older adults. *Health Qual Life Outcomes* [Internet]. 2011 Mar 9 [cited
632 2020 Oct 20];9(1):1–10. Available from:
633 <https://link.springer.com/articles/10.1186/1477-7525-9-13>
- 634 28. Christopher KA, Kulig JC. Determinants of Psychological Well-Being in Irish

Immigrants. *West J Nurs Res*. 2000 Mar;22(2):123–43.

29. Gerino E, Rollè L, Sechi C, Brustia P. Loneliness, resilience, mental health, and quality of life in old age: a structural equation model. *Front Psychol* [Internet]. 2017 Nov 14 [cited 2020 Oct 29];8(NOV):2003. Available from: www.frontiersin.org
30. Gooding PA, Hurst A, Johnson J, Tarrier N. Psychological resilience in young and older adults. *Int J Geriatr Psychiatry* [Internet]. 2012 Mar 1 [cited 2021 Feb 19];27(3):262–70. Available from: <http://doi.wiley.com/10.1002/gps.2712>
31. Montross LP, Depp C, Daly J, Reichstadt J, Golshan S, Moore D, et al. Correlates of self-rated successful aging among community-dwelling older adults. *Am J Geriatr Psychiatry* [Internet]. 2006 [cited 2020 Oct 29];14(1):43–51. Available from: <https://www.sciencedirect.com/science/article/pii/S1064748112615917>
32. Nygren B, Aléx L, Jonsén E, Gustafson Y, Norberg A, Lundman B. Resilience, sense of coherence, purpose in life and self-transcendence in relation to perceived physical and mental health among the oldest old. *Aging Ment Health* [Internet]. 2005 Jul [cited 2020 Aug 13];9(4):354–62. Available from: <http://www.tandfonline.com/doi/abs/10.1080/1360500114415>
33. Smith JL, Hollinger-Smith L. Savoring, resilience, and psychological well-being in older adults. *Aging Ment Health* [Internet]. 2015 Mar 4 [cited 2020 Jun 5];19(3):192–200. Available from: <http://www.tandfonline.com/doi/abs/10.1080/13607863.2014.986647>
34. Zeng Y, Shen K. Resilience significantly contributes to exceptional longevity. *Curr Gerontol Geriatr Res*. 2010;2010.
35. Shen K, Zeng Y. The association between resilience and survival among chinese elderly. *Demogr Res* [Internet]. 2010 Apr [cited 2020 Oct 30];23(5):105–16. Available from: [/pmc/articles/PMC2990503/?report=abstract](http://pmc/articles/PMC2990503/?report=abstract)

- 660 36. Jeste D V., Savla GN, Thompson WK, Vahia I V., Glorioso DK, Martin AS, et al.
661 Association between older age and more successful aging: Critical role of resilience
662 and depression. *Am J Psychiatry*. 2013 Feb 1;170(2):188–96.
- 663 37. Borrás C, Ingles M, Mas-Bargues C, Dromant M, Sanz-Ros J, Román-Domínguez A,
664 et al. Centenarians: An excellent example of resilience for successful ageing. Vol. 186,
665 *Mechanisms of Ageing and Development*. Elsevier Ireland Ltd; 2020. p. 111199.
- 666 38. Bonaccio M, Di Castelnuovo A, Costanzo S, Pounis G, Persichillo M, Cerletti C, et al.
667 Mediterranean-type diet is associated with higher psychological resilience in a general
668 adult population: findings from the Moli-sani study. *Eur J Clin Nutr [Internet]*. 2018
669 Jan 27 [cited 2020 Oct 30];72(1):154–60. Available from:
670 <http://www.nature.com/articles/ejcn2017150>
- 671 39. Perna L, Mielck A, Lacruz ME, Emeny RT, Holle R, Breitfelder A, et al.
672 Socioeconomic position, resilience, and health behaviour among elderly people. *Int J*
673 *Public Health [Internet]*. 2012 Sep 13 [cited 2020 Oct 30];57(2):341–9. Available
674 from: <http://www.helmholtz-muenchen.de/igm/>
- 675 40. Tiainen A-MK, Männistö S, Lahti M, Blomstedt PA, Lahti J, Perälä M-M, et al.
676 Personality and Dietary Intake – Findings in the Helsinki Birth Cohort Study.
677 Rustichini A, editor. *PLoS One [Internet]*. 2013 Jul 18 [cited 2020 Oct
678 30];8(7):e68284. Available from: <https://dx.plos.org/10.1371/journal.pone.0068284>
- 679 41. Reichstadt J, Depp CA, Palinkas LA, Folsom DP, Jeste D V. Building blocks of
680 successful aging: A focus group study of older adults perceived contributors to
681 successful aging. *Am J Geriatr Psychiatry [Internet]*. 2007 [cited 2020 Oct
682 19];15(3):194–201. Available from:
683 <https://www.sciencedirect.com/science/article/pii/S1064748112604369>
- 684 42. Bosnes I, Nordahlid HM, Stordalid E, Bosnes O, Myklebust TÅ, Almkvist O.

Lifestyle predictors of successful aging: A 20-year prospective HUNT study. 2019
[cited 2020 Mar 17]; Available from: <https://doi.org/10.1371/journal.pone.0219200>

43. Davey M, Eaker DG, Walters LH. Resilience processes in adolescents: Profiles, self-worth, and coping. Vol. 18, *Journal of Adolescent Research*. 2003. p. 347–62.

44. Riolli L, Savicki V, Cepani A. Resilience in the face of catastrophe: Optimism, personality, and coping in the Kosovo crisis. *J Appl Soc Psychol* [Internet]. 2002 Aug 1 [cited 2020 Oct 23];32(8):1604–27. Available from: <https://onlinelibrary.wiley.com/doi/full/10.1111/j.1559-1816.2002.tb02765.x>

45. Schmeichel BJ, Volokhov RN, Demaree HA. Working memory capacity and the self-regulation of emotional expression and experience. *J Pers Soc Psychol* [Internet]. 2008 Dec [cited 2020 Oct 13];95(6):1526–40. Available from: <https://www.researchgate.net/publication/23489273>

46. Schmeichel BJ, Demaree HA. Working memory capacity and spontaneous emotion regulation: high capacity predicts self-enhancement in response to negative feedback. *Emotion* [Internet]. 2010 [cited 2020 Oct 13];10(5):739–44. Available from: <https://www.researchgate.net/publication/47642303>

47. van den Berg AE, Maas J, Verheij RA, Groenewegen PP. Green space as a buffer between stressful life events and health. *Soc Sci Med*. 2010 Apr 1;70(8):1203–10.

48. Wells M. Resilience in Older Adults Living in Rural, Suburban, and Urban Areas. *Online J Rural Nurs Heal Care*. 2010;10(2):45–54.

49. Windle G, Bennett KM, Noyes J. A methodological review of resilience measurement scales. *Health Qual Life Outcomes* [Internet]. 2011 Feb 4 [cited 2020 Mar 26];9(1):8. Available from: <http://hqlo.biomedcentral.com/articles/10.1186/1477-7525-9-8>

50. Chmitorz A, Kunzler A, Helmreich I, Tüscher O, Kalisch R, Kubiak T, et al. Intervention studies to foster resilience – A systematic review and proposal for a

resilience framework in future intervention studies. Vol. 59, Clinical Psychology Review. Elsevier Inc.; 2018. p. 78–100.

51. Kalisch R, Veer I, Yuen KS, Hendler T, Myin-Germeys, I. Walter, H. Riepenhausen A. DynaCORE-C: The DynaMORE cross-sectional survey study on psychological resilience to the mental health consequences of the Corona crisis [Internet]. 2020. Available from: osf.io/5xq9p
52. Netuveli G, Wiggins RD, Montgomery SM, Hildon Z, Blane D. Mental health and resilience at older ages: bouncing back after adversity in the British Household Panel Survey. J Epidemiol Community Health. 2008 Nov 1;62(11):987–91.
53. Fritz J, Stochl J, Fried EI, Goodyer IM, Van Borkulo CD, Wilkinson PO, et al. Unravelling the complex nature of resilience factors and their changes between early and later adolescence. BMC Med [Internet]. 2019 Nov 14 [cited 2021 Apr 14];17(1):203. Available from: <https://bmcmmedicine.biomedcentral.com/articles/10.1186/s12916-019-1430-6>
54. Fritz J, Fried EI, Goodyer IM, Wilkinson PO, van Harmelen AL. A Network Model of Resilience Factors for Adolescents with and without Exposure to Childhood Adversity. Sci Rep [Internet]. 2018 [cited 2020 Sep 9];8(1). Available from: <https://www.nature.com/articles/s41598-018-34130-2>
55. Scheffer M, Elizabeth Bolhuis J, Borsboom D, Buchman TG, Gijzel SMW, Goulson D, et al. Quantifying resilience of humans and other animals [Internet]. Vol. 115, Proceedings of the National Academy of Sciences of the United States of America. National Academy of Sciences; 2018 [cited 2021 Apr 14]. p. 11883–90. Available from: www.pnas.org/lookup/suppl/doi:10.1073/pnas.1810630115/-/DCSupplemental.www.pnas.org/cgi/doi/10.1073/pnas.1810630115
56. Borsboom D. A network theory of mental disorders. World Psychiatry. 2017 Feb

1;16(1):5–13.

57. Kalisch R, Cramer AOJ, Binder H, Fritz J, Leertouwer Ij, Lunansky G, et al. Deconstructing and reconstructing resilience: a dynamic network approach. *Perspect Psychol Sci* [Internet]. 2019 Sep [cited 2020 Mar 24];14(5):765–77. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/31365841>
58. Lunansky G, van Borkulo C, Borsboom D. Personality, Resilience, and Psychopathology: A Model for the Interaction between Slow and Fast Network Processes in the Context of Mental Health. *Eur J Pers* [Internet]. 2020 Dec 4 [cited 2021 Mar 17];34(6):969–87. Available from: <http://journals.sagepub.com/doi/10.1002/per.2263>
59. Zendedel R, Schouten BC, van Weert JCM, van den Putte B. Informal interpreting in general practice: Are interpreters' roles related to perceived control, trust, and satisfaction? *Patient Educ Couns* [Internet]. 2018 [cited 2020 Sep 21];101(6):1058–65. Available from: <https://www.sciencedirect.com/science/article/pii/S073839911830034X>
60. Abdel-Khalek AM. Measuring happiness with a single-item scale [Internet]. Vol. 34, *Social Behavior and Personality*. 2006 [cited 2020 Jun 26]. p. 139–50. Available from: <http://openurl.ingenta.com/content/xref?genre=article&issn=0301-2212&volume=34&issue=2&spage=139>
61. Ikink JGM, Lamers SM, Boliuer JM. De Warwick-Edinburgh Mental Well-being Scale (WEMWBS) als meetinstrument voor mentaal welbevinden in Nederland. 2012;(november).
62. Tennant R, Hiller L, Fishwick R, Platt S, Joseph S, Weich S, et al. The Warwick-Dinburgh mental well-being scale (WEMWBS): Development and UK validation. *Health Qual Life Outcomes* [Internet]. 2007 Nov 27 [cited 2020 Apr 14];5(1):63.

Available from: <http://hqlo.biomedcentral.com/articles/10.1186/1477-7525-5-63>

63. Andresen EM, Malmgren JA, Carter WB, Patrick DL. Screening for depression in well older adults: evaluation of a short form of the CES-D (Center for Epidemiologic Studies Depression Scale). *Am J Prev Med* [Internet]. 1994 Jan 1 [cited 2020 Jul 27];10(2):77–84. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/8037935>
64. Gobbens RJJ, van Assen MALM. Psychometric properties of the Dutch WHOQOL-OLD. *Health Qual Life Outcomes*. 2016 Jul 15;14(1).
65. Power M, Quinn K, Schmidt S. Development of the WHOQOL-Old module. Springer [Internet]. 2005 Dec [cited 2020 Apr 6];14(10):2197–214. Available from: <https://link.springer.com/article/10.1007/s11136-005-7380-9>
66. Zigmond AS, Snaith RP. The Hospital Anxiety and Depression Scale. *Acta Psychiatr Scand*. 1983;67(6):361–70.
67. Spinhoven P, Ormel J, Sloekers PPA, Kempen GIJM, Speckens AEM, Van Hemert AM. A validation study of the hospital anxiety and depression scale (HADS) in different groups of Dutch subjects. *Psychol Med* [Internet]. 1997 [cited 2020 Apr 28];27(2):363–70. Available from: <https://doi.org/10.1017/S0033291796004382>
68. Leontjevas R, de Beek WO, Lataster J, Jacobs N. Resilience to affective disorders: A comparative validation of two resilience scales. *J Affect Disord* [Internet]. 2014 Oct 15 [cited 2020 Aug 3];168:262–8. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0165032714004418>
69. Smith BW, Dalen J, Wiggins K, Tooley E, Christopher P, Bernard J. The Brief Resilience Scale: Assessing the Ability to Bounce Back. *Int J Behav Med* [Internet]. 2008 Jul [cited 2020 Apr 1];15(3):194–200. Available from: https://idp.springer.com/authorize/casa?redirect_uri=https://link.springer.com/article/10.1080/10705500802222972&casa_token=wT4YpwuEJMAAAAAA:qUJTOjj2CWych

7c3wjcP8V9w0eLWz8sKZ0-

n12FGpenAuGMHEqltr2tEY8oXBimGacJt74AVEaPfNCjoMbs

70. Kampa M, Schick A, Yuen K, Sebastian A, Chmitorz A, Saase V, et al. A Combined Behavioral and Neuroimaging Battery to Test Positive Appraisal Style Theory of Resilience in Longitudinal Studies. *bioRxiv* [Internet]. 2018 [cited 2020 Aug 3];470435. Available from: <https://doi.org/10.1101/470435>
71. Kampa M, Schick A, Sebastian A, Wessa M, Tüscher O, Kalisch R, et al. Replication of fMRI group activations in the neuroimaging battery for the Mainz Resilience Project (MARP). *Neuroimage* [Internet]. 2020 [cited 2020 Aug 3];204. Available from: <https://www.sciencedirect.com/science/article/pii/S1053811919308146>
72. Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. *J Health Soc Behav.* 1983;24(4):385–96.
73. Korten NCM, Comijs HC, Penninx BWJH, Deeg DJH. Perceived stress and cognitive function in older adults: which aspect of perceived stress is important? *Int J Geriatr Psychiatry* [Internet]. 2017 Apr 1 [cited 2020 Aug 3];32(4):439–45. Available from: <https://pubmed.ncbi.nlm.nih.gov/27059116/>
74. Gobbens RJJ, van Assen MALM, Luijckx KG, Wijnen-Sponselee MT, Schols JMGA. The tilburg frailty indicator: Psychometric properties. *J Am Med Dir Assoc* [Internet]. 2010 [cited 2020 Mar 30];11(5):344–55. Available from: <https://www.researchgate.net/publication/44638590>
75. Holmes TH, Rahe RH. The social readjustment rating scale. *J Psychosom Res* [Internet]. 1967 [cited 2020 Aug 3];11(2):213–8. Available from: <https://psycnet.apa.org/record/1968-03998-001>
76. Gosling SD, Rentfrow PJ, Swann WB. A very brief measure of the Big-Five personality domains. *J Res Pers* [Internet]. 2003 [cited 2020 Aug 27];37(6):504–28.

- 810 Available from:
- 811 <https://www.sciencedirect.com/science/article/pii/S0092656603000461>
- 812 77. Hofmans J, Kuppens P, Allik J. Is short in length short in content? An examination of
813 the domain representation of the Ten Item Personality Inventory scales in Dutch
814 language. *Pers Individ Dif*. 2008 Dec 1;45(8):750–5.
- 815 78. Teeuw B, Schwarzer R, Jerusalem M. Dutch adaptation of the general self-efficacy
816 scale. Berlin, Ger [Internet]. 1994; Available from:
817 https://meetinstrumentenzorg.nl/wp-content/uploads/instrumenten/328_3.pdf
- 818 79. Schwarzer R, Jerusalem M. General Self--efficacy Scale (GES) [Internet]. diabetes-
819 psychologie.de. 1995 [cited 2020 Aug 3]. Available from: [https://diabetes-](https://diabetes-psychologie.de/downloads/Beschreibung_GSE.pdf)
820 [psychologie.de/downloads/Beschreibung_GSE.pdf](https://diabetes-psychologie.de/downloads/Beschreibung_GSE.pdf)
- 821 80. Rosenberg M. Conceiving the self. [Internet]. New York: Basic Books; 1979 [cited
822 2020 Aug 3]. Available from:
823 [https://www.google.com/search?q=Rosenberg%2C+M.+\(1979\).+Conceiving+the+self.](https://www.google.com/search?q=Rosenberg%2C+M.+(1979).+Conceiving+the+self.+New+York%3A+Basic+Books&rlz=1C1CHBD_nlNL904NL904&oq=Rosenberg%2C+M.+(1979).+Conceiving+the+self.+New+York%3A+Basic+Books&aqs=chrome..69i57.314j0j4&sourceid=chrome&ie=UTF-8)
824 [+New+York%3A+Basic+Books&rlz=1C1CHBD_nlNL904NL904&oq=Rosenberg%2](https://www.google.com/search?q=Rosenberg%2C+M.+(1979).+Conceiving+the+self.+New+York%3A+Basic+Books&rlz=1C1CHBD_nlNL904NL904&oq=Rosenberg%2C+M.+(1979).+Conceiving+the+self.+New+York%3A+Basic+Books&aqs=chrome..69i57.314j0j4&sourceid=chrome&ie=UTF-8)
825 [C+M.+\(1979\).+Conceiving+the+self.+New+York%3A+Basic+Books&aqs=chrome..6](https://www.google.com/search?q=Rosenberg%2C+M.+(1979).+Conceiving+the+self.+New+York%3A+Basic+Books&rlz=1C1CHBD_nlNL904NL904&oq=Rosenberg%2C+M.+(1979).+Conceiving+the+self.+New+York%3A+Basic+Books&aqs=chrome..69i57.314j0j4&sourceid=chrome&ie=UTF-8)
826 [9i57.314j0j4&sourceid=chrome&ie=UTF-8](https://www.google.com/search?q=Rosenberg%2C+M.+(1979).+Conceiving+the+self.+New+York%3A+Basic+Books&rlz=1C1CHBD_nlNL904NL904&oq=Rosenberg%2C+M.+(1979).+Conceiving+the+self.+New+York%3A+Basic+Books&aqs=chrome..69i57.314j0j4&sourceid=chrome&ie=UTF-8)
- 827 81. Franck E, De Raedt R, Barbez C, Rosseel Y. Psychometric properties of the Dutch
828 Rosenberg self-esteem scale. *Psychol Belg*. 2008;48(1):25–35.
- 829 82. Cramm JM, Strating MM, de Vreede PL, Steverink N, Nieboer AP. Validation of the
830 self-management ability scale (SMAS) and development and validation of a shorter
831 scale (SMAS-S) among older patients shortly after hospitalisation. *Health Qual Life*
832 *Outcomes* [Internet]. 2012 Jan 24 [cited 2020 Jul 14];10(1):9. Available from:
833 <http://hqlo.biomedcentral.com/articles/10.1186/1477-7525-10-9>
- 834 83. Manapat PD, Edwards MC, MacKinnon DP, Poldrack RA, Marsch LA. A

Psychometric Analysis of the Brief Self-Control Scale. Assessment [Internet]. 2019
[cited 2020 Aug 3]; Available from: <https://doi.org/10.1177/1073191119890021>

84. Tangney JP, Baumeister RF, Boone AL. High Self-Control Predicts Good Adjustment,
Less Pathology, Better Grades, and Interpersonal Success. J Pers [Internet]. 2004 Apr 1
[cited 2020 Aug 4];72(2):271–324. Available from:
/pmc/articles/PMC7178234/?report=abstract

85. Verweijen EHS. Ongezonder snackgedrag Plannen uitvoeren of uitstellen ? 2011 [cited
2020 Aug 3];(3272303). Available from:
[https://dspace.library.uu.nl/bitstream/handle/1874/236421/E.H.S. Verweijen
3272303.pdf?sequence=1](https://dspace.library.uu.nl/bitstream/handle/1874/236421/E.H.S.Verweijen3272303.pdf?sequence=1)

86. De Jong-Gierveld J, Van Tilburg T. Manual of the Loneliness Scale. fsw.vu.nl
[Internet]. 1999 [cited 2020 Apr 1];1–26. Available from:
http://home.fsw.vu.nl/TG.van.Tilburg/manual_loneliness_scale_1999.html

87. Van Sonderen E. Sociale steun lijst-Interacties (SSL-i) en Sociale steun lijst-
Discrepanties (SSL-d). Noord Cent voor gezondheidsvraagstukken. 1993;

88. Aaronson NK, Muller M, Cohen PDA, Essink-Bot ML, Fekkes M, Sanderman R, et al.
Translation, validation, and norming of the Dutch language version of the SF-36 Health
Survey in community and chronic disease populations. J Clin Epidemiol. 1998 Nov
1;51(11):1055–68.

89. Ware, J. E., Sherbourne CD. The MOS 36-Item Short-Form Health Survey (SF-36): I .
Conceptual Framework and Item Selection. Med Care. 1992;30(6):473–83.

90. Buysse DJ, Reynolds CF, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh sleep
quality index: A new instrument for psychiatric practice and research. Psychiatry Res
[Internet]. 1989 [cited 2020 Mar 3];28(2):193–213. Available from:
<https://psychiatry.pitt.edu/sites/default/files/inline-files/PSQI Article.pdf>

- 860 91. Babor TF, Higgins-Biddle JC, Saunders JB, Monteiro MG. The Alcohol Use Disorders
861 Identification Test Guidelines for Use in Primary Care [Internet]. 2001 [cited 2020 Aug
862 4]. Available from: <http://oki-kyo.jp/who-audit-jp.pdf>
- 863 92. Bush K, Kivlahan DR, McDonell MB, Fihn SD, Bradley KA. The AUDIT alcohol
864 consumption questions (AUDIT-C): An effective brief screening test for problem
865 drinking. Arch Intern Med [Internet]. 1998 [cited 2020 Aug 4];158(16):1789–95.
866 Available from: [https://jamanetwork.com/journals/jamainternalmedicine/article-](https://jamanetwork.com/journals/jamainternalmedicine/article-abstract/208954)
867 [abstract/208954](https://jamanetwork.com/journals/jamainternalmedicine/article-abstract/208954)
- 868 93. Saunders JB, AASLAND OG, BABOR TF, DE LA FUENTE JR, GRANT M.
869 Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO
870 Collaborative Project on Early Detection of Persons with Harmful Alcohol
871 Consumption-II. Addiction. 1993;88(6):791–804.
- 872 94. Slotman A, Cramm JM, Nieboer AP. Validation of the Dutch aging perceptions
873 questionnaire and development of a short version. Health Qual Life Outcomes. 2015
874 May 12;13(1).
- 875 95. Crawford JR, Smith G, Maylor EA, Della Sala S, Logie RH. The Prospective and
876 Retrospective Memory Questionnaire (PRMQ): Normative data and latent structure in
877 a large non-clinical sample. Memory [Internet]. 2003 May [cited 2020 Aug
878 4];11(3):261–75. Available from: <https://pubmed.ncbi.nlm.nih.gov/12908675/>
- 879 96. Gondo Y, Renge N, Ishioka Y, Kurokawa I, Ueno D, Rendell P. Reliability and
880 validity of the Prospective and Retrospective Memory Questionnaire (PRMQ) in young
881 and old people: A Japanese study. Jpn Psychol Res [Internet]. 2010 Sep 1 [cited 2020
882 Aug 4];52(3):175–85. Available from: [http://doi.wiley.com/10.1111/j.1468-](http://doi.wiley.com/10.1111/j.1468-5884.2010.00433.x)
883 [5884.2010.00433.x](http://doi.wiley.com/10.1111/j.1468-5884.2010.00433.x)
- 884 97. Smith G, Sala D, Logie RH, Maylor EA. Prospective and retrospective memory in

normal ageing and dementia: A questionnaire study. Taylor Fr [Internet]. 2000 [cited 2020 Aug 4];8(5):311–21. Available from:

<https://www.tandfonline.com/action/journalInformation?journalCode=pmem20>

98. Zimprich D, Kliegel M, Rast P. The factorial structure and external validity of the prospective and retrospective memory questionnaire in older adults. *Eur J Ageing* [Internet]. 2011 [cited 2020 Aug 4];8(1):39–48. Available from:

http://www.psy.ed.ac.uk/research/hcn/PRMQclick/document_view.

99. Epskamp S, Borsboom D, Fried EI. Estimating psychological networks and their accuracy: A tutorial paper. *Behav Res Methods*. 2018 Feb 1;50(1):195–212.

100. Haslbeck JMB, Waldorp LJ. Structure estimation for mixed graphical models in high-dimensional data. 2015 Oct 19 [cited 2020 Sep 22]; Available from:

<http://arxiv.org/abs/1510.05677>

101. Haslbeck JMB, Waldorp LJ. MGM: Estimating time-varying mixed graphical models in high-dimensional data. *J Stat Softw*. 2020;93.

102. Hastie T, Tibshirani R, Wainwright M. Statistical learning with sparsity: The lasso and generalizations [Internet]. *Statistical Learning with Sparsity: The Lasso and Generalizations*. 2015 [cited 2020 Aug 5]. 1–337 p. Available from:

[https://books.google.nl/books?hl=nl&lr=&id=f-](https://books.google.nl/books?hl=nl&lr=&id=f-A_CQAAQBAJ&oi=fnd&pg=PP1&dq=Hastie+T,+Tibshirani+R,+Wainwright+M+(2015).+Statistical+learning+with+sparsity.+CRC+press.&ots=G4_PFZnYT3&sig=42GE9z__6jzB_Vs42tKYWugrevA)

[A_CQAAQBAJ&oi=fnd&pg=PP1&dq=Hastie+T,+Tibshirani+R,+Wainwright+M+\(2015\).+Statistical+learning+with+sparsity.+CRC+press.&ots=G4_PFZnYT3&sig=42GE9z__6jzB_Vs42tKYWugrevA](https://books.google.nl/books?hl=nl&lr=&id=f-A_CQAAQBAJ&oi=fnd&pg=PP1&dq=Hastie+T,+Tibshirani+R,+Wainwright+M+(2015).+Statistical+learning+with+sparsity.+CRC+press.&ots=G4_PFZnYT3&sig=42GE9z__6jzB_Vs42tKYWugrevA)

103. Nelder J, Baker R. Generalized linear models. In: *Encyclopedia of statistical sciences* [Internet]. 1972 [cited 2020 Aug 5]. Available from:

<https://rss.onlinelibrary.wiley.com/doi/abs/10.2307/2344614>

104. Epskamp S, Cramer AOJ, Waldorp LJ, Schmittmann VD, Borsboom D. Qgraph:

910 network visualizations of relationships in psychometric data. J Stat Softw [Internet].
 911 2012 [cited 2020 Sep 22];48(4):1–18. Available from: <http://www.jstatsoft.org/>

912 105. Bollen KA, Stine RA. Bootstrapping Goodness-of-Fit Measures in Structural Equation
 913 Models. Sociol Methods Res. 1992;21(2):205–29.

914 106. Efron B. Bootstrap methods: another look at the jackknife. Ann Stat. 1979;7(1):1–26.

915 107. Hastie T, Tibshirani R, Wainwright M. Statistical learning with sparsity: the lasso and
 916 generalizations. CRC Press; 2015.

917 108. Haslbeck JMB, Waldorp LJ. How well do network models predict observations? On
 918 the importance of predictability in network models. Behav Res Methods [Internet].
 919 2018 Apr 1 [cited 2020 Sep 22];50(2):853–61. Available from: <https://jmbh.github.io/>

920 109. Haslbeck JMB, Fried EI. How predictable are symptoms in psychopathological
 921 networks? A reanalysis of 18 published datasets. Psychol Med [Internet]. 2017 Dec 1
 922 [cited 2021 Mar 8];47(16):2267–76. Available from:
 923 <https://pubmed.ncbi.nlm.nih.gov/28625186/>

924 110. Fortunato S. Community detection in graphs [Internet]. Vol. 486, Physics Reports.
 925 2010 [cited 2020 Sep 21]. p. 75–174. Available from:
 926 <https://www.sciencedirect.com/science/article/pii/S0370157309002841>

927 111. Lange J. CliquePercolation: Clique Percolation for Networks [Internet]. 2019.
 928 Available from: <https://cran.r-project.org/package=CliquePercolation>

929 112. Farkas I, Ábel D, Palla G, Vicsek T. Weighted network modules. New J Phys.
 930 2007;9(6):180.

931 113. Palla G, Derényi I, Farkas I, Vicsek T. Uncovering the overlapping community
 932 structure of complex networks in nature and society. Nature [Internet]. 2005 Jun 9
 933 [cited 2020 Sep 23];435(7043):814–8. Available from: <http://angel.elte.hu/clustering/>

934 114. Blanken TF, Deserno MK, Dalege J, Borsboom D, Blanken P, Kerkhof GA, et al. The

- role of stabilizing and communicating symptoms given overlapping communities in psychopathology networks. *Sci Rep* [Internet]. 2018 Dec 1 [cited 2021 Apr 30];8(1):5854. Available from: www.nature.com/scientificreports
115. Haslbeck J, Borsboom D, Waldorp L. Moderated Network Models. *Multivariate Behav Res* [Internet]. 2018 Jul 8 [cited 2020 Aug 5]; Available from: <http://arxiv.org/abs/1807.02877>
116. Van Borkulo CD, Borsboom D, Epskamp S, Blanken TF, Boschloo L, Schoevers RA, et al. A new method for constructing networks from binary data. *Sci Rep* [Internet]. 2014 Aug 1 [cited 2021 Apr 26];4(1):1–10. Available from: <http://cran.r-project.org/web/packages/>
117. Epskamp S, Fried EI. A tutorial on regularized partial correlation networks. *Psychol Methods*. 2018 Dec 1;23(4):617–34.
118. Leys C, Ley C, Klein O, Bernard P, Licata L. Detecting outliers: Do not use standard deviation around the mean, use absolute deviation around the median. *J Exp Soc Psychol* [Internet]. 2013 [cited 2020 Sep 23];49(4):764–6. Available from: <https://www.sciencedirect.com/science/article/pii/S0022103113000668>
119. Bowling A. Aspirations for older age in the 21st century: What is successful aging? *Int J Aging Hum Dev*. 2007;64(3):263–97.
120. Depp CA, Jeste D V. Definitions and Predictors of Successful Aging: A Comprehensive Review of Larger Quantitative Studies. *Focus (Madison)*. 2009 Jan;7(1):137–50.
121. Domènech-Abella J, Mundó J, Haro JM, Rubio-Valera M. Anxiety, depression, loneliness and social network in the elderly: Longitudinal associations from The Irish Longitudinal Study on Ageing (TILDA). *J Affect Disord*. 2019 Mar 1;246:82–8.
122. Tseng FM, Petrie D, Leon-Gonzalez R. The impact of spousal bereavement on

- subjective wellbeing: Evidence from the Taiwanese elderly population. *Econ Hum Biol*. 2017 Aug 1;26:1–12.
123. Halpin SN. Declining with purpose: Resilience despite Cognitive Decline. *Life Span Disabil* [Internet]. 2018 [cited 2020 Oct 30];21(1):89–108. Available from: <https://www.researchgate.net/publication/326331328>
124. Peels DA, Bolman C, Golsteijn RHJ, Vries H De, Mudde AN, Van Stralen MM, et al. Differences in reach and attrition between web-based and print-delivered tailored interventions among adults over 50 years of age: Clustered randomized trial. *J Med Internet Res*. 2012;14(6).
125. Lavrakas P. Self-Selection Bias. In: *Encyclopedia of Survey Research Methods*. Sage Publications, Inc.; 2013.
126. CBS. Bevolking; leeftijd, migratieachtergrond, geslacht, regio, 1 jan. 1996-2020 [Dataset] [Internet]. 2020. Available from: <https://opendata.cbs.nl/statline/#/CBS/nl/dataset/37713/table?dl=3E6BC>
127. Bonevski B, Randell M, Paul C, Chapman K, Twyman L, Bryant J, et al. Reaching the hard-to-reach: A systematic review of strategies for improving health and medical research with socially disadvantaged groups. *BMC Med Res Methodol* [Internet]. 2014 Mar 25 [cited 2021 Jan 12];14(1):1–29. Available from: <http://www.biomedcentral.com/1471-2288/14/42>
128. Van Loon AJM, Tijhuis M, Picavet HSJ, Surtees PG, Ormel J. Survey non-response in the Netherlands: Effects on prevalence estimates and associations. *Ann Epidemiol* [Internet]. 2003 Feb [cited 2021 Feb 5];13(2):105–10. Available from: <https://pubmed.ncbi.nlm.nih.gov/12559669/>
129. Asmundson GJG, Paluszek MM, Landry CA, Rachor GS, McKay D, Taylor S. Do pre-existing anxiety-related and mood disorders differentially impact COVID-19 stress

985 responses and coping? J Anxiety Disord. 2020 Aug 1;74:102271.

986 130. Grossman ES, Hoffman YSG, Palgi Y, Shrira A. COVID-19 related loneliness and
987 sleep problems in older adults: Worries and resilience as potential moderators. Pers
988 Individ Dif. 2021 Jan 1;168:110371.

989 131. Pearman A, Hughes ML, Smith EL, Neupert SD. Age Differences in Risk and
990 Resilience Factors in COVID-19-Related Stress. Journals Gerontol Ser B [Internet].
991 2020 Aug 3 [cited 2020 Nov 18];XX:1–7. Available from:
992 [https://academic.oup.com/psychsocgerontology/advance-](https://academic.oup.com/psychsocgerontology/advance-article/doi/10.1093/geronb/gbaa120/5879986)
993 [article/doi/10.1093/geronb/gbaa120/5879986](https://academic.oup.com/psychsocgerontology/advance-article/doi/10.1093/geronb/gbaa120/5879986)

994 132. Kalisch R, Baker DG, Basten U, Boks MP, Bonanno GA, Brummelman E, et al. The
995 resilience framework as a strategy to combat stress-related disorders [Internet]. Vol. 1,
996 Nature Human Behaviour. 2017 [cited 2020 Apr 1]. p. 784–90. Available from:
997 <https://www.nature.com/articles/s41562-017-0200-8>

998 133. Narici M, De Vito G, Franchi M, Paoli A, Moro T, Marcolin G, et al. Impact of
999 sedentarism due to the COVID-19 home confinement on neuromuscular,
1000 cardiovascular and metabolic health: Physiological and pathophysiological
1001 implications and recommendations for physical and nutritional countermeasures. Eur J
1002 Sport Sci. 2020;

1003 134. Pruchno R, Heid AR, Genderson MW. Resilience and Successful Aging: Aligning
1004 Complementary Constructs Using a Life Course Approach. Vol. 26, Psychological
1005 Inquiry. Routledge; 2015. p. 200–7.

1006

1007

1008 Figure 1. Visualization of the edge-parameters of a fictitious MGM. Green edges indicate
1009 positive relationships and red edges indicate negative relationships. Grey edges indicate
1010 pairwise interactions involving categorical variables. The width of the edges indicate the
1011 absolute strength of the connection.

1012

1013

1014 Figure 2. Visualization of a fictitious MGM with several nodes, including health condition
1015 (HC), stress (S), depression (D) and coping ability (C). Green edges indicate positive
1016 relationships; red edges indicate negative relationships. Left: C and C are conditionally
1017 dependent. Right: C and D are conditionally independent.

1018

1019

1020