**The Psychological Impact of Threat & Lockdowns during the COVID-19 Pandemic:
Exacerbating Factors and Mitigating Actions**

This paper has now been accepted for publication in Translational Behavioral Medicine

Valerie van Mulukom¹ Ph.D., Barbara Muzzulini² Ph.D., Bastiaan T. Rutjens³ Ph.D., Caspar J. van Lissa⁴ Ph.D., and Miguel Farias¹ Ph.D.

¹ Centre for Trust, Peace, and Social Relations, Coventry University, United Kingdom

² Centre for the Study of Social Cohesion, University of Oxford, United Kingdom

³ Psychology Research Institute, University of Amsterdam, Netherlands

⁴ Department of Methodology & Statistics, Utrecht University, Netherlands

*Corresponding author contact information: Valerie van Mulukom, Centre for Trust, Peace and Social Relations, Innovation Village IV5, Cheetah Road, Coventry CV1 2TL, United Kingdom (e-mail: valerie.vanmulukom@coventry.ac.uk).

Abstract

Background

In spring 2020, the COVID-19 pandemic was declared. The threat the pandemic poses as well as associated lockdown measures created challenging times for many.

Purpose

This study aimed to investigate the individual and social factors associated with low mental health, particularly perceived threat and lockdown measures, and factors associated with psychological wellbeing, particularly sense of control.

Methods

An online survey was completed by participants (N=8,229) recruited from 79 countries.

Results

In line with pre-registered hypotheses, participants showed elevated levels of anxiety and depression worldwide. This poor mental health was predicted by perceived threat. The effect of threat on depression was further moderated by social isolation, but there was no effect of sense of control. Sense of control was low overall, and was predicted negatively by maladaptive coping, but positively by adaptive coping and the perception that the government is dealing with the outbreak. Social isolation increased with quarantine duration, but was mitigated by frequent communication with close ones. Engaging in individual actions to avoid contracting the virus was associated with higher anxiety, except when done professionally.

Conclusions

We suggest that early lockdown of the pandemic may have had detrimental psychological effects, which may be alleviated by individual actions such as maintaining frequent social contact and adaptive coping, and by governmental actions which demonstrate support in a public health crisis. Citizens and governments can work together to adapt better to restrictive but necessary measures during the current and future pandemics.

Keywords

COVID-19, mental health, quarantine, sense of control, social isolation, government action

Introduction

On March 11th, 2020, the COVID-19 outbreak was declared a pandemic by the World Health Organisation (WHO). Billions of people were suddenly faced with the threat of catching the virus and the threat of financial and job uncertainty, and potential social isolation as a result of lockdown measures. How might this combination of threat and lockdown measures impact mental health and what factors might contribute to alleviate their negative impact during the pandemic? To address this question, we asked participants ($N=8,236$) worldwide who had been living in social isolation (range 0-110 days, $M = 23.0$, $SD = 9.5$ days), to answer a pre-registered online survey (from March 28th until April 24th, 2020) which included measures of perceived personal threat, anxiety, and depression. Pre-registered analyses sought to assess how various factors, particularly exacerbating (e.g., threat) and mitigating factors (e.g., social contact), may be associated with mental health outcomes. In addition, we investigated what may be associated with a sense of personal control during the pandemic, a factor which we predicted would be associated with better psychological wellbeing.

Threat, Lockdown, & Mental Health Model. The pandemic brings with it high levels of uncertainty: No one can accurately predict how the pandemic will progress. Thus, the level of perceived threat includes a projection of one's fears concerning the likelihood of becoming ill or dying (for oneself or one's family), or of losing one's job, among other threatening possibilities. For example, levels of perceived threat may be based on number of diagnoses in one's community, one's perceived risk of contracting the virus, and one's personal financial situation, given that the pandemic has substantially increased unemployment (e.g., in the U.S.A., the unemployment rate went from a range of 3.5-3.7% to 4.4% in March, and then shot to 14.7% in April and 13.3% in May [1]).

However, perceived threat does not necessarily dictate psychological ill health. Individuals may engage with various behaviours in order to help them deal with the negative aspects of lockdown; here, we suggest that these mitigating factors will moderate the effect of perceived threat on psychological well-being. Such behaviours include staying in contact with close others, taking actions to avoid contracting the virus (e.g., washing one's hands), and engaging in adaptive coping

mechanisms (e.g., reframing the situation in a positive light). Together, these behaviours can give a sense of control and togetherness, and manifest resilience in the face of a threat, thus avoiding anxiety and depression – see Figure 1 for the Threat, Lockdown & Mental Health Model, and Table 1 at the end of the introduction for an overview of the pre-registered hypotheses. We also considered external factors that might contribute to people’s mental health during lockdown. For example, the number of days spent in isolation potentially affects stress levels, though not necessarily in a linear fashion. In the early days of lockdown, a ‘novelty effect’ of lockdown may be expected, reflecting initial low levels of stress, which after this initial adjustment phase may be followed by a spike of the negative effects of lockdown. This in turn may decrease in time due to habituation.

Figure 1. The Threat, Lockdown & Mental Health Model; explaining the effect of personal threat and lockdown consequences (social isolation & reduced sense of control) on depression and anxiety, and factors that may mitigate or exacerbate the effect.

Antecedents of Control Model. The state of emergency issued by many countries, which severely restricted freedom of movement, has led to an inevitable decrease in direct control over one's environment. Therefore, we hypothesised that sense of control would be particularly affected, and that people would engage in actions to regain this control. There are two main processes for achieving control: primary or personal control, which is the ability to influence the environment in response to personal needs, and secondary or compensatory control, which is the ability to adapt the self in response to the environment [2, 3]. Having a sense of control over one's environment is considered a fundamental motivation [4], and a perceived lack of control is directly associated with poor mental health outcomes [5].

Although the level of direct control over a pandemic is, for most of us, negligible, we hypothesise that there are various behaviours which will impact one's sense of control, through both primary and secondary pathways: individuals can take actions on their own to try reducing the risk of contracting the virus (e.g., avoiding physical contact with others) and to gather information about the disease (through checking news sources or through communicating with others), but they can also gain a sense of control without attempting to change the current situation, such as by trusting the government to take effective action in order to control the pandemic [6]; see Figure 2 for the Antecedents of Control Model, and Table 1 for the hypotheses).

Figure 2. Antecedents of Control Model; explaining how a sense of control may be achieved through primary and secondary control mechanisms during the COVID-19 pandemic.

Table 1*Overview of pre-registered hypotheses and models*

Model	Hypotheses
Threat, Lockdown & Mental Health Model	Personal threat will be predicted by the perceived risk of contracting the virus, which in turn is predicted by contagion rates in one's environment. Finances will further negatively impact threat. Levels of social connectedness are predicted by the social actions one takes; levels of sense of control are predicted by actions to regain control (at the individual or governmental level). Days in isolation will have a quadratic effect on social connectedness and perceived sense of control: initially, levels will be high as isolation does not yet take grip, and later on they will be low as individuals will have adapted to isolation.
Antecedents of Control Model	A perceived sense of control is predicted by (a) frequency of behaviours taken to avoid contracting the virus, (b) communicating with close others, and (c) other types of coping behaviour. In addition, sense of control is predicted by how often an individual consults information sources, and this relationship is moderated by the level of trust in these sources. A perceived sense of control is further predicted by the extent to which governments take actions to deal with the COVID-19 pandemic, and this relationship is moderated by the level of trust in the government.

Methods

Pre-registration

We pre-registered our models and analyses at the Open Science Framework (<https://doi.org/10.17605/OSF.IO/TSJNB>) while data collection was ongoing but prior to examining the data and running analyses.

Ethics

University ethics approval was obtained prior to data collection. All participants read the participant information page and signed an online consent form.

Procedure

Data was collected via an online survey hosted by Qualtrics. The survey was online from 28th of March to 24th of April, 2020, and was advertised on social media, including international COVID-19 groups on Facebook and Reddit. In a snowballing method, the survey link was also e-mailed through the social networks of the authors.

Participants

We collected data from 8,236 participants. Of these, seven individuals were excluded on the basis of pre-registered criteria (see SM.2). The resulting data set contained responses from 8,229 participants, of whom 3245 male (39.4%), 4888 female (59.4%), and 91 other/non-binary (1.1%). Age ranged from 18 to 88 years old ($M = 38.3$, $SD = 13.5$). In descending order of total number of responses, we obtained responses from the following countries: United States ($n=2167$), United Kingdom ($n=1082$), Italy ($n=1029$), Brazil ($n=884$), Australia ($n=489$), the Netherlands ($n=489$), Portugal ($n=367$), Germany ($n=282$), France ($n=237$), Finland ($n=219$), Croatia ($n=209$), and New Zealand ($n=194$), with remaining responses divided among 67 countries with fewer than 100 responses each (total participants $n=377$). See SM.3 for an overview of the descriptive statistics (gender, age, education

levels, days in isolation) and situation (e.g., mandatory self-isolation, key worker) of the participants from the top 12 countries.

Measures

See SM2.3 for example items of the scales. Scoring options were kept similar across scales as much as possible. Standard quantity options were: “Not at all” (0), “A little” (1), “A moderate amount” (2), “A lot” (3), to “A great deal” (4). Standard frequency options were: “Never” (0), “Rarely” (1), “Occasionally” (2), “Sometimes, more than occasionally” (3), “Often” (4), “All the time” (5)¹. For income, government action, and institutional trust, an option “I’d rather not say” was also provided, which was excluded from the analyses. For frequency of communication about COVID-19, the option “We don’t communicate/NA” was excluded from the analyses.

Demographics. Demographics included: Country of residence; Age (in years); Gender (female, male, or other/non-binary); level of education in nine tiers, from “No schooling completed, or less than 1 year” (1) to “Doctorate degree (PhD, EdD, etc.)” (9); household income from “Far below average” (-2) to “Far above average” (2); essential worker job (medical profession; delivery or post personnel, supermarket or shop personnel, garbage or recycling collection; driver in public transport; nursery/childcare/education employee; construction worker; police or armed forces; other; none of the above).

Diagnoses. Participants were asked to indicate whether there had been diagnoses of COVID-19 in any of their environment(s). Options were: no, or global: in my country, in my state/province/county, in my city, and local: in my workplace, school, or children’s nursery, my acquaintances, my close friends, my family, my household, and I’m infected. Answers (yes=1, no=0) were summed in global and local scores.

Days of self-isolation. Participants were asked how many days they had been in self-isolation (voluntary or mandatory).

---

¹ To ensure measurement invariance, we estimated configural and metric invariance models for all scales across countries with >100 respondents. Although most chi-square difference tests were significant, these are overpowered with a large sample size. According to BIC, the metrically invariant models actually had superior fit to the configurally invariant models for all most scales except institutional trust. This means that the scales measure the same construct across countries (although mean levels might vary).

Anxiety and depression. The Hospital Anxiety and Depression Scale [7] was used to measure symptoms of anxiety and depression, with 7 items each. Example of items for anxiety are: “I get a sort of frightened feeling as if something awful is about to happen.”; examples for depression are: “I feel as if I am slowed down.” The score per item is 0-3, total 0-21. We summed the scores to compare to Zigmond & Snaith [7]’s criteria levels: a score of 7 or less reflected ‘non-cases’, scores of 8-10 were ‘doubtful cases’ and scores of 11 or more were ‘definite cases’. In addition, scores of 11-14 were considered ‘moderate’, and 15-21 ‘severe’ [8]. We also computed averages for anxiety ($\alpha = .861$) and depression ($\alpha = .778$), as sum scores are not robust to missing values in structural equation models.

Threat. We measured threat by asking participants how severe they thought the effects of COVID-19 would be on their personal lives (personal threat) and on global society (global threat). Options for answers were: “Not at all” (0), “A little” (1), “Somewhat” (2), “Considerably” (3), “Very” (4), and “Extremely” (5). We used personal threat as a single-item measure, meaning we cannot compute reliability.

Perceived risk. We asked participants to rate, for themselves and their family/close friends, the risk of becoming infected, of experiencing a severe version of the infection, and of dying from the infection. These questions were answered on a scale from “Very low” (-2), “Low” (-1), “Neither low nor high” (0), “High” (1), to “Very high” (2). These scores were averaged ($\alpha = .787$).

Sense of control. We used two items to measure sense of control: “Currently, do you feel that you are the actor in, or the director of, your life?” and “Is the amount of control you experience in life right now lower or higher than you would want, or is it just right?”. Both questions were measured on a sliding scale from -10 to 10 to create an average score ($\alpha = .664$).

Social isolation. Four items of the Social Connectedness Scale [9] measured social isolation. Example: “I catch myself losing all sense of connectedness with society.” These scores were averaged ($\alpha = .885$).

Social actions. Social actions were measured by the frequency with which participants were in contact with their family, friends, and colleagues/fellow students, either through the exchange of messages, calls, or video chats. Options were: “not once/NA” (0), “few times a month” (1), “weekly”

(2), “few times a week” (3), “once a day” (4), to “few times a day” (5). These scores were averaged ($\alpha = .824$).

Avoidance actions. Avoidance actions were measured by COVID-19 safety precautions, adapted from a concurrent Stanford survey [10]. Participants were asked: “In the past week, how much have you made it a priority to...” followed by each of 12 actions, including ‘Wash your hands for at least 20 seconds’, ‘Stay home’, and ‘Wear a mask of any kind’. Options were: “Not at all” (0), “A little bit” (1), “Some” (2), “A lot” (3), to “As much as possible” (4). Scores were averaged ($\alpha = .757$).

PPE availability for essential workers. Key workers in the sample were asked whether they had access to sufficient personal protective equipment (PPE), on a slider from “Not sufficient; nothing” (-10) to “Sufficient; plenty” (10).

Information actions. Participants were asked how often they consulted, and how much they trusted, nine different sources for information about COVID-19, including: the World Health Organization (WHO), national health services, government website(s), newspaper website(s), Facebook, Twitter, Instagram, websites with maps or tables with COVID-19 cases counts, and websites following Google searches about COVID-19. The response options were: “Never” (0), “Rarely” (1), “Occasionally” (2), “Sometimes, more than occasionally” (3), “Often” (4), “All the time” (5) for frequency; and “Not at all” (0), “A little” (1), “A moderate amount” (2), “A lot” (3), to “A great deal” (4) for quantity of trust.

Frequency of communication about COVID-19. Participants were asked how often they spoke about COVID-19 with family, friends, and colleagues/fellow students by exchanging messages, calling, and videochatting. The response options were “Never” (0), “Rarely” (1), “Occasionally” (2), “Sometimes, more than occasionally” (3), “Often” (4), “All the time” (5), and scores were averaged ($\alpha = .824$).

Coping actions. The Brief COPE scale [11] was used, which consists of 28 items reflecting adaptive (e.g., using emotional support; $\alpha = .785$) and maladaptive coping (e.g., substance use; $\alpha = .714$).

Government actions. We included six government actions that help in making quarantine experiences tolerable [12], including: Telling people what is happening and why; Explaining how long it will continue; Providing meaningful activities for people to do while in quarantine; Providing clear communication; Ensuring basic supplies (such as food, water, and medical supplies) are available; Reinforcing a sense of togetherness. Participants were asked to indicate how much they think the government of the country they reside in has accomplished these actions. The response options were: “Not at all” (0), “A little” (1), “A moderate amount” (2), “A lot” (3), to “A great deal” (4). These scores were averaged ($\alpha = .906$).

Institutional trust. Institutional trust was measured through three factors: capability, integrity, and benevolence [6], measured by two items each. Participants were asked “To what degree do you perceive that..” followed by the items, including “...the government is capable of stopping the COVID-19 outbreak?” (capability, $\alpha = .783$); “...the actions of the government in regard to COVID-19 outbreak are driven by sound values and principles?” (benevolence, $\alpha = .885$); “...the government tries to counteract COVID-19 outbreak?” (integrity, $\alpha = .885$). Together, these items constituted institutional trust ($\alpha = .913$). The response options were: “Not at all” (0), “A little” (1), “A moderate amount” (2), “A lot” (3), to “A great deal” (4). For USA respondents, questions were also asked about state capability ($\alpha = .811$), state benevolence ($\alpha = .921$), and state integrity ($\alpha = .924$), together forming state institutional trust ($\alpha = .933$).

Perceived knowledge. Participants were asked how informed they felt with regards to four aspects of COVID-19: The risk of contracting COVID-19, symptoms of COVID-19, how COVID-19 spreads, how to prevent COVID-19 from spreading. Participants were also asked how informed they felt about the treatment of COVID-19, but this option was removed as it was considered too medically niche. Participants were asked to include in their judgment how much they felt they knew, and how much they trusted that information. The response options were: “Not at all” (0), “A little” (1), “A moderate amount” (2), “A lot” (3), to “A great deal” (4). These scores were averaged ($\alpha = .880$).

Actual knowledge To examine how informed the participants were, four facts were gleaned from the WHO’s Whatsapp list of common COVID-19 myths, and participants were asked to indicate if they thought them to be true or false, or to indicate that they did not know. The answer to all fact

statements was “False”: Only people of a certain age can get COVID-19; Pets can transmit COVID-19; Vaccines against pneumonia provide protection against COVID-19. (We also had a fact statement “Taking antibiotics is effective against COVID-19”, but this was removed when it was pointed out that antibiotics do help with the recovery of pneumonic infection following COVID-19). The sum of all ‘false’ (i.e., correct) answers was used as an indicator of actual knowledge.

One of the main hypotheses of this study was that having information in times of threat would increase psychological well-being. Having information was measured through either actual or perceived knowledge (i.e., being or feeling informed). Upon closer inspection, it appeared that the actual knowledge variable has too little variance to meaningful test in this sample. None of the participants got all four myths wrong: correctly judging each of the 4 myths ($n=6924$, 84.3%), 3 myths ($n=1212$, 14.8%), 2 myths ($n=71$, 0.9%), 1 myth ($n=8$, 0.1%), 0 myths ($n=0$). As such, this variable has been excluded from the analyses.

Analyses

All analyses were conducted in R [R Team, 13], using the lme4 package for linear regressions [14], lavaan for structural equation modelling [15], ggplot2 for plotting figures [16], and tidySEM for plotting SEM graphs [17]. This project follows the Workflow for Open Reproducible Code in Science [18]. Regression analyses allow countries (of residence) to have random intercepts, and conditional R^2 and ICC which take these effects into account [19] have been reported. All analysis scripts are publicly available on our project on Open Science Framework (<https://doi.org/10.17605/OSF.IO/TSJNB>).

Power analyses

Using the semPower package in R [20], we conducted a post-hoc power analysis for the Threat, Lockdown & Mental Health Model, and found that a sample size of $N = 8229$ is associated with a power larger than $> 99.99\%$ (more precisely, $1 - 9.812621e-116$) to reject a wrong model (with $df = 27$) with an amount of misspecification corresponding to $RMSEA = .06$ on $\alpha = .05$. Using G*Power [21], post-hoc power analyses for our main two regressions, predicting sense of control and

reduced anxiety, confirmed sufficient power ($>.99$) for the effect sizes $f^2 = .24$ and $.64$ for the regressions respectively (f^2 calculated from regression R^2 with the formula $R^2/(1-R^2)$), given $N = 8229$, an alpha-level of $.05$, and 9 predictors.

Translations

The survey was translated from English into Italian, French, Dutch, German, Portuguese, Brazilian Portuguese, and Spanish by native speakers (an author or a colleague familiar with surveys), and checked by another native speaker. See our project on Open Science (<https://doi.org/10.17605/OSF.IO/TSJNB>) for all full versions of the survey.

Results

First, we examined levels of anxiety ($M = 7.96$, $SD = 4.49$) and depression ($M = 6.97$, $SD = 4.01$) for the 12 main countries, see Figure 3 (see also Figure SM4.2 and SM.5 for comparisons). See Supplementary Materials for an overview of descriptive statistics of the participants (SM.3) and model variables (SM.3).

Figure 3. Scatterplot of average anxiety symptoms by depression symptoms per country (averages of sums of symptom scores).

Threat, Lockdown, & Mental Health Model. We pre-registered a basic model including anxiety, depression, perceived threat, social isolation and sense of control, which was significant (see SM.6). Our second pre-registered model (see Figure 1) added a number of important variables to this basic model. We considered $RMSEA \leq 0.08$, combined with $CFI > 0.90$ and $SRMR \leq 0.08$, to be indicative of an acceptable model fit [22]. Using these criteria, we were able to accept this model ($RMSEA$ (0.06), CFI (0.94), and $SRMR$ (0.04)¹, see Figure 4 and Table SM7.1).

Most of our pre-registered predictions were confirmed: Social isolation positively predicted depression, a reduced perception of control predicted anxiety, and perceived personal threat positively predicted both anxiety and depression. The interaction of social isolation by personal threat was significant. Social isolation also significantly predicted anxiety, and a reduced sense of control predicted depression. In addition, as expected, the more local diagnosed infections, the higher the perceived risk of contracting COVID-19. Perceived risk in turn predicted a higher perceived threat of

COVID-19. Contrary to expectation, though, income level did not predict perceived personal threat, and the quadratic function of days in isolation did not significantly predict sense of control (nor the linear function), though the linear effect did predict social isolation (see SM.8 for scatterplots).

Figure 4. Threat, Lockdown & Mental Health Model results. Standardised coefficients are shown; (residual) variances omitted for clarity, non-significant paths printed in grey, and the dotted line reflects correlation. “Days2” is included to test the quadratic function of days in isolation on social isolation and sense of control. * $p < .05$, ** $p < .01$, *** $p < .001$.

Antecedents of Control Model. Our next pre-registered analysis examined the predictors for sense of control (see Figure 2). Fit indices RMSEA (0.06), CFI (0.54), and SRMR (0.05) of this model were examined, and it was found that these indicated an unacceptable model fit, given our criteria (see above), and we thus rejected the model. See SM.9 for the figure and table of the unexpected model results. One explanation may be that the hypothesised latent variable did not bear out and reduced the power of the analysis. We therefore ran an exploratory regression analysis predicting sense of control

by the main hypothesised mechanisms of control without the hypothesised latent variable, see Figure 5 (and Table SM10.1 for a full regression table).

Next we investigated whether anxiety could be predicted from the same factors. While we found that a lower sense of control predicted anxiety, the level of sense of control had no influence on the effect of personal threat over anxiety. Therefore, we investigated whether the factors influencing control would affect anxiety; see Figure 5 (and Table SM10.2 for a full regression table).

The analyses showed a good fit in predicting sense of control (cond. $R^2 = 0.203$) and anxiety (cond. $R^2 = 0.384$). While sense of control was predicted first and foremost by coping — maladaptive coping (negatively) and adaptive coping (positively) — it was also predicted by frequency of communication about COVID-19 (negatively) and government actions and perceived knowledge (positively). We also found that avoidance actions, institutional trust, and their interaction did not significantly predict sense of control. Anxiety, on the other hand, was predicted by all factors, except for the interaction between government actions and institutional trust.

Figure 5. Plot of regression coefficients for regressions predicting sense of control (blue circles) and reduced anxiety (orange squares) with 95% confidence intervals. Predictor variables are ordered by

magnitude from the sense of control regression, with the exception of the interaction variable (placed last). Estimates are standardised.

Primary control. We were surprised by the lack of efficacy of primary control mechanisms: avoidance actions did not predict sense of control; unexpectedly they predicted an increase rather than a decrease in anxiety. We therefore ran additional regressions predicting control and anxiety with all of avoidance behaviours individually. As the conditional R^2 of the control regression was only 0.025, we will only discuss the anxiety regression (cond. $R^2=.086$, see Table SM11.1). We found that wearing a N-95 or higher (health grade) mask was the only predictor of lower levels of anxiety, whilst wearing a mask of any kind significantly predicted higher levels of anxiety. In addition, washing one's hands for at least 20 seconds, cleaning and disinfecting one's home, using antibacterial products, and avoiding food prepared by someone you do not know were also all predictive of higher levels of anxiety. The same regression was run for participants who indicated that they are essential workers ($n = 942$; Table SM11.2) and it showed a similar pattern, except that the use of antibacterial products in this group was overall not predictive of anxiety. Indeed, essential keyworkers engaged in more actions to avoid contracting the virus ($M = 2.57$, $SD = 0.65$) than participants who were not essential workers virus ($M = 2.51$, $SD = 0.61$, $F(1,8133)=10.09$, $p=.001$, $\eta_p^2=.001$), and also had higher levels of anxiety ($M = 8.28$, $SD = 4.56$, compared to $M = 7.92$, $SD = 4.48$, $F(1,8219)=5.23$, $p=0.2.$, $\eta_p^2= .001$)². Essential workers were also asked whether they felt they had sufficient PPE (see Figure SM11.1 for overall frequencies and Figure SM11.2 for access to PPE per country) and access to sufficient PPE was significantly correlated with lower levels of anxiety symptoms ($r=-.17$, $p<.001$) and depressive symptoms ($r=-.12$, $p<.001$).

---

² Essential workers were also found to have lower levels of social isolation ($M = 1.02$, $SD = 1.02$, compared to $M = 1.23$, $SD = 1.08$, $F(1,8212)=30.12$, $p<.001.$, $\eta_p^2=.004$), despite being in less contact with friends, family, and colleagues ($M = 2.21$, $SD = 0.99$) than participants who were not essential workers ($M = 2.29$, $SD = 1.00$, $F(1,8167)=5.35$, $p=.02$, $\eta_p^2=.001$). They did communicate about COVID-19 more ($M = 3.35$, $SD = 0.99$) than participants who were not essential workers ($M = 3.10$, $SD = 1.01$, $F(1,6503)=45.29$, $p<.001$, $\eta_p^2=.007$), with no further significant differences between the groups in the main variables, including depression symptoms (F 's < 2.04 , p 's $< .15$).

Secondary control. Government actions were found to be predictive of both sense of control and reduced anxiety, whereas institutional trust was only associated with reduced anxiety. See Figure 6 for averages of government actions and institutional trust per country (and SM.12 for further breakdowns, including of USA federal and state governments³). Additional exploratory regressions (see SM.13) indicated that positive government actions both contributed to a greater sense of control (cond. $R^2 = 0.062$), and predicted lower levels of anxiety (cond. $R^2 = 0.101$).

Figure 6. Scatterplot of government actions averages and institutional trust averages per country.

Discussion

The global crisis around COVID-19 stands out because of the combination of two main factors: it involves (1) a practically constant threat and (2) global lockdowns. We hypothesised in a pre-registration that many perceive that COVID-19 will have severe effects on their lives, and that this

would be negatively associated with their psychological wellbeing. Moreover, we hypothesised that lockdown measures would be associated with depression symptoms in people who are socially disconnected and anxiety symptoms in those experiencing a sense of loss of personal control. Finally, we hypothesised that a range of behaviours, such as communicating with others, or governmental actions, would be associated with positive mental health outcomes.

Data from over 8,000 participants worldwide showed that overall rates of anxiety ($M = 7.96$) and depression ($M = 6.97$) were at quasi-clinical levels [8]. While the Hospital Anxiety and Depression Scale is typically used with patients, some studies have used the scale in the general population [23]. This study showed averages ranging from 3.9 (57 years and over) to 5.1 (18-65 years old), suggesting that though we were not able to obtain a pre-pandemic baseline, there is a marked elevation in anxiety and depressive symptoms, in line with other recent findings [24, 25]. The United States, the United Kingdom, and Brazil had especially elevated levels of anxiety and depression compared to other countries, with Italy additionally scoring highest on depression, at clinical levels of mild symptoms [8]. This is an alarming finding.

Our main pre-registered hypotheses were confirmed: Perceived personal threat (i.e., thinking that COVID-19 will have severe effects on one's life) predicted anxiety and depression symptoms in our model. The effect of personal threat on depressive symptoms was stronger when one felt socially isolated. A low sense of control predicted anxiety symptoms, but did not further increase the effect of personal threat on anxiety symptoms. The level of local COVID-19 diagnoses increased the perceived risk of contracting the virus, and this significantly increased perceived personal threat, in line with recent COVID-19 research on risk and resilience [26]. Both perceived risk and threat predicted levels of anxiety. Days of lockdown were, as expected, a significant predictor of social isolation, but not of reduced control. We did not find evidence for a hypothesised U-curve for either factor, where initial low levels of stress would increase but then reduce due to habituation.

The COVID-19 pandemic has involved lockdown, social distancing, and quarantine for people across the globe. This has led, amongst other things, to a severe reduction in social connectedness between people. Social connectedness is of pivotal importance to psychological wellbeing: feelings of loneliness and social isolation have been associated with higher mortality [27],

depression [28], and other negative mental and physical health outcomes [29]. The current results show that engaging in frequent communication with close ones (family, friends, colleagues/fellow students) through a variety of sources (messages, calls, and video chats) was generally associated with reduced feelings of social isolation.

Secondary control mechanisms (e.g., coping, government actions) significantly contributed to a higher sense of control, but primary control mechanisms (e.g., actions to avoid contracting COVID-19) did not. When a threat is deemed controllable individuals will adapt their behaviours to address the threat [30]. However, when threat cannot be controlled, defensive reactions may set in, leading to feelings of helplessness [31], which is often associated with depression [32]. In support of the idea that participants in our sample felt there was little they could do to regain control, we found that sense of control predicted both depressive and anxiety symptoms.

We further investigated the puzzling association between avoidance behaviours and increased anxiety, and found that of the listed avoidance behaviours, only wearing high-grade medical masks (e.g., N-95) was associated with lower levels of anxiety. Other common and ostensibly sensible actions, such as washing your hands for at least 20 seconds, were associated with higher levels of anxiety. These results may reflect that individuals who are particularly anxious engage in these activities more often. In line with that idea, the results showed that essential workers with high levels of anxiety engaged in those activities more often too. Moreover, for this group, wearing masks of any kind was associated with higher levels of anxiety, whereas wearing a high health-grade mask (N-95) was associated with lower levels of anxiety. Having access to sufficient PPE was also associated with lower levels of anxiety and depressive symptoms. Thus, whilst professional actions to reduce the chance of contracting the virus reduce anxiety, other avoidance actions are likely performed more often (at home) by individuals with high levels of anxiety.

As expected, adaptive coping mechanisms such as acceptance and humour were associated with higher levels of sense of control and lower levels of anxiety, whereas maladaptive coping mechanisms such as denial and substance use were associated with higher levels of anxiety, in line with previous findings [33]. Interestingly, frequency of COVID-19 communication –hypothesised to reflect the seeking out of information about COVID-19 – was associated with lower levels of sense of

control. It may be that low levels of sense of control and high levels of anxiety feed the need to talk more frequently about COVID-19.

A recent review [12] listed what governments can do to alleviate suffering due to quarantine, and in our survey we found that the perception that the government had undertaken many of these actions was associated with higher levels of sense of control and lower levels of anxiety. Institutional trust — the belief in the government’s capability, benevolence, and integrity to deal with the COVID-19 outbreak —independently predicted lower levels of anxiety. Trust in the government is extremely important; during the Ebola outbreak, trust in the Liberian government was correlated with taking precautions and compliance with government-instituted control policies [34] and the use of health services during the outbreak [35]. Therefore, it is paramount that the leaders of national governments and officials of national health institutions put forward factual and reliable information.

There are some limitations to this research, such as the use of a cross-sectional survey, which means that causal inferences cannot be made, and that we did not assess levels of pre-existing anxiety and depression. We made this decision on the basis of debates about the quality of, and biases in, the retrospective recall of emotional states [36]. It should be noted that the participants in our sample were, on average, relatively highly educated and predominantly —though not exclusively- from Western countries. This might limit the generalizability of our findings. Due to the employed study design and samples, we were not able to directly compare countries. However, we hypothesised that people’s psychological responses to the pandemic would be generally uniform given the universal themes of lack of social contact and feelings of threat due to the pandemic. Another limitation was that we could not use our measurement of actual COVID-19 knowledge due to too little variance. Luckily, this reflects that four common myths about COVID-19 were not believed widely in the current large sample. Moreover, recent research has shown that individuals’ perceived knowledge about COVID-19 infection was more predictive of emotional well-being than actual knowledge [24].

Conclusions

Disasters, whether natural or human-made, are becoming more common occurrences [37, 38], with such traumatic events being experienced by people almost every day [39]. Often those who

experience and survive these disasters show a variety of psychological problems, including generalised anxiety disorder and major depression disorder [40]. These psychological issues are often exacerbated –if not caused – by additional problems, such as loss of income and disruption of social networks and daily activities [41]. It is therefore paramount that factors that might mitigate these issues are identified. The results of our study indicate that some individual and social actions, such as maintaining frequent social contact and governments taking positive actions to manage a public health crisis, are associated with psychological wellbeing during lockdown. This means that citizens and governments could together engage in helpful actions to adapt better to restrictive measures during the current and future pandemics.

Funding Sources: This study was performed without funding.

Conflicts of Interest: All listed authors declare that they have no conflicts of interest.

Human Rights: All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Informed Consent: Informed consent was obtained from all individual participants included in the study.

Welfare of Animals: This article does not contain any studies with animals performed by any of the authors.

Acknowledgements: We thank the following external translators for their work: Lotte Pummerer (German), Francesca Mellano (French), Ingrid Ruesink (French, Dutch), Everton Maraldi (Brazilian Portuguese), and Lluís Oviedo (Spanish). We received no funding for this work.

References

1. Statistica. (2020). *Monthly unemployment rate in the United States from May 2019 to May 2020*. Retrieved from <https://www.statista.com/statistics/273909/seasonally-adjusted-monthly-unemployment-rate-in-the-us/>: <https://www.statista.com/statistics/273909/seasonally-adjusted-monthly-unemployment-rate-in-the-us/>
2. Rothbaum, F., Weisz, J. R., & Snyder, S. S. (1982). Changing the world and changing the self: A two-process model of perceived control. *Journal of personality and social psychology*, 42(1), 5-37.
3. Kay, A. C., Gaucher, D., Napier, J. L., Callan, M. J., & Laurin, K. (2008). God and the government: testing a compensatory control mechanism for the support of external systems. *Journal of personality and social psychology*, 95(1), 18-35.
4. Heckhausen, J., & Schulz, R. (1995). A life-span theory of control. *Psychological review*, 102(2), 284.
5. Skinner, E. A. (1996). A guide to constructs of control. *Journal of personality and social psychology*, 71(3), 549-570.
6. Mayer, R. C., Davis, J. H., & Schoorman, F. D. (1995). An integrative model of organizational trust. *Academy of management review*, 20(3), 709-734.
7. Zigmond, A. S., & Snaith, R. P. (1983). The hospital anxiety and depression scale. *Acta psychiatrica scandinavica*, 67(6), 361-370.
8. Stern, A. F. (2014). The hospital anxiety and depression scale. *Occupational Medicine*, 64(5), 393-394.
9. Lee, R. M., & Robbins, S. B. (1995). Measuring belongingness: The social connectedness and the social assurance scales. *Journal of counseling psychology*, 42(2), 232-241.
10. Zion, S., Heathcote, L. C., Horii, R., Leibowitz, K., Louis, K., & Crum, A. (2020). The role of mindsets in the context of the COVID-19 viral outbreak. Retrieved March 17
11. Carver, C. S. (1997). You want to measure coping but your protocol's too long: Consider the brief COPE. *International journal of behavioral medicine*, 4(1), 92-100.
12. Brooks, S. K., Webster, R. K., Smith, L. E., Woodland, L., Wessely, S., Greenberg, N., & Rubin, G. J. (2020). The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *The Lancet*, 395(10227), 912-920.
13. R Core Team. (2020). R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from <https://www.R-project.org/>
14. Bates, D., Mächler, M., Bolker, B., & Walker, S. (2014). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, 67(1), 1-48. doi:10.18637/jss.v067.i01
15. Rosseel, Y. (2012). lavaan: An R Package for Structural Equation Modeling. *Journal of Statistical Software*, 48(2), 1-36. Retrieved from <http://www.jstatsoft.org/v48/i02/>
16. Wickham, H. (2016). *ggplot2: elegant graphics for data analysis*. New York: Springer-Verlag.
17. Van Lissa, C. J. (2020). tidySEM: Tidy Structural Equation Modeling. Retrieved from <https://cjvanlissa.github.io/tidySEM>
18. Van Lissa, C. J., Brinkman, L., Vreede, B., van de Schoot, R., & Brandmaier, A. M. (2020). WORCS: A Workflow for Open Reproducible Code in Science.
19. Nakagawa, S., Johnson, P. C., & Schielzeth, H. (2017). The coefficient of determination R² and intra-class correlation coefficient from generalized linear mixed-effects models revisited and expanded. *Journal of the Royal Society Interface*, 14(134), 20170213.
20. Moshagen, M., & Erdfelder, E. (2016). A new strategy for testing structural equation models. *Structural Equation Modeling: A Multidisciplinary Journal*, 23(1), 54-60.
21. Faul, F., Erdfelder, E., Buchner, A., & Lang, A.-G. (2009). Statistical power analyses using G* Power 3.1: Tests for correlation and regression analyses. *Behavior research methods*, 41(4), 1149-1160.
22. Kline, R. B. (2011). *Principles and practice of structural equation modeling*: Guilford publications.
23. Spinhoven, P., Ormel, J., Sloekers, P., Kempen, G., Speckens, A., & Van Hemert, A. (1997). A validation study of the Hospital Anxiety and Depression Scale (HADS) in different groups of Dutch subjects. *Psychological medicine*, 27(2), 363-370.

24. Yang, H., & Ma, J. (2020). How an epidemic outbreak impacts happiness: Factors that worsen (vs. protect) emotional well-being during the coronavirus pandemic. *Psychiatry research*, 289, 113045.
25. Shevlin, M., McBride, O., Murphy, J., Miller, J. G., Hartman, T. K., Levita, L., . . . Stocks, T. V. (2020). Anxiety, Depression, Traumatic Stress, and COVID-19 Related Anxiety in the UK General Population During the COVID-19 Pandemic. Retrieved from <https://psyarxiv.com/hb6nq>.
26. Dryhurst, S., Schneider, C. R., Kerr, J., Freeman, A. L., Recchia, G., Van Der Bles, A. M., . . . van der Linden, S. (2020). Risk perceptions of COVID-19 around the world. *Journal of Risk Research*. 1-13.
27. Holt-Lunstad, J., Smith, T. B., Baker, M., Harris, T., & Stephenson, D. (2015). Loneliness and social isolation as risk factors for mortality: a meta-analytic review. *Perspectives on psychological science*, 10(2), 227-237.
28. Luo, Y., Hawkey, L. C., Waite, L. J., & Cacioppo, J. T. (2012). Loneliness, health, and mortality in old age: A national longitudinal study. *Social Science & Medicine*, 74(6), 907-914.
29. Hawkey, L. C., & Cacioppo, J. T. (2010). Loneliness matters: A theoretical and empirical review of consequences and mechanisms. *Annals of behavioral medicine*, 40(2), 218-227.
30. Witte, K., & Allen, M. (2000). A meta-analysis of fear appeals: Implications for effective public health campaigns. *Health education & behavior*, 27(5), 591-615.
31. Alloy, L. B., & Abramson, L. Y. (1982). Learned helplessness, depression, and the illusion of control. *Journal of personality and social psychology*, 42(6), 1114-1126.
32. Maier, S. F., & Seligman, M. E. (2016). Learned helplessness at fifty: Insights from neuroscience. *Psychological review*, 123(4), 349-367.
33. Mahmoud, J. S. R., Staten, R. T., Hall, L. A., & Lennie, T. A. (2012). The relationship among young adult college students' depression, anxiety, stress, demographics, life satisfaction, and coping styles. *Issues in mental health nursing*, 33(3), 149-156.
34. Blair, R. A., Morse, B. S., & Tsai, L. L. (2017). Public health and public trust: Survey evidence from the Ebola Virus Disease epidemic in Liberia. *Social Science & Medicine*, 172, 89-97.
35. Morse, B., Grépin, K. A., Blair, R. A., & Tsai, L. (2016). Patterns of demand for non-Ebola health services during and after the Ebola outbreak: panel survey evidence from Monrovia, Liberia. *BMJ global health*, 1(1), e000007.
36. Levine, L. J., & Safer, M. A. (2002). Sources of bias in memory for emotions. *Current Directions in Psychological Science*, 11(5), 169-173.
37. Warsini, S., West, C., Ed, G. D., Res Meth, G. C., Mills, J., & Usher, K. (2014). The psychosocial impact of natural disasters among adult survivors: An integrative review. *Issues in mental health nursing*, 35(6), 420-436.
38. Beck, U., Lash, S., & Wynne, B. (1992). *Risk society: Towards a new modernity* (Vol. 17): sage.
39. Norris, F. H., Friedman, M. J., & Watson, P. J. (2002). 60,000 disaster victims speak: Part II. Summary and implications of the disaster mental health research. *Psychiatry: Interpersonal and biological processes*, 65(3), 240-260.
40. Norris, F. H., Friedman, M. J., Watson, P. J., Byrne, C. M., Diaz, E., & Kaniasty, K. (2002). 60,000 disaster victims speak: Part I. An empirical review of the empirical literature, 1981–2001. *Psychiatry: Interpersonal and biological processes*, 65(3), 207-239.
41. Nikapota, A. (2006). After the tsunami: A story from Sri Lanka. *International Review of Psychiatry*, 18(3), 275-279.

**The Psychological Impact of Threat & Lockdowns during the COVID-19 Pandemic:
Exacerbating Factors and Mitigating Actions**

Supplementary Materials

SM.2 Methods elaborations

SM2.1 Ethics elaboration

All participants read the participant information page and signed an online consent form to confirm that they had read and had understood the participant information, that they were voluntarily taking part in the survey, that they understood that they were able to stop participating at any time without any consequences, and that they were aged 18 or over.

SM2.2 Participant exclusion elaboration

As per the pre-registration, we investigated the completed responses to our survey, and all non-serious responses were removed from the dataset. We started with checking how many days everyone had been self-isolating: we excluded all participants who said they had been self-isolating for longer than 200 days, because by the time the survey stopped (24th of April), 115 days had passed since the coronavirus was first flagged up in Wuhan, China (31st of December 2019 was the first time a cluster of cases of pneumonia was reported in Wuhan, China, from which the novel coronavirus was eventually identified; WHO). This means the participants were either not responding seriously to the survey (one participant claimed they used a crystal ball to find information about COVID-19), or were in some extraordinary circumstances outside of COVID-19 (one participant admitted to “I do not engage with life ordinarily and have been self-isolating for a few years”). Following these data exclusion criteria meant to exclusion of one woman from Afghanistan, two men from Germany, two men from the United States, and one individual from Monaco. One more participant indicated they were 150 years old, which is less probable score, and upon investigation it appeared that this data was not entered in seriousness, and so this male from the United States was also removed. This means that the data of a total of seven individuals was removed from the data set of completed responses.

SM2.3 Measures elaboration

Removed item: Avoid Asian food/restaurants

Of the avoidance actions, one item (“Avoid Asian food/restaurants”) was considered racist by several participants and as such was excluded from the analyses.

SM.3 Descriptive statistics for participants

Out of all participants, 8219 filled out their country of residence (n=10 missing data), and 8,223 filled in their gender (n=6 missing).

Table SM3.1

Descriptive statistics of participants by country

Country	N	Gender			Age range	M (SD)
		Male	Female	Other		
United States	2167	658 (30.4%)	1481 (68.3%)	27 (1.2%)	18-88	37.3 (12.1)
United Kingdom	1082	457 (42.2%)	607 (56.1%)	18 (1.7%)	18-81	35.0 (11.4)
Italy	1029	314 (30.6%)	711 (69.2%)	2 (0.2%)	18-82	45.6 (14.7)
Brazil	884	526 (59.5%)	347 (39.3%)	11 (1.2%)	18-73	33.3 (11.3)
Australia	683	247 (36.2%)	427 (62.5%)	9 (1.3%)	18-76	43.6 (12.6)
Netherlands	489	214 (43.8%)	271 (55.4%)	4 (0.8%)	18-77	36.4 (13.2)
Portugal	367	162 (44.1%)	205 (55.9%)	0 (0.0%)	18-82	46.0 (17.9)
Germany	282	124 (44.0%)	154 (54.6%)	4 (1.4%)	18-80	38.8 (12.2)
France	237	117 (49.4%)	119 (50.2%)	1 (0.4%)	18-85	38.9 (16.4)
Finland	219	116 (53.2%)	98 (45.0%)	4 (1.8%)	18-83	34.9 (11.4)
Croatia	209	105 (50.2%)	103 (49.3%)	1 (0.5%)	18-68	31.0 (9.5)
New Zealand	194	60 (30.9%)	132 (68.0%)	2 (1.0%)	18-80	38.0 (12.4)
All	8229	3246 (39.4%)	4888 (59.4%)	90 (1.1%)	18-88	38.3 (13.5)

Note. Countries approximating 200 participants or more included (all other country samples comprised of fewer than 60 participants each), ordered by participant numbers. The 'Other' gender included non-binary.

A question about USA states was added after 591 USA responses. The top responding states were Texas (193M, 275F, 5N; 474 total), Massachusetts (160M, 213F, 5N; 378 total), Washington (33M, 91F, 5N; 129 total), and New Mexico (37M, 67F, 2N; 106 total), with all other states fewer than 100 participants reported each.

Table SM3.2*Percentages of participants for each education level per country and on average (of total)*

Country	No schooling completed, or less than 1 year	Nursery, kindergarten, and primary school	Some secondary school	Completed secondary school	College/A- levels	Associate's degree (including occupational or academic degrees)	Bachelor's degree (BA, BSc, AB, etc.)	Master's degree (MA, MSc, MENG, MSW, etc.)	Doctorate degree (PhD, EdD, etc.)
United States	0.3	0.4	3.2	7.9	13.5	5.7	35.4	19.1	14.6
United Kingdom	0.1	0.0	1.4	6.6	16.1	10.0	34.8	22.1	8.9
Italy	0.0	0.0	6.6	13.6	24.8	15.1	13.1	22.9	3.8
Brazil	0.0	0.1	0.2	4.7	14.8	8.6	46.2	17.2	8.1
Australia	0.0	0.0	3.8	9.6	3.7	22.0	37.3	17.3	6.2
Netherlands	0.2	0.0	0.8	8.5	8.5	12.5	35.1	28.9	5.4
Portugal	0.0	0.0	1.1	4.4	11.3	2.8	51.4	24.6	4.4
Germany	0.0	0.0	2.9	6.9	15.6	19.3	20.0	27.3	8.0
France	0.9	0.0	2.1	3.9	5.2	11.6	17.2	51.1	8.2
Finland	0.0	0.9	4.7	10.2	11.2	12.1	31.2	26.5	3.3
Croatia	0.0	0.0	1.9	20.7	10.1	2.9	25.0	34.1	5.3
New Zealand	0.0	0.0	4.2	11.5	7.9	20.9	38.7	9.9	6.8
Average	0.1	0.1	2.8	8.2	13.4	10.6	32.2	22.1	8.7

Table SM3.3

Number of participants for each essential worker category per country and on % of total participants (ppts) and keyworkers

Country	Medical profession	Delivery / post personnel	Supermarket / shop personnel	Garbage / recycling collection	Driver in public transport	Nursery / childcare / education employee	Construction worker	Police or armed forces	Other	% of total ppts
United States	72	9	19	0	1	4	10	7	128	11.5
United Kingdom	28	2	12	0	3	9	1	1	39	8.8
Italy	151	2	10	0	1	2	2	4	60	22.5
Brazil	33	1	7	0	1	0	1	4	19	7.5
Australia	42	0	9	1	1	10	2	5	32	14.9
Netherlands	9	2	7	0	1	6	0	1	15	8.4
Portugal	21	0	3	0	0	0	0	0	13	10.1
Germany	12	1	3	0	0	1	0	1	17	12.4
France	7	1	0	0	0	0	0	0	4	5.1
Finland	9	0	0	0	0	1	0	1	7	8.2
Croatia	10	1	1	0	0	0	0	0	6	8.6
New Zealand	5	2	2	0	0	0	0	0	6	7.7
% of total keyworkers	43.3	2.3	7.9	0.1	0.9	3.6	1.7	2.6	37.6	11.7

Table SM3.4*Percentages for ‘current situations’ of participants in each country at time of survey completion*

	Mandatory self-isolation	Voluntary self-isolation	Restricted normal	Normal	Essential worker
United States	32.5	52.8	3.0	0.1	11.5
United Kingdom	30.7	57.4	3.0	0.1	8.8
Italy	34.8	29.2	13.0	0.0	22.5
Brazil	27.5	62.2	2.8	0.0	7.5
Australia	14.7	56.8	13.2	0.3	14.9
Netherlands	10.7	74.0	7.0	0.0	8.4
Portugal	27.3	57.1	5.5	0.0	10.1
Germany	15.0	55.0	16.8	0.7	12.4
France	83.5	10.1	0.8	0.4	5.1
Finland	8.7	71.2	11.0	0.9	8.2
Croatia	13.9	71.3	6.2	0.0	8.6
New Zealand	76.8	14.4	1.0	0.0	7.7
Average	30.2	51.8	6.1	0.2	11.5

Note. Restricted normal means people go about their daily business (work, school, etc.), but do not visit restaurants, cinemas, etc., whereas normal includes all of these.

SM.4 Descriptive statistics and figures for model variables**Table SM4.1***Descriptive statistics for variables in the pre-registered models*

	Type	Range	Mean	SD	nr ppts
<i>Model 1 (simple model) variables</i>					
Perceived personal threat	Avg	0-5	2.84	1.19	8212
Anxiety	Sum	0-21	7.96	4.49	8222
Depression	Sum	0-21	6.97	4.01	8222
Sense of control	Avg	-10-10	-2.07	4.77	8071
Social isolation	Avg	0-4	1.20	1.08	8215
<i>Model 2 (main model) variables</i>					
Income	Avg	-2-2	.24	1.08	7939
Perceived risk	Avg	0-4	1.84	.77	8230
Local diagnoses	Sum	0-6	.58	.84	8230
Social actions	Avg	0-5	2.28	1.00	8170
Days in isolation	Avg	0-115	23.0	9.5	6720
<i>Model 3 (perceived control) variables</i>					
Avoidance actions	Avg	0-4	2.51	.62	8136
Communicating about COVID	Avg	0-5	3.13	1.01	6506
Coping actions	Avg	0-4	1.06	.32	8181
Frequency of checking sources	Avg	0-5	1.83	.81	8127
Trust in checked sources	Avg	0-4	1.59	.53	8066
Government actions	Avg	0-4	1.62	1.08	8003
Institutional trust	Avg	0-4	1.97	1.15	7994

Note: Columns indicate type of calculation (average or sum), range, mean and standard deviation (SD), and number of participants who completed the variable items in the survey (nr ppts).

Table SM4.2*Table with lockdown start and days in isolation by country*

Country	Lockdown start	Comparison to project start	Range days in isolation	Average days in isolation (SD)
Australia	March 23	-5 days	0-110	28.0 (11.1)
Brazil		-	0-90	23.5 (8.1)
Santa Catarina	March 17	-11 days	-	-
Sao Paolo	March 24	-4 days	-	-
Croatia	March 18	-10 days	0-60	15.2 (6.5)
Finland	March 16**	-12 days	0-80	30.5 (12.1)
France	March 17	-11 days	2-42	20.4 (5.2)
Germany	March 23	-5 days	4-87	21.1 (9.3)
Italy	March 11*	-19 days	0-70	26.2 (9.1)
Netherlands	March 23**	-5 days	0-70	20.4 (7.4)
New Zealand	March 19	-9 days	3-60	23.5 (7.5)
Portugal	March 19	-9 days	0-60	21.2 (5.7)
United Kingdom	March 23	-5 days	0-100	19.1 (9.6)
USA		-	0-79	23.7 (8.8)
Texas	April 7	+11 days	3-79	25.1 (8.8)
Massachusetts	March 23	-5 days	0-63	25.4 (6.9)
Washington	March 23	-5 days	7-50	27.4 (8.3)
New Mexico	March 30	+2 days	7-46	25.9 (7.2)

*Note. Survey start and end dates: March 28th - April 24th .***Some areas had lockdown implemented from February 23rd****A less strict lockdown was implemented (e.g., 'intelligent lockdown' in the Netherlands)*

Figure SM4.1: Frequency histograms of perceived threat by country. Perceived threat reflects how severe participants think the effects of COVID-19 will be on their personal lives from 0 “Not at all” to 5 “Extremely”. Dotted blue line indicates country average. Order of countries is alphabetical.

Figure SM4.2. Density plots of anxiety symptoms (pink) and depressive symptoms (turquoise). Score is sum (range from 0 to 21) by country. Dotted lines reflect country averages. Order of countries is alphabetical.

Figure SM4.3: Ridgeline plots of perceived sense of control by country. Black lines are the median. Order of countries is descending average sense of control.

Figure SM4.4: Ridgeline plots of social isolation by country. Black lines are the median. Order of countries is descending average social isolation.

SM.5 Anxiety and depression per country**Table SM5.1***Means and standard deviations for anxiety and depression by country*

	Anxiety		Depression	
	M	SD	M	SD
Australia	6.63	4.14	6.31	3.81
Brazil	8.86	4.57	7.22	3.92
Croatia	7.57	4.22	6.52	3.73
Finland	5.81	3.95	5.49	3.33
France	6.35	4.11	5.33	3.62
Germany	7.09	4.35	6.36	3.93
Italy	6.99	3.59	8.22	4.10
Netherlands	7.00	4.23	6.03	3.81
New Zealand	6.29	4.44	5.95	3.60
Portugal	6.87	3.90	5.45	3.38
United Kingdom	8.23	4.67	7.36	4.17
United States	9.59	4.57	7.39	4.01
Average	8.00	4.50	6.99	4.01

Note. Both variables were measured on a scale from 0-21.

SM.6 Basic Model of threat, anxiety and depression

Figure SM6.1: The basic model; reflecting the effect of personal threat on mental health and its potential mitigating factors during the COVID-19 pandemic

Table SM6.1

Pre-registered hypotheses for the Basic Model

Basic Model	A higher perceived personal threat will increase levels of anxiety and depression.
	The relationship between personal threat and well-being will be moderated by levels of social connectedness and sense of control.

Results for the Basic Models

The model has 0 degrees of freedom, so it has perfect fit (CFI = 1.00). Most of our pre-registered predictions were confirmed: Social isolation positively predicts depression, a reduced perception of control predicts anxiety, and perceived personal threat significantly and positively predicts both anxiety and depression. The interaction of social isolation by personal threat is significant, but the interaction of sense of control by personal threat was not. Social isolation also significantly predicts anxiety, and a reduced sense of control predicts depression.

Figure SM6.2. Basic Model results. Standardised coefficients are shown; (residual) variances omitted for clarity, and non-significant paths printed in grey, the dotted line reflects a correlation.

Table SM6.2*Full model parametrization of the Basic Model*

Pathway	Estimate	CI	SE	<i>p</i>-value
Depression on perceived threat	0.15***	[0.14, 0.17]	0.01	<.001
Depression on sense of control	-0.30***	[-0.32, -0.28]	0.01	<.001
Depression on social isolation	0.33***	[0.31, 0.35]	0.01	<.001
Depression on perceived threat moderated by sense of control	0	[-0.02, 0.02]	0.01	0.95
Depression on perceived threat moderated by social isolation	0.03**	[0.01, 0.05]	0.01	<.001
Anxiety on perceived threat	0.28***	[0.26, 0.30]	0.01	<.001
Anxiety on sense of control	-0.27***	[-0.29, -0.26]	0.01	<.001
Anxiety on social isolation	0.25***	[0.23, 0.27]	0.01	<.001
Anxiety on perceived threat moderated by sense of control	-0.02	[-0.04, 0.00]	0.01	0.10
Anxiety on perceived threat moderated by social isolation	0.01	[-0.01, 0.03]	0.01	0.30
Depression with anxiety	0.43***	[0.42, 0.45]	0.01	<.001

Note. All values are standardized; standardized coefficients shown, CI= 95% confidence intervals.

*** $p < .001$

SM.7 Full model parametrization of Threat, Lockdown, and Mental Health Model**Table SM7.1***Full model parametrization of the Threat, Lockdown, and Mental Health Model*

Pathway	Estimate	CI	SE	<i>p-value</i>
Perceived risk on local diagnoses	0.05***	[0.02, 0.07]	0.01	<.001
Perceived threat on income	-0.01	[-0.03, 0.02]	0.01	0.51
Perceived threat on perceived risk	0.30***	[0.28, 0.32]	0.01	<.001
Social isolation on social actions	-0.22***	[-0.24, -0.20]	0.01	<.001
Social isolation on days in isolation	0.06***	[0.03, 0.08]	0.01	0.01
Social isolation on days in isolation^2	-0.01	[-0.04, 0.01]	0.01	0.39
Sense of control on days isolation	<0.01	[-0.02, 0.03]	0.01	0.82
Sense of control on days in isolation^2	0.01	[-0.01, 0.04]	0.01	0.31
Perceived threat with social isolation	0.18***	[0.16, 0.20]	0.01	<.001
Social isolation with sense of control	-0.42***	[-0.44, -0.40]	0.01	<.001
Perceived threat with sense of control	-0.19***	[-0.21, -0.17]	0.01	<.001
Depression on social actions	0.04***	[0.02, 0.06]	0.01	<.001
Depression on days isolation	0.02*	[0.00, 0.04]	0.01	0.02
Depression on local diagnoses	-0.05***	[-0.06, -0.03]	0.01	<.001
Depression on perceived risk	0.06***	[0.04, 0.08]	0.01	<.001
Depression on perceived threat, moderated by social isolation	0.03*	[0.01, 0.05]	0.01	0.01
Depression on social isolation	0.35***	[0.33, 0.37]	0.01	<.001
Depression on perceived threat	0.13***	[0.11, 0.16]	0.01	<.001
Depression on sense of control	-0.31***	[-0.33, -0.29]	0.01	<.001

Depression on perceived threat, moderated by sense of control	0.01	[-0.02, 0.03]	0.01	0.62
Anxiety on social actions	0.08***	[0.06, 0.10]	0.01	<.001
Anxiety on days isolation	-0.03**	[-0.05, -0.01]	0.01	0.01
Anxiety on local diagnoses	0.03**	[0.01, 0.05]	0.01	<.001
Anxiety on perceived risk	0.17***	[0.15, 0.19]	0.01	<.001
Anxiety on perceived threat, moderated by social isolation	0.01	[-0.01, 0.04]	0.01	0.21
Anxiety on social isolation	0.27***	[0.25, 0.29]	0.01	<.001
Anxiety on perceived threat	0.22***	[0.20, 0.24]	0.01	<.001
Anxiety on sense of control	-0.27***	[-0.29, -0.25]	0.01	<.001
Anxiety on perceived threat, moderated by sense of control	-0.01	[-0.03, 0.01]	0.01	0.32

Note. All values are standardized; standardized coefficients shown, CI= 95% confidence intervals.

*p<.05, **p<.01, ***p<.001

SM.8 Days in isolation

Figure SM8.1 Scatterplot of days in isolation by sense of control with a curvilinear function fitted

Figure SM8.2 Scatterplot of days in isolation by feelings of social isolation with a curvilinear function fitted

SM.9 Antecedents of Control Model

Figure SM9.1: Antecedents of Control Model results. Standardized coefficients are shown; (residual) variances omitted for clarity, and non-significant coefficients printed in gray. * $p < .05$, ** $p < .01$, *** $p < .001$. Media is a latent variable, 'Int' stands for interaction (between frequency and trust of the sources), fb=Facebook, gov=government websites, ig=Instagram, maps=COVID-19 maps, who=WHO, tw=Twitter, nhs=websites of country-specific national health services. Freqcomcov=frequency of which communications with close ones is about COVID-19.

Note that the interaction terms between frequency and trust for each source (i.e., the frequency with which you check sources that you trust) was computed and used those as latent variable indicators for the latent variable 'Media'.

Table SM9.1

Fit indices of Antecedents of Control Model

	parameters	chisq	df	AIC	BIC	RMSEA	CFI	SRMR
model	40	2588.72	103	336496.8	336496.8	0.055	0.543	0.047

Table SM9.2*Full model parametrization of Antecedents of Control Model*

Pathway	Estimate	CI	SE	<i>p-value</i>
Media by interaction of Facebook	0.07***	[0.04, 0.09]	0.01	<.001
Media by interaction of Google	0.11***	[0.09, 0.14]	0.01	<.001
Media by interaction of government websites	0.46***	[0.43, 0.48]	0.01	<.001
Media by interaction of Instagram	0.04**	[0.01, 0.07]	0.01	0.01
Media by interaction of COVID maps	0.13***	[0.11, 0.16]	0.01	<.001
Media by interaction of newspapers	0.18***	[0.15, 0.20]	0.01	<.001
Media by interaction of NHS websites	0.74***	[0.71, 0.77]	0.02	<.001
Media by interaction of Twitter	0.05***	[0.03, 0.08]	0.01	<.001
Media by interaction of WHO	0.49***	[0.47, 0.52]	0.01	<.001
Sense of control on feeling informed	0.06***	[0.04, 0.09]	0.01	<.001
Sense of control on avoidance actions	0.01	[-0.02, 0.03]	0.01	0.60
Sense of control on media	0.06***	[0.03, 0.09]	0.01	<.001
Sense of control on communicating about COVID	-0.12***	[-0.15, -0.10]	0.01	<.001
Sense of control on coping actions	0.04***	[0.02, 0.06]	0.01	<.001
Sense of control on government actions	0.14***	[0.10, 0.18]	0.02	<.001
Sense of control on institutional trust	0.03	[-0.01, 0.07]	0.02	0.13
Sense of control on government actions, moderated by institutional trust	0.01	[-0.02, 0.03]	0.01	0.61
Feeling informed on media	0.05***	[0.02, 0.08]	0.01	<.001

Note. All values are standardized; standardized coefficients shown, CI= 95% confidence intervals.

*p<.05 **p<.01, ***p<.001.

SM.10 Regressions predicting sense of control and anxiety**Table SM10.1***Regression of core factors predicting sense of control*

<i>Fixed effects</i>	β	95% CI	<i>t</i>	<i>p</i>
Intercept	0.02	[-0.04 , 0.09]	0.70	0.49
Adaptive coping	0.21	[0.19 , 0.23]	17.23	<.001
Government actions	0.07	[0.03 , 0.12]	3.26	<.001
Perceived knowledge	0.03	[<0.01 , 0.05]	2.19	0.03
Institutional trust	-0.01	[-0.05 , 0.03]	-0.42	0.68
Avoidance actions	-0.02	[-0.05 , <-0.01]	-1.94	0.05
Frequency of COVID-19 communication	-0.07	[-0.10 , -0.05]	-6.27	<.001
Maladaptive coping	-0.37	[-0.39 , -0.35]	-29.71	<.001
Government actions* Institutional trust	0.02	[<-0.01, 0.05]	1.83	0.07
<i>Random effects</i>	Variance	SD		
Country	0.012	0.11		
Residual	0.770	0.88		

Note. Participants (countries): 6034 (70). All variables (predictor; outcome) were standardized before regressing. * $p < .05$, ** $p < .01$, *** $p < .001$.

Fit indices of the sense of control regression

R² (cond.)	ICC (cond.)	AIC	BIC
0.203	0.012	15647.39	15721.15

Table SM10.2*Regression of core factors predicting reduced anxiety*

<i>Fixed effects</i>	β	95% CI	<i>t</i>	<i>p</i>
Intercept	0.04	[-0.01 , 0.10]	1.52	0.14
Adaptive coping	0.07***	[0.05 , 0.09]	6.79	<.001
Government actions	0.08***	[0.04 , 0.12]	4.13	<.001
Perceived knowledge	0.03**	[0.01 , 0.05]	2.64	0.01
Institutional trust	0.07***	[0.03 , 0.11]	3.55	<.001
Avoidance actions	-0.17***	[-0.19 , -0.15]	-16.06	<.001
Frequency of COVID-19 communication	-0.10***	[-0.12 , -0.08]	-10.03	<.001
Maladaptive coping	-0.50***	[-0.52 , -0.48]	-46.76	<.001
Government actions* Institutional trust	-0.01	[-0.04 , 0.01]	-1.30	0.19
<i>Random effects</i>	Variance	SD		
Country	0.010	0.10		
Residual	0.575	0.76		

Note. Participants (countries): 6128 (71). All variables (predictor; outcome) were standardized before regressing. * $p < .05$, ** $p < .01$, *** $p < .001$.

Fit indices of the sense of control regression

R^2 (cond.)	ICC (cond.)	AIC	BIC
0.384	0.010	14105.18	14179.11

SM.11 Avoidance actions, PPE, and anxiety**Table SM11.1***Regression of avoidance actions predicting anxiety*

	β	95% CI	<i>t</i>	<i>p</i>
Intercept	-0.15**	[-0.24 , -0.06]	-3.35	<.001
Wash your hands	0.02	[<-0.01 , 0.05]	1.65	0.10
Wash your hands for at least 20 seconds	0.04**	[0.02 , 0.07]	3.22	<.001
Stay home	0.01	[-0.02 , 0.03]	0.52	0.60
Clean and disinfect your home	0.06***	[0.03 , 0.09]	4.18	<.001
Use antibacterial products	0.05***	[0.02 , 0.07]	3.33	<.001
Not touch your face	0.02	[<-0.01 , 0.05]	1.72	0.09
Avoid shaking hands with other people	-0.00	[-0.03 , 0.02]	-0.27	0.79
Avoid any physical contact with other people	0.00	[-0.02 , 0.03]	0.32	0.75
Wear a mask of any kind	0.08***	[0.05 , 0.10]	5.49	<.001
Wear an N-95 or higher (health grade) mask	-0.03*	[-0.05 , -0.01]	-2.39	0.02
Avoid food prepared by someone you do not know	0.06***	[0.04 , 0.09]	5.59	<.001
<i>Random effects</i>	Variance	SD		
Country	0.044	0.209		
Residual	0.893	0.945		

Note. Participants (countries): 8122 (77). All variables (predictor; outcome) were standardized before regressing. * $p < .05$, ** $p < .01$, *** $p < .001$.

Fit indices of the sense of control regression

R² (cond.)	ICC (cond.)	AIC	BIC
0.086	0.045	22279.47	22377.5

Table SM11.2*Regression of avoidance actions predicting anxiety for essential workers only*

	β	95% CI	t	p
Intercept	0.05	[-0.23 , 0.12]	-0.56	0.58
Wash your hands	<0.01	[-0.09 , 0.08]	-0.10	0.92
Wash your hands for at least 20 seconds	0.11**	[0.03 , 0.19]	2.70	0.01
Stay home	-0.06	[-0.12 , 0.01]	-1.79	0.07
Clean and disinfect your home	0.12**	[0.04 , 0.20]	2.87	<.001
Use antibacterial products	-0.05	[-0.13 , 0.02]	-1.38	0.17
Not touch your face	0.07	[<-0.01, 0.14]	1.84	0.07
Avoid shaking hands with other people	-0.02	[-0.1 , 0.06]	-0.51	0.61
Avoid any physical contact with other people	0.01	[-0.07 , 0.08]	0.13	0.89
Wear a mask of any kind	0.14**	[0.05 , 0.22]	3.15	<.001
Wear an N-95 or higher (health grade) mask	-0.12***	[-0.19 , -0.05]	-3.49	<.001
Avoid food prepared by someone you do not know	0.14***	[0.07 , 0.20]	4.18	<.001
<i>Random effects</i>	Variance	SD		
Country	0.084	0.29		
Residual	0.856	0.93		

Note. Participants (countries): 937 (24). All variables (predictor; outcome) were standardized before regressing. * $p < .05$, ** $p < .01$, *** $p < .001$.

Fit indices of the sense of control regression with essential workers

R^2 (cond.)	ICC (cond.)	AIC	BIC
0.170	0.081	2609.115	2676.913

Figure SM11.1: Access to personal protective equipment (PPE) for essential workers (from -10 “Not sufficient; nothing” to 10 “Sufficient; plenty”). Dotted black line indicates overall mean. Question answered by 895 essential workers worldwide.

Figure SM11.2: Density plots of access to PPE per country. Dotted blue lines indicate country averages.

SM.12 Country differences in government actions and institutional trust

Figure SM12.1. Density plots of perceived government actions to fight the COVID-19 outbreak (pink) and institutional trust (turquoise) per country. Dotted lines indicate country average. Order of countries is alphabetical.

Table SM12.1*Means and standard deviations for government actions and institutional trust by country*

	Government actions		Institutional trust	
	M	SD	M	SD
Australia	2.14	0.98	2.74	0.95
Brazil	0.97	0.82	0.88	0.80
Croatia	2.45	0.84	2.99	0.76
Finland	2.41	0.88	2.88	0.96
France	1.84	0.98	2.49	0.96
Germany	2.16	0.86	2.63	1.03
Italy	2.01	0.79	2.32	0.85
Netherlands	2.32	0.90	2.65	1.01
New Zealand	3.18	0.74	3.50	0.73
Portugal	2.55	0.83	2.62	0.79
United Kingdom	1.53	0.83	1.89	0.94
United States	0.78	0.78	1.21	0.87
Average	1.58	1.08	1.93	1.15

Note. Both variables were measured on a scale from 0-4.

Figure SM12.2: Density plots of institutional trust components – government capability (pink), government benevolence (green), and government integrity (blue) – per country. Dotted lines indicate country averages. Order of countries is alphabetical.

Figure SM12.3: Frequency histograms of institutional trust components (capability, benevolence, and trust) for federal government and state government in the U.S.A. Dotted blue lines indicate country averages.

Capability of the state government ($M = 1.95$, $SD = 1.23$) was estimated significantly higher than that of the federal government ($M = 1.25$, $SD = 1.14$), $t(1366)=-19.25$, $p<.001$, as was the benevolence of the state government ($M = 2.19$, $SD = 1.45$) as compared to that of the federal government ($M = .61$, $SD = .83$), $t(1365)=-37.69$, $p<.001$, as well as the integrity of the state government ($M = 2.55$, $SD = 1.37$) as compared to the federal government ($M = 1.46$, $SD = 1.11$), $t(1347)=-26.40$, $p<.001$.

SM.13 Government actions predicting sense of control and anxiety

Table SM13.1

Regression of the six government actions predicting sense of control

<i>Fixed effects</i>	β	95% CI	<i>t</i>	<i>p</i>
Intercept	0.03	[-0.04, 0.11]	0.71	0.49
Telling people what is happening and why	-0.02	[-0.06, 0.02]	-1.15	0.25
Explaining how long it will continue	0.12***	[0.08, 0.15]	6.86	<.001
Providing meaningful activities for people to do while in quarantine	0.05***	[0.02, 0.08]	3.62	<.001
Providing clear communication	-0.01	[-0.06, 0.03]	-0.69	0.49
Ensuring basic supplies (such as food, water, and medical supplies) are available	0.03	[<0.01, 0.07]	1.92	0.05
Reinforcing a sense of togetherness	0.07***	[0.04, 0.11]	3.80	<.001
<i>Random effects</i>	Variance	SD		
Country	0.020	0.14		
Residual	0.951	0.97		

Note. Participants (countries): 7853 (77). All variables (predictor; outcome) were standardized before regressing. * $p < .05$, ** $p < .01$, *** $p < .001$.

Fit indices of the sense of control regression

R² (cond.)	ICC (cond.)	AIC	BIC
0.062	0.020	21983.76	22046.47

Table SM13.2

Regression of the six government actions predicting anxiety symptoms

<i>Fixed effects</i>	β	95% CI	<i>t</i>	<i>p</i>
Intercept	-0.06	[-0.15, 0.02]	-1.55	0.13
Telling people what is happening and why	-0.08***	[-0.12, -0.04]	-4.17	<.001
Explaining how long it will continue	-0.08***	[-0.11, -0.05]	-4.82	<.001
Providing meaningful activities for people to do while in quarantine	0.03*	[<0.01, 0.06]	1.96	0.05
Providing clear communication	-0.03	[-0.07, 0.01]	-1.60	0.11
Ensuring basic supplies (such as food, water, and medical supplies) are available	-0.09***	[-0.12, -0.06]	-5.51	<.001
Reinforcing a sense of togetherness	-0.04*	[-0.08, <-0.01]	-2.22	0.03
<i>Random effects</i>	Variance	SD		
Country	0.033	0.18		
Residual	0.888	0.94		

Note. Participants (countries): 7989 (78). All variables (predictor; outcome) were standardized before regressing. * $p < .05$, ** $p < .01$, *** $p < .001$.

Fit indices of the sense of control regression

R^2 (cond.)	ICC (cond.)	AIC	BIC
0.101	0.034	21823.89	21886.76