Brief Report: Making Experience Personal: Internal States Language in the Memory Narratives
of Children with and without Asperger's Disorder

Benjamin T. Brown, Georgia Gwinnett College; Gwynn Morris, Meredith College; Robert E.
Nida, Canisius College; & Lynne Baker-Ward, North Carolina State University

Brown, B.T., Morris, G.D., Nida, R., & Baker-Ward, L. (2011). Brief report: Making experience
personal: Internal states language in the memory narratives of children with and without
Asperger's Disorder. *Journal of Autism and Developmental Disorders*. DOI:
10.1007/s10803-011-1246-5

The final publication is available at www.springerlink.com
<http://www.springerlink.com/content/e76743t81h723432/>

Abstract

The development of the personal past is complex, requiring the operation of multiple components of cognitive and social functioning. Because many of these components are affected by autism spectrum disorders, it is likely that autobiographical memory (ABM) in children with Asperger's Disorder (AD) will be impaired. We predicted that the memory narratives of children with AD, in comparison to typically-developing peers, would reflect less personal interpretation as evidenced by internal states language. Thirty children with AD and 20 typically-developing children aged 6 to 14 reported their earliest memories and two emotional experiences (one positive and one negative). Consistent with our predictions, children with AD included fewer emotional, cognitive, and perceptual terms than the comparison sample.

Keywords: Asperger's Disorder, autobiographical memory, narratives, internal states language

Brief Report: Making Experience Personal: Internal States Language in the Memory Narratives
of Children with and without Asperger's Disorder

Given its importance in establishing and maintaining social relationships, autobiographical memory (ABM) can be expected to play a role in Asperger's Disorder (AD), a developmental disorder characterized in part by difficulties in interactions with others (American Psychiatric Association [APA], 2000). The purpose of this investigation was to extend the analysis of the quality of event reports in children with AD. Although adults with Asperger syndrome show impaired memory performance, specifically recall for semantically-related word lists (Bowler, Matthews, & Gardiner, 1997), relatively little work examines the impact of the condition on autobiographical memory which is generally shared through personal narratives. Recent work by Goldman (2008) documented limited inclusion of high points in the narratives of children with high-functioning autism in comparison to language delayed and typically-developing children. We augmented this work by examining the inclusion of terms that connote the individual's interpretation of the experience, including words indicating emotion and cognitive processing. According to Fivush (2009), "it is such internal states language that gives an event its personal significance, linking the experience most closely to self and enabling linkages between self and other" (p. 292). Moreover, Spreng, Mar, and Kim (2009) found neurological evidence of a functional overlap in the processes of theory of mind and autobiographical memory. Less inclusion of internal states language is thus consistent with the limitations in theory of mind and emotion understanding that characterize autism spectrum disorders. Although Collee and colleagues (Colle, Baron-Cohen, Wheelwright, & van der Lely, 2008) did not report a difference in internal states language between adults with and without autism spectrum disorders, Crane, Goddard and Pring (2010) recently reported that adults with

AD extracted less meaning from narratives of personal experiences than did neurotypical adults, as reflected in part by their inclusion of fewer terms indicating insight.

At present, no investigation has examined internal states language in children with autism spectrum disorders. Indeed, little information is available about the development of internal states language in general (Pennebaker & Stone, 2003). Research by Losh and Capps (2006) supports the prediction of less internal states language in children with AD in one category in particular, emotional terms. They presented groups of 7-13 year old children with and without high-functioning autism with emotion cues and asked them to describe experiences consistent with the emotion. The authors described the narratives of the participants with high-functioning autism as “script-like emotional accounts” (p. 816). We extended their investigation by examining specific inclusions of words from multiple categories of internal states language.

Method

Participants

Children with AD. Male children, ages 6 to 14 years, with AD were recruited from a treatment study for children with high-functioning autism spectrum disorders. Female participants with Asperger Syndrome were not available for recruitment, which is consistent with the finding that the incidence of Asperger Syndrome is four times greater in males than females (Ehlers & Gillberg, 1993). Prior to treatment, all of the children met inclusion criteria using a multiple-gate screening process (see Lopata, Thomeer, Volker, Nida, & Lee, 2008; Volker et al., 2010). First, written documentation of a formal diagnosis of Asperger Syndrome was provided by a licensed psychiatrist, physician, or psychologist. Second, parents were required to submit prior testing and evaluation records. A comprehensive records review by a licensed clinical psychologist took place to determine if the testing and evaluation results were consistent with the

diagnostic criteria of AD, as specified in the DSM-IV. Children who met the requirements of the first two gates were then invited to participate in the third gate, which involved formal assessment of intelligence (WISC-IV short-form, Wechsler, 2003) and social-emotional functioning (an Asperger symptomatology scale, a behavior rating scale, and an adaptive scale) by licensed psychological examiners. Parental permission was obtained for 42 children with a confirmed diagnosis of a high-functioning autism spectrum disorder (i.e., AD, autism [high functioning], or Pervasive Developmental Disorder - Not Otherwise Specified (PDD-NOS) [high functioning]), representing about 81% of the children in the treatment study.

While there is ongoing debate regarding whether AD can be reliably distinguished from other ASDs (see Klin, McPartland, & Volkmar, 2005), diagnostic criteria for AD require significant impairment in social interaction and restricted and repetitive behavior and interests, in the absence of significant cognitive or language impairment (APA, 2000). Of the 42 potential participants who met inclusion criteria for the treatment study (i.e., prior diagnosis of an ASD, documented absence of a current language delay, IQ score > 70, evidence of social interaction impairments [≥ 2] and restricted and repetitive behaviors and interests [≥ 1 per diagnostic criteria; APA, 2000]), 11 were excluded due to developmental histories positive for early language delays. In addition to being consistent with diagnostic criteria for AD, this exclusionary criterion was instituted to minimize any potential influence of early language delays, which can affect ABM (see Nelson & Fivush, 2004). An additional participant did not complete the interview. Thus, the final clinical sample consisted of 30 children. The scores of the participants with Asperger syndrome included in the final sample ranged from 63-131 ($M = 107.5$, $SD = 15.6$) on the Asperger Syndrome Diagnostic Scale (Myles, Bock, & Simpson, 2000), indicating a "likely" diagnosis. Based on parents' reports, the AD sample was 93% Caucasian,

6% Hispanic, and 1% biracial and consisted of children from middle and upper-middle income families.

Typically-developing children. Twenty typically-developing male children from middle and upper-middle class families were recruited from the same geographical area as the clinical group. There were no significant differences in the mean age, IQ, or verbal comprehension index (as measured by the WISC-IV short-form) of the sample of typically-developing children [Mean age= 8.95 (2.35) years, Mean IQ = 111.00 (11.53), Mean verbal comprehension = 112.69 (11.83)] and the AD group [Mean age= 9.73 (2.17) years, Mean IQ = 104.64 (11.76). Mean verbal comprehension = 104.71 (14.59)]. Based on parents' reports, the typically developing sample was 97% Caucasian, 2% Hispanic, and 1% Asian American and from middle and upper-middle income families.

Procedure

Children were assessed individually by a psychological examiner. After providing assent, the children with AD were administered two autobiographical memory interviews in a testing session lasting about one hour. The emotional experiences interview was based on a procedure developed by Fivush, Haden, and Adam (1995). Children were asked to provide narratives for two autobiographical memories, one positive and one negative event, of their own choosing. To elicit reports of their earliest memories, the children were asked, "I want you to think way back and tell me the first thing you ever remember, something that happened when you were really little." More prompts were given if needed. In both the emotional and early memory interviews, children's responses were followed up with empty prompts (e.g. "Can you tell me more about that?") until the child could spontaneously provide no more information about the event.

The typically-developing children received a battery of tests that included the IQ assessment and the Memory Interviews in a single testing session lasting four to five hours, with frequent breaks. The IQ assessment was administered prior to other measures. The Autobiographical Memory Interviews were audio-recorded for subsequent coding, and the presentation of the emotional and earliest memory interviews was counterbalanced across participants. In addition, within the emotional memory interview, the order of request for positive and negative events was counterbalanced.

All memories were transcribed verbatim. Identical coding procedures were applied to the narrative accounts of the earliest and the two emotional memories. Internal states language was measured using the Linguistic Inquiry and Word Count software (LIWC, Pennebaker, Francis, & Booth, 2001), which tallies the percentage of words per narrative in each designated category. We examined the emotional (e.g., happy, scared), cognitive (e.g., thought, believed), perceptual (e.g., saw, heard), physiological (e.g., cold, sick), and social content (e.g., we, our) of the children's narratives.

Results

Children provided narratives containing at least two unique pieces of information in response to 85.7% of the prompts, resulting in a total of 126 memory narratives for analysis. All 30 children with AD provided at least one narrative whereas one of the 20 typically-developing children provided no usable narrative. As examined in separate χ^2 analyses, diagnostic group was not associated with the frequency of reports to any of the prompts [$\chi^2(1, N = 50) \leq .87, ps \geq .77$]. A mixed model MANOVA with diagnosis as the between-subject factor and narrative type as the within subject factor demonstrated that all children talk more about recent rather than earliest events (Wilks' lambda=0.56, $F[2, 102]=12.34, p<.01, \eta^2=0.44$), and that narrative length

of earliest, negative, or positive experiences did not differ between children with and without AD (Wilks' $\lambda=0.85$, $F[2, 102]=2.70$, $p > .05$, $\eta^2=0.15$).

Figure 1 depicts the usage of internal states language terms by children with and without AD.

[place figure 1 about here]

MANOVA was applied in order to assess differences in the percentages of words classified into each of several internal states language categories (emotional, cognitive, perceptual, physiological, and social) in the narratives of children with and without AD by memory prompt (earliest, positive, or negative). Two significant main effects emerged from this model: diagnosis (Wilks' $\lambda=0.86$, $F[5, 116]=3.76$, $p<.01$, $\eta^2=0.14$) and Memory Prompt (Wilks' $\lambda=0.82$, $F[5, 115]=2.48$, $p<.01$, $\eta^2=0.10$). The interaction was not significant. Follow up univariate tests localized the difference in ISL usage between children with and without AD to only the use of emotional, perceptual, and cognitive terms; typically-developing children included over one and a half times as many emotional words ($F[1, 126]= 9.11$, $p < .01$, $\eta^2=0.07$) and perceptual words ($F[1, 126]= 5.18$, $p < .03$, $\eta^2=0.04$) and slightly more cognitive words ($F[1, 126]= 4.12$, $p < .05$, $\eta^2=0.03$) in their memory narratives as children with AD. Univariate tests also localized the ISL differences between memory prompts to only the usage of physiological terms, with these terms being reported three times more frequently in earliest and negative memories than in positive memories ($F[2, 126]= 4.73$, $p = .01$, $\eta^2=0.07$).

Because it was not possible to determine if the main effect of memory prompt found above could be attributed to memory characteristics of very early childhood experiences or later affective experiences, a follow-up analysis was conducted in which we combined positive and negative memories into one group. MANOVA was again used to examine differences in the

percentages of internal states language used by children with and without AD by memory type (emotional or earliest). The only significant main effect that emerged from this model was diagnosis (Wilks' lambda=0.88, $F[5, 118]=3.17$ $p=.01$, $\eta^2=0.12$). No other main effect or interaction was significant, indicating that the use of internal states language was related to diagnosis but not the distinction between very early childhood memories and reports of later emotional events. Follow up univariate tests localized the difference in ISL usage between children with and without AD to only the use of emotional and cognitive terms; typically-developing children included over one and a half times as many emotional words ($F[1, 126]=6.49$, $p = .01$, $\eta^2=0.05$) and slightly more cognitive words ($F[1, 126]= 5.20$, $p < .03$, $\eta^2=0.04$) in their memory narratives as children with AD.

Discussion

Our prediction that children with AD would include less internal states language in ABM narratives when compared to typically-developing peers was supported. More specifically, children with AD were less likely to include emotional, perceptual and cognitive terms in their memory narratives. Because the minimal inclusion of emotion characterized earliest memory reports as well as accounts of affective experiences, it appears that it reflects an overall deficit in narrative contents among children with AD.

Perhaps surprisingly, children with AD did not differ from their typically developing peers in their inclusion of social terms in their memory narratives. Social terms, however, are not a part of internal state language (Stone & Pennebaker, 2002) and thus do not reflect interpretation of events. Instead, the high levels of social terms included in these narratives may reflect the extent to which the events that comprise children's lives are experienced in conjunction with other people, including family members and classmates.

Taken together, the limited inclusion of cognitive, perceptual, and emotional terms when describing autobiographical memories is consistent with the possibility that children with AD lack awareness of their own reactions to personal experiences. Future investigations should examine this possibility. Less cognitive term usage in children with AD is consistent with Crane, Goddard and Pring's (2010) findings that adults with AD included fewer insight terms in their personal narratives than did a comparison sample of neurotypical adults. Similarly, Goldman (2008) suggested that high-functioning children with autism provide lists of facts but few interpretations of behavior when reporting personal experiences. This finding fits well with previous research demonstrating limitations that individuals with AD have in theory of mind development and perspective-taking (Baron-Cohen, 1995; Baron-Cohen et al., 2005). Individuals with AD may lack an understanding of self in relation to others; instead, they may experience an intense egocentrism (Frith & De Vignemont, 2005). Because of their difficulties taking the perspective of others, children with AD may not realize the importance of including information about their emotional, cognitive and perceptual experiences in their memory narratives. This interpretation is consistent with classic developmental research that established that egocentric children often engage in collective monologues (Piaget, 1926) ineffectively communicating because of their difficulty in taking the perspectives of others.

Additionally, children with AD may be including fewer emotional terms than their typically-developing peers because of their previously documented difficulties with emotional understanding (Losh & Capps, 2006). If children with AD do not fully comprehend their emotional experiences, they would likely encode and store less emotional information in memory.

Because of the nature of the present clinical sample, we can attribute the differences we found in children's early memory narratives to the social and emotional deficits characteristic of autism spectrum disorders rather than contributions of language development and IQ. In order to qualify for a diagnosis of AD, participants had to display typical intellectual functioning and no delays in language development. Previous research that has examined memory functioning in children with autism spectrum disorders (e.g., Bruck, London, Landa, & Goldman, 2007; Losh & Capps, 2006; Millward, Powell, Messer, & Jordan, 2000) did not distinguish between the performances of children with AD and other autism spectrum disorders. Both intellectual ability and language development contribute to the development of autobiographical memory (Nelson & Fivush, 2004) and are thus both potential confounds in this previous research.

While the present investigation established differences in interpreting and sharing one's personal past in children with and without AD, several limitations warrant mention. One limitation involved the demographic characteristics of the sample. Future studies should increase the generalizability of findings by recruiting samples that include greater diversity with regard to gender, SES, and ethnicity. The present sample size also restricted the number of research questions that could be reasonably examined. Larger scale studies would allow for examination of additional aspects of autobiographical memory, the manner in which children with AD construct these memories, and factors that affect these constructions. Lastly, future studies should examine the manner in which the differences exhibited by children with AD impacts their social lives and/or influences their continuing development.

Individuals who provide memory narratives that include more discussion of their internal states tend to have more fully developed senses of self, more coherent understandings of their lives, and higher levels of physical and psychological health (see McLean & Pasupathi, 2009). If

children with AD do in fact use less internal state language in their memory narratives, it would be expected this limitation would have a negative influence on their identity development and general well-being. Thus, researchers may wish to explore potential interventions designed to increase the use of internal state language in children with AD.

References

- American Psychiatric Association (2000). *Diagnostic and Statistical Manual of Mental Disorders* (4th edition). Washington, DC.
- Baron-Cohen, S. (1995). *Mindblindness: An essay on autism and theory of mind*. MIT Press: Cambridge, MA.
- Baron-Cohen, S., Wheelwright, S., Lawson, J., Griffin, R., Ashwin, C., Billington, J., & Chakrabarti, B. (2005). Empathizing and Systemizing in Autism Spectrum Conditions. In F. R. Volkmar, P. Rhea, A. Klin, & D. Cohen (Eds.), *Handbook of autism and pervasive development disorders, Vol 1: Diagnosis, development, neurobiology, and behavior* (3rd ed.) (pp. 628-639). Hoboken, NJ US: John Wiley & Sons Inc.
- Bowler, D., Matthews, N., & Gardiner, J. (1997). Asperger's syndrome and memory: Similarity to autism but not amnesia. *Neuropsychologia*, 35, 65-70. doi: 10.1016/S0028-3932(96)00054-1
- Bruck, M., London, K., Landa, L., & Goodman, J. (2007). Autobiographical memory and suggestibility in children with autism spectrum disorder. *Development and Psychopathology*, 19, 73-95. doi:10.1017/S0954579407070058
- Colle, L., Baron-Cohen, S., Wheelwright, S., van der Lely, H. K. J. (2008). Narrative discourse in adults with high-functioning autism or Asperger Syndrome. *Journal of Autism and Developmental Disorders*, 38, 28-40. doi:10.1007/s10803-007-0357-5
- Crane, L., Goddard, L., & Pring, L. (2010). Self-defining and everyday autobiographical memories in adults with autism spectrum disorders. *Journal of Autism and Developmental Disorders*, 40, 383-391. doi:10.1007/s10803-009-0875-4

- Ehlers S. & Gillberg C. (1993). The epidemiology of Asperger's syndrome: a total population study, *Journal of Child Psychology and Psychiatry*, 34, 1327-1350. doi: 10.1111/j.1469-7610.1993.tb02094.x
- Fivush, R. (2009). Sociocultural perspectives on autobiographical memory. In M. L. Courage & N. Cowan (Eds.), *The development of memory in infancy and childhood* (2nd ed.). (pp. 283-301). New York, NY: Psychology Press.
- Fivush, R., Haden, C. A., & Adam, S. (1995). Structure and coherence of preschoolers' personal narratives over time: Implications for childhood amnesia. *Journal of Experimental Child Psychology*, 60, 32-56.
- Frith, U., & De Vignemont, F. (2005). Egocentrism, allocentrism, and Asperger syndrome. *Consciousness and Cognition: An International Journal*, 14, 719-738.
doi:10.1016/j.concog.2005.04.006
- Goldman, S. (2008). Brief report: Narratives of personal events in children with autism and developmental language disorders: Unshared memories. *Journal of Autism and Developmental Disorders*, 38, 1982-1988. doi: 10.1007/s10803-008-0588-0.
- Klin, A., McPartland, J., & Volkmar, F. R. (2005). Asperger syndrome. In F. R. Volkmar, R. Paul, A. Klin, & D. Cohen (Eds.), *Handbook of autism and pervasive developmental disorders: Vol 1. Diagnosis, development, neurobiology, and behavior* (3rd ed., pp. 88-125). Hoboken, NJ: John Wiley & Sons.
- Lopata, C., Thomeer, M. L., Volker, M. A., Nida, R. E., & Lee, G. K. (2008). Effectiveness of a manualized summer social treatment program for high-functioning children with autism spectrum disorders. *Journal of Autism and Developmental Disorders*, 38(5), 890-904.

- Losh, M., & Capps, L. (2006). Understanding of emotional experience in autism: Insights from the personal accounts of high-functioning children with autism. *Developmental Psychology*, 42, 809-818. doi:10.1037/0012-1649.42.5.809
- McLean, K. & Pasupathi, M. (2009) Narrative Development in Adolescence: Creating the Storied Self. NY: Springer.
- Millward, C., Powell, S., Messer, D., & Jordan, R. (2000). Recall for self and other in autism: Children's memory for events experienced by themselves and their peers. *Journal of Autism and Developmental Disorders*, 30, 15-28. doi:10.1023/A:1005455926727
- Myles, B. S., Bock, S. J., & Simpson, R. L. (2000). Asperger syndrome diagnostic scale. Austin, TX: PRO-ED.
- Nelson, K., & Fivush, R. (2004). The emergence of autobiographical memory: A social cultural developmental theory. *Psychological Review*, 111, 486-511.
- Pennebaker, J.W., Francis, M.E., & Booth, R.J. (2001). Linguistic Inquiry and Word Count: LIWC 2001. Mahwah, NJ: Erlbaum Publishers
- Pennebaker, J. W., & Stone, L. D. (2003). Words of wisdom: Language use over the life span. *Journal of Personality and Social Psychology*, 85(2), 291-301. doi: 10.1037/0022-3514.85.2.291
- Piaget, J. (1926). The language and thought of the child. New York: Harcourt, Brace, Jovanovich.
- Spreng, R.N., Mar, R.A., & Kim, A.S. (2009). The common neural basis of autobiographical memory, prospection, navigation, theory of mind, and the default mode: a quantitative meta-analysis. *Journal of Cognitive Neuroscience*, 21, 489-510. doi: 10.1162/jocn.2008.21029

- Stone, L. D., & Pennebaker, J. W. (2002). Trauma in real time: Talking and avoiding online conversations about the death of Princess Diana. *Basic and Applied Social Psychology*, 24, 172–182. doi: 10.1111/j.1475-6811.2008.00207.x
- Volker, M. A., Lopata, C., Vujnovic, R. K., Smerbeck, A. M., Toomey, J. A., Rodgers, J. D., Schiavo, A. & Thomeer, M. L. (2010). Comparison of the Bender Gestalt-II and VMI-V in samples of typical children and children with high-functioning autism spectrum disorders. *Journal of Psychoeducational Assessment*, 28(3), 187-200. doi: 10.1177/0734282909348216
- Wechsler, D. (2003). Wechsler Intelligence Scale for Children—Fourth Edition. San Antonio, TX: Psychological Corporation.