Title: Sympathetic nervous system dominance during stress recovery mediates associations between stress sensitivity and social anxiety symptoms in female adolescents

Authors: Tiffany C. Ho, Ph.D.^{a,b*†}, Holly T. Pham^{c†}, Jonas G. Miller, Ph.D.^b, Katharina Kircanski, Ph.D.^d, & Ian H. Gotlib, Ph.D.^b

^aUniversity of California, San Francisco, San Francisco, CA

^bStanford University, Stanford, CA

^cThe Pennsylvania State University, University Park, PA

^dNational Institute of Mental Health, Bethesda, MD

[†]Equal contribution

*Corresponding author details:

Tiffany Ho, Ph.D.

Department of Psychiatry and Weill Institute for Neurosciences

University of California, San Francisco

401 Parnassus Avenue, San Francisco, CA 94143 USA

E-mail: tiffany.ho@ucsf.edu

Phone: +1 (415) 476-7519

Short title: Stress sensitivity, autonomic flexibility, and social anxiety

Keywords: stress, social anxiety disorder, adolescence, respiratory sinus arrhythmia, skin conductance level

Acknowledgments: This research was supported by the National Institutes of Health (NIH; R37MH101495 to IHG, K01MH117442 to TCH, T32MH019908 to JGM) and the Brain and Behavior Research Foundation (Young Investigator Award 20814 to KK). The funding agencies played no role in the preparation of this article. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH. We thank Monica Ellwood-Lowe, M. Catalina Camacho, Sophie Schouboe, Alexandria Price, and Anna Cichocki for their help with data collection, and Lucy King, Kathryn Humphreys, Natalie Colich, and Sarah Ordaz for their help with data organization and coding. Finally, we thank the families who participated in this study.

Abstract

Social anxiety disorder (SAD) is one of the most common anxiety disorders diagnosed during adolescence. SAD is associated with both psychological stress reactivity and heightened physiological arousal; however, no study has systematically examined which aspects of autonomic nervous system function mediate the likely links between stress sensitivity and symptoms of SAD in adolescents. In this study, we assessed 163 adolescents (90 females; 12.29 ± 1.39 years) with respect to severity of life stress and social anxiety symptoms and measured respiratory sinus arrhythmia (RSA) and skin conductance levels (SCL) during a well-validated psychosocial stress paradigm composed of baseline, reactivity, and recovery periods. We operationalized stress sensitivity as the residual variance in subjective stress severity after accounting for objective stress severity and changes in autonomic regulation (relative to baseline) during reactivity and recovery using standardized change scores in RSA and SCL. We found that in females only, stress sensitivity and symptoms of SAD were significantly correlated with each other ($p < 0.001$) and with autonomic regulation during both reactivity and recovery (all $ps < 0.04$). Further, sympathetic nervous system dominance during recovery specifically mediated associations between stress sensitivity and symptoms of SAD ($B = 1.06$, 95% CI: 0.02–2.64). In contrast, in males, stress sensitivity, autonomic regulation during reactivity or recovery, and symptoms of SAD were not significantly associated (all $ps > 0.1$). We interpret these results in the context of psychobiological models of SAD and discuss implications for interventions targeting autonomic processes in this disorder.

Introduction

Social anxiety disorder (SAD) is characterized by a persistent fear of judgment from others that often drives avoidance of social situations, resulting in behaviors that adversely affect individuals' functioning across multiple domains (e.g., social, academic, and occupational; American Psychiatric Association, 2013). The median age of onset for SAD is 13 years (Stein, 2006), making it one of the most common anxiety disorders diagnosed during adolescence (Beesdo, Pine, Lieb, & Wittchen, 2010). Because of the impairments experienced by individuals suffering from SAD and because the presence of social anxiety problems during adolescence is associated with an increased risk for concurrent and subsequent behavioral problems and other severe disorders—including depression, concentration difficulties, and conduct problems (Beesdo et al., 2007; Stein et al., 2001; Van Roy, Kristensen, Groholt, & Clench-Aas, 2009)—it is critical to understand the psychobiological mechanisms that contribute to the emergence of SAD during adolescence. Elucidating these mechanisms will inform the development of more effective, mechanism-driven, evidence-based treatments.

In this context, contemporary models have posited that SAD is characterized by an increased sensitivity to social stress resulting from maladaptive cognitive and emotion regulatory processing (e.g., fear of social rejection, negative self-evaluation). In particular, a growing body of literature has suggested that individuals vulnerable to SAD and other anxiety disorders tend to respond to stressors with pronounced negative affect—that is, they have greater *stress sensitivity* to negative events (Farmer & Kashdan, 2015). Heightened responses to life stressors may reflect a propensity to perceive interpersonal events negatively, thereby contributing to behaviors that are intended to minimize distress for the individual but that often generate or perpetuate interpersonal conflict (e.g., avoidance, withdrawal), leading to sustained symptoms of SAD

(Hammen, 2015). Indeed, researchers have demonstrated that adolescents with a history of SAD symptoms report experiencing stress in response to social-evaluative threat, which, in turn, predicts greater SAD symptoms in late adolescence (Nelemans et al., 2017).

Several researchers have also documented that individuals with, or at risk for, SAD are *physiologically* more sensitive to stress (Crişan, Vulturar, Miclea, & Miu, 2016; Farmer & Kashdan, 2015; Nelemans et al., 2017), suggesting that in addition to subjective stress reactivity, physiological function is an important domain to consider in understanding the psychobiology of SAD. Indeed, perceived stress is associated with physiological changes related to the autonomic nervous system (ANS), including changes in arousal, cardiovascularity, and respiration, which collectively mobilize individuals to engage in adaptive behavioral responses. Specifically, when encountering a stressor, an organism shifts from a “rest and digest” mode governed by parasympathetic nervous system (PNS) activity to a “fight or flight” mode governed by heightened sympathetic nervous system (SNS) activity (Berntson, Norman, Hawkley, & Cacioppo, 2008; Porges, 2007). During recovery from a stressor, healthy autonomic regulation is characterized by a shift back to PNS dominance (Kahle, Miller, Lopez, & Hastings, 2016; Mezzacappa, Kelsey, Katkin, & Sloan, 2001). These patterns of physiological coordination between SNS to PNS activity during reactivity and recovery are posited to reflect the effectiveness of individuals’ autonomic regulation to stressors (Berntson et al., 2008; Brosschot, Pieper, & Thayer, 2005). Interestingly, recent work suggests that flexible changes in the PNS and SNS (i.e., increased reactivity and recovery) in response to emotionally evocative events are correlated significantly with adaptive emotion regulation (Kahle, Miller, Helm, & Hastings, 2018; Miller et al., 2013); in contrast, blunted PNS reactivity and recovery have been documented in internalizing disorders (Kircanski, Waugh, Camacho, & Gotlib, 2016). Thus,

dysfunction in ANS regulation, measured by disruptions in the ability to appropriately shift autonomic balance during or following a stressor, may represent two plausible—and not mutually exclusive—biological indicators of the maladaptive emotion regulatory processes that underlie SAD.

Finally, it is imperative that researchers consider sex when examining links among stress sensitivity (i.e., the tendency to perceive a stressor as distressing), autonomic regulation, and increasing rates of anxiety problems across adolescence. Sex-differentiated exposure to stressors (i.e., more experiences of interpersonal stress in females) may lead to differences in stress sensitivity that, in turn, contribute to sex differences in related physiological stress regulation and, ultimately, in symptoms of SAD and related disorders (Hamilton, Stange, Abramson, & Alloy, 2015). Indeed, females report higher levels of perceived stress than do males (Bergdahl & Bergdahl, 2002; Brougham, Zail, Mendoza, & Miller, 2009), and are twice as likely both to report internalizing symptoms and to be diagnosed with anxiety disorders in adolescence (Angold & Costello, 1995). In fact, sex differences in physiological responses to stress have been documented across the lifespan, with females exhibiting significantly greater SNS reactivity than do males starting in adolescence (Kudielka, Buske-Kirschbaum, Hellhammer, & Kirschbaum, 2004; Kudielka & Kirschbaum, 2005; Ordaz & Luna, 2012). Thus, the well documented sex differences in the onset of SAD from adolescence through adulthood (i.e., females are more likely than are males to be diagnosed with SAD; Aune & Stiles, 2009; Caballo et al., 2008; Merikangas et al., 2010) may arise from sex differences in both subjective stress sensitivity and autonomic reactivity and recovery to social stress.

In the present investigation we examined the associations among stress sensitivity, autonomic regulation in response to a well-validated laboratory social stress paradigm, and social

anxiety symptoms in a community sample of female and male adolescents. Importantly, we recruited female and male adolescents to be matched on pubertal status, thereby allowing us to examine sex-specific effects without puberty-related confounds. Moreover, we conducted and coded comprehensive interviews in all participants to assess cumulative objective life stress as well as subjective stress severity. Thus, we were able to derive a measure of stress sensitivity that reflects the subjective severity of stressful experiences after accounting for objective severity. We hypothesized that: 1) stress sensitivity will be associated with symptoms of social anxiety; 2) this association will be mediated by dysfunction in ANS regulation (more inflexible responding; that is, a smaller shift from PNS to SNS dominance during physiological reactivity to an acute stressor and a smaller shift from SNS to PNS dominance during physiological recovery to the stressor); and 3) these associations will be stronger in females than in males.

Methods

Participants

Potential participants were recruited from the San Francisco Bay Area through posted printed flyers, online advertisements placed in local parenting magazines, and social media platforms. Inclusion criteria for participants were English fluency and being 9–13 years of age. Given that the larger study was focused on the effects of early life stress on brain development across the pubertal transition, exclusion criteria for participants were post-pubertal status as measured by self-report Tanner staging (see “Pubertal Staging” below for more details), any contraindications to MRI (e.g., claustrophobia, braces, metal implants), a history of neurological disorder or any major medical illness, serious cognitive or physical challenges that might interfere with their ability to understand or complete procedures, or, for female participants, the onset of menses. Males and females were recruited to be matched on pubertal stage, measured by

self-report Tanner staging. Participants and their parent(s)/legal guardian(s) signed assent and consent forms, respectively, to participate in this study, which was approved by the Stanford University Institutional Review Board. All participants were compensated for their time.

At the baseline assessment (T1) of the study, we recruited 214 adolescents (121 females) ages 9–13 years ($M \pm SD = 11.38 \pm 1.05$). Of these participants, 170 (92 females) ages 11–16 years ($M \pm SD = 13.40 \pm 1.06$) returned for a follow-up assessment (T2) approximately 24 months later ($M \pm SD = 23.98 \pm 4.28$ months). One hundred and sixty-nine participants completed a laboratory social stress test; 96 participants completed the task at T1 and 73 participants completed the task at T2 (because participants were debriefed after the task, they could only complete it at one timepoint). Of these 169 participants, 6 did not complete this task within 90 days of completing the measure of social anxiety and were subsequently excluded from our analyses. Thus, a total of 163 participants (90 females) were included in the current study: 94 participants (51 females) from T1 and 69 participants (39 females) from T2. As we note, below, participants who underwent the TSST at T1 did not differ from participants who received the TSST at T2 in sex, stress sensitivity, social anxiety, or physiological measures (see “Results”).

Stress Sensitivity

Participants were interviewed at both timepoints using the modified version of the Traumatic Screening Inventory for Children (Ribbe, 1996), during which they were asked whether they had experienced any of over 30 different types of stressful events (e.g., “Have you ever experienced a severe illness or injury?”, “Have your parents ever separated or divorced?”, “Have you ever been left alone or with a caregiver who was not able to take care of you?”). At T1, participants were interviewed about events that happened in their lifetime; at T2, participants were interviewed about events that happened since their T1 visit. Thus, for participants whose T2

data we used for the present investigation, we combined data from both of their stress interviews to obtain a single measurement of their cumulative stress exposure.

If a participant endorsed having experienced a stressful event, interviewers followed up with specific questions to contextualize the event (e.g., timing, duration, frequency, involvement of others). Further, for each type of stressor that a participant endorsed, s/he provided subjective severity ratings of how helpless, confused, or scared s/he felt at the time of the experience on a 4-point scale (0=*not scared*; 3=*extremely scared*). Next, for each event that a participant endorsed, a panel of three coders, blind to the participant's subjective severity ratings and reactions and behaviors during the interview, rated the objective severity of the event using a modified version of the UCLA Life Stress Interview coding system (Rudolph et al., 2000) based on a 5-point scale (0=*non-event or no impact*; 4=*extremely severe impact*). After this coding, we z-scored and summed the maximum subjective and objective severity scores for each type of stressor endorsed (we chose this method in order to not overweight reports of frequent but less severe events; see King et al., 2017), thereby creating standardized indices of subjective stress severity and objective stress severity.

As in prior work (Ho et al., 2017), we operationalized stress sensitivity as the residual variance in subjective stress severity after accounting for objective stress severity using a linear regression model on the standardized scores. Thus, stress sensitivity represented participants' perceived severity of the stressful events beyond the events' cumulative objective severity, such that positive values represented higher stress sensitivity than would be expected objectively, and negative values represented lower stress sensitivity than would be expected objectively.

Autonomic Regulation During Stress

Participants were randomized to undergo a modified version of the Trier Social Stress Test (TSST; Kirschbaum, Pirke, & Hellhammer, 1993) at either T1 or T2. Electrodes were placed on the participants' abdomen and non-dominant hand in order to record electrocardiogram (ECG) and skin conductance level (SCL), respectively. Respiratory sinus arrhythmia (RSA; i.e., heart rate variability corresponding to respiration) is a measure of parasympathetic nervous system activity and was computed from the ECG data (Berntson, Cacioppo, & Quigley, 1993). SCL is an index of sympathetic nervous system activity and measures skin conductivity through changes in electrical potentials in the skin (Critchley & Nagai, 2013). The electrodes were attached to wireless devices that transmitted data to receiver modules placed in the session room (Biopac Systems, Goleta, CA). A computer in the session room recorded the data at a sampling rate of 500 Hz using the AcqKnowledge data acquisition and analysis program (Biopac Systems, Goleta, CA).

In our TSST protocol, participants were asked to sit quietly by themselves for 5 minutes in order to collect baseline data. After this 5-minute baseline period, the experimenter read aloud a story prompt to the participants and told them that they had 5 minutes to prepare a 5-minute ending to the story. Participants were told that they would tell this story ending in front of a judge, who would be videotaping them and evaluating the quality of their story. After the 5-minute preparation period, participants gave their speech, while seated, to a confederate judge, who was trained to maintain a neutral demeanor and to not react to participants' actions unless it was to keep a participant on task. The judge also acted as if s/he was taking notes throughout the interaction with the participant. Participants were given 5 minutes to complete their story. If participants finished earlier, the judge prompted them to continue. If participants refused to continue speaking for the rest of the 5-minute story period, the judge and participant sat quietly

until the period was over. After the story period, the judge notified the participant that his/her next task was to complete a 5-minute mental math task during which s/he was given an original number and told to continuously mentally subtract a given number. Participants were interrupted if they made a mistake during this task and were told to start over. After the 5-minute math period, participants were shown two 15-minute videos depicting calming scenes of nature. These two 15-minute recovery periods were combined to form one full recovery period used in analyses. Following the task, participants were debriefed and were informed that the judge was not actually evaluating them.

Autonomic Nervous System Laboratory (ANSLAB; Wilhelm & Peyk, 2005) software was used to preprocess and extract RSA and SCL data from activity recorded by the electrodes. RSA values were computed from ECG data using a high-frequency bandpass ranging from 0.15 to 0.40 Hz during the story and math periods and were averaged and subtracted from baseline to yield RSA reactivity. SCL reactivity was calculated in the same manner. RSA and SCL recovery were calculated by averaging values across both recovery periods and subtracting from the average of the story and math scores (Miller, Gillette, Manczak, Kircanski, & Gotlib, 2019).

Autonomic regulation was operationalized by combining standardized RSA and SCL change values, thereby representing physiological shifts from sympathetic nervous system (SNS) activity to parasympathetic nervous system (PNS) activity (Miller et al., 2019). RSA and SCL change values were z-scored for standardization. Because RSA is a measure of PNS activity, RSA values were multiplied by -1 so that higher RSA scores represented lower PNS activity. Inverse standardized RSA values and standardized SCL values were summed to yield an index of a shift from PNS activity to SNS activity. Thus, for stress reactivity, positive values represent a shift to SNS dominance and negative values represent a shift to PNS dominance, from the

baseline to the TSST. For stress recovery, positive and negative values represent a shift to SNS and PNS dominance, respectively, from the TSST to the recovery period.

Anxiety Symptoms

The Multidimensional Anxiety Scale for Children 2nd edition (MASC; March, Parker, Sullivan, Stallings, & Conners, 1997) was used to assess anxiety problems at both timepoints. The MASC is a reliable measure of anxiety in children and adolescents (Baldwin & Dadds, 2007; March, Sullivan, & Parker, 1999). Because of our interest in symptoms related to SAD, we focused our analyses on the Social Anxiety subscale of the MASC.

Pubertal Staging

Pubertal staging at both T1 and T2 was measured with self-report Tanner staging (Marshall & Tanner, 1968; Marshall & Tanner, 1970; Morris & Udry, 1980), which has been shown to be strongly correlated with physicians' physical examinations of pubertal development (Shirtcliff, Dahl, & Pollak, 2009). Female and male participants were shown sets of schematic drawings, dependent on their sex, depicting different developmental stages of pubic hair and breast/testes development ranging from 1 (prepubertal) to 5 (postpubertal). As in prior investigations from our group (Colich et al., 2017; King et al., 2017), pubic hair and breast/testes ratings were averaged to compute an index of overall pubertal development (Dorn, Dahl, Woodward, & Biro, 2006).

Medication Usage

Use of medications that could affect physiological stress reactivity (e.g., vasodilators, beta-blockers, calcium channel blockers) was coded as a dichotomous variable (0=*no medications*, 1=*taking medications*).

Statistical Analysis

All statistical analyses were conducted using R version 3.3 (<https://www.r-project.org/>). Two-tailed t-tests and χ^2 tests, where appropriate, were conducted to examine sex differences in demographics and variables of interest. Two-way repeated measures ANOVAs with Greenhouse-Geisser sphericity correction were conducted to evaluate the main and interactive effects of sex and TSST condition on RSA and SCL. Linear regression models were conducted to test for associations among stress sensitivity, social anxiety symptoms, and autonomic regulation (during recovery and reactivity, separately), whether sex moderated these associations, and whether there were significant effects in males and females, separately, given sex-specific effects for several of these variables in prior literature (e.g., Brougham et al., 2009; Merikangas et al., 2010; Ordaz & Luna, 2012). Finally, based on our findings (see “Results”), we tested whether autonomic regulation (in separate models for recovery and reactivity) statistically mediated the associations between stress sensitivity and social anxiety in females by conducting a series of bootstrapped regressions. We estimated the 95% confidence interval for the indirect effect (i.e., the mediation effect) using Monte Carlo simulations with 10,000 chains using the “RMediation” package in R (Tofighi & MacKinnon, 2011). In all of our statistical models, age, Tanner stage, and medication usage were included as covariates.

Results

Descriptive Statistics

Descriptive statistics by sex are presented in Table 1. By design, females and males did not differ in pubertal status but did differ significantly in age, such that females were, on average, younger than males ($t(161)=3.36, p<.001$). While males and females did not differ in objective stress severity ($t(161)=0.20, p=.842$) or in subjective stress severity ($t(159)=-1.46, p=.146$), females exhibited significantly greater stress sensitivity than did males ($t(159)=-2.89,$

$p=0.004$). There were no sex differences on any other measures of interest. In Table 2 we present descriptive statistics by timepoint. As expected, participants differed across timepoint by age and pubertal status; there were no other significant differences. Zero-order correlations among all key variables are presented in Tables 3 (all participants) and 4 (separated by sex).

Effects of TSST

We first examined whether the TSST was effective in eliciting autonomic reactivity by increasing SNS activity. Figure 1 depicts RSA (panel A) and SCL (panel B) values across the TSST session (separated by sex). There was a main effect of time during TSST on RSA ($F(2.76,623.35)=4.465, p<.001$) and SCL ($F(3.42,414.24)=39.107, p=1.17 \times 10^{-24}$) but no main or interactive effects of sex (all $ps>.244$). We also tested whether RSA and SCL during reactivity and recovery differed from baseline. Participants had lower RSA during reactivity compared to baseline, an effect that was driven by females (all: mean difference=-0.33, $t(322)=-3.08, p=.002$; females: mean difference=-0.43, $t(177)=-3.28, p=.001$; males: mean difference=-0.22, $t(143)=-1.20, p=.231$). Similarly, participants had greater SCL during reactivity compared to baseline (all: mean difference=3.59, $t(311)=5.14, p<.001$; females: mean difference=4.06, $t(171)=4.82, p<.001$; males: mean difference=3.03, $t(138)=2.62, p=.010$). RSA during recovery did not differ significantly from baseline (all: mean difference=0.07, $t(297)=-0.56, p=.573$; females: mean difference=-0.16, $t(163)=-1.21, p=.227$; males: mean difference=0.06, $t(132)=0.28, p=.778$). Participants also had greater SCL during recovery compared to baseline, an effect that was also driven by females (mean difference=1.50, $t(282)=2.34, p=.020$; females: mean difference=2.27, $t(259)=2.90, p=.004$; males: mean difference=0.56, $t(121)=0.52, p=.602$).

Stress Sensitivity and Social Anxiety

Across all participants, greater stress sensitivity was significantly associated with more severe social anxiety symptoms ($\beta=.25, p=.004$). This association was significantly moderated by sex ($\beta=.30, p=.022$): it was significantly positive in females ($\beta=.43, p<.001$) and was non-significant in males ($p=0.90$). See Figure 2 for more details.

Stress Sensitivity and Regulation of the Autonomic Nervous System

Across all participants, greater stress sensitivity was significantly associated with PNS dominance during reactivity ($\beta=-.19, p=.037$) and SNS dominance during recovery ($\beta=.25, p=.013$). These associations were not driven by changes in RSA activity from baseline to reactivity or from reactivity to recovery (all $ps>.05$). Rather, stress sensitivity was significantly associated with change in SCL activity from baseline to reactivity ($\beta=-.205, p=.022$) and from reactivity to recovery ($\beta=-.036, p=.003$). While sex did not significantly moderate the association between stress sensitivity and autonomic regulation during reactivity ($p=.314$), this association was significantly negative in females ($\beta=-.26, p=.031$) and was non-significant in males ($p=.487$). See Figure 3A for details. Similarly, while sex did not significantly moderate the association between stress sensitivity and autonomic regulation during recovery ($p=.932$), this association was significantly positive in females ($\beta=.30, p=.023$) and was non-significant in males ($p=.095$). See Figure 3B for details.

Regulation of the Autonomic Nervous System and Social Anxiety

Across all participants, social anxiety symptoms were not significantly associated with autonomic regulation during reactivity or recovery (all $ps>.113$). While sex did not significantly moderate the association between social anxiety and autonomic regulation during reactivity ($p=.051$), this association was significantly negative in females ($\beta=-.235, p=.040$) and non-significant in males ($p=.485$). See Figure 4A for details. Similarly, while sex did not significantly

moderate the association between social anxiety and autonomic regulation during recovery ($p=.076$), this association was significantly positive in females ($\beta=.277, p=.026$) and non-significant in males ($p=.861$). See Figure 4B for details. Specifically, for females, social anxiety was significantly associated with change in RSA activity from baseline to reactivity ($\beta=.272, p=.016$) and with change in RSA activity from reactivity to recovery ($\beta=-.251, p=.041$) but not with comparable changes in SCL ($ps>.05$). Neither changes in RSA nor in SCL during reactivity and recovery were significantly associated with social anxiety symptoms for males ($ps>.05$).

Mediation Models

Because the associations among stress sensitivity, autonomic regulation during reactivity and recovery, and social anxiety symptoms were all statistically significant in females, we explored whether autonomic regulation during reactivity and recovery (separately) mediated the association between stress sensitivity and social anxiety. While the mediation effect for autonomic regulation during reactivity was not significantly different from 0 (95% CI: -0.03–2.12), the mediation effect for autonomic regulation during recovery was significant ($B=1.06$, 95% CI: 0.02–2.64). Specifically, SNS dominance during recovery significantly explained variance in the association between stress sensitivity and social anxiety symptoms in females.

Discussion

Adolescence is a formative period of development marked by social reorienting, including a desire for social approval from peers (Blakemore, 2008; Kilford, Garrett, & Blakemore, 2016). Not surprisingly, therefore, adolescence is also a vulnerable time for the development of conditions characterized by fear of judgment or social humiliation, particularly social anxiety disorder (SAD). While heightened stress sensitivity is a central feature in conceptual models of psychopathology—and while there is evidence for greater stress sensitivity

in individuals with SAD—the specific psychobiological mechanisms by which stress sensitivity contributes to the emergence of SAD in adolescents is not well understood.

Previous work from our group examining a subset of the adolescents reported here focused on neural correlates underlying emotion regulation (i.e., white matter connectivity between the amygdala and ventromedial prefrontal cortex) as putative mediators of the link between stress sensitivity and SAD. Importantly, we did not find evidence that this neural circuit mediated links between stress sensitivity and SAD, and, critically, we did not examine sex-specific associations (Ho et al., 2017). Given that a growing body of literature has implicated autonomic nervous system (ANS) dysfunction both in stress reactivity generally (Spear, 2009) and in SAD specifically (Farmer & Kashdan, 2015), in this study we sought to examine whether the links between stress sensitivity and social anxiety symptoms were mediated by dysfunction in autonomic regulation. Specifically, we operationalized ANS dysfunction as the failure to shift from parasympathetic nervous system (PNS) to sympathetic nervous system (SNS) during stress reactivity and as the failure to shift from SNS to PNS during stress recovery. While both processes are indicative of autonomic dysregulation, the former implicates a dominant PNS whereas the latter implicates a dominant SNS; thus, these two accounts not only yield distinct contributions to existing psychobiological models of SAD but also have distinct treatment implications. Finally, because we were limited in our statistical power to detect significant moderating effects of sex (Perugini, Gallucci, & Costantini, 2018), we stratified our analyses by sex given evidence of sex-specific effects in these processes (e.g., Brougham et al., 2009; Merikangas et al., 2010; Ordaz & Luna, 2012).

Consistent with our hypotheses, we found that females showed heightened stress sensitivity compared to males; further, among females only, sustained SNS activity during

recovery from a stressor explained the extent to which stress sensitivity—that is, the tendency to perceive and experience stressors as more distressing—was associated with social anxiety symptoms in adolescents. While we found significant associations across all participants between stress sensitivity and autonomic regulation, and between stress sensitivity and social anxiety, our sex-specific analyses revealed that these effects were driven by females; stress sensitivity, autonomic regulation, and social anxiety were not significantly associated with each other in males.

Our finding that dominant SNS activity during stress recovery is a key physiological mechanism explaining the contribution of psychological stress reactivity to social anxiety symptoms in females has important clinical implications. Interventions that target SNS function (either by dampening activity in this system or by promoting PNS activity) through breathing exercises, in conjunction with cognitive and emotion regulatory strategies, may be particularly effective for female adolescents who are suffering from or at risk for SAD. Such interventions may draw from existing mindfulness-based therapies, such as mindfulness-based stress reduction (MBSR), mindfulness-based cognitive therapy (MBCT), and acceptance and commitment therapy (ACT), that are appropriately modified for adolescents. Indeed, MBSR, MBCT, and ACT have been demonstrated to help reduce avoidance behaviors and emotion reactivity to negative self-beliefs in both adults (Goldin & Gross, 2010) and adolescents with SAD (Azadeh, Kazemi-Zahrani, & Besharat, 2016; Swain, Hancock, Dixon, Koo, & Bowman, 2013).

We did not find statistically significant associations among stress sensitivity, autonomic regulation, and SAD in male adolescents. Thus, a clear direction for future research is to investigate the psychobiological mechanisms that underlie the onset and maintenance of SAD in male adolescents. Additional work is also needed to address some of the limitations of the

present study. For example, the design of this investigation required us to rely on retrospective reports of stressful life events; however, recent work suggests that there are differences between retrospective and prospective reports of adverse experiences in youth (Baldwin, Reuben, Newbury, & Danese, 2019, although see Scott, McLaughlin, Smith, & Ellis, 2012, who found that both types of report predict mental health outcomes). In addition, prospective studies starting at earlier ages will help to elucidate processes that shape the development of heightened stress sensitivity. One possibility is that this tendency is a consequence of biological sensitivity to stressors combined with excessive exposure to events that elicit stress responses (Ellis, Boyce, Belsky, Bakermans-Kranenburg, & Van IJzendoorn, 2011). As a related point, there are additional aspects of the experience of life stress during development that warrant further investigation. For example, consistent with a stress sensitization model, repeated or chronic exposure to threatening stimuli may affect stress sensitivity and associated autonomic systems (McLaughlin, Conron, Koenen, & Gilman, 2010); however, life stress characterized by an absence of exposure to positive stimuli (i.e., experiences of deprivation or neglect) may affect the development of stress sensitivity and autonomic regulation differently than would life stress characterized by threat (McLaughlin, Sheridan, & Nelson, 2017). Finally, while we focused in this study on symptoms of SAD, exposure to life stressors during childhood and adolescence is associated with a greater risk of developing a range of psychopathology, including depression, substance dependence, and schizophrenia in both adolescence (Green et al., 2010; LeMoult et al., 2019) and adulthood (Kessler et al., 2010; McLaughlin et al., 2010). Because our study sample was a community sample of adolescents who were exposed to normative stressors, and because SAD is one of the most commonly diagnosed disorders during this developmental period, we limited the present investigation to symptoms in this domain. It is clear, however, that future

work should examine whether dysfunction in autonomic regulation is a transdiagnostic mechanism through which stress sensitivity leads to the emergence of psychopathology in general, or alternatively, if this mechanism is specific to disorders that are characterized by maladaptive responses to stress (e.g., depression, anxiety, PTSD).

Conclusions

In the present study, we explored associations among stress sensitivity, autonomic stress responses during reactivity and recovery phases of a social stressor, and symptoms of social anxiety in a large community sample of female and male adolescents. Guided by a stress sensitivity framework, we found that inappropriately prolonged sympathetic nervous system activity following stressful events may explain how stress sensitivity—that is, the tendency to perceive and experience stressors as more distressing—leads to social anxiety symptoms in adolescent girls. Our findings also suggest that interventions targeting sympathetic nervous system function, either by dampening activity in this system or by promoting parasympathetic nervous system activity, will be particularly effective for female adolescents who are experiencing SAD.

References

- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Washington, DC: Publisher.
- Angold, A., & Costello, E. J. (1995). Developmental epidemiology. *Epidemiologic Reviews*, 17(1), 74-82.
- Aune, T., & Stiles, T. C. (2009). The effects of depression and stressful life events on the development and maintenance of syndromal social anxiety: Sex and age differences. *Journal of Clinical Child & Adolescent Psychology*, 38(4), 501-512.
- Azadeh, S. M., Kazemi-Zahrani, H., & Besharat, M. A. (2016). Effectiveness of acceptance and commitment therapy on interpersonal problems and psychological flexibility in female high school students with social anxiety disorder. *Global Journal of Health Science*, 8(3), 131.
- Baldwin, J. S., & Dadds, M. R. (2007). Reliability and validity of parent and child versions of the multidimensional anxiety scale for children in community samples. *Journal of the American Academy of Child & Adolescent Psychiatry*, 46(2), 252-260.
- Baldwin, J. R., Reuben, A., Newbury, J. B., & Danese, A. (2019). Agreement between prospective and retrospective measures of childhood maltreatment: A systematic review and meta-analysis. *JAMA Psychiatry*, 76(6), 584-593.
- Beesdo, K., Bittner, A., Pine, D. S., Stein, M. B., Höfler, M., Lieb, R., & Wittchen, H. U. (2007). Incidence of social anxiety disorder and the consistent risk for secondary depression in the first three decades of life. *Archives of General Psychiatry*, 64(8), 903-912.
- Beesdo, K., Pine, D. S., Lieb, R., & Wittchen, H. U. (2010). Incidence and risk patterns of

- anxiety and depressive disorders and categorization of generalized anxiety disorder. *Archives of General Psychiatry*, 67(1), 47-57.
- Bergdahl, J., & Bergdahl, M. (2002). Perceived stress in adults: Prevalence and association of depression, anxiety and medication in a Swedish population. *Stress and Health: Journal of the International Society for the Investigation of Stress*, 18(5), 235-241.
- Berntson, G. G., Cacioppo, J. T., & Quigley, K. S. (1993). Respiratory sinus arrhythmia: Autonomic origins, physiological mechanisms, and psychophysiological implications. *Psychophysiology*, 30(2), 183-196.
- Berntson, G. G., Norman, G. J., Hawkley, L. C., & Cacioppo, J. T. (2008). Cardiac autonomic balance versus cardiac regulatory capacity. *Psychophysiology*, 45(4), 643-652.
- Blakemore, S. J. (2008). The social brain in adolescence. *Nature Reviews Neuroscience*, 9(4), 267.
- Brosschot, J. F., Pieper, S., & Thayer, J. F. (2005). Expanding stress theory: Prolonged activation and perseverative cognition. *Psychoneuroendocrinology*, 30(10), 1043-1049.
- Brougham, R. R., Zail, C. M., Mendoza, C. M., & Miller, J. R. (2009). Stress, sex differences, and coping strategies among college students. *Current Psychology*, 28(2), 85-97.
- Caballo, V. E., Salazar, I. C., Irtia, M. J., Arias, B., Hofmann, S. G., & CISO-A Research Team. (2008). Social anxiety in 18 nations: Sex and age differences. *Behavioral Psychology/Psicología Conductual*, 16(2), 163-187.
- Colich, N. L., Williams, E. S., Ho, T. C., King, L. S., Humphreys, K. L., Price, A. N., ... & Gotlib, I. H. (2017). The association between early life stress and prefrontal cortex activation during implicit emotion regulation is moderated by sex in early adolescence. *Development and Psychopathology*, 29(5), 1851-1864.

- Crişan, L. G., Vulturar, R., Miclea, M., & Miu, A. C. (2016). Reactivity to social stress in subclinical social anxiety: Emotional experience, cognitive appraisals, behavior, and physiology. *Frontiers in Psychiatry*, 7, 5.
- Critchley, H., & Nagai, Y. (2013). Electrodermal activity (EDA). *Encyclopedia of Behavioral Medicine*, 666-669.
- Dorn, L. D., Dahl, R. E., Woodward, H. R., & Biro, F. (2006). Defining the boundaries of early adolescence: A user's guide to assessing pubertal status and pubertal timing in research with adolescents. *Applied Developmental Science*, 10(1), 30-56.
- Ellis, B. J., Boyce, W. T., Belsky, J., Bakermans-Kranenburg, M. J., & Van IJzendoorn, M. H. (2011). Differential susceptibility to the environment: An evolutionary–neurodevelopmental theory. *Development and Psychopathology*, 23(1), 7-28.
- Farmer, A. S., & Kashdan, T. B. (2015). Stress sensitivity and stress generation in social anxiety disorder: A temporal process approach. *Journal of Abnormal Psychology*, 124(1), 102.
- Goldin, P. R., & Gross, J. J. (2010). Effects of mindfulness-based stress reduction (MBSR) on emotion regulation in social anxiety disorder. *Emotion*, 10(1), 83.
- Green, J. G., McLaughlin, K. A., Berglund, P. A., Gruber, M. J., Sampson, N. A., Zaslavsky, A. M., & Kessler, R. C. (2010). Childhood adversities and adult psychiatric disorders in the national comorbidity survey replication I: Associations with first onset of DSM-IV disorders. *Archives of General Psychiatry*, 67(2), 113-123.
- Hammen, C. (2015). Stress sensitivity in psychopathology: Mechanisms and consequences. *Journal of Abnormal Psychology*, 124(1), 152.
- Hamilton, J. L., Stange, J. P., Abramson, L. Y., & Alloy, L. B. (2015). Stress and the

- development of cognitive vulnerabilities to depression explain sex differences in depressive symptoms during adolescence. *Clinical Psychological Science*, 3(5), 702-714.
- Ho, T. C., King, L. S., Leong, J. K., Colich, N. L., Humphreys, K. L., Ordaz, S. J., & Gotlib, I. H. (2017). Effects of sensitivity to life stress on uncinate fasciculus segments in early adolescence. *Social Cognitive and Affective Neuroscience*, 12(9), 1460-1469.
- Kahle, S., Miller, J. G., Helm, J. L., & Hastings, P. D. (2018). Linking autonomic physiology and emotion regulation in preschoolers: The role of reactivity and recovery. *Developmental Psychobiology*, 60(7), 775-788.
- Kahle, S., Miller, J. G., Lopez, M., & Hastings, P. D. (2016). Sympathetic recovery from anger is associated with emotion regulation. *Journal of Experimental Child Psychology*, 142, 359-371.
- Kessler, R. C., McLaughlin, K. A., Green, J. G., Gruber, M. J., Sampson, N. A., Zaslavsky, A. M., ... & Benjet, C. (2010). Childhood adversities and adult psychopathology in the WHO World Mental Health Surveys. *The British Journal of Psychiatry*, 197(5), 378-385.
- Kilford, E. J., Garrett, E., & Blakemore, S. J. (2016). The development of social cognition in adolescence: An integrated perspective. *Neuroscience & Biobehavioral Reviews*, 70, 106-120.
- King, L. S., Colich, N. L., LeMoult, J., Humphreys, K. L., Ordaz, S. J., Price, A. N., & Gotlib, I. H. (2017). The impact of the severity of early life stress on diurnal cortisol: The role of puberty. *Psychoneuroendocrinology*, 77, 68-74.
- Kircanski, K., Waugh, C. E., Camacho, M. C., & Gotlib, I. H. (2016). Aberrant parasympathetic stress responsivity in pure and co-occurring major depressive disorder and generalized anxiety disorder. *Journal of Psychopathology and Behavioral Assessment*, 38(1), 5-19.

- Kirschbaum, C., Pirke, K. M., & Hellhammer, D. H. (1993). The 'Trier Social Stress Test'—a tool for investigating psychobiological stress responses in a laboratory setting. *Neuropsychobiology*, 28(1-2), 76-81.
- Kudielka, B. M., Buske-Kirschbaum, A., Hellhammer, D. H., & Kirschbaum, C. (2004). HPA axis responses to laboratory psychosocial stress in healthy elderly adults, younger adults, and children: Impact of age and gender. *Psychoneuroendocrinology*, 29(1), 83-98.
- Kudielka, B. M., & Kirschbaum, C. (2005). Sex differences in HPA axis responses to stress: A review. *Biological Psychology*, 69(1), 113-132.
- LeMoult, J., Humphreys, K. L., Tracy, A., Hoffmeister, J. A., Ip, E., & Gotlib, I. H. (2019). Meta-analysis: Exposure to early life stress and risk for depression in childhood and adolescence. *Journal of the American Academy of Child & Adolescent Psychiatry*.
- March, J. S., Parker, J. D., Sullivan, K., Stallings, P., & Conners, C. K. (1997). The Multidimensional Anxiety Scale for Children (MASC): Factor structure, reliability, and validity. *Journal of the American Academy of Child & Adolescent Psychiatry*, 36(4), 554-565.
- March, J. S., Sullivan, K., & Parker, J. (1999). Test-retest reliability of the Multidimensional Anxiety Scale for Children. *Journal of Anxiety Disorders*, 13(4), 349-358.
- Marshall, W. A., & Tanner, J. M. (1968). Growth and physiological development during adolescence. *Annual Review of Medicine*, 19(1), 283-300.
- Marshall, W. A., & Tanner, J. M. (1970). Variations in the pattern of pubertal changes in boys. *Archives of Disease in Childhood*, 45(239), 13-23.
- McLaughlin, K. A., Conron, K. J., Koenen, K. C., & Gilman, S. E. (2010). Childhood adversity,

- adult stressful life events, and risk of past-year psychiatric disorder: A test of the stress sensitization hypothesis in a population-based sample of adults. *Psychological Medicine*, 40(10), 1647-1658.
- McLaughlin, K. A., Green, J. G., Gruber, M. J., Sampson, N. A., Zaslavsky, A. M., & Kessler, R. C. (2010). Childhood adversities and adult psychiatric disorders in the national comorbidity survey replication II: Associations with persistence of DSM-IV disorders. *Archives of General Psychiatry*, 67(2), 124-132.
- McLaughlin, K. A., Sheridan, M. A., & Nelson, C. A. (2017). Neglect as a violation of species-expectant experience: Neurodevelopmental consequences. *Biological Psychiatry*, 82(7), 462-471.
- Merikangas, K. R., He, J. P., Burstein, M., Swanson, S. A., Avenevoli, S., Cui, L., ... & Swendsen, J. (2010). Lifetime prevalence of mental disorders in US adolescents: Results from the National Comorbidity Survey Replication–Adolescent Supplement (NCS-A). *Journal of the American Academy of Child & Adolescent Psychiatry*, 49(10), 980-989.
- Mezzacappa, E. S., Kelsey, R. M., Katkin, E. S., & Sloan, R. P. (2001). Vagal rebound and recovery from psychological stress. *Psychosomatic Medicine*, 63(4), 650-657.
- Miller, J. G., Choccol, C., Nuselovici, J. N., Utendale, W. T., Simard, M., & Hastings, P. D. (2013). Children's dynamic RSA change during anger and its relations with parenting, temperament, and control of aggression. *Biological Psychology*, 92(2), 417-425.
- Miller, J. G., Gillette, J. S., Manczak, E. M., Kircanski, K., & Gotlib, I. H. (2019). Fine particle air pollution and physiological reactivity to social stress in adolescence: The moderating role of anxiety and depression. *Psychosomatic Medicine*, 81(7), 641-648.

- Morris, N. M., & Udry, J. R. (1980). Validation of a self-administered instrument to assess stage of adolescent development. *Journal of Youth and Adolescence*, 9(3), 271-280.
- Nelemans, S. A., Hale III, W. W., Branje, S. J., van Lier, P. A. C., Koot, H. M., & Meeus, W. H. J. (2017). The role of stress reactivity in the long-term persistence of adolescent social anxiety symptoms. *Biological Psychology*, 125, 91-104.
- Ordaz, S., & Luna, B. (2012). Sex differences in physiological reactivity to acute psychosocial stress in adolescence. *Psychoneuroendocrinology*, 37(8), 1135-1157.
- Perugini, M., Gallucci, M., & Costantini, G. (2018). A practical primer to power analysis for simple experimental designs. *International Review of Social Psychology*, 31(1).
- Porges, S. W. (2007). The polyvagal perspective. *Biological Psychology*, 74(2), 116-143.
- Ribbe, D. (1996). Psychometric review of traumatic event screening instrument for children (TESI-C). *Measurement of Stress, Trauma, and Adaptation*, 386-387.
- Rudolph, K.D., Hammen, C., Burge, D., Lindberg, N., Herzberg, D., & Daley, S.E. (2000). Toward an interpersonal life-stress model of depression: The developmental context of stress generation. *Developmental Psychopathology*, 12, 215-234.
- Scott, K. M., McLaughlin, K. A., Smith, D. A., & Ellis, P. M. (2012). Childhood maltreatment and DSM-IV adult mental disorders: Comparison of prospective and retrospective findings. *The British Journal of Psychiatry*, 200(6), 469-475.
- Shirtcliff, E. A., Dahl, R. E., & Pollak, S. D. (2009). Pubertal development: Correspondence between hormonal and physical development. *Child Development*, 80(2), 327-337.
- Spear, L. P. (2009). Heightened stress responsivity and emotional reactivity during pubertal maturation: Implications for psychopathology. *Development and Psychopathology*, 21(1), 87-97.

- Stein, M. B. (2006). An epidemiologic perspective on social anxiety disorder. *The Journal of Clinical Psychiatry*, 67, 3-8.
- Stein, M. B., Fuetsch, M., Müller, N., Höfler, M., Lieb, R., & Wittchen, H. U. (2001). Social anxiety disorder and the risk of depression: A prospective community study of adolescents and young adults. *Archives of General Psychiatry*, 58(3), 251-256.
- Swain, J., Hancock, K., Dixon, A., Koo, S., & Bowman, J. (2013). Acceptance and Commitment Therapy for anxious children and adolescents: Study protocol for a randomized controlled trial. *Trials*, 14(1), 140.
- Tofighi, D., & MacKinnon, D. P. (2011). RMediation: An R package for mediation analysis confidence intervals. *Behavior Research Methods*, 43(3), 692-700.
- Van Roy, B., Kristensen, H., Groholt, B., & Clench-Aas, J. (2009). Prevalence and characteristics of significant social anxiety in children aged 8–13 years. *Social Psychiatry and Psychiatric Epidemiology*, 44(5), 407-415.
- Wilhelm, F. H., & Peyk, P. (2005). ANSLab: Autonomic nervous system laboratory (Version 2.4) (Computer software). Retrieved from the SPR Software Repository: <http://www.sprweb.org>.

Figure 1. Mean values of RSA and SCL during the TSST for each sex. Each value on the x-axis represents a five-minute period during the TSST. 1= baseline, 2= preparation, 3=story, 4=mental math, 5–10=recovery.

Figure 2. Associations between stress sensitivity and social anxiety by sex. Across all participants, greater stress sensitivity was significantly associated with greater social anxiety ($\beta=.25, p=.004$). Sex significantly moderated the association between stress sensitivity and social anxiety ($\beta=.30, p=.022$), such that this association was significantly positive in females ($\beta=.43, p<.001$) and non-significant in males ($p=0.90$). All statistical models adjusted for age, pubertal status, and medication usage. Data and trends are plotted without adjustment for covariates for the purposes of visualization.

Figure 3. Associations between stress sensitivity and regulation of the autonomic nervous system (ANS) during reactivity (A) and recovery (B) by sex. Given that the measure of autonomic regulation was based on aggregating standardized scores, the zero value represents the mean shift to sympathetic nervous system dominance (i.e., decreasing heart rate variability and increasing skin conductance). Across all participants, stress sensitivity was significantly associated with parasympathetic nervous system dominance during reactivity ($\beta=-.19, p=.037$) and sympathetic nervous system dominance during recovery ($\beta=.25, p=.013$). While sex did not significantly moderate the association between stress sensitivity and flexibility of ANS during reactivity ($p=.314$), this association was significantly negative in females ($\beta=-.26, p=.031$) and non-significant in males ($p=.487$). Similarly, while sex did not significantly moderate the association between stress sensitivity and autonomic regulation during recovery ($p=.932$), this association was significantly positive in females ($\beta=.30, p=.023$) and non-significant in males ($p=.095$). All statistical models adjusted for age, pubertal status, and medication usage. Data and trends are plotted without adjustment for covariates for the purposes of visualization.

Figure 4. Associations between social anxiety symptoms and regulation of autonomic nervous system (ANS) during reactivity (A) and recovery (B) by sex. Given that the measure of autonomic regulation was based on aggregating standardized scores, the zero value represents the mean shift to sympathetic nervous system dominance (i.e., decreasing heart rate variability and increasing skin conductance). Across all participants, social anxiety symptoms were not significantly associated with autonomic regulation during reactivity ($p=.294$) or recovery ($p=.113$). While sex did not significantly moderate the association between social anxiety and flexibility of ANS during reactivity ($p=.051$), this association was significantly negative in females ($\beta=-.235$, $p=.040$) and non-significant in males ($p=.485$). While sex did not significantly moderate the association between social anxiety and flexibility of ANS during recovery ($p=.076$), this association was significantly positive in females ($\beta=.277$, $p=.026$) and non-significant in males ($p=.861$). All statistical models adjusted for age, pubertal status, and medication usage. Data and trends are plotted without adjustment for covariates for the purposes of visualization.

Table 1. Descriptive statistics by sex.

	All (M±SD; Range)	Male (M±SD; Range)	Female (M±SD; Range)	Statistic
<i>n</i>	163	73	90	
<i>Timepoint</i>				$\chi^2(1) = 0.08$
<i>T1</i>	94	43	51	
<i>T2</i>	69	30	39	
Age in years	12.29 ± 1.39 (9.56–15.82)	12.69 ± 1.25 (10.16–15.65)	11.97 ± 1.42 (9.56–15.82)	$t(161) = 3.36^{***}$
Pubertal stage	2.70 ± 1.11 (1–5)	2.64 ± 1.14 (1–5)	2.76 ± 1.11 (1–5)	$t(161) = -0.68$
Stress sensitivity	0 ± 0.55 (-2.07–1.85)	-0.13 ± 0.54 (-2.07–1.85)	0.11 ± 0.54 (-1.11–1.47)	$t(159) = -2.89^*$
Objective stress (z-scores)	-0.02 ± 0.93 (-1.34–3.46)	-0.01 ± 0.88 (-1.18–2.82)	-0.04 ± 0.97 (-1.34–3.46)	$t(161) = 0.20$
Subjective stress (z-scores)	-0.02 ± 0.96 (-1.23–2.94)	0.14 ± 0.92 (-.123–2.94)	0.08 ± 0.99 (-1.23–2.18)	$t(159) = 1.46$
MASC social anxiety	9.52 ± 6.86 (0–27)	8.78 ± 6.19 (0–26)	10.12 ± 7.33 (0–27)	$t(161) = -1.24$
RSA reactivity	-0.34 ± 0.86 (-2.46–3.75)	-0.23 ± 0.70 (-2.16–1.98)	-0.43 ± 0.97 (-2.46–3.75)	$t(159) = 1.44$
RSA recovery	0.22 ± 0.88 (-3.21–4.69)	0.17 ± 0.83 (-1.74–4.69)	0.26 ± 0.91 (-3.21–3.27)	$t(135) = -0.57$

SCL reactivity	3.69 ± 3.59 (-17.32–19.32)	3.26 ± 4.28 (-17.32–19.32)	4.03 ± 2.91 (-7.72–12.67)	$t(153) = -1.34$
SCL recovery	-1.02 ± 2.48 (-17.77–3.72)	-1.07 ± 3.28 (-17.77–3.72)	-0.99 ± 1.75 (-9.16–2.28)	$t(125) = -0.19$
Autonomic flexibility during reactivity	0 ± 1.47 (-7.14–3.80)	-0.25 ± 1.47 (-7.14–3.80)	0.19 ± 1.45 (-4.28–3.48)	$t(153) = -1.87$
Autonomic flexibility during recovery	0.05 ± 1.33 (-6.04–3.41)	0.18 ± 1.46 (-6.04–2.20)	-0.04 ± 1.24 (-3.94–3.41)	$t(122) = 0.90$
Child race				$\chi^2(5) = 3.92$
<i>White</i>	72	31	41	
<i>Black</i>	12	6	6	
<i>Hispanic/Latino</i>	14	4	10	
<i>Asian</i>	19	7	12	
<i>Biracial</i>	39	21	18	
<i>Other</i>	7	3	4	
Family income				$\chi^2(9) = 6.02$
<i><\$5000</i>	1	0	1	
<i>\$5001-10,000</i>	1	1	0	
<i>\$10,001-15,000</i>	3	0	3	
<i>\$15,001-25,000</i>	3	1	2	

<i>\$25,001-35,000</i>	4	2	2	
<i>\$35,001-50,000</i>	7	4	3	
<i>\$50,001-75,000</i>	16	7	9	
<i>\$75,001-100,000</i>	19	10	9	
<i>\$100,001-150,000</i>	49	20	29	
<i>\$150,001+</i>	53	25	28	
<i>No response</i>	7	3	4	
<i>Parental education</i>				$\chi^2(7) = 10.18$
<i>No GED/High school diploma</i>	1	0	1	
<i>GED/High school diploma</i>	2	0	2	
<i>Some college</i>	33	14	19	
<i>2-Year college degree</i>	12	6	6	
<i>4-Year college degree</i>	58	22	36	
<i>Master's degree</i>	45	28	17	

<i>Professional degree</i>	6	2	4	
<i>Doctorate</i>	4	1	3	
<i>No response</i>	2	0	2	

* indicates $p < .05$, ** indicates $p < .01$, and *** indicates $p < .001$

Table 2. Descriptive statistics by timepoint.

	All	T1	T2	Statistic
<i>n</i>	163	94	69	
Sex				$\chi^2(1) = 0.08$
<i>Male</i>	73	43	30	
<i>Female</i>	90	51	39	
Age in years	12.29 ± 1.39 (9.56–15.82)	11.53 ± 1.10 (9.56–14.04)	13.33 ± 1.02 (11.08–15.82)	$t(161) = -10.64^{***}$
Pubertal stage	2.70 ± 1.11 (1–5)	2.01 ± 0.71 (1–4)	3.64 ± 0.84 (1–5)	$t(161) = -13.49^{***}$
Stress sensitivity	0 ± 0.55 (-2.07–1.85)	-0.05 ± 0.50 (-0.99–1.34)	0.07 ± 0.62 (-2.07–1.85)	$t(159) = -1.40$
Objective stress (z-scores)	-0.02 ± 0.93 (-1.34 –3.46)	0 ± 0.87 (-1.27 –2.82)	-0.05 ± 1.01 (-1.34 –3.46)	$t(161) = 0.33$
Subjective stress (z-scores)	-0.02 ± 0.96 (-1.23–2.94)	-0.05 ± 0.92 (-1.15–2.73)	0.03 ± 1.03 (-1.23–2.94)	$t(159) = -0.53$
MASC social anxiety scores	9.52 ± 6.86 (0–27)	9.98 ± 6.60 (0–27)	8.90 ± 7.20 (0–27)	$t(161) = 0.99$
RSA reactivity	-0.34 ± 0.86 (-2.46–3.75)	-0.38 ± 0.72 (-2.37–1.98)	-0.29 ± 1.02 (-2.46–3.75)	$t(159) = -0.63$
RSA recovery	0.22 ± 0.88 (-3.21–4.69)	0.25 ± 0.74 (-1.74–3.27)	0.19 ± 1.01 (-3.21–4.69)	$t(135) = 0.39$

SCL reactivity	3.69 ± 3.59 (-17.32–19.32)	3.84 ± 4.33 (-17.32–19.32)	3.48 ± 2.16 (-3.12–9.92)	$t(153) = 0.61$
SCL recovery	-1.02 ± 2.48 (-17.77–3.72)	-1.44 ± 3.18 (-17.77–3.63)	-0.60 ± 1.38 (-4.07–3.72)	$t(125) = -1.91$
ANS regulation (reactivity)	0 ± 1.47 (-7.14–3.80)	0.07 ± 1.54 (-7.14–3.80)	-0.11 ± 1.36 (-4.28–3.40)	$t(153) = 0.78$
ANS regulation (recovery)	0.05 ± 1.33 (-6.04–3.41)	-0.10 ± 1.39 (-6.04–1.87)	0.19 ± 1.27 (-4.46–3.41)	$t(122) = -1.23$
Child race				$\chi^2(5) = 8.14$
<i>White</i>	72	37	35	
<i>Black</i>	12	6	6	
<i>Hispanic/Latino</i>	14	11	3	
<i>Asian</i>	19	15	4	
<i>Biracial</i>	39	22	17	
<i>Other</i>	7	3	4	
Family income				$\chi^2(9) = 6.83$
<i><\$5000</i>	1	1	0	
<i>\$5001-10,000</i>	1	1	0	
<i>\$10,001-15,000</i>	3	1	2	
<i>\$15,001-25,000</i>	3	3	0	
<i>\$25,001-35,000</i>	4	1	3	
<i>\$35,001-50,000</i>	7	3	4	
<i>\$50,001-75,000</i>	16	9	7	

<i>\$75,001-100,000</i>	19	10	9	
<i>\$100,001-150,000</i>	49	27	22	
<i>\$150,001+</i>	53	31	22	
<i>No response</i>	7	7	0	
Parental education				$\chi^2(7) = 4.43$
<i>No GED/High school diploma</i>	1	1	0	
<i>GED/High school diploma</i>	2	2	0	
<i>Some college</i>	33	22	11	
<i>2-Year college degree</i>	12	7	5	
<i>4-Year college degree</i>	58	30	28	
<i>Masters degree</i>	45	25	20	
<i>Professional degree</i>	6	3	3	
<i>Doctorate</i>	4	2	2	
<i>No response</i>	2	2	0	

* indicates $p < .05$, ** indicates $p < .01$, and *** indicates $p < .001$

Table 3. Correlation matrix among key variables across all participants.

	1	2	3	4	5	6	7	8	9	10	11	12	13
1. Age		-0.26**	0.66	-0.10	0.02	-0.03	-0.08	0.07	-0.01	-0.07	0.09	-0.09	0.04
2. Sex			0.05	0.22**	-0.01	0.12	0.10	- 0.11	0.05	0.11	0.02	0.15	-0.08
3. Pubertal stage				0.04	0.21**	0.20*	-0.05	0.06	-0.01	-0.07	0.12	-0.08	0.05
4. Stress sensitivity					0.06	0.62***	0.26**	0.04	-0.07	-0.17*	0.27**	-0.14	0.21*
5. Objective stress						0.82***	0.20*	0.02	0.00	-0.06	0.03	-0.05	0.01
6. Subjective stress							0.30***	0.04	-0.03	-0.18*	0.17	-0.15	0.12
7. MASC social anxiety scores								0.12	-0.07	0.09	0.02	-0.02	0.06
8. RSA reactivity									-0.62***	-0.06	-0.08	-0.73***	0.43***

9. RSA recovery										-0.19*	0.10	0.39***	-0.71***
10. SCL reactivity											-0.38***	0.73***	-0.08
11. SCL recovery												-0.13	0.63***
12. Physiological reactivity shift													-0.40***
13. Physiological recovery shift													

* indicates $p < .05$, ** indicates $p < .01$, and *** indicates $p < .001$

Table 4. Correlation matrix among key variables by sex. Correlations above the diagonal are among males. Correlations below the diagonal are among females.

	1	2	3	4	5	6	7	8	9	10	11	12
1. Age		0.72***	-0.09	0.07	0.02	-0.06	0.20	-0.08	-0.02	0.18	-0.12	0.09
2. Pubertal stage	0.69***		-0.02	0.16	0.11	-0.08	0.17	0.00	-0.07	0.19	-0.14	0.08
3. Stress sensitivity	-0.02	0.07		0.05	0.64***	0.10	-0.16	0.05	-0.20	0.34*	-0.06	0.18
4. Objective stress	-0.01	0.25*	0.08		0.80***	0.05	-0.20	0.32*	-0.01	-0.02	0.11	-0.29*
5. Subjective stress	-0.01	0.26*	0.60***	0.84***		0.10	-0.24*	0.30*	-0.21	0.18	-0.01	-0.15
6. MASC social anxiety scores	-0.06	-0.04	0.34**	0.29**	0.41***		0.02	0.08	0.17	0.02	0.13	-0.09
7. RSA reactivity	-0.03	0.01	0.20	0.13	0.20	0.19		-0.30*	-0.05	-0.09	-0.59***	0.08
8. RSA recovery	0.06	-0.01	-0.18	-0.19	-0.25*	-0.18	-0.80***		-0.30*	0.14	-0.03	-0.57***
9. SCL reactivity	-0.07	-0.08	-0.18	-0.12	-0.19	-0.01	-0.07	-0.07		-0.50***	0.83***	-0.12

10. SCL recovery	0.02	0.04	0.22	0.11	0.19	0.03	-0.08	0.06	-0.19		-0.26	0.73***
11. Physiological reactivity shift	-0.01	-0.06	-0.26*	-0.16	-0.26*	-0.16	- 0.83***	0.65***	0.61***	-0.02		-0.14
12. Physiological recovery shift	-0.04	0.02	0.28*	0.23*	0.33**	0.18	0.65***	-0.82***	-0.04	0.52***	-0.58***	

* indicates $p < .05$, ** indicates $p < .01$, and *** indicates $p < .001$