Assessing dysfunctional expectations in posttraumatic stress disorder – Development and validation of the Posttraumatic Expectations Scale (PTES)

Philipp Herzog^{*a,b,c}, Tim Kaiser^b, Winfried Rief^a, Eva-Lotta Brakemeier^{a,b †}, & Tobias Kube^{a,d,e †}

^a Philipps-University of Marburg, Department of Psychology, Division of Clinical Psychology and Psychological Treatment Gutenbergstraße 18, D-35032 Marburg, Germany

^b University of Greifswald, Department of Psychology, Clinical Psychology and Psychotherapy, Franz-Mehring-Straße 47, D-17489 Greifswald, Germany

^c Department of Psychiatry and Psychotherapy, University of Lübeck, Ratzeburger Allee 160, D-23562 Lübeck, Germany

^d University of Koblenz-Landau, Pain and Psychotherapy Research Lab, Ostbahnstr. 10, 76829 Landau, Germany

^e Harvard Medical School, Program in Placebo Studies, Beth Israel Deaconess Medical Center, Brookline Avenue 330, Boston, MA 02115, USA

*Correspondence address: Philipp Herzog, Philipps-University of Marburg, Department of Clinical Psychology and Psychotherapy, Gutenbergstraße 18, D-35032 Marburg, Germany, Email: philipp.herzog@staff.uni-marburg.de, Telephone: +49 6421 2824061, Fax: +49 6421 2828904

[†] Shared senior authorship: Eva-Lotta Brakemeier and Tobias Kube contributed equally.

Short title:

Assessing dysfunctional expectations in PTSD

24 **Conflict of interest statement**

25 The authors declare no conflict of interest.

26 **Funding**

27 This research did not receive any specific grant from funding agencies in the public,
28 commercial, or not-for-profit sectors.

29 **Author Contributions**

30 PH, ToKu, and ELB conceptualized the study. PH collected data and originally drafted and
31 edited the manuscript, ToKu and ELB supervised, reviewed, and edited the manuscript, TiKa
32 computed the statistical analyses, WR revisited the manuscript critically for important
33 intellectual content, and gave substantial feedback for the conception and design of this
34 study.

35 **Acknowledgments**

36 The authors would especially like to thank Dr. Kerstin Kühl for her valuable feedback at
37 generating the item pool, Anna-Lena Eckert for her support in translating the questionnaire
38 into English, and all bachelor and master graduate students as well as practitioners
39 (especially at the Schön Clinic, Parkland-Clinic, and the Outpatient Clinic Marburg) for their
40 support in recruiting patients for our study.

Abstract

Dysfunctional expectations are a particularly important subset of cognitions that influence the development and maintenance of various mental disorders. This study aimed to develop and validate a scale to assess dysfunctional expectations in posttraumatic stress disorder (PTSD), the “Posttraumatic Expectations Scale” (PTES). In a cross-sectional study, 70 PTSD patients completed the PTES, the Posttraumatic Cognitions Inventory (PTCI), as well as measures of the severity of symptoms of PTSD and depression. The results show that the PTES has excellent internal consistency and correlates significantly with the PTCI and PTSD symptom severity. A regression analysis revealed that the PTES explained variance of PTSD symptom severity above the PTCI, supporting the incremental validity of the PTES. While the original version of the PTES comprises 81 items, short scales were constructed using the BISCUIT method. The current findings suggest that the PTES is a reliable and valid novel self-report measure in patients with PTSD.

Keywords:

Expectations, posttraumatic stress disorder, PTSD, assessment, scale, validation

Introduction

While the psychopathology of Posttraumatic Stress Disorder (PTSD) is multifaceted and complex, the role of dysfunctional cognitions has regained attention in PTSD research in the last decade (Woud et al., 2019). In cognitively focused PTSD research, it is well known that the view of the world changes dramatically after exposure to a traumatic event (Bernardi et al., 2019; Brown et al., 2018; LoSavio et al., 2017). More specifically, research indicates that people's "belief in a just world" is violated by a traumatic event (Dalglish, 2004) and as a result, negative beliefs about the world, other people, and oneself are formed (Woud et al., 2019). Importantly, research found that negative appraisals of the traumatic event and its consequences are strongly associated with PTSD symptom severity; in addition, the persistence of such negative reappraisals predicts later symptom severity (Ehring et al., 2008; Kleim et al., 2013; ter Heide et al., 2017). Accordingly, in the DSM-5, cognitive changes after the experience of a traumatic event have been included as part of the new symptom cluster D "negative alterations in cognition and mood" (American Psychiatric Association, 2013), highlighting the importance of further research on the role of dysfunctional cognitions in PTSD.

Several insightful theories have been developed to elucidate how cognitive factors contribute to the development and maintenance of PTSD symptoms, such as the *emotional processing theory* (Foa & Kozak, 1986; Rauch & Foa, 2006) and the *cognitive processing approach* by Resick & Schnicke (1992). One of the most influential cognitive approaches to PTSD is the Cognitive Model of PTSD, developed by Ehlers & Clark (2000). This model assumes that PTSD symptoms persist when people continue to perceive the trauma with a sense of threat, resulting from negative appraisals of the traumatic event and distortions in autobiographic memory. Numerous empirical studies support the assumptions of the model

by Ehlers & Clark (2000): it has been shown that negative cognitions are associated with PTSD symptom severity and contribute to their exacerbation and maintenance (Dunmore et al., 1999; Ehling et al., 2008; Halligan et al., 2003; Kleim et al., 2013; ter Heide et al., 2017). Drawing on such cognitive accounts, psychological treatments focusing on the modification of dysfunctional cognitive processes have been developed, such as the cognitive processing therapy (CPT; Resick & Schnicke, 1992) and the cognitive therapy for PTSD (CT-PTSD; Ehlers, 1999; Ehlers & Clark, 2000).

In recent years, expectations have received particular attention in research on aberrant cognitive processes across mental disorders (Rief et al., 2015). Expectations have been defined as a subgroup of cognitions, relating to the likelihood of the occurrence of future events or experiences (Kube et al., 2016; Laferton et al., 2017; Olson et al., 1996). Unlike more general negative appraisals and dysfunctional cognitions, expectations offer the advantage of being amenable to a relatively straightforward empirical confirmation vs. disconfirmation due to their “if, then” structure (Rief et al., 2015). Thus, dysfunctional expectations might be a promising target in the cognitive-behavioral treatment of PTSD, as they may provide the basis of behavioral experiments in which patients sample new information that helps them revise their initial expectations (Craske et al., 2014; Murray & El-Leithy, 2020; Rief et al., 2015). The importance of expectations in PTSD has also recently been spelled out in neuroscientific and computational approaches that conceive PTSD in terms of hyper-precise prior predictions referring to the anticipation of threat, hence shaping perception towards the continued experience of threat, much like a self-fulfilling prophecy (Kube et al., 2020; Linson & Friston, 2019).

In order to systematically investigate how dysfunctional expectations influence the course of PTSD, it is important to be able to assess several types of expectations that are core

to PTSD. Although we are cognizant of some assessment tools that aim to examine dysfunctional cognitions in PTSD, such as the Posttraumatic Cognition Inventory (PTCI; Dunmore et al., 1999; Ehlers, 1999), there is, to our knowledge, no measure available that focuses specifically on expectations in PTSD, in terms of testable future-directed cognitions that are put in an “if-then” structure. When aiming to assess relevant types of expectations in PTSD, one may consider symptom-specific expectations referring to the memory of the traumatic event (e.g., “When I am reminded of the traumatic event, I will feel like being at the mercy of these memories.”) and the intense emotions related to the trauma (e.g., “When I experience unpleasant emotions such as fear or guilt, I will feel helpless coping with these emotions.”). Because such disorder-specific expectations have been shown to incrementally contribute to the development of depressive symptoms, beyond other types of cognitions (Kube et al., 2018; Kube et al., 2019), we believe that it might be valuable to examine them in PTSD as well. Besides disorder-specific expectations, it might also be worthwhile to consider patients’ treatment expectations (e.g., “If I confront myself with the traumatic event during therapy, I will feel much better in the longer run.”), as it has been shown that such outcome expectations predict treatment response for combat veterans suffering from PTSD (Price et al., 2015).

Accordingly, the present study sought to develop a new self-report questionnaire that allows examining both disorder-specific expectations in PTSD and expectations that might be relevant to cognitive-behavioral treatments of PTSD. To this end, we examined a clinical sample of individuals being treated for their PTSD symptoms, and we aimed to evaluate the psychometric properties of our new scale, the “Posttraumatic Expectations Scale” (PTES). In addition to reliability analyses, we particularly aimed to assess the incremental validity of our scale by examining whether it is able to predict PTSD symptom severity above the well-

128 established PTCL. Moreover, we aimed to build a short version of the scale in order to
129 facilitate its use in clinical practice.

Methods

This study was approved by the local ethics committee of the Faculty of Psychology at the Philipps-University of Marburg (reference number 2018-6k) and was conducted in accordance with the ethical standards as laid down in the Declaration of Helsinki (1964) and its later amendments.

Patients and Procedure

Data was collected between May 2018 and July 2019. Participation in this study was voluntary and no financial benefit was offered. Before inclusion in this study, all patients gave their informed consent. To be included, patients had to have the primary diagnosis of PTSD according to ICD-10, as verified by structured clinical interviews. Also, patients had to be at least 18 years old and fluent in German. Patients were recruited in several local outpatient and inpatient clinics in the surrounding of the study site. In both outpatient and inpatient settings, patients were enrolled after the diagnosis of PTSD was confirmed, but before treatment began. Because some of the expectations assessed in our scale relate to the anticipation of specific aspects of the therapy, it was crucial to complete them prior to the actual therapy. A total of 70 PTSD patients participated in this paper-pencil study.

Measures

Scale development and procedure: Posttraumatic Expectations Scale (PTES)

First, an extensive literature review of dysfunctional cognitions and expectations in PTSD was done. In doing so, we particularly focused on research on the cognitive model of PTSD (Ehlers & Clark, 2000) and the cognitive processing theory (Resick & Schnicke, 1992) to get a sense of what might be relevant domains where individuals with PTSD hold dysfunctional expectations. Second, various experts in the field of PTSD research and who frequently work with PTSD patients were consulted to ask them for domains of

dysfunctional expectations that, in their experience, are core to PTSD. As a result, we defined the following domains for item development: general treatment expectations (subscale A; e.g., “In psychotherapy I will feel supported.”), exposure to the traumatic event during psychotherapy (subscale B; e.g., “If I face the traumatic event in psychotherapy, I will lose control.”), talking about the traumatic event in psychotherapy (subscale C; e.g., “If I tell my psychotherapist about the traumatic event, she/he will blame me for it.”), reexperiencing symptoms (subscale D; e.g., “When I am reminded of the traumatic event, I will feel helpless.”), emotion regulation (subscale E; e.g., “When I experience unpleasant feelings (e.g., blame, anger, sadness, disgust...) in relation to the traumatic experience *outside* of the psychotherapy, there will be nothing I can do to feel better.”), openness to other people (subscale F; e.g., “When I confide in other people (e.g. significant others), they will misuse my trust.”), encounters with strangers (subscale G; e.g., “If I meet strangers, I will be able to make new acquaintances.” (inversely scored)), and avoidance behavior (subscale H; e.g., “If I stop avoiding certain things that remind me of the traumatic experience(s) (e.g., places, activities, people), I will reexperience a traumatic event.”). While the first three subscales (i.e., subscales A-C) refer to expectations regarding the psychological treatment of PTSD, subscales D-H were developed to assess expectations that relate to various disorder-specific aspects. In terms of domain sampling (Nunnally, 1978), we used these nine categories to derive specific items that were formulated in accordance with the existing expectation literature and theories (Rief et al., 2015; Rief & Glombiewski, 2017); that is, we paid particular attention to the fact that items are put in a testable, future-directed manner. Specifically, when developing each item, we carefully thought about whether it would allow an empirical test of the respective expectation in a behavioral experiment. Furthermore, we used an additional ninth category named “general disorder-specific expectations” (subscale

I), where additional expectations were assessed that we considered important but could not clearly be linked to one the aforementioned domains (e.g., “I will be able to live a happy life, despite the traumatic experience I had.”).

As a result of this item generation process, an 81-item self-report scale was developed. The items are rated on a 5-point Likert-Scale ranging from 1 (‘I don’t agree at all’) to 5 (‘I totally agree’). Lower sum scores indicate fewer dysfunctional expectations. As we aimed to balance negative expectations with positive expectations, about half of the items need to be inversely scored. Originally, the scale was developed in German, as the questionnaire was planned to be administered in German in the present validation study; for this article, we translated the scale into English. All 81 items of the English version of the PTES are presented in the supplemental material (see supplemental material 1). Upon request, the German version can be made available by the corresponding author.

Dysfunctional cognitions in PTSD: Posttraumatic Cognitions Inventory

The Posttraumatic Cognitions Inventory (PTCI) is a 33-item self-report questionnaire measuring maladaptive cognitions in relation to a traumatic event and/or its consequences (Dunmore et al., 1999; Ehlers, 1999). Participants are asked to rate their agreement with the statements on a 7-point Likert-scale ranging from 1 (‘totally disagree’) to 7 (‘totally agree’). The PTCI consists of three subscales: negative cognitions about the self, the world, and self-blame (Foa et al., 1999). Beyond its use in the original English version, the PTCI has also been used and validated in a German sample (Müller et al., 2010). Along with the original 33-item version of the PTCI, the Interpretation of PTSD Symptoms Inventory (IPSI; Dunmore et al., 1999) was used in this study. This scale consists of 11 items that measures the distress by intrusions, problems in remembering the traumatic event and the consequences of a traumatic event. The PTCI and IPSI was shown to have an excellent

reliability across studies (Foa et al., 1999; Halligan et al., 2003; Müller et al., 2010).

Symptom severity: Impact-of-event-Scale-revised

The severity of PTSD symptoms was assessed with the Impact-of-event-Scale-Revised (IES-R). The IES-R is a 22-item self-report questionnaire measuring three areas of PTSD symptoms, namely, intrusion, avoidance, and hyperarousal (Maercker & Schützwohl, 1998). Patients suffering from PTSD are asked to rate the degree of distress of each item during the past week on a 4-point-Likert scale ranging from 0 ('not at all') to 3 ('many times'). The IES-R was shown to have high internal consistency (Creamer et al., 2003).

Other measures

PHQ-9. Depressive symptoms were assessed with the Patient Health Questionnaire-9 (Gräfe et al., 2004; Kroenke et al., 2001), and has been extensively validated previously and shown excellent psychometric properties (Kroenke et al., 2010).

Sociodemographics. Socio-demographic questions were asked in a self-report questionnaire including age, sex, education level, and marital status.

Statistical Analyses

Preprocessing. In general, data screening was performed according to the recommendations of Tabachnick & Fidell (2013). Outliers, extreme values and missing values were systematically inspected. The squared Mahalanobis distance was computed to determine multivariate outliers indicating no significant outliers in our sample (Jacob Cohen et al., 1983; Stevens, 2009). The MCAR-Test (Little, 1988) was used to identify if missing values were missing not at random (MNAR) or missing completely at random (MCAR), respectively, missing at random (MAR). The MCAR-Test showed no significant result ($\chi^2(5675, N = 70) < .0001$ $p = 1.00$), indicating a MCAR distribution of missing values.

Missing values were imputed with multiple imputations with chained equations (MICE; van

Buuren & Groothuis-Oudshoorn, 2011). Since 27% of the cases had missing values, 30 imputed data sets with 10 iterations were generated. Descriptive item and scale analyses were performed. Bivariate Pearson correlations between the scales and p-values were computed. The p-values were corrected according to the Holm-Method for multiple comparisons.

Reliability and validity analyses. Reliability was determined by computing internal consistency (Cronbach's alpha). To determine the incremental validity of the PTES, beyond the PTCL, a multiple hierarchical regression analysis was performed for the complete scale, with symptom severity (IES-R) being the dependent variable. On a subscale level, due to the small sample size, a cross-validated LASSO regression was computed that sets regularized parameters to 0. The coefficients are standardized (beta) and all coefficients not set to 0 predict the respective scale.

Building a criterion-keyed short scale. BISCUIT (best items scales that are cross validated, unit weighted, informative and transparent) analyses were performed (Elleman et al., 2020). From a statistical point of view, BISCUIT build scales of a predetermined number of items that show the strongest pairwise correlations with a certain criterion and cross-validates or bootstraps these scales. The cross-validated items showing the highest pairwise correlation with the criterion were then combined to a scale. For combination, all retained items are weighted equally (i.e., unit weighted). This way, a realistic estimate of out-of-sample generalizability is obtained, as recommended by Cohen (1992). BISCUIT is particularly robust to high proportions of missing data and limitations in sample (Elleman et al., 2020). The short scale derived from the complete PTES was validated against the IES-R with an upper bound of a maximum of 15 items. In addition, short scales were developed for

249 the subscales of the IES-R with an upper bound of a maximum of 30 items (10 items for
250 each of the three subscales).

Results

Patient characteristics

Of all 70 PTSD patients, 82.6% were female with a mean age of $M = 42.06$ years ($SD = 11.82$). 45.7% had children, with a mean number of $M = 2.22$ ($SD = 1.09$) children per subject, and 32.9% were married while 30% were single. The educational level was relatively equally distributed; while 61.4% had completed an apprenticeship or vocational training, 1.4% had a higher vocational qualification (e.g., master craftsman), 11.4% had completed a university degree, 4.3% had another vocational training qualification, 20.0% had no vocational training.

The mean age of onset of PTSD was $M = 23.70$ years ($SD = 15.45$) with a mean of $M = 17.96$ years of manifestation ($SD = 14.64$). According to the LEC-5, patients reported on average to have experienced five traumatic events ($M = 5.31$; $SD = 2.88$), with sexual assaults being the most frequently reported trauma (54%). The average sum score of the IES-R was 79.24 ($SD = 16.58$), indicating that the sample was severely impaired on average (intrusion subscale: 26.92 ($SD = 6.72$); avoidance subscale: 24.17 ($SD = 7.59$); hyperarousal subscale: 28.11 ($SD = 6.68$)). The average sum score of the PTCI was 155.58 ($SD = 35.05$) and the mean value of the IPSI was $M = 4.03$ ($SD = 1.34$), both speaking to presence of highly dysfunctional cognitions as compared to other PTSD samples (Ehlers, 1999; Müller et al., 2010). In line with these clinical characteristics of the current sample, the mean PHQ-9 sum score was $M = 20.70$ ($SD = 5.52$), indicating severe comorbid depressive symptoms. Table 1 portrays the relevant sociodemographic and clinical characteristics of the sample.

Insert Table 1 here.

Scale analyses, construct validity and reliability of the PTES

The items statistics of the PTES and the descriptive statistics of all subscales can be found in the supplements (see supplemental material 2 and 3). For the entire PTES and the respective subscales, the reliabilities are displayed in Table 2, along with the inter-correlations of all scales used in the present study.

Insert Table 2 here.

Correlational analyses showed that the sum score of the PTES was significantly correlated with all other scales used in the present study. Specifically, the correlation of the PTES with the PTCI was $r = .66$; its correlation with the IES-R was $r = .54$; and the correlation with the PHQ-9 was $r = .63$. The PTCI was also significantly correlated with the IES-R ($r = .57$) and the PHQ-9 ($r = .60$), with similar correlation coefficients as the PTES.

The internal consistency of the total scale was $\alpha = .96$ (95% CI [.94, .97]) indicating a very high reliability. The internal consistency of the PTES subscales ranged between $\alpha = .52$ (95% CI [.34, .70]) for the scale A, assessing general treatment expectations, and $\alpha = .92$ (95% CI [.89, .95]) for the scale F, assessing openness to other people. Among all subscales, only two subscales did not reach the critical value of $\alpha = .70$ as an acceptable reliability for newly developed scales as proposed by DeVellis (2012); namely, subscale A (general treatment expectations) and subscale C (talking about the traumatic event in therapy).

Incremental validity analyses

The results of the regression analyses revealed that the PTCI as the sole predictor explained 31.4% of the variance of the IES-R (Fit $R^2 = .314$; R^2 95% CI [.12, .52]). When including the PTES sum score as an additional predictor, a total of 36.4% of the variance was explained (Fit $R^2 = .364$; R^2 95% CI [.18, .55]). The difference in the explained outcome variance of 5.0% (difference $\Delta R^2 = -.050$; 95% CI [-.06, -.03]) was significant; thus, the

PTES had incremental value over the PTCI. When examining the predictive value of the subscales of the PTES, the results indicated a distinct predictor profile as displayed in Table 3. In both prediction models (IES-R and PHQ-9), the PTCI was the strongest predictor, additionally there were some PTES subscales found to be important predictors; e.g., the subscale A (general treatment expectations) ($\beta = -.037$), subscale E (emotion regulation) ($\beta = .129$) and subscale I (general disorder-specific expectations) ($\beta = .239$).

Insert Table 3 here.

Building a short scale

The BISCUIT method extracted 15 PTES items when validated on the IES-R sum score as the criterion building a 15-items ultra-short scale. Separately, the BISCUIT method extracted 10 PTES items when validated on each of the three IES-R subscales (i.e., intrusion, avoidance, hyperarousal) as criteria leading together to a 30-items short scale. These items are displayed in Table 4.

Using the IES-R sum score as the criterion, the best symptom-specific items seemed to be Item 74, Item 40, Item 48, Item 71, Item 39, Item 70, Item 78, Item 77, Item 58, Item 5, Item 32, Item 81, Item 72, Item 43, and Item 33. These items are part of the 15-items ultra-short scale that yields a more favorable validity (see appendix for the ultra-short PTES). When applying the IES-R subscales as criteria, more general expectations were selected as criteria for the intrusion subscale and the hyperarousal subscale, whereas expectations regarding negative emotions caused by the traumatic event (scale E) and expectations regarding openness to other people (scale F) were selected as criteria for the avoidance subscale in particular. As the subscales are interrelated, some items were selected repeatedly leading to less favorable validity results.

Insert Table 4 here.

Discussion

In this study, the Posttraumatic Expectations Scale (PTES) was developed in an iterative and theory-driven process to assess both disorder-specific expectations and treatment expectations in a clinical sample of patients with PTSD. The results of the reliability and validity analyses showed that the PTES is a reliable measure and its validity appears promising. More specifically, the results of this validation study showed excellent internal consistency of the entire scale and most of the subscales. Moreover, in cross-sectional analyses, the PTES was significantly correlated with the well-established PTCI, as a measure for more general dysfunctional cognitions, as well as the severity of PTSD symptoms, speaking to the concurrent validity of the PTES. With respect to its incremental validity, the results of the regression analyses revealed that the PTES incrementally explained variance of the severity of PTSD symptoms over the PTCI, suggesting that the PTES might be a useful addition to the PTCI. To enhance the applicability for practitioners, the present study used a state-of-the-art approach (Elleman et al., 2020) to construct short scales that can easily be administered in clinical practice, with 15 items for the total symptom-specific criterion and 10 items for each symptom-specific criterion. Collectively, the results of the reliability and validity analyses suggest that the PTES might be a useful tool to assess several types of dysfunctional expectations in PTSD patients reliably.

Theoretical implications

While general cognitions, such as negative appraisals of the traumatic event and its consequences have been well-studied in PTSD research (Dunmore et al., 1999; Ehrling et al., 2008; Halligan et al., 2003; Kleim et al., 2013; ter Heide et al., 2017; Woud et al., 2019), the present study is the first to systematically focus on expectations as a particularly important subset of cognitions. The results of this study showed that expectations (as measured by the

PTES) indeed explained incremental variance of symptoms of PTSD beyond general cognitions (as measured by the PTCI); therefore, the PTES seems to have value in shifting the focus to patients' expectations, i.e., expectations resulting from core features of the disorder (such as expectations about perceived threat) and expectations concerning psychological treatment. Thus, the present findings might have the potential to extend the cognitive model of PTSD (Ehlers & Clark, 2000) by differentiating between different types of cognitions and therefore refining the association between negative appraisals of the trauma and/or its sequelae and the experience as a current threat (intrusions, arousal symptoms, strong emotions) within the cognitive model. In particular, expectations as assessed with our newly developed scale reflect a person's prediction of future events and experiences. Recent advances in cognitive neuroscience suggest people's predictions significantly affect their perception; in other words, people interpret new information in light of their prior predictions (Bastos et al., 2012; Friston, 2009; Kanai et al., 2015). It has been suggested that in people with PTSD, predictions are tied to the anticipation of threat (due to the (life-)threatening nature of the traumatic event) that lead to the perception of the world as being insecure and dangerous (Kube et al., 2020; Linson & Friston, 2019). Consistent with these theories from computational psychiatry, merit of the PTES can be seen in its systematic assessment of people's predictions that, consequently, might offer insights into how – and why – people with PTSD perceive their world. Drawing on our approach, future research may aim to investigate how patients' expectations in various domains predict the course of PTSD symptoms in longitudinal designs.

Clinical implications

Although meta-analyses and systematic reviews of randomized-controlled trials (Bisson et al., 2007; Cusack et al., 2016; Ehlers et al., 2010) and practice-based studies (Herzog et

al., 2020) indicate the general efficacy of psychotherapy (especially individual trauma-focused cognitive-behavioral therapy (CBT)) in PTSD patients with moderate to large effect sizes across various settings, non-response rates and dropout rates are relatively high (Bradley et al., 2005; Schottenbauer et al., 2008). Therefore, there is space for optimizing the effects of psychotherapy in the treatment of PTSD. Since empirical hypothesis-testing is the backbone of CBT for PTSD in general, and of CT-PTSD (Ehlers, 1999; Ehlers & Clark, 2000) and CPT (Resick & Schnicke, 1992) in particular, the PTES might have the potential to facilitate this approach by allowing a structured assessment of patients' expectations that can subsequently be tested in interventions such as behavioral experiments. Indeed, changes in posttraumatic cognitions predict changes in CT (Kleim et al., 2013) and CPT (Schumm et al., 2015) for PTSD. Drawing on the work by Kleim et al. (2013) and Schumm et al. (2015), the special focus on dysfunctional expectations in patients with PTSD might offer several advantages for practitioners: Knowledge about situation-specific expectations might be useful in designing effective interventions such as behavioral experiments. When considering dysfunctional expectations instead of more general cognitions, the empirical testing of hypotheses may be more straight-forward, i.e., dysfunctional beliefs might be easier disconfirmed by new information as gained in a behavioral experiment. To illustrate this argument, we discuss below some examples of how PTES might be used to optimize psychotherapy by the use of behavioral experiments in PTSD.

For instance, the dysfunctional expectation as assessed with the PTES, "When I confide in other people (e.g., significant others), they will misuse my trust.", might be easier to test in behavioral experiment than the more global cognition, "People can't be trusted", as assessed with the PTCI. We believe so because the latter is hard to evaluate empirically, whereas the former can be examined in a relatively straight forward manner: The therapist

may encourage the patient to talk to a significant other (e.g., a member of the family) and observe whether the other person takes advantage of that situation. Obviously, to be an effective intervention, the therapist and the patient in this example would also have to discuss what “misuse” or “take advantage” could mean exactly, and under which circumstances the behavioral experiment would – in the patients’ perspective – reveal useful and trustworthy information, such that the patient would use it to revise the initial expectation. To give another example, the PTES might also allow an easier disconfirmation of avoidance-related expectations. For instance, while the expectation, “If I stop avoiding certain things that remind me of the traumatic experience(s) (e.g., places, activities, people), I will re-experience a traumatic event.” might be easier to test in behavioral experiment or an exposure session (e.g., by exposure the patient to triggering cues) than the more global cognition, “The world is a dangerous place” from the PTCI.

A structured assessment of patients’ individual dysfunctional expectations with the PTES might help optimize case conceptualization and, accordingly, developing a treatment plan, at an early treatment stage and targeting these expectations subsequently. Furthermore, the PTES might be completed repeatedly throughout the therapy, such that it can be examined whether the interventions to modify patients’ expectations were actually effective. As an additional advantage, addressing expectation violation through the use of idiosyncratic behavioral experiments could be promising in remotely delivered treatments which recently gained attention in times of Covid-19 and its related challenges in delivering evidence-based practice (Murray & El-Leithy, 2020).

Limitations and future research directions

This study serves as a preliminary validation of the PTES as a newly developed questionnaire assessing dysfunctional expectations of PTSD patients. By interpreting the results of this study, the following limitations must be taken into account:

This study used a cross-sectional design and therefore does not allow any conclusions about causality. Future studies could incorporate the shortened version of the PTES in longitudinal studies (for example in RCTs) using larger samples, which would offer several advantages: first, it can provide information on the causal pathways between expectations change and its relation to symptom severity throughout treatment; second, it can be used to determine test-retest-reliability; and third, it would allow more sophisticated statistical procedures to determine the factor validity (e.g., CFA). A further drawback of this study might be seen in the characteristics of the sample; specifically, PTSD patients in the current sample reported relatively high impairment and severe symptoms as compared to previous studies using similar measures (Müller et al., 2010). At least in part, this might be due to the fact that the most frequent traumatic events were sexual assaults and other unwanted or unpleasant sexual experiences as well as other interpersonal traumas, whereas the experience of an accidental trauma was rarely reported. In fact, the experience of an interpersonal trauma has been shown to cause more dysfunctional cognitions and greater symptom severity than an accidental trauma (Foa et al., 1999; Müller et al., 2010; Shakespeare-Finch & Armstrong, 2010; Startup et al., 2007).

Moreover, this study used only self-rating questionnaires for the assessment of PTSD-specific cognitions and expectations as well as PTSD symptom severity. Regarding the potential risk of common method bias, future studies on the PTES may additionally use an observer-rated instrument such as the Clinician-Administered PTSD Scale for DSM-5

(CAPS-5; Weathers et al., 2018). In terms of the psychometric properties of the PTES, it must be noted that while the total scale and most of the subscales of the PTES were shown to be reliable and valid, the subscale A (i.e., treatment-specific expectations) had lower values of Cronbach's alpha. Also, this subscale seemed to be relatively unrelated to the total scale; therefore, it might be considered to analyze this scale independent of the other subscales of the PTES. An additional limitation of the present study is that the English version of the PTES (see supplemental material 1) was not tested. The authors encourage other researchers to further validate the PTES in English-speaking countries as the data we used in our study was a German-speaking sample of PTSD patients.

Conclusions

In the present study, we used a sample of 70 patients with PTSD to develop the Posttraumatic Expectations Scale (PTES) as a new questionnaire assessing dysfunctional expectations in relation to PTSD. The results suggest that the PTES is a highly reliable assessment tool that incrementally explains variance of PTSD symptom severity, above and beyond the well-established Posttraumatic Cognition Inventory (PTCI). The PTES might have the potential to further improve cognitive-behavioral treatment of PTSD as it allows a structured assessment of various domains of dysfunctional expectations, which can be used as a basis for interventions such as behavioral experiments or exposure-based interventions, with the aim of disconfirming patients' expectations and facilitating their modification.

References

- American Psychiatric Association. (2013). *Diagnostic and Statistical Manual of Mental Disorders*. American Psychiatric Association.
<https://doi.org/10.1176/appi.books.9780890425596>
- Bastos, A. M., Usrey, W. M., Adams, R. A., Mangun, G. R., Fries, P., & Friston, K. J. (2012). Canonical Microcircuits for Predictive Coding. In *Neuron* (Vol. 76, Issue 4, pp. 695–711). Neuron. <https://doi.org/10.1016/j.neuron.2012.10.038>
- Bernardi, J., Engelbrecht, A., & Jobson, L. (2019). The impact of culture on cognitive appraisals: Implications for the development, maintenance, and treatment of posttraumatic stress disorder. *Clinical Psychologist*, 23(2), 91–102.
<https://doi.org/10.1111/cp.12161>
- Bisson, J. I., Ehlers, A., Matthews, R., Pilling, S., Richards, D., & Turner, S. (2007). Psychological treatments for chronic post-traumatic stress disorder: Systematic review and meta-analysis. In *British Journal of Psychiatry* (Vol. 190, Issue FEB., pp. 97–104).
<https://doi.org/10.1192/bjp.bp.106.021402>
- Bradley, R., Ph, D., Greene, J., Russ, E., Dutra, L., Westen, D., Ph, D., & Al, E. T. (2005). Reviews and Overviews A Multidimensional Meta-Analysis of Psychotherapy for PTSD. *American Journal of Psychiatry*, 162(February), 214–227.
<https://doi.org/10.1093/clipsy/bpg024>
- Brown, L. A., Belli, G. M., Asnaani, A., & Foa, E. B. (2018). A Review of the Role of Negative Cognitions About Oneself, Others, and the World in the Treatment of PTSD. *Cognitive Therapy and Research*. <https://doi.org/10.1007/s10608-018-9938-1>
- Cohen, J. (1992). Things I have learned (so far). In A. E. Kazdin (Ed.), *Methodological*

- 482 *issues & strategies in clinical research* (pp. 315–333). American Psychological
483 Association.
- 484 Cohen, J., Cohen, P., West, S., & Aiken, L. (1983). *Applied Multiple Regression /*
485 *Correlation Analysis for the Behavioral Sciences* (2nd ed) by. 2nd Ed Hillsdale NJ
486 *Lawrence Erlbaum Associates, Third Edit*, 703 S.
487 <http://books.google.com/books?hl=de&lr=&id=fuq94a8C0ioC&pgis=1>
- 488 Craske, M. G., Treanor, M., Conway, C. C., Zbozinek, T., & Vervliet, B. (2014).
489 Maximizing exposure therapy: an inhibitory learning approach. *Behaviour Research*
490 *and Therapy*, 58, 10–23. <https://doi.org/10.1016/j.brat.2014.04.006>
- 491 Creamer, M., Bell, R., & Failla, S. (2003). Psychometric properties of the Impact of Event
492 Scale - Revised. *Behaviour Research and Therapy*, 41(12), 1489–1496.
493 <https://doi.org/10.1016/j.brat.2003.07.010>
- 494 Cusack, K., Jonas, D. E., Forneris, C. A., Wines, C., Sonis, J., Middleton, J. C., Feltner, C.,
495 Brownley, K. A., Olmsted, K. R., Greenblatt, A., Weil, A., & Gaynes, B. N. (2016).
496 Psychological treatments for adults with posttraumatic stress disorder: A systematic
497 review and meta-analysis. *Clinical Psychology Review*, 43(290), 128–141.
498 <https://doi.org/10.1016/j.cpr.2015.10.003>
- 499 Dalglish, T. (2004). Cognitive Approaches to Posttraumatic Stress Disorder: The Evolution
500 of Multirepresentational Theorizing. *Psychological Bulletin*, 130(2), 228–260.
501 <https://doi.org/10.1037/0033-2909.130.2.228>
- 502 DeVellis, R. F. (2012). *Scale development : theory and applications*. SAGE.
503 [https://books.google.de/books/about/Scale_Development.html?id=vmwBHYuchfAC&r](https://books.google.de/books/about/Scale_Development.html?id=vmwBHYuchfAC&redir_esc=y)
504 [edir_esc=y](https://books.google.de/books/about/Scale_Development.html?id=vmwBHYuchfAC&redir_esc=y)

- 505 Dunmore, E., Clark, D. M., & Ehlers, A. (1999). Cognitive factors involved in the onset and
506 maintenance of posttraumatic stress disorder (PTSD) after physical or sexual assault.
507 *Behaviour Research and Therapy*, 37, 809–829. [https://doi.org/10.1016/S0005-](https://doi.org/10.1016/S0005-7967(98)00181-8)
508 7967(98)00181-8
- 509 Ehlers, A. (1999). *Posttraumatische Belastungsstörung*. Hogrefe, Verl. für Psychologie.
- 510 Ehlers, A., Bisson, J., Clark, D. M., Creamer, M., Pilling, S., Richards, D., Schnurr, P. P.,
511 Turner, S., & Yule, W. (2010). Do all psychological treatments really work the same in
512 posttraumatic stress disorder? *Clinical Psychology Review*, 30(2), 269–276.
513 <https://doi.org/10.1016/j.cpr.2009.12.001>
- 514 Ehlers, A., & Clark, D. M. (2000). A cognitive model of posttraumatic stress disorder.
515 *Behaviour Research and Therapy*, 38(4), 319–345. [https://doi.org/10.1016/S0005-](https://doi.org/10.1016/S0005-7967(99)00123-0)
516 7967(99)00123-0
- 517 Ehring, T., Ehlers, A., & Glucksman, E. (2008). Do cognitive models help in predicting the
518 severity of posttraumatic stress disorder, phobia, and depression after motor vehicle
519 accidents? A prospective longitudinal study. *Journal of Consulting and Clinical*
520 *Psychology*, 76(2), 219–230. <https://doi.org/10.1037/0022-006X.76.2.219>
- 521 Elleman, L. G., McDougald, S. K., Condon, D. M., & Revelle, W. (2020). That takes the
522 BISCUIT : A comparative study of predictive accuracy and parsimony of four statistical
523 learning techniques in personality data , with data missingness conditions. *Preprint*, 1–
524 23. <https://doi.org/10.31234/osf.io/tuqap>
- 525 Foa, E. B., & Kozak, M. J. (1986). Emotional Processing of Fear. Exposure to Corrective
526 Information. In *Psychological Bulletin* (Vol. 99, Issue 1, pp. 20–35).
527 <https://doi.org/10.1037/0033-2909.99.1.20>

- 528 Foa, E. B., Tolin, D. F., Ehlers, A., Clark, D. M., & Orsillo, S. M. (1999). The Posttraumatic
529 Cognitions Inventory (PTCI): Development and validation. *Psychological Assessment*,
530 11(3), 303–314. <https://doi.org/10.1037/1040-3590.11.3.303>
- 531 Friston, K. (2009). The free-energy principle: a rough guide to the brain? *Trends in*
532 *Cognitive Sciences*, 13(7), 293–301. <https://doi.org/10.1016/j.tics.2009.04.005>
- 533 Gräfe, K., Zipfel, S., Herzog, W., & Löwe, B. (2004). Screening psychischer störungen mit
534 dem “Gesundheitsfragebogen für Patienten (PHQ-D)”. Ergebnisse der Deutschen
535 validierungsstudie. *Diagnostica*, 50(4), 171–181. [https://doi.org/10.1026/0012-](https://doi.org/10.1026/0012-1924.50.4.171)
536 1924.50.4.171
- 537 Halligan, S. L., Michael, T., Clark, D. M., & Ehlers, A. (2003). Posttraumatic stress disorder
538 following assault: The role of cognitive processing, trauma memory, and appraisals.
539 *Journal of Consulting and Clinical Psychology*, 71(3), 419–431.
540 <https://doi.org/10.1037/0022-006X.71.3.419>
- 541 Herzog, P., Voderholzer, U., Gärtner, T., Osen, B., Svitak, M., Doerr, R., Rolvering-
542 Dijkstra, M., Feldmann, M., Rief, W., & Brakemeier, E. L. (2020). Predictors of
543 outcome during inpatient psychotherapy for posttraumatic stress disorder: a single-
544 treatment, multi-site, practice-based study. *Psychotherapy Research*.
545 <https://doi.org/10.1080/10503307.2020.1802081>
- 546 Kanai, R., Komura, Y., Shipp, S., & Friston, K. (2015). Cerebral hierarchies: predictive
547 processing, precision and the pulvinar. *Philosophical Transactions of the Royal Society*
548 *B: Biological Sciences*, 370(1668), 20140169. <https://doi.org/10.1098/rstb.2014.0169>
- 549 Kleim, B., Grey, N., Wild, J., Nussbeck, F. W., Stott, R., Hackmann, A., Clark, D. M., &
550 Ehlers, A. (2013). Cognitive change predicts symptom reduction with cognitive therapy

- 551 for posttraumatic stress disorder. *Journal of Consulting and Clinical Psychology*, 81(3),
552 383–393. <https://doi.org/10.1037/a0031290>
- 553 Kroenke, K., Spitzer, R. L., & Williams, J. B. W. (2001). The PHQ-9. *Journal of General*
554 *Internal Medicine*, 16(9), 606–613. [https://doi.org/10.1046/j.1525-](https://doi.org/10.1046/j.1525-1497.2001.016009606.x)
555 1497.2001.016009606.x
- 556 Kroenke, K., Spitzer, R. L., Williams, J. B. W., & Löwe, B. (2010). The Patient Health
557 Questionnaire Somatic, Anxiety, and Depressive Symptom Scales: A systematic
558 review. *General Hospital Psychiatry*, 32(4), 345–359.
559 <https://doi.org/10.1016/j.genhosppsych.2010.03.006>
- 560 Kube, T., Siebers, V. H. A., Herzog, P., Glombiewski, J. A., Doering, B. K., & Rief, W.
561 (2018). Integrating situation-specific dysfunctional expectations and dispositional
562 optimism into the cognitive model of depression – A path-analytic approach. *Journal of*
563 *Affective Disorders*, 229. <https://doi.org/10.1016/j.jad.2017.12.082>
- 564 Kube, T., Berg, M., Kleim, B., & Herzog, P. (2020). Rethinking post-traumatic stress
565 disorder – A predictive processing perspective. *Neuroscience & Biobehavioral Reviews*,
566 113(April), 448–460. <https://doi.org/10.1016/j.neubiorev.2020.04.014>
- 567 Kube, T., D’Astolfo, L., Glombiewski, J. A., Doering, B. K., & Rief, W. (2016). Focusing
568 on situation-specific expectations in major depression as basis for behavioural
569 experiments - Development of the Depressive Expectations Scale. *Psychology and*
570 *Psychotherapy: Theory, Research and Practice*. <https://doi.org/10.1111/papt.12114>
- 571 Kube, T., Herzog, P., Michalak, C. M., Glombiewski, J. A., Doering, B. K., & Rief, W.
572 (2019). Further Specifying the Cognitive Model of Depression: Situational
573 Expectations and Global Cognitions as Predictors of Depressive Symptoms. *Clinical*

- 574 *Psychology in Europe*, 1(4). <https://doi.org/10.32872/cpe.v1i4.33548>
- 575 Laferton, J. A. C., Kube, T., Salzmann, S., Auer, C. J., & Shedden-Mora, M. C. (2017).
576 Patients' expectations regarding medical treatment: A critical review of concepts and
577 their assessment. In *Frontiers in Psychology* (Vol. 8, Issue FEB, p. 233). Frontiers
578 Research Foundation. <https://doi.org/10.3389/fpsyg.2017.00233>
- 579 Linson, A., & Friston, K. (2019). Reframing PTSD for computational psychiatry with the
580 active inference framework. *Cognitive Neuropsychiatry*, 24(5), 347–368.
581 <https://doi.org/10.1080/13546805.2019.1665994>
- 582 Little, R. J. A. (1988). A Test of Missing Completely at Random for Multivariate Data with
583 Missing Values. *Journal of the American Statistical Association*, 83(404), 1198.
- 584 LoSavio, S. T., Dillon, K. H., & Resick, P. A. (2017). Cognitive factors in the development,
585 maintenance, and treatment of post-traumatic stress disorder. *Current Opinion in*
586 *Psychology*, 14, 18–22. <https://doi.org/10.1016/j.copsyc.2016.09.006>
- 587 Maercker, A., & Schützwohl, M. (1998). Erfassung von psychischen Belastungsfolgen: Die
588 Impact of Event Skala-revidierte Version. *Diagnostica*, 44(3), 130–141.
589 <https://doi.org/10.1037/t01210-000>
- 590 Müller, J., Wessa, M., Rabe, S., Dörfel, D., Knaevelsrud, C., Flor, H., Maercker, A., & Karl,
591 A. (2010). Psychometric properties of the Posttraumatic Cognitions Inventory (PTCI) in
592 a German sample of individuals with a history of trauma. *Psychological Trauma:*
593 *Theory, Research, Practice, and Policy*, 2(2), 116–125.
594 <https://doi.org/10.1037/a0018603>
- 595 Murray, H., & El-Leithy, S. (2020). Behavioural Experiments in Cognitive Therapy for

- 596 Posttraumatic Stress Disorder: Why, When, and How? *Verhaltenstherapie*, 1–11.
597 <https://doi.org/10.1159/000511921>
- 598 Nunnally, J. C. (1978). *Psychometric Theory*. McGraw-Hill.
- 599 Olson, J. M., Roese, N. J., & Zanna, M. P. (1996). Expectancies. In E. T. Higgins & A. W.
600 Kruglanski (Eds.), *Social psychology: Handbook of basic principles* (pp. 211–238). The
601 Guilford Press.
- 602 Price, M., Maples, J. L., Jovanovic, T., Norrholm, S. D., Heekin, M., & Rothbaum, B. O.
603 (2015). An investigation of outcome expectancies as a predictor of treatment response
604 for combat veterans with PTSD: Comparison of clinician, self-report, and biological
605 measures. *Depression and Anxiety*, 32(6), 392–399. <https://doi.org/10.1002/da.22354>
- 606 Rauch, S., & Foa, E. (2006). Emotional processing theory (EPT) and exposure therapy for
607 PTSD. *Journal of Contemporary Psychotherapy*, 36(2), 61–65.
608 <https://doi.org/10.1007/s10879-006-9008-y>
- 609 Resick, P. A., & Schnicke, M. K. (1992). Cognitive processing therapy for sexual assault
610 victims. *Journal of Consulting and Clinical Psychology*, 60(5), 748–756.
611 <https://doi.org/10.1037//0022-006x.60.5.748>
- 612 Rief, W., & Anna Glombiewski, J. (2017). The role of expectations in mental disorders and
613 their treatment. *World Psychiatry*, 16(2), 210–211. <https://doi.org/10.1002/wps.20427>
- 614 Rief, W., Glombiewski, J. A., Gollwitzer, M., Schubö, A., Schwarting, R., & Thorwart, A.
615 (2015). Expectancies as core features of mental disorders. *Current Opinion in*
616 *Psychiatry*, 28(5), 378–385. <https://doi.org/10.1097/YCO.0000000000000184>
- 617 Schottenbauer, M. A., Glass, C. R., Arnkoff, D. B., Tendick, V., & Gray, S. H. (2008).

- 618 Nonresponse and Dropout Rates in Outcome Studies on PTSD: Review and
619 Methodological Considerations. *Psychiatry: Interpersonal and Biological Processes*,
620 71(2), 134–168. <https://doi.org/10.1521/psyc.2008.71.2.134>
- 621 Schumm, J. A., Dickstein, B. D., Walter, K. H., Owens, G. P., & Chard, K. M. (2015).
622 Changes in posttraumatic cognitions predict changes in posttraumatic stress disorder
623 symptoms during cognitive processing therapy. *Journal of Consulting and Clinical*
624 *Psychology*, 83(6), 1161–1166. <https://doi.org/10.1037/ccp0000040>
- 625 Shakespeare-Finch, J., & Armstrong, D. (2010). Trauma type and posttrauma outcomes:
626 Differences between survivors of motor vehicle accidents, sexual assault, and
627 bereavement. *Journal of Loss and Trauma*, 15(2), 69–82.
628 <https://doi.org/10.1080/15325020903373151>
- 629 Startup, M., Makgekenene, L., & Webster, R. (2007). The role of self-blame for trauma as
630 assessed by the Posttraumatic Cognitions Inventory (PTCI): A self-protective
631 cognition? *Behaviour Research and Therapy*, 45(2), 395–403.
632 <https://doi.org/10.1016/j.brat.2006.02.003>
- 633 Stevens, J. (James P. (2009). *Applied multivariate statistics for the social sciences*.
634 Routledge.
- 635 Tabachnick, B. G., & Fidell, L. S. (2013). *Using multivariate statistics*. Pearson Education.
- 636 ter Heide, F. J. J., Sleijpen, M., & van der Aa, N. (2017). Posttraumatic world assumptions
637 among treatment-seeking refugees. *Transcultural Psychiatry*, 54(5–6), 824–839.
638 <https://doi.org/10.1177/1363461517741811>
- 639 van Buuren, S., & Groothuis-Oudshoorn, K. (2011). mice: Multivariate imputation by

- 640 chained equations in R. *Journal of Statistical Software*, 45(3), 1–67.
641 <https://doi.org/10.18637/jss.v045.i03>
- 642 Weathers, F. W., Bovin, M. J., Lee, D. J., Sloan, D. M., Schnurr, P. P., Kaloupek, D. G.,
643 Keane, T. M., & Marx, B. P. (2018). The clinician-administered ptsd scale for DSM-5
644 (CAPS-5): Development and initial psychometric evaluation in military veterans.
645 *Psychological Assessment*, 30(3), 383–395. <https://doi.org/10.1037/pas0000486>
- 646 Woud, M. L., Kleim, B., & Cwik, J. C. (2019). Editorial for the Special Issue on Negative
647 Appraisals in Trauma: Current Status and Future Directions for Research. *Cognitive*
648 *Therapy and Research*, 43(1), 139–142. <https://doi.org/10.1007/s10608-018-09992-5>
- 649