Momentary Interpersonal Processes of Suicidal Surges in Borderline Personality Disorder

Aleksandra Kaurin^a, Alexandre Y. Dombrovski^b, Michael N. Hallquist^c & Aidan G.C. Wright^a

This is an unedited manuscript accepted for publication at Psychological Medicine.

The manuscript will undergo further copyediting and typesetting.

Word count: 4480

^a Department of Psychology, University of Pittsburgh

^b Department of Psychiatry, University of Pittsburgh School of Medicine

^c Department of Psychology and Neuroscience, University of North Carolina, Chapel Hill

Correspondence concerning this article should be addressed to Aleksandra Kaurin, Department of Psychology, University of Pittsburgh, 3213 Sennott Square, 210 S. Bouquet St., Pittsburgh, PA, 15260. E-Mail: kaurinaleksandra@pitt.edu

Abstract

Background: Suicide rates are high in borderline personality disorder (BPD) where interpersonal problems trigger intense affective dysregulation and impulses to act on suicidal thoughts. To date, however, no study has examined how interpersonal stressors contribute to momentary within-person links among affect and impulsivity with suicidal ideation, and how those links vary over time in people's daily lives. **Methods:** 153 individuals diagnosed with BPD and 52 healthy controls completed a 21-day ecological momentary assessment protocol. Of these 153 BPD individuals with BPD, 105 had a history of suicide attempts. Multilevel structural equation modeling (MSEM) was used to examine dynamic links among interpersonal perceptions, affect, state impulsivity, and suicidal intent. **Results:** Aggregated across interactions, lower perceived warmth in others was associated with suicidal ideation. This direct relationship, however, did not extend to momentary within-person associations. Instead, interpersonal conflicts were linked to suicidal ideation indirectly via greater negative affect and lower positive affect. While a robust within-person link between interpersonal perceptions and impulsivity emerged, impulsivity did not account for the relationship between interpersonal perceptions and suicidal ideation. **Conclusion:** This intensive longitudinal study illustrates momentary interpersonal signatures of an emerging suicidal crisis. Among people with BPD at high risk for suicide, interpersonal triggers initiate a cascade of affective dysregulation, which in turn gives rise to suicidal ideation.

Key words: suicidal ideation; ecological momentary assessment; borderline personality disorder; multilevel structural equation modelling; affective dysregulation; impulsivity;

People diagnosed with borderline personality disorder (BPD) are at high risk of dying by suicide: almost all report chronic suicidal ideation, 84% of patients with BPD engage in suicidal behavior, 70% attempt suicide, with a mean of 3.4 lifetime attempts per individual, and 5–10% die by suicide (Black et al., 2004; McGirr et al., 2007; Soloff et al., 1994).

Although correlates of suicidal behavior have been studied extensively in the past 30 years (Turecki et al., 2019), prediction of suicide attempts remains close to chance (Franklin et al., 2017; Nock, 2009). One important reason for this lack of progress is that most studies have focused on between-person differences, or have employed long follow-up intervals, missing the temporal dynamics of suicide risk that require more frequent assessments.

Given the high rate and potential lethality of suicide attempts in BPD, we need a better understanding of proximal and potentially modifiable factors that predict short-term surges in ideation and catalyze suicidal crises (Galynker, 2017). Accordingly, we investigate how interpersonal triggers elicit momentary surges in suicidal ideation – either directly or indirectly via affective or impulsive processes – among people diagnosed with BPD and a history of suicide attempts.

Two key observations emerge from recent studies using ambulatory assessment of the real-time occurrence of suicidal ideation in daily life. First, suicidal ideation severity varies considerably from hour to hour (Ben-Zeev et al., 2012; Hallensleben et al., 2017; Husky et al., 2017; Witte et al., 2006), and covaries with well-known precipitants, which are limited in their ability to predict prospective changes in suicidal ideation (Kleiman et al., 2017; Victor et al., 2019). Second, episodes of suicidal ideation tend to be brief, with participants reporting most episodes to be shorter than an hour (Nock et al., 2009). This descriptive work aligns well with prior theory. The *Three Step-Theory*, for instance, posits that negative affective states motivate

suicidal desire contemporaneously or over seconds to minutes (Klonsky et al., 2017). More specifically, the theory suggests that pain, hopelessness, and suicidal desire are reciprocally influential within a situation, rather than unfolding over longer periods of time. Similarly, *Fluid Vulnerability Theory*'s concept of a suicidal mode suggests that suicidal crises represent a sudden and intense departure from a baseline state, resulting from complex interactions of mutually influential affective, cognitive, behavioral, and physiological processes (Rudd, 2006).

In BPD, interpersonal stressors often exacerbate emotional instability (Tragesser et al., 2008), yet it is unclear how interpersonal stressors contribute to suicidal crises as they emerge in daily life. On a momentary and daily timescale, interpersonal perceptions of hostility, disaffiliation, and rejection are linked to increases in negative affect (Kaurin et al., 2020; Lazarus et al., 2018; Sadikaj et al. 2013; Sadikaj et al., 2010), aversive tension (Stiglmayr et al., 2005), affective instability, severe anger (Miskewicz et al., 2015), and impulsivity (Coifman et al., 2012; Koenigsberg et al., 2001). Although previous studies have largely suggested that the amplification of affect in BPD is specific to negative affect, some evidence indicates that individuals with BPD also report smaller increases in positive affect in response to agreeable behavior of interaction partners (Sadikaj et al., 2010). Overall, this pattern of reactivity is particularly striking in comparison to healthy controls (Berenson et al., 2011) or depressed patients (Hepp et al., 2017).

Because impulsivity accelerates the transition from thoughts to action, it has long been conceptualized as a central risk factor for suicide attempts in general (Bryan & Rudd, 2006), and in BPD in particular (Brodsky et al., 2006; Soloff et al., 2000). Meta-analytic evidence, however, suggests that the predictive validity of impulsivity for suicide attempts is paradoxically rather modest (Anestis et al., 2014), with little discriminatory value for the differentiation of suicide

ideation and attempt (Klonsky & May 2010; May & Klonsky, 2016). However, some evidence derived from a 6-day ecological momentary assessment protocol suggests that impulsivity is related to the likelihood of suicidal behavior and particularly to within-person fluctuations in suicidal ideation, but not to the general tendency to think about suicide (Hadzic et al., 2020). More importantly, previous work has predominantly focused on self-reports of impulsiveness, which are likely relevant to determining long-term suicide risk. State-sensitive indices of impulsivity, in contrast, have the potential to reveal more proximal risk factors for suicidal ideation as it emerges in daily life (Liu et al., 2017).

In contrast, negative affectivity has been more consistently related to the emergence of suicidal ideation (Hallquist & Pilkonis, 2012; Linehan et al., 1993; McGirr et al., 2007; Soloff et al., 2005; Wedig et al., 2012; Yen et al., 2004; 2009). Links et al. (2007) found that in participants with BPD and a history of suicidal behavior, negative mood intensity was significantly related to the intensity of self-reported suicide ideation and to the number of suicidal behaviors during the past year. More recent work further suggests that the within-person link between negative affect and suicidal ideation is enhanced among patients with BPD compared to those without (Mou et al., 2018). These findings are consistent with the notion that suicidal behaviors reflect efforts to escape intense aversive arousal or distress in response to acute stressors (Gratz et al., 2006; Millner et al., 2019), and that the BPD-specific association among both is reinforced by reductions in negative affect (Kleiman et al., 2018; Selby et al., 2009).

Though negative emotions are clearly related to suicidal ideation, positive affect may have incremental predictive value. For instance, Yen et al. (2013) found that over a period of six months low positive affectivity was a stronger prospective predictor for suicidal acts than

negative affectivity. Similarly, Hirsch et al.'s (2007) cross-sectional study found that dispositional positive affectivity differentiated older primary care patients with increased levels of suicidal ideation from those without, independently of trait negative affect. In BPD, positive affect may be essential for research on interpersonal precipitants of suicidal ideation, because perceptions of others' warmth do not necessarily translate to the same amount of increases in positive affect in BPD as in community controls (Sadikaj et al., 2010).

Though a vast body of work has consistently reported momentary contingencies among interpersonal perceptions and affect or state impulsivity in BPD, no study has illustrated how those processes propel real-life suicidal crises. The evidence reviewed above suggests at least two different pathways through which suicidal ideation in BPD may emerge in daily interpersonal interactions. First, interpersonal perceptions of disaffiliation may be directly linked to increased momentary suicidal ideation (Brodsky et al., 2006). Alternatively, suicidal ideation may arise via a cascade of socio-affective processes, where the effect of interpersonal perceptions on momentary suicidal ideation is indirect, depending on enhanced affective reactivity (both positive and negative) or increased impulsivity in response to interpersonal stressors (Sadikaj et al., 2013; Selby et al., 2009).

We test these pathways from the viewpoint of transactional models of suicidal surges that emphasize within-person links among interpersonal perception, affect, impulsivity, and suicidal ideation. This approach can delineate momentary processes that potentially conduce to suicide. Serious suicidal thoughts are rare, episodic events, and will manifest infrequently during a 21-day EMA protocol, even in a high-risk population, which makes achieving adequate power particularly difficult. Therefore, our analyses are based on a case-control sample enriched for history of high-lethality suicide attempts, which are among the most powerful long-term

predictors of lethal suicidal behavior (Christiansen & Jensen, 2007; Gibb et al., 2005; Haw et al., 2007; Suominen et al., 2004). Thus, for the purpose of the current study, healthy controls, non-attempters, and attempters were pooled to represent a range of suicidal ideation severity consistent with dimensional conceptualizations of psychopathology (Stanton et al., 2020).

Methods

All study procedures were approved by the Institutional Review Boards of the University of Pittsburgh (STUDY19050210).

Sample. Participants were drawn from a longitudinal study (Soloff et al., 2017), and recruited from in- and outpatient clinics or the nearby community by advertisement. At enrollment, participants had to be between 18-45 years. Exclusion criteria included a lifetime diagnosis of any psychotic or bipolar disorder, clinical evidence of organic brain disease, physical disorders or treatments with known psychiatric consequence, and $IQ < 70$ measured by the WTAR (Wechsler, 2001).

The sample comprised 153 individuals diagnosed with BPD and 52 healthy control participants (M_{age} : 33.71, $SD=9.43$; 80% female). Of these 153 individuals with BPD, 105 had a history of non-zero lethality suicide attempts; 48 reported no past suicide attempts. To increase the reliability of our EMA measurements, participants with fewer than ten reported interactions ($N=19$) were excluded. This resulted in a final sample size of 186 participants. The majority of the sample identified as white/Caucasian (76%), followed by Black/African American (15%), Asian (4%), Pacific Islander (3%), or other/did not report racial demographics (2%).

BPD diagnoses were based on the ICD-10-based International Personality Disorder Examination (Loranger et al., 1994). Non-suicidal participants with BPD had no lifetime history

of suicide attempts, healthy control participants had no lifetime history of psychiatric disorders, nor suicide attempts, as determined by the SCID/DSM-IV.

Suicide attempters had a history of a self-injurious act with the intent to die within a one-month period prior to completing the study assessments or had a history of a past suicide attempt with strong current SI at the time of study enrollment. Attempters were required to have medically significant attempts, defined by a score of >1 on the Beck Lethality Scale (Beck et al., 1975). High-lethality attempts were defined by a score of >3 . For participants with multiple attempts, data for the highest-lethality attempt were used. High-lethality attempts resulted in coma, need for resuscitation, unstable vital signs, penetrating wounds of abdomen or chest, third-degree burns, or major bleeding.

Power estimations. Power calculations were based on Monte Carlo simulations of fully specified MSEM models with plausible values taken from previous 21-day studies with similar sampling schemes (Sadikaj et al., 2013; Lazarus et al., 2018). These simulations indicated that power would exceed .80 at an alpha level of .05 for each individual within-person effect with person-level sample sizes of $N > 50$, including tests of main effects and variance components.

Momentary Assessments. Participants completed a 21-day EMA protocol within pre-defined time windows, using the MetricWire smartphone application, which reminded them to complete surveys via push notifications. If participants indicated that an interaction occurred, they were asked to report on the behavior of one of their interaction partners along with features of the situation. This resulted in $N=9009$ reported interactions, with an average number of 48 interactions per participant, ranging from 10 to 116 observations overall. On 12% of days participants reported suicidal ideation at least once during the day, which corresponds to a total of 4% of observations that were characterized by suicidal ideation. Participants rated the

behavior of their interaction partner/s regarding dominance or warmth, on a sliding scale from - 50 to +50. The *dominance* scale ranged from “Accommodating/Submissive/Timid” to “Assertive/Dominant/Controlling” and the *warmth* scale ranged from “Cold/Distant/Hostile” to “Warm/Friendly/Caring.”

Participants also rated the degree to which they felt negative (i.e., nervous, sad, irritated, angry) or positive emotions (i.e., happy, content, excited) derived from the Positive and Negative Affect Schedule (Watson et al., 1988). Items read “How [ADJECTIVE] did you feel during the interaction?”, and ratings were made on a slider scale from 0 (“Not at All”) to 100 (“Extremely”) for each adjective, and an additional item asking about their impulsivity (i.e., “How would you describe your behavior during the interaction?, 0 (“In Control”) to 100 (“Impulsive”).

Suicidal ideation was assessed with two dichotomous items: “Since the interaction,” “Have you wished you were dead or wished you could go to sleep and not wake up?” and “Have you actually had any thoughts of killing yourself?”, derived from the *suicidal ideation subscale* of the Columbia-Suicide Severity Rating Scale (C-SSRS; Posner et al., 2008).

Data Analysis

Because we repeatedly sampled social interactions from participants, our data had a hierarchical structure, such that interactions (within-person level) were nested within individuals (between-person level). Therefore, we used multilevel structural equation modeling (MSEM), which accommodates this data structure. Variables can be between-person (e.g., gender), which only include variance at the between-person level, or within-person (e.g., interpersonal interaction variables), which include variance at both levels. The total variability in interaction variables are partitioned into the between-person variance, reflecting individual differences in average (or trait-level) responses, and within-person variance, reflecting moment-to-moment

fluctuations from an individual's average level. At each level, MSEM can be used to examine associations among variables (Sadikaj et al., 2019). The *between-person* portion of the model estimates associations among individual differences in each observed variable. For instance, the correlation or regression path between how suicidal a person tends to be with how much negative affect (NA) they typically tend to experience. In contrast, *within-person* associations reflect how strongly variables are coupled together as they fluctuate from interaction to interaction. Thus, within-person associations reflect *dynamic processes* (e.g., how much NA is arises at time of an interpersonal stressor). MSEM also allows for the estimation of random slopes (i.e., slopes that vary across individuals), such that individuals can differ in the strength of their within-person associations among variables that are repeatedly assessed in the EMA protocol. That is, some individuals may have a strong within-person link among variables, but others may have a weaker one, or one of an opposite sign. The fixed effects of these slopes represent the average association in the sample, and at the random effects represent individual differences in the extent to which those situational features co-occur across participants.

Importantly, the within-person coefficients reported here represent contemporaneous associations (i.e., within the same wave), although the suicidality items were referenced to "since the interaction." This modelling decision was based on theoretical and empirical work suggesting that, when the variables assessed fluctuate substantially over the frequency in which they are being monitored, prioritizing lagged associations is not more valuable (Granger, 1969).

Figure 1 provides a schematic overview of the estimated models, along with path annotations as referenced in the results section. Three sets of models were estimated with perceptions of other behavior (i.e., warmth/dominance) predicting suicidal ideation. Model 1 regressed momentary suicidal ideation on perceptions of interaction partner's behavior at the

within-person level (i.e., fluctuations in momentary interpersonal perceptions predicting fluctuations in momentary suicidal ideation) and the between-person level (i.e., individual differences in average perceptions and suicidal ideation). Model 2 introduced negative or positive affect or impulsivity as additional predictors of momentary suicidal ideation at each level, each variable was also regressed on interpersonal perceptions. Finally, Model 3 simultaneously introduced two of the three intermediate variables (i.e., negative and positive affect, impulsivity) as correlated predictors/covariates of momentary suicidal ideation at each level, which permitted the evaluation of unique associations of each predictor with suicidal ideation.

Table 1 summarizes pooled within-person correlations among the variables along with correlations among the random intercepts at the between-person level. Sex (0=female; 1=male) and age (centered on mean age) were also included as covariates in all models at the between-person level. Within-person variables were adjusted for observation number (time centered on mean of observations) and whether the interaction occurred on a weekend to account for possible changes over time and weekly cycles. Along with other parameters not reported in the tables (e.g., residual variances, coefficients for covariates) full specifications and detailed output from all models can be found online at <https://osf.io/zpc3u/>. All models were estimated in Mplus (version 8.4; Muthén & Muthén, 1998-2019). Bayesian estimation was used because it provides a latent decomposition into within- and between-person variance for both the predictors and outcome variables when random effects are specified. Significance for all model parameters was based on 95% Credibility Intervals (CIs), with CIs that excluded zero indicative of a parameter that differed significantly from zero.

Results

Model Set 1

For models examining the effects of others' perceived warmth on suicidal ideation (Model 1), we found no significant fixed effect at the within-person level ($c = .187$, CI $[-.017, .404]$), but did at the between-person level ($\beta_{SI,W} = -.184$, CI $[-.357, .000]$). Similarly, perceiving interaction partners as more dominant was not associated with suicidal ideation at the within-person ($c = -.014$, CI $[-.167, .145]$), but was at the between-person level ($\beta_{SI,D} = .185$, CI $[.029, .331]$). Note that for these and all subsequently reported within-person paths, we found significant random effects indicative of individual differences in these associations.

Model Set 2

Next, momentary affect and impulsivity were added as additional intervening predictors of suicidal ideation to our model. Table 2 provides a detailed overview of estimates for models including perceptions of interpersonal warmth, and Table 3 for estimates based on models including perceptions of dominance.

Negative Affect. At the within-person level, the fixed effect of perceptions of others' *warmth* negatively predicted negative affect ($a-NA$), and negative affect predicted suicidal ideation ($b-NA$), while warmth now predicted SI (c ; Table 2A). The indirect effect (ab) was also significant, suggesting that negative affect accounted for the link between perceived coldness and momentary suicidal ideation. At the between-person level, lower average perceptions of warmth were associated with experiencing more negative affect on average, and higher average levels of negative affect were associated with an increased likelihood for suicidal ideation; the link between perceptions of warmth and suicidal ideation, however, was no longer significant, suggesting individual differences in negative affect accounted for this association.

Relationships with perceived *dominance* were similar but somewhat weaker (Table 3A): interpersonal perceptions predicted negative affect (*a-NA*), and negative affect predicted suicidal ideation (*b-NA*). Although the total of the *c* path was not significant, an indirect effect (*ab*) emerged, suggesting that negative affect accounted for the link between perceived dominance and momentary suicidal ideation. The same pattern emerged at the between-person level.

Positive Affect. A very similar pattern with opposing signs emerged for the model including positive affect (Table 2B), such that at the within-person level interpersonal *warmth* was positively associated with positive affect (*a-PA*), and positive affect negatively with suicidal ideation (*b-PA*). Again, the indirect effect (*ab*) was significant, while the *c* path did not reach significance. The same pattern emerged at the between-person level.

Models including perceived *dominance* (Table 3B) revealed a similar pattern of results, with the only exceptions being an expectedly negative association with positive affect (*a-PA*), and no between-person level association between perceived dominance and positive affect.

Impulsivity. For models that included impulsivity (Table 2C), we found a significant negative within-person link between perceptions of *warmth* and impulsivity (*a-Imp*), while the effects for the *b-Imp* path and the *c* path were not significant. At the between-person level, lower average perceptions of warmth were associated with feeling more impulsive on average, and that higher average levels of impulsivity were associated with an increased likelihood for suicidal ideation.

In models where impulsivity was included to account for the link between *dominance* and suicidal ideation (Table 3C), only a significant link between interpersonal perceptions and impulsivity emerged (*a-Imp*). At the between-person level, however, higher average perceptions

of dominance were associated with experiencing more impulsivity, and higher average levels of impulsivity were associated with an increased likelihood for suicidal ideation.

Model Set 3

Finally, to test the robustness of our indirect effects, we added negative affect and impulsivity (and negative with positive affect, respectively) simultaneously as covarying predictors of suicidal ideation to our models. Table 4 provides estimates for models including negative affect and impulsivity, and Table 5 summarizes estimates for models including negative and positive affect.

Negative Affect and Impulsivity. At the within-person level, perceptions of others' warmth negatively predicted negative affect ($a-NA$), and impulsivity ($a-Imp$). Moreover, negative affect ($b-NA$), but not impulsivity ($b-Imp$), was related to suicidal ideation. The path between perceived warmth and suicidal ideation, however, was not significant (c). At the between person level, the same pattern of results emerged, and impulsivity and negative affect were positively associated with each other (Table 4A).

Perceptions of others' dominance were positively related to negative affect ($a-NA$), and impulsivity ($a-Imp$) at the within-person level (Table 4B). Again, negative affect ($b-NA$), but not impulsivity ($b-Imp$), was related to suicidal ideation, while the path between perceived dominance and suicidal ideation was not significant (c). At the between-person level, the same pattern of results emerged, and impulsivity and negative affect were positively associated with each other.

Negative and Positive Affect. Finally, when both positive and negative affect were simultaneously added as predictors of suicidal ideation (Table 5A), we found that perceptions of others' warmth negatively predicted negative affect ($a-NA$), and positively positive affect ($a-PA$).

Both were uniquely related to suicidal ideation (*b-NA*; *b-PA*), albeit in different directions, and positive and negative affect were negatively associated with each other. The path between perceived warmth and suicidal ideation, was not significant. At the between person level, the same pattern of results emerged. For the *a-NA* and *a-PA* paths, a converse pattern emerged for our model including perceptions of *dominance* (Table 5B).

Discussion

We administered intensive longitudinal assessments to BPD patients, many of whom had a history of high-lethality suicide attempts, over several weeks to evaluate relationships between interpersonally stressful encounters, and suicidal surges (i.e., within-person links of affective dysregulation and suicidal ideation) in the moment-to-moment stream of individuals' daily lives. When aggregated across all interactions, an association between perceived coldness and suicidal ideation emerged, supporting the notion that interpersonal experiences are meaningfully related to suicidal ideation in BPD (Brodsky et al., 2006; Brown et al., 2002). However, this between-person association did not extend to the within-person momentary timescale. Instead, the triggering effect of interpersonal conflicts was indirect, such that greater NA or lower PA in the context of perceived conflict or withdrawal was associated with suicidal ideation. The present study suggests that in people diagnosed with BPD at high risk for suicide, situational interpersonal triggers do not directly elicit suicidal thoughts. Instead, the effect of negative interpersonal experiences on suicidal ideation unfolds indirectly via affective dysregulation.

Moreover, positive and negative affect each uniquely accounted for the link between interpersonal stressors and suicidal ideation, even after adjusting for their covariation, both at the within- and between-person level. This pattern of incremental effects supports theoretical assumptions that affective manifestations of BPD encompass positive as well as negative mood

states (Linehan et al., 1993; Russel et al., 2007). It also underlines the notion that suicidal ideation in BPD is driven not only by distress (NA) giving rise to escape motivations (Millner et al., 2019), but also by low PA, which may undermine a persons' ability to find alternative solutions or put the current crisis in perspective (Baumeister, 1990).

Ratings of impulsivity were unassociated with ideation within-person, and as such they did not explain the indirect effect of negative affect. However, impulsivity did exhibit a momentary relationship with interpersonal perception, consistent with previous work (Coifman et al., 2012). Average impulsivity was associated with average suicidal ideation, but not after adjusting for negative affect consistent with a vast body of work indicating that maladaptive interpersonal behaviors and impulsive coping are related to affective instability among individuals with BPD (Bradley et al., 2007) though longitudinally affective instability is a stronger predictor of suicidal behavior than impulsivity (Yen et al., 2004). Although widely in line with previous work, we note that characteristics of our sample (e.g., mean age) may have attenuated the effect, because impulsive suicidal acts become less common as BPD patients transition into midlife (Wedig et al., 2012; Zanarini et al., 2016).

Additional considerations of our study include that, the direct link between interpersonal perceptions and suicidal ideation was not significant at the within-person level, though the indirect effect through affect was. Although this circumstance has traditionally been discussed to preclude indirect-effect models, more recent writings note that inferences on intervening relationships are justified if the indirect effect carried by the $X \rightarrow M$ and $M \rightarrow Y$ paths is significant (MacKinnon et al., 2007), regardless of whether the bivariate association between X and Y was significant.

Additionally, our study primarily included people diagnosed with BPD, and follow-up

studies replicating our findings of the socio-affective dysregulation-suicidal ideation nexus in clinically more diverse samples are needed to corroborate our findings. Yet, BPD is a particularly informative population for the study of suicidal ideation, because it reflects a confluence of internalizing and externalizing psychopathology (Eaton et al., 2011; Crowell et al., 2009; James & Taylor, 2008): Diagnostic criteria such as affective instability relate more strongly to internalizing, while others, such as marked impulsivity, relate more strongly to externalizing forms of psychopathology. Such heterogeneity may imply generalizability to other clinical disorders.

A key question arising from our analyses is whether suicidal surges are a cause, in and of themselves, of suicide attempts. High negative and low positive affect could provide the emotional substrate for suicidal ideation as a proximal risk factor, potentially informing treatment. Alternatively, affective dysregulation may represent a reliable correlate, but not a causal component of suicidal ideation. It could be, for example, that both suicidal ideation and negative affect share risks. For instance, a growing literature suggests that Pavlovian escape biases (from negative affect) invigorate suicidal behavior (Millner et al., 2019) while disrupted decision processes undermine the consideration of deterrents and alternatives (Dombrovski et al., 2013; Brown et al., 2020). Future computational studies could help disentangle the role of affective states from dispositional deficits.

Future intensive longitudinal research on suicidal processes should also consider both the assessment schedule and statistical modeling alternatives. Suicidal ideation is infrequent and episodic, severely complicating its assessment. Although EMA-based sampling strategies seem uniquely capable of capturing meaningful variation in suicidal thought and behavior (Gratch et al., 2020), how best to balance intensity and duration of assessment to capture this meaningful

variation is challenging. Relatedly, a recurring debate is whether lagged associations should be prioritized over contemporaneous associations. Although contemporaneous and cross-sectional associations are often treated as the same, in intensive longitudinal data contemporaneous associations imply dynamic processes, which refer to the systematic covariation of variables as they fluctuate from measurement occasion to occasion. Contemporaneous associations capture processes that are assumed to be shorter than the time between assessments, while lagged associations provide a statistical model for processes that are likely to be longer than the sampling frequency (Granger, 1969). If the process assessed is faster than the assessment interval, prioritizing lagged associations may not be inherently more valuable, particularly when measurements were not evenly spaced in time (e.g., when assessments are randomized throughout the day). Thus, the choice of an assessment and data analytic approach requires theoretical justification, and should be based on estimates of how long episodes of suicidal thinking may last. Research has just begun to estimate the true timescale of an emergent suicidal crisis, and future studies should, therefore, systematically examine theoretically informed real-time sampling and modeling strategies of suicidal ideation.

This is the first reported study using EMA to characterize differential associations among affect and impulsivity with suicidal ideation, contextualized within stressful interpersonal interactions. Our study uncovers how momentary links among socio-affective processes and suicidal thoughts may drive the progression of suicidal crises in daily life: In people diagnosed with BPD at high risk for suicide, situational triggers do not directly elicit suicidal thoughts, but unfold indirectly via affective dysregulation.

References

- Anestis, M. D., Soberay, K. A., Gutierrez, P. M., Hernández, T. D., & Joiner, T. E. (2014). Reconsidering the link between impulsivity and suicidal behavior. *Personality and Social Psychology Review*, 18(4), 366-386.
- Baumeister, R. F. (1990). Suicide as escape from self. *Psychological Review*, 97(1), 90–113.
- Beck, A. T., Beck, R., & Kovacs, M. (1975). Classification of suicidal behaviors: I. Quantifying intent and medical lethality. *The American Journal of Psychiatry*, 132(3), 285–287.
- Ben-Zeev, D., Young, M. A., & Depp, C. A. (2012). Real-time predictors of suicidal ideation: mobile assessment of hospitalized depressed patients. *Psychiatry Research*, 197(1-2), 55-59.
- Berenson, K. R., Downey, G., Rafaeli, E., Coifman, K. G., & Paquin, N. L. (2011). The rejection–rage contingency in borderline personality disorder. *Journal of Abnormal Psychology*, 120(3), 681–690.
- Black, D. W., Blum, N., Pfohl, B., & Hale, N. (2004). Suicidal behavior in borderline personality disorder: prevalence, risk factors, prediction, and prevention. *Journal of Personality Disorders*, 18(3), 226-239.
- Bradley, R., Conklin, C. Z., & Westen, D. (2007). Borderline personality disorder. In W. O'Donohue, K. A. Fowler, & S. O. Lilienfeld (Eds.), *Personality disorders: Toward the DSM-V* (p. 167–201). Sage Publications, Inc.
- Brodsky, B. S., Groves, S. A., Oquendo, M. A., Mann, J. J., & Stanley, B. (2006). Interpersonal precipitants and suicide attempts in borderline personality disorder. *Suicide and Life-Threatening Behavior*, 36(3), 313-322.
- Brown, M. Z., Comtois, K. A., & Linehan, M. M. (2002). Reasons for suicide attempts and

- 432 nonsuicidal self-injury in women with borderline personality disorder. *Journal of*
433 *Abnormal Psychology*, 111(1), 198–202.
- 434 Brown, V. M., Wilson, J., Hallquist, M. N., Szanto, K., & Dombrovski, A. Y. (2020).
435 Ventromedial prefrontal value signals and functional connectivity during decision-
436 making in suicidal behavior and impulsivity. *Neuropsychopharmacology*, 45(6), 1034-
437 1041.
- 438 Bryan, C. J., & Rudd, M. D. (2006). Advances in the assessment of suicide risk. *Journal of*
439 *Clinical Psychology*, 62(2), 185-200.
- 440 Christiansen, E., & Frank Jensen, B. (2007). Risk of repetition of suicide attempt, suicide or all
441 deaths after an episode of attempted suicide: a register-based survival analysis. *Australian*
442 *and New Zealand Journal of Psychiatry*, 41(3), 257-265.
- 443 Coifman, K. G., Berenson, K. R., Rafaeli, E., & Downey, G. (2012). From negative to positive
444 and back again: Polarized affective and relational experience in borderline personality
445 disorder. *Journal of Abnormal Psychology*, 121(3), 668.
- 446 Crowell, S. E., Beauchaine, T. P., & Linehan, M. M. (2009). A biosocial developmental model
447 of borderline personality: Elaborating and extending linehan's theory. *Psychological*
448 *Bulletin*, 135(3), 495–510.
- 449 Dombrovski, A. Y., Szanto, K., Clark, L., Reynolds, C. F., & Siegle, G. J. (2013). Reward
450 signals, attempted suicide, and impulsivity in late-life depression. *JAMA Psychiatry*,
451 70(10), 1020-1030.
- 452 Eaton, N. R., Krueger, R. F., Keyes, K. M., Skodol, A. E., Markon, K. E., Grant, B. F., & Hasin,

- 453 D. S. (2011). Borderline personality disorder co-morbidity: relationship to the
454 internalizing–externalizing structure of common mental disorders. *Psychological*
455 *Medicine*, 41(5), 1041-1050.
- 456 Franklin, J. C., Ribeiro, J. D., Fox, K. R., Bentley, K. H., Kleiman, E. M., Huang, X.,
457 Musacchio, K. M., Jaroszewski, A. C., Chang, B. P., & Nock, M. K. (2017). Risk factors
458 for suicidal thoughts and behaviors: A meta-analysis of 50 years of research.
459 *Psychological Bulletin*, 143(2), 187–232.
- 460 Galynker, I., Yaseen, Z. S., Cohen, A., Benhamou, O., Hawes, M., & Briggs, J. (2017).
461 Prediction of suicidal behavior in high risk psychiatric patients using an assessment of
462 acute suicidal state: The suicide crisis inventory. *Depression and anxiety*, 34(2), 147-158.
- 463 Gibb, S. J., Beautrais, A. L., & Fergusson, D. M. (2005). Mortality and further suicidal
464 behaviour after an index suicide attempt: a 10-year study. *Australian & New Zealand*
465 *Journal of Psychiatry*, 39(1-2), 95-100.
- 466 Granger, C. W. (1969). Investigating causal relations by econometric models and cross-spectral
467 methods. *Econometrica: Journal of the Econometric Society*, 424-438.
- 468 Gratch, I., Choo, T. H., Galfalvy, H., Keilp, J. G., Itzhaky, L., Mann, J. J., ... & Stanley, B.
469 (2020). Detecting suicidal thoughts: The power of ecological momentary assessment.
470 *Depression and Anxiety*, 1-9.
- 471 Gratz, K. L., Rosenthal, M. Z., Tull, M. T., Lejuez, C. W., & Gunderson, J. G. (2006). An
472 experimental investigation of emotion dysregulation in borderline personality disorder.
473 *Journal of Abnormal Psychology*, 115(4), 850-855.
- 474 Hadzic, A., Spangenberg, L., Hallensleben, N., Forkmann, T., Rath, D., Strauß, M., ... &

- 475 Glaesmer, H. (2020). The association of trait impulsivity and suicidal ideation and its
476 fluctuation in the context of the Interpersonal Theory of Suicide. *Comprehensive*
477 *Psychiatry*, 98, 152158.
- 478 Hallensleben, N., Glaesmer, H., Forkmann, T., Rath, D., Strauss, M., Kersting, A., &
479 Spangenberg, L. (2019). Predicting suicidal ideation by interpersonal variables,
480 hopelessness and depression in real-time. An ecological momentary assessment study in
481 psychiatric inpatients with depression. *European Psychiatry*, 56(1), 43-50.
- 482 Hallquist, M. N., & Pilkonis, P. A. (2012). Refining the phenotype of borderline personality
483 disorder: Diagnostic criteria and beyond. *Personality Disorders: Theory, Research, and*
484 *Treatment*, 3(3), 228–246.
- 485 Haw, C., Bergen, H., Casey, D., & Hawton, K. (2007). Repetition of deliberate self-harm: a
486 study of the characteristics and subsequent deaths in patients presenting to a general
487 hospital according to extent of repetition. *Suicide and Life-Threatening Behavior*, 37(4),
488 379-396.
- 489 Hepp, J., Lane, S. P., Wycoff, A. M., Carpenter, R. W., & Trull, T. J. (2018). Interpersonal
490 stressors and negative affect in individuals with borderline personality disorder and
491 community adults in daily life: A replication and extension. *Journal of Abnormal*
492 *Psychology*, 127(2), 183–189.
- 493 Hirsch, J. K., Duberstein, P. R., Chapman, B., & Lyness, J. M. (2007). Positive affect and suicide
494 ideation in older adult primary care patients. *Psychology and Aging*, 22(2), 380–385.
- 495 Husky, M., Swendsen, J., Ionita, A., Jaussent, I., Genty, C., & Courtet, P. (2017). Predictors of
496 daily life suicidal ideation in adults recently discharged after a serious suicide attempt: a
497 pilot study. *Psychiatry research*, 256, 79-84.

- 498 James, L. M., & Taylor, J. (2008). Associations between symptoms of borderline personality
499 disorder, externalizing disorders, and suicide-related behaviors. *Journal of*
500 *Psychopathology and Behavioral Assessment*, 30(1), 1-9.
- 501 Kaurin, A., Wright, A. G., Pilkonis, P. A., Scott, L., Woods, W. C., Stepp, S., & Beeney, J. E.
502 (Manuscript Submitted for Publication). Attachment and Borderline Personality
503 Disorder: Differential Effects on Daily Socio-Affective Processes.
504 <https://doi.org/10.31234/osf.io/fncjg>
- 505 Kleiman, E. M., Turner, B. J., Fedor, S., Beale, E. E., Huffman, J. C., & Nock, M. K. (2017).
506 Examination of real-time fluctuations in suicidal ideation and its risk factors: Results
507 from two ecological momentary assessment studies. *Journal of Abnormal Psychology*,
508 126(6), 726.
- 509 Kleiman, E. M., Coppersmith, D. D., Millner, A. J., Franz, P. J., Fox, K. R., & Nock, M. K.
510 (2018). Are suicidal thoughts reinforcing? A preliminary real-time monitoring study on
511 the potential affect regulation function of suicidal thinking. *Journal of Affective*
512 *Disorders*, 232, 122-126.
- 513 Klonsky, E. D., Saffer, B. Y., & Bryan, C. J. (2018). Ideation-to-action theories of suicide: a
514 conceptual and empirical update. *Current Opinion in Psychology*, 22, 38-43.
- 515 Klonsky, E. D., & May, A. (2010). Rethinking impulsivity in suicide. *Suicide and Life-*
516 *Threatening Behavior*, 40(6), 612-619.
- 517 Koenigsberg, H. W., Harvey, P. D., Mitropoulou, V., New, A. S., Goodman, M., Silverman, J.,
518 ... & Siever, L. J. (2001). Are the interpersonal and identity disturbances in the borderline
519 personality disorder criteria linked to the traits of affective instability and impulsivity?.
520 *Journal of Personality Disorders*, 15(4), 358-370.

- 521 Lazarus, S., Scott, L., Beeney, J., Wright, A.G.C., Stepp, S., & Pilkonis, P. (2018). Borderline
522 Personality Disorder Symptoms and Affective Responding to Perceptions of Rejection
523 and Acceptance From Romantic Versus Nonromantic Partners. *Personality Disorders:
524 Theory, Research, and Treatment*, 9(3), 197-206.
- 525 Linehan, M. M., Heard, H. L., & Armstrong, H. E. (1993). Naturalistic follow-up of a behavioral
526 treatment for chronically parasuicidal borderline patients. *Archives of General
527 Psychiatry*, 50(12), 971-974.
- 528 Links, P. S., Eynan, R., Heisel, M. J., Barr, A., Korzekwa, M., McMain, S., & Ball, J. S. (2007).
529 Affective instability and suicidal ideation and behavior in patients with borderline
530 personality disorder. *Journal of Personality Disorders*, 21(1), 72-86.
- 531 Liu, R. T., Trout, Z. M., Hernandez, E. M., Cheek, S. M., & Gerlus, N. (2017). A behavioral and
532 cognitive neuroscience perspective on impulsivity, suicide, and non-suicidal self-injury:
533 Meta-analysis and recommendations for future research. *Neuroscience & Biobehavioral
534 Reviews*, 83, 440-450.
- 535 Loranger, A. W., Sartorius, N., Andreoli, A., Berger, P., Buchheim, P., Channabasavanna, S. M.,
536 ... & Jacobsberg, L. B. (1994). The international personality disorder examination: The
537 World Health Organization/Alcohol, Drug Abuse, and Mental Health Administration
538 international pilot study of personality disorders. *Archives of General Psychiatry*, 51(3),
539 215-224.
- 540 MacKinnon, D. P., Fairchild, A. J., & Fritz, M. S. (2007). Mediation analysis. *Annual Review of
541 Psychology*, 58, 593-614.
- 542 May, A. M., & Klonsky, E. D. (2016). "Impulsive" suicide attempts: What do we really mean?
543 *Personality Disorders: Theory, Research, and Treatment*, 7(3), 293-302.

- 544 McGirr, A., Paris, J., Lesage, A., Renaud, J., & Turecki, G. (2007). Risk factors for suicide
545 completion in borderline personality disorder: a case-control study of cluster B
546 comorbidity and impulsive aggression. *The Journal of Clinical Psychiatry*.
- 547 Millner, A. J., den Ouden, H. E. M., Gershman, S. J., Glenn, C. R., Kearns, J. C., Bornstein, A.
548 M., Marx, B. P., Keane, T. M., & Nock, M. K. (2019). Suicidal thoughts and behaviors
549 are associated with an increased decision-making bias for active responses to escape
550 aversive states. *Journal of Abnormal Psychology*, 128(2), 106–118.
- 551 Miskewicz, K., Fleson, W., Arnold, E. M., Law, M. K., Mneimne, M., & Furr, R. M. (2015). A
552 contingency-oriented approach to understanding borderline personality disorder:
553 Situational triggers and symptoms. *Journal of Personality Disorders*, 29(4), 486-502.
- 554 Mou, D., Kleiman, E. M., Fedor, S., Beck, S., Huffman, J. C., & Nock, M. K. (2018). Negative
555 affect is more strongly associated with suicidal thinking among suicidal patients with
556 borderline personality disorder than those without. *Journal of Psychiatric Research*, 104,
557 198-201.
- 558 Muthén, L., & Muthén, B. (2019). Mplus User's Guide (1998–2019). *Muthén & Muthén: Los*
559 *Angeles, CA*.
- 560 Nock, M. K., Hwang, I., Sampson, N., Kessler, R. C., Angermeyer, M., Beautrais, A., ... & De
561 Graaf, R. (2009). Cross-national analysis of the associations among mental disorders and
562 suicidal behavior: findings from the WHO World Mental Health Surveys. *PLoS*
563 *Medicine*, 6(8).
- 564 Nock, M. K., Prinstein, M. J., & Sterba, S. K. (2009). Revealing the form and function of self-
565 injurious thoughts and behaviors: A real-time ecological assessment study among
566 adolescents and young adults. *Journal of Abnormal Psychology*, 118(4), 816–827.

- 567 Posner, K., Brent, D., Lucas, C., Gould, M., Stanley, B., Brown, G., ... & Mann, J. (2008).
568 *Columbia-suicide severity rating scale (C-SSRS)*. New York, NY: Columbia University
569 Medical Center, 2008.
- 570 Rudd, M. D. (2006). *Fluid vulnerability theory: A cognitive approach to understanding the*
571 *process of acute and chronic risk*. In T. E. Ellis (Ed.), *Cognition and suicide: Theory,*
572 *research, and therapy* (pp. 355–368). Washington, DC: American Psychological
573 Association.
- 574 Russell, J. J., Moskowitz, D. S., Zuroff, D. C., Sookman, D., & Paris, J. (2007). Stability and
575 variability of affective experience and interpersonal behavior in borderline personality
576 disorder. *Journal of Abnormal Psychology, 116*(3), 578.
- 577 Sadikaj, G., Wright, A. G., Dunkley, D., Zuroff, D., & Moskowitz, D. (2019, October 8).
578 Multilevel Structural Equation Modeling for Intensive Longitudinal Data: A Practical
579 Guide for Personality Researchers. <https://doi.org/10.31234/osf.io/hwj9r>
- 580 Sadikaj, G., Moskowitz, D. S., Russell, J. J., Zuroff, D. C., & Paris, J. (2013). Quarrelsome
581 behavior in borderline personality disorder: Influence of behavioral and affective
582 reactivity to perceptions of others. *Journal of Abnormal Psychology, 122*(1), 195-207.
- 583 Sadikaj, G., Russell, J. J., Moskowitz, D. S., & Paris, J. (2010). Affect dysregulation in
584 individuals with borderline personality disorder: Persistence and interpersonal triggers.
585 *Journal of Personality Assessment, 92*(6), 490-500.
- 586 Selby, E. A., Anestis, M. D., Bender, T. W., & Joiner, T. E., Jr. (2009). An exploration of the
587 emotional cascade model in borderline personality disorder. *Journal of Abnormal*
588 *Psychology, 118*(2), 375–387.
- 589 Soloff, P. H., & Chiappetta, L. (2017). Suicidal behavior and psychosocial outcome in borderline

- 590 personality disorder at 8-year follow-up. *Journal of Personality Disorders*, 31(6), 774-
591 789.
- 592 Soloff, P. H., Fabio, A., Kelly, T. M., Malone, K. M., & Mann, J. J. (2005). High-lethality status
593 in patients with borderline personality disorder. *Journal of Personality Disorders*, 19(4),
594 386-399.
- 595 Soloff, P. H., Lis, J. A., Kelly, T., Cornelius, J. R., & Ulrich, R. (1994). Risk factors for suicidal
596 behavior in borderline personality disorder. *The American Journal of Psychiatry*, 151(9),
597 1316–1323.
- 598 Soloff, P. H., Lynch, K. G., Kelly, T. M., Malone, K. M., & Mann, J. J. (2000). Characteristics of
599 suicide attempts of patients with major depressive episode and borderline personality
600 disorder: a comparative study. *American Journal of Psychiatry*, 157(4), 601-608.
- 601 Stanton, K., McDonnell, C. G., Hayden, E. P., & Watson, D. (2020). Transdiagnostic approaches
602 to psychopathology measurement: Recommendations for measure selection, data
603 analysis, and participant recruitment. *Journal of Abnormal Psychology*, 129(1), 21–28.
- 604 Stiglmayr, C. E., Grathwol, T., Linehan, M. M., Ihorst, G., Fahrenberg, J., & Bohus, M. (2005).
605 Aversive tension in patients with borderline personality disorder: A computer-based
606 controlled field study. *Acta Psychiatrica Scandinavica*, 111(5), 372-379.
- 607 Suominen, K., Isometsä, E., Suokas, J., Haukka, J., Achte, K., & Lönnqvist, J. (2004).
608 Completed suicide after a suicide attempt: a 37-year follow-up study. *American Journal*
609 *of Psychiatry*, 161(3), 562-563.
- 610 Tragesser, S. L., Lippman, L. G., Trull, T. J., & Barrett, K. C. (2008). Borderline personality
611 disorder features and cognitive, emotional, and predicted behavioral reactions to teasing.
612 *Journal of Research in Personality*, 42(6), 1512-1523.

- 613 Turecki, G., Brent, D. A., Gunnell, D., O'Connor, R. C., Oquendo, M. A., Pirkis, J., & Stanley,
614 B. H. (2019). Suicide and suicide risk. *Nature Reviews Disease Primers*, 5(1), 1-22.
- 615 Victor, S. E., Scott, L. N., Stepp, S. D., & Goldstein, T. R. (2019). I Want You to Want Me:
616 Interpersonal Stress and Affective Experiences as Within-Person Predictors of
617 Nonsuicidal Self-Injury and Suicide Urges in Daily Life. *Suicide and Life-Threatening*
618 *Behavior*, 49(4), 1157-1177.
- 619 Watson, D., Clark, L. A., & Tellegen, A. (1988). Development and validation of brief measures
620 of positive and negative affect: the PANAS scales. *Journal of Personality and Social*
621 *Psychology*, 54(6), 1063-1070.
- 622 Wechsler, D. (2001). *Wechsler Test of Adult Reading (WTAR)*. San Antonio, TX: The
623 Psychological Corporation.
- 624 Wedig, M. M., Silverman, M. H., Frankenburg, F. R., Reich, D. B., Fitzmaurice, G., & Zanarini,
625 M. C. (2012). Predictors of suicide attempts in patients with borderline personality
626 disorder over 16 years of prospective follow-up. *Psychological medicine*, 42(11), 2395-
627 2404.
- 628 Yen, S., Shea, M. T., Sanislow, C. A., Grilo, C. M., Skodol, A. E., Gunderson, J. G., ... & Morey,
629 L. C. (2004). Borderline personality disorder criteria associated with prospectively
630 observed suicidal behavior. *American Journal of Psychiatry*, 161(7), 1296-1298.
- 631 Yen S, Shea MT, Sanislow CA, Skodol AE, Grilo CM, Edelen MO, et al. Personality traits as
632 prospective predictors of suicide attempts. (2009). *Acta Psychiatrica Scandinavica*,
633 120(3):222–229.
- 634 Yen, S., Weinstock, L. M., Andover, M. S., Sheets, E. S., Selby, E. A., & Spirito, A. (2013).

- 635 Prospective predictors of adolescent suicidality: 6-month post-hospitalization follow-up.
636 *Psychological Medicine*, 43(5), 983-993.
- 637 Zanarini, M. C., Frankenburg, F. R., Reich, D. B., & Fitzmaurice, G. M. (2016). Fluidity of the
638 subsyndromal phenomenology of borderline personality disorder over 16 years of
639 prospective follow-up. *American Journal of Psychiatry*, 173(7), 688-694.
- 640

641 **Acknowledgements**

642 We are grateful to the patients who took part in this study and to the clinical and research
643 teams who made this work possible.

644 **Financial Support**

645 The study was supported by the National Institute of Mental Health (R01 MH048463).

646 **Conflict of Interest**

647 The authors certify that they have no affiliations with or involvement in any organization
648 or entity with any financial interest, or non-financial interest in the subject matter or materials
649 discussed in this manuscript.

650 **Ethical Standards**

651 The authors assert that all procedures contributing to this work comply with the ethical
652 standards of the relevant national and institutional committees on human experimentation and
653 with the Helsinki Declaration of 1975, as revised in 2008.

Figure 1. Overview of model sets used for all analyses of suicidal surges including decomposition of observed momentary variables into between- (subscript i) and within-person (subscript t) variance. Note, not all parameters are diagrammed (e.g., residual variances are not depicted). In Model Set 3, at the within-person level, a process was modelled in which perceived warmth or dominance was associated with negative affect and impulsivity *or* negative and positive affect. In these models, all three variables were independently associated with momentary suicidal ideation, and were allowed to covary. All models controlled for age and gender at the between-person level, and for time and week day at the within-person level. Coefficients for covariates are not presented for parsimony. Single headed arrows indicate regression paths. Filled dots represent random effects. Perc = interpersonal perceptions of warmth or dominance; NA = negative affect; PA = positive affect; Imp = impulsivity; SI = suicidal ideation.

Table 1. Correlations among study variables at within- and between-person levels.

	SI	warmth	dominance	impulsivity	NA	PA
within-person						
suicidal ideation	1					
warmth	-.14	1				
dominance	.06	-.36	1			
impulsivity	.12	-.23	.17	1		
NA	.25	-.49	.28	.38	1	
PA	-.20	.43	-.20	-.16	-.43	1
<i>Variance</i>	.62	.73	.64	.46	.60	.67
between-person						
suicidal ideation	1					
warmth	-.26	1				
dominance	.11	-.15	1			
impulsivity	.41	-.20	.30	1		
NA	.57	-.30	.26	.51	1	
PA	-.41	.46	-.15	-.15	-.38	1
<i>M (SD) / %</i>	4%	26.06	-.74	22.72	1.53	2.36
<i>Variance</i>	.38	.27	.36	.54	.40	.34

Note. $N=189$ (between), $N= 9100$ (within); Between-person variance was calculated as the intraclass correlation coefficient (ICC), which can be interpreted as the proportion of total variance accounted for at the between-person level. Within-person variance is therefore calculated as $1.0-ICC$. Bolded values indicate the credibility interval does not contain zero. NA = negative affect, PA = positive affect, SI = suicidal ideation.

Table 2. Key estimates from three individual multi-level structural equation models showing associations among suicidal ideation, negative affect, positive affect, or impulsivity, and perceived warmth of others during interactions (Model Set 2).

	(A) Negative Affect		(B) Positive Affect		(C) Impulsivity	
	<i>Est</i> (CI)	β	<i>Est</i> (CI)	β	<i>Est</i> (CI)	β
within-person estimates						
warmth $\rightarrow$ SI [<i>c</i>]	.007 [.000, .014]	.120	.001 [-.005, .010]	.062	-.002 [-.009, .007]	.014
warmth $\rightarrow$ NA [<i>a-NA</i>]	-.014 [-.016, -.012]	-.402	-	-	-	-
warmth $\rightarrow$ PA [<i>a-PA</i>]	-	-	.022 [.019, .024]	.422	-	-
warmth $\rightarrow$ Imp [<i>a-Imp</i>]	-	-	-	-	-.198 [-.242, -.154]	-.188
NA $\rightarrow$ SI [<i>b-NA</i>]	.466 [.324, .593]	.204	-	-	-	-
PA $\rightarrow$ SI [<i>b-PA</i>]	-	-	-.294 [-.458, -.181]	-.230	-	-
Imp $\rightarrow$ SI [<i>b-Imp</i>]	-	-	-	-	.006 [-.001, .013]	.068
indirect Effect [β_{ab}]	-.007 [-.009, -.004]		-.006 [-.010, -.004]		-.001 [-.003, .001]	
between-person estimates						
warmth $\rightarrow$ SI	-.006 [-.028, .012]	-.057	-.004 [-.030, .020]	-.033	-.021 [-.046, .003]	-.146
warmth $\rightarrow$ NA	-.014 [-.020, -.008]	-.326	-	-	-	-
warmth $\rightarrow$ PA	-	-	.025 [.018, .033]	.474	-	-
warmth $\rightarrow$ Imp	-	-	-	-	-.345 [-.599, -.090]	-.206
NA $\rightarrow$ SI	1.262 [.813, 1.664]	.472	-	-	-	-
PA $\rightarrow$ SI	-	-	-1.037 [-1.489, -.591]	-.418	-	-
Imp $\rightarrow$ SI	-	-	-	-	.028 [.014, .043]	.323

Note. $N=186$ (between), $N= 9009$ (within); $\rightarrow$ indicates regression; β parameter estimates are standardized. 95% credibility intervals of unstandardized parameter estimates are in parentheses. Bolded values indicate the credibility interval does not contain zero; warmth = perceived warmth of interaction partner, NA = negative affect, PA = positive affect, Imp = impulsivity, SI = suicidal ideation.

Table 3. Key estimates from three individual multi-level structural equation models showing associations among suicidal ideation, negative affect, positive affect, or impulsivity, and perceived dominance of others during interactions (Model Set 2).

	(A) Negative Affect		(B) Positive Affect		(C) Impulsivity	
	<i>Estimate</i> (CI)	β	<i>Estimate</i> (CI)	β	<i>Estimate</i> (CI)	β
within-person estimates						
dominance → SI [<i>c</i>]	-.004 [-.011, .004]	-.034	-.000 [-.007, .007]	.010	.002 [-.005, .008]	.028
dominance → NA [<i>a-NA</i>]	.009 [.007, .012]	.228	-	-	-	-
dominance → PA [<i>a-PA</i>]	-	-	-.010 [-.013, -.007]	-.178	-	-
dominance → Imp [<i>a-Imp</i>]	-	-	-	-	.158 [.112, .201]	.133
NA → SI [<i>b-NA</i>]	.423 [.285, .533]	.196	-	-	-	-
PA → SI [<i>b-PA</i>]	-	-	-.331 [-.452, -.228]	-.233	-	-
Imp → SI [<i>b-Imp</i>]	-	-	-	-	.004 [-.003, .011]	.040
indirect Effect [β_{ab}]	.004 [.002, .006]		.003 [.002, .005]		.001 [-.001, .002]	
between-person estimates						
dominance → SI	-.003 [-.020, .014]	-.031	.005 [-.014, .026]	.043	.002 [-.017, .023]	.019
dominance → NA	.009 [.0034, .015]	.251	-	-	-	-
dominance → PA	-	-	-.007 [-.014, .001]	-.138	-	-
dominance → Imp	-	-	-	-	.432 [.210, .625]	.298
NA → SI	1.281 [.901, 1.669]	.513	-	-	-	-
PA → SI	-	-	-1.078 [-1.48, -.665]	-.449	-	-
Imp → SI	-	-	-	-	.025 [.014, .037]	.347

Note. *N*=186 (between), *N*= 9009 (within); → indicates regression; β parameter estimates are standardized. 95% credibility intervals of unstandardized parameter estimates are in parentheses. Bolded values indicate the credibility interval does not contain zero; dominance = perceived dominance of interaction partner, NA = negative affect, PA = positive affect, Imp = impulsivity, SI = suicidal ideation.

Table 4. Key estimates from multi-level structural equation models simultaneously estimating associations among suicidal ideation, negative affect and impulsivity, and perceptions of others during interactions (Model Set 3).

	Negative Affect ↔ Impulsivity			
	(A) warmth		(B) dominance	
	<i>Estimate (CI)</i>	β	<i>Estimate (CI)</i>	β
within-person estimates				
perception → SI [c]	.005 [-.002, .013]	.116	-.003 [-.011, .006]	-.016
perception → NA [<i>a-NA</i>]	-.014 [-.016, -.012]	-.407	.009 [.007, .012]	.230
perception → Imp [<i>a-Imp</i>]	-.199 [-.244, -.155]	-.188	.159 [.114, .204]	.133
NA → SI [<i>b-NA</i>]	.467 [.305, .660]	.194	.470 [.323, .621]	.199
Imp → SI [<i>b-Imp</i>]	.000 [-.009, .007]	.000	.001 [-.008, .008]	-.011
NA ↔ Imp	2.640 [2.45, 2.83]	.301	3.27 [3.06, 3.50]	.331
between-person estimates				
perception → SI	-.010 [-.031, .016]	-.073	-.007 [-.028, .017]	-.054
perception → NA	-.014 [-.020, -.007]	-.316	.009 [.004, .015]	.248
perception → Imp	-.362 [-.606, -.101]	-.219	.432 [.213, .642]	.296
NA → SI	1.264 [.701, 1.809]	.416	1.31 [.67, 1.97]	.403
Imp → SI	.007 [-.005, .022]	.095	.014 [-.006, .030]	.160
NA ↔ Imp	4.60 [3.19, 6.25]	.494	4.50 [3.12, 6.22]	.482

Note. $N=186$ (between), $N= 9009$ (within); → indicates regression, ↔ indicates correlation between variables; β parameter estimates are standardized. 95% credibility intervals of unstandardized parameter estimates are in parentheses. Bolded values indicate the credibility interval does not contain zero; perception = degree to which interaction partners' behavior is perceived to be warm / dominant, NA = negative affect, Imp = impulsivity, SI = suicidal ideation.

Table 5. Key estimates from multi-level structural equation models simultaneously estimating associations among suicidal ideation, negative and positive affect, and perceptions of others during interactions (Model Set 3).

	Negative Affect ↔ Positive Affect			
	(A) warmth		(B) dominance	
	<i>Estimate (CI)</i>	β	<i>Estimate (CI)</i>	β
within-person estimates				
perception → SI [c]	.007 [.000, .015]	.136	-.003 [-.011, .004]	-.032
perception → NA [<i>a-NA</i>]	-.014 [-.016, -.013]	-.402	.009 [.007, .012]	.228
perception → PA [<i>a-PA</i>]	.022 [.019, .024]	.422	-.010 [-.013, -.007]	-.177
NA → SI [<i>b-NA</i>]	.376 [.241, .514]	.178	.361 [.221, .487]	.168
PA → SI [<i>b-PA</i>]	-.188 [-.333, -.056]	-.163	-.177 [-.328, -.021]	-.127
NA ↔ PA	-.119 [-.127, -.110]	-.299	-.189 [-.200, -.178]	-.389
between-person estimates				
perception → SI	.007 [-.016, .031]	.052	-.007 [-.025, .012]	-.060
perception → NA	-.014 [-.021, -.008]	-.322	.010 [.004, .015]	.249
perception → PA	.026 [.018, .033]	.477	-.007 [-.014, .000]	-.142
NA → SI	1.248 [.778, 1.74]	.405	1.304 [.854, 1.796]	.430
PA → SI	-.770 [-1.257, -.306]	-.310	-.695 [-1.20, -.302]	-.288
NA ↔ PA	-.075 [-.123, -.035]	-.278	-.112 [-.170, -.066]	-.360

Note. $N=186$ (between), $N= 9009$ (within); → indicates regression, ↔ indicates correlation between variables; β parameter estimates are standardized. 95% credibility intervals of unstandardized parameter estimates are in parentheses. Bolded values indicate the credibility interval does not contain zero; perception = degree to which interaction partners' behavior is perceived to be warm / dominant, NA = negative affect, PA = positive affect, SI = suicidal ideation.