Type: Preprint (PsyArXiv) v1 (4/11/21, word version); v2 (4/11/21, pdf version); v3 (8/11/21, minor grammar editing); v4 (15/3/22, minor content revision, minor title editing, supplement adding)

Full-title

Predicting self-harm and suicide ideation during the COVID-19 pandemic in Indonesia:
A nationwide survey report

Short-title

Self-harm & suicide ideation during the pandemic

Type

Original research article

Authors and affiliations

1. Andrian Liem*

Jeffrey Cheah School of Medicine and Health Sciences, Monash University
Malaysia, Selangor, Malaysia; andrian.liem@monash.edu; <https://orcid.org/0000-0002-1746-7235>.

2. Benny Prawira*

Into the Light Indonesia, DKI Jakarta, Indonesia; prawirabenny89@gmail.com;
<https://orcid.org/0000-0003-4417-9674>.

3. Selvi Magdalena

Into the Light Indonesia, DKI Jakarta, Indonesia; selvi.eru@gmail.com.

4. Monica Jenifer Siandita

Into the Light Indonesia, DKI Jakarta, Indonesia; jensiandita@gmail.com.

5. Joevarian Hudiyana

Faculty of Psychology, Universitas Indonesia, Jawa Barat, Indonesia;
joevarian91@ui.ac.id; <https://orcid.org/0000-0001-5507-0573>.

**corresponding authors*

Please cite

Liem, A., Prawira, B., Magdalena, S., Siandita, M. J., & Hudiyana, J. (2021). Predicting self-harm and suicide ideation during the COVID-19 pandemic in Indonesia: A brief report of nationwide survey. *PsyArXiv* (preprint). <https://doi.org/10.31234/osf.io/f3c8w>

Predicting self-harm and suicide ideation during the COVID-19 pandemic in Indonesia: A nationwide survey report

Abstract

Background: It is estimated that 77.0% of suicide cases occurred in low-and-middle-income countries (LMICs), which would increase because of the COVID-19 pandemic and socioeconomic inequity. However, there is lack of reports on this topic from LMICs, especially during the pandemic. Therefore, this nationwide study aimed to explore self-harm and suicide ideation and its predictive variables during the pandemic in Indonesia as a MIC with the highest COVID-19 fatality rate in Asia.

Methods: Non-random sampling online survey was conducted nationwide between 25 May and 16 June 2021. The collected data were demographic variables (*i.e.* age group), loneliness from social isolation using The UCLA Loneliness Scale Six Items (ULS-6), and self-harm and suicide ideation using item 9 of The Patient Health Questionnaire-9 (PHQ-9). Predictive model was analyzed using hierarchical logistic regression.

Results: A total of 5,211 participants from all 34 provinces in Indonesia completed the survey. Among 39.3% of them reported self-harm and suicide ideation during the pandemic, which significantly correlated with loneliness. The predictive variables associated with the likelihood of self-harm and suicide ideation were age, residence, job, religion, sex-gender, sexual orientation, HIV status, disability status, and loneliness. The predictive model showed a significant goodness-of-fit to the observed data ($\chi^2(15) = 1,803.46, p < .001$), $R_N^2 = .40$.

Conclusion: Four out of 10 Indonesians experienced self-harm and suicide ideation during the COVID-19 pandemic, particularly people within the age range of 18-24, living in the Java Island, unemployed/student/retired and freelancer, women, members of minority and marginalized communities, and experience of loneliness during the pandemic.

Keywords: suicide ideation, health inequity, mental health surveillance, loneliness, marginalized group, minority, LGBTIQ

Highlights

- 4 in 10 Indonesians experienced self-harm and suicide ideation during the pandemic.
- Loneliness from social isolation correlated with self-harm and suicide ideation.
- Socioeconomic inequity may contribute to self-harm and suicide ideation.

Predicting self-harm and suicide ideation during the COVID-19 pandemic in Indonesia: A nationwide survey report

BACKGROUND

Approximately 800,000 people worldwide died by suicide in 2019, 77.0% of whom occurred in low-and-middle-income countries (LMICs) (1), which was anticipated to increase due to the Coronavirus disease 2019 (COVID-19) pandemic. Life problems during this pandemic have been escalating and leading to mental disorders and self-harm, which is linked to suicide plan and suicide attempt (2-4). A systemic review and meta-analysis on the prevalence of the symptoms of depression, anxiety, and stress during the COVID-19 pandemic in another LMIC like Bangladesh found the pooled prevalence of these three conditions were 47%, 47%, and 45% respectively (5). Despite high suicide prevalence in LMICs, studies on this topic are clustered in high-income countries (HICs) and upper MICs. Therefore, the results might be less relevant to LMICs like Indonesia due to incomparable factors such as socioeconomic, resources, and health infrastructure between the two groups (6-8). In addition, Indonesia was downgraded back as a lower-middle income country again after a year classified as a upper-middle income country in 2020 because the economy was affected by the COVID-19 pandemic (9).

Previous nationwide study on anxiety conducted early in the pandemic in Indonesia was found about 20% of participants reported clinical anxiety symptoms, mostly among younger people, unemployed, and women (10). This finding is comparable to a nationwide survey in Bangladesh where 33% of the population were screened for depression, particularly among younger people (4). A community study in Indonesia among men who have sex with men (MSM) and transgender individuals in July-September 2020 found that nearly 70% of participants reported worse psychological distress during the COVID-19 pandemic than pre-pandemic (11). One of the reasons was discrimination faced by LGBTIQ people, which has existed long before the pandemic (12, 13). In other studies, vulnerable and marginalized communities (*e.g.* people with HIV positive and people with disability) reported more severe mental problems than general population due to multiple barriers they faced during the

pandemic (14-16). Additionally, the burden was also experienced by the caregivers as found in a qualitative study among family members of schizophrenia patients who care them on a daily basis and felt more difficult to provide care during the pandemic, which also affected caregivers' quality of life (17). Despite the importance of the findings, these nationwide quantitative studies did not investigate suicidal behavior.

There are studies on suicide behavior in certain age groups and specific geographical locations in Indonesia in pre-pandemic (18-20). However, there are no studies on self-harm and suicide ideation for wider population and more diverse groups, especially during the pandemic. Therefore, the current study aimed to explore self-harm and suicide ideation during the COVID-19 pandemic in Indonesia that had the highest COVID-19 fatality rate in Asia (21) and previous systematic analysis of suicide mortality burden found that the rates of suicide mortality in Indonesia was very high and influenced the average suicide rate the Southeast Asia region even before the pandemic (22). Moreover, self-harm was identified as a predictor of suicides among young people in LMICs (23). In addition, loneliness was consistently found to increase during the pandemic, which was associated with mental health problems and suicide attempt in diverse socioeconomic backgrounds globally (24-29). This study will also give attention to demographic variables, minority status (*i.e.* sexual orientation), and loneliness from social isolation, as possible predictors of self-harm and suicide ideation during the COVID-19 pandemic in Indonesia. It is expected that findings from this study will contribute to suicide intervention plan and program, which more than half of the WHO member states are lacking, including Indonesia. (1).

METHODS

Participants and procedure

The participant inclusion criteria were Indonesian, aged 18 or older, and living in Indonesia. The survey was set up using Typeform online survey software. The software recorded 15,184 view times (only at the cover page) and 10,430 attempts to fill it in. At the end, a total of 5,211 participants completed the survey data collection and incomplete responses were automatically deleted by the software. The response and

completion rates could not be calculated as view times and attempts to fill in the online survey were not representing the real number of people. For example, one person could visit the cover page multiple times then filled in the survey but discontinued or paused it. When the person tried to fill it in again, previous data was deleted and the software counted it as a new attempt. The participants were predominantly aged 18-24 (45.9%), had a diploma or higher educational background (54.4%), were Muslims (70.9%), and identified as women (67.3%). Details of the participants were provided in the Results and total participants in each province is provided in Supplement 1. The data were collected online from 25 May to 16 June 2021 using non-randomized sampling method. The link to complete the online survey was posted on social media accounts of Into The Light (ITL) Indonesia and Change.org Indonesia¹. The survey link was also sent through the mailing lists of ITL Indonesia and Change.org. To improve the survey inclusivity, personalized messages were sent to social media accounts of the marginalized communities (LGBTIQ people, people with disability, and people with HIV positive). This study was approved by the Research Ethics Committee of Research and Community Service Center (LPPM)-Universitas Katolik Indonesia Atma Jaya (0519A/III/LPPM-PM.10.05/05/2021).

Measures

All items were presented in Bahasa Indonesia, including the demographic information: age groups, current province of residence, ethnicity, educational background, job status/type, income category, religion/belief, and sex-gender. There were also three questions that asked participants about their minority and marginalized status, namely sexual orientation, HIV status, and disability status that covered both mental and physical disabilities. The demographic variables were grouped in analysis by considering the sociocultural situation in Indonesia. For example, 34 provinces were categorized into two groups, namely “Outside Java Island” and “Within Java Island”

¹ ITL Indonesia is a youth-based community with the focus as a center of advocacy, research, and education related to suicide prevention and mental health issues in Indonesia. Change.org Indonesia is the Indonesian office of Change.org, an American petition website that offers the public the ability to promote the petitions they care about to potential signers.

because more than half of total population resided in six provinces (Banten, West Java, DKI Jakarta, Central Java, DI Yogyakarta, and East Java) in Java Island (30).

Loneliness from social isolation was measured using The UCLA Loneliness Scale Six Items (ULS-6)-Indonesian version (31), which has an α of .81 in this study. The ULS-6 is a cross-cultural measure of loneliness, and it primarily assesses the perception of objective social isolation. Four intensity options were provided for each item from Never (1) to Often (4). The total score ranged from 6 to 24, where higher score indicated more intense loneliness in the last four weeks. Participants who answered 'Never' on all six items were categorized as "never felt loneliness", while participants who gave any other responses were grouped in "felt loneliness" for the descriptive analysis. The loneliness intensity was used for the correlational test with self-harm and suicide ideation intensity in the logistic regression analysis.

Self-harm and suicide ideation was measured using item number 9 of The Patient Health Questionnaire-9 (PHQ-9)-Indonesian version (32), which is "Over the last two weeks, how often have you been bothered by thoughts that you would be better off dead or of hurting yourself in some way?" with response options ranging from Not at all (0) to Nearly every day (3). Previous study found that higher score means higher risk for suicide in mental health screening for the community sample (33). For the descriptive and logistic regression analyses, participants who answered 'Not at all' were categorized in "no ideation", while participants who answered any other responses were categorized in "with ideation". The PHQ-9 has been validated among Indonesians, and the item 9 was classified into cognitive/affective factor of the two-factor model structure (34).

Statistical analysis

Data analyses were conducted in Statistical Package for the Social Sciences (SPSS) version 26. Descriptive analysis was performed, followed by a Pearson's test to analyze the correlation between loneliness intensity and self-harm and suicide ideation. A hierarchical logistic regression analysis was conducted to determine variables that predict self-harm and suicide ideation. Three blocks were built with the demographic variables in Block 1, minority conditions (sexual orientation, HIV status, and disability

status) in Block 2, and loneliness intensity in Block 3. In order to explore the best model, all variables in three blocks were entered using the “Backward-Stepwise” method and the first category of each categorical variable was set as the indicator reference. Adjusted odds ratios with the associated 95% confidence intervals were reported. The significance value (p) was set at .05 (two-sided).

RESULTS

Table 1 showed that almost all participants (98.0%) reported feeling lonely during the pandemic. The proportion was relatively lower among participants who were aged 35 or older (95.1%), worked as civil servants/police/military (95.2%), and had monthly income of Rp 9,000,000-11,999,999 ($\approx$ USD 632-842) (95.5%). The intensity of loneliness was relatively less intense among groups of participants aged 35 or older and worked as civil servants/police/military than other groups. In contrast, participants with disability and participants who were HIV positive reported higher intensity.

More than one-third of the total participants (39.3%) reported self-harm and suicide ideation during the pandemic, which was relatively higher in some groups, including participants with disability (64.5%), participants with HIV positive (62.8%), and non-heterosexual participants (56.8%). Conversely, the percentage was relatively lower in some other groups, such as participants who were aged 35 or older (14.2%), worked as civil servants/police/military (17.9%), and had monthly income of USD 843 or higher (21.5%). In general, participants reported having self-harm and suicide ideation intensity nearly several days in the last two weeks (score 1 on the scale), which was relatively more intense among participants who identified themselves as intersex, transgender, or non-binary and participants with disability compared to other groups. Correlational test showed loneliness intensity was significantly and positively correlated with self-harm and suicide ideation intensity ($r = .51, p < .001$).

Demographic variables	%	Loneliness experience (% yes)	Loneliness intensity (M, SD)	Self-harm & suicide ideation (% yes)	Self-harm & suicide ideation intensity (M, SD)
All participants	100	98.0	14.0 (4.6)	39.3	0.5 (0.8)
Age group					
18-24	45.9	98.9	15.2 (4.4)	50.4	0.7 (0.9)
25-34	34.0	98.5	13.8 (4.4)	39.1	0.5 (0.8)
≥ 35	20.1	95.1	11.3 (3.9)	14.2	0.2 (0.5)
Province					
Outside Java Island	24.2	97.5	13.7 (4.5)	36.1	0.5 (0.8)
Within Java Island	75.8	98.1	14.0 (4.6)	40.3	0.6 (0.8)
Ethnicity					
Non-Javanese	55.6	98.1	13.8 (4.5)	38.5	0.5 (0.8)
Javanese	44.4	97.9	14.1 (4.6)	40.3	0.6 (0.8)
Educational background					
High school and lower	45.6	98.6	14.8 (4.5)	45.2	0.7 (0.9)
Diploma and higher	54.4	97.5	13.2 (4.4)	34.3	0.5 (0.7)
Job status/type					
Unemployed/student/retired	41.1	98.6	15.3 (4.4)	50.4	0.7 (0.9)
Freelancer	13.9	98.1	13.8 (4.7)	39.4	0.5 (0.8)
Private employer/blue collar/driver/domestic worker	27.6	98.0	13.2 (4.4)	33.4	0.4 (0.7)
Civil servants/police/military	5.6	95.2	11.9 (4.2)	17.9	0.2 (0.6)
Entrepreneur	7.7	97.3	12.3 (4.1)	25.1	0.4 (0.7)
Others	4.1	96.2	12.7 (4.6)	22.6	0.3 (0.7)
Average monthly income group					
Other/have no income	9.8	98.8	15.2 (4.4)	50.9	0.7 (0.9)
<Rp 3,000,000 ($\approx$ USD 210)	48.6	98.5	14.7 (4.5)	44.9	0.7 (0.9)
Rp 3,000,000-5,999,999 ($\approx$ USD 210-421)	21.2	97.6	13.3 (4.4)	34.2	0.4 (0.7)
Rp 6,000,000-8,999,999 ($\approx$ USD 422-631)	8.8	97.0	12.6 (4.4)	28.6	0.4 (0.7)
Rp 9,000,000-11,999,999 ($\approx$ USD 632-842)	4.3	95.5	12.0 (4.0)	26.0	0.3 (0.6)
$\geq$ Rp 12,000,000 ($\approx$ USD 843)	7.3	97.4	12.0 (4.1)	21.5	0.3 (0.6)
Religion/belief					
Non-Islam	29.1	97.9	13.6 (4.4)	39.6	0.5 (0.8)
Islam	70.9	98.0	14.1 (4.6)	39.2	0.5 (0.8)
Sex-gender					
Male	31.9	96.8	12.9 (4.5)	28.6	0.4 (0.7)
Female	67.3	98.5	14.4 (4.5)	44.1	0.6 (0.8)
Intersex/transgender/non-binary	0.8	97.7	14.7 (4.7)	55.8	1.0 (1.0)
Sexual orientation					
Non-heterosexual	10.4	99.1	15.3 (4.5)	56.8	0.8 (0.9)
Heterosexual	89.6	97.9	13.8 (4.5)	37.2	0.5 (0.8)
HIV status					
Negative/do not know	99.2	98.0	13.9 (4.6)	39.1	0.5 (0.8)
Positive	0.8	100	15.5 (4.3)	62.8	0.9 (1.0)
Disability status					
Without disability	77.7	97.5	13.3 (4.4)	32.0	0.4 (0.7)
With disability	22.3	99.7	16.4 (4.3)	64.5	1.0 (1.0)

220 *Notes.* M = Mean. SD = Standard Deviation.

222 The final predictive model for self-harm and suicide ideation (Model 3) showed
 223 a significant goodness-of-fit to the observed data ($\chi^2 (15) = 1,803.46, p < .001$).
 224 Nagelkerke's pseudo R^2 indicated the significant variables in the third model explained
 225 approximately 40% of the variability. Variables and groups that were significantly
 226 associated with higher odds of experiencing self-harm and suicide ideation were age
 227 (18-24 years old), current residence (provinces in Java Island), job status/type
 228 (unemployed/student/retired and freelancer), religion (non-Islam), sex-gender (female),
 229 sexual orientation (non-heterosexual), disability status (with disability), and loneliness
 230 intensity (see Table 2).

231

232 **Table 2.** Hierarchical logistic regression models predicting self-harm and suicide
 233 ideation ($N = 5,211$)

Variables	Model 1			Model 2			Model 3		
	Adj OR	95% CI	p	Adj OR	95% CI	p	Adj OR	95% CI	p
Age group									
18-24	1 (ref)			1 (ref)			1 (ref)		
25-34	0.81	0.70-0.94	***	0.80	0.68-0.93	**	0.91	0.77-1.09	.30
≥ 35	0.24	0.19-0.29	***	0.26	0.21-0.33	***	0.39	0.31-0.50	***
Province									
Outside Java Island	1 (ref)			1 (ref)			1 (ref)		
In Java Island	1.33	1.16-1.53	***	1.29	1.12-1.50	**	1.24	1.06-1.46	.01
Job status/type									
Unemployed/student/retired	1 (ref)			1 (ref)			1 (ref)		
Freelancer	0.92	0.76-1.11	.38	0.88	0.72-1.08	.22	0.99	0.80-1.24	.98
Private employer/blue collar/driver/domestic worker	0.68	0.58-0.81	***	0.70	0.59-.83	***	0.82	0.68-0.99	.04
Civil servants/police/military	0.39	0.28-0.54	***	0.41	0.29-0.59	***	0.53	0.36-.77	**
Entrepreneur	0.62	0.48-0.81	***	0.64	0.48-0.84	**	0.85	0.63-1.14	.27
Others	0.42	0.30-0.60	***	0.42	0.29-0.61	***	0.47	0.31-.71	***
Religion/belief									
Non-Islam	1 (ref)			1 (ref)			1 (ref)		
Islam	0.79	0.69-0.91	**	0.84	0.73-0.96	.01	0.75	0.65-0.88	***
Sex-gender									
Male	1 (ref)			1 (ref)			1 (ref)		
Female	1.55	1.35-1.77	***	1.70	1.47-1.97	***	1.55	1.32-1.82	***
Intersex/transgender/non-binary	2.49	1.30-4.74	**	1.43	0.73-2.79	.29	1.53	0.73-3.17	.26
Sexual orientation									
Non-heterosexual				1 (ref)			1 (ref)		
Heterosexual				0.49	0.39-0.60	***	0.58	0.46-0.73	***
HIV status									
Negative/do not know				1 (ref)			1 (ref)		
Positive				2.31	1.16-4.60	.02	1.90	0.92-3.95	.08
Disability status									
Without disability				1 (ref)			1 (ref)		
With disability				3.25	2.81-3.75	***	2.18	1.86-2.56	***
Loneliness intensity							1.27	1.25-1.29	***
Constant	0.76		**	0.99		.93	0.03		***

Variables	Model 1			Model 2			Model 3		
	Adj OR	95% CI	p	Adj OR	95% CI	p	Adj OR	95% CI	p
Overall correct predicted percentage		63.5			68.1			75.2	
R_N^2		.14			.22			.40	
ΔR_N^2		.14			.08			.18	
Omnibus test / Chi-square (df)		566.99 (10)***			921.23 (14)***			1,803.46 (15)***	

Notes. Adj OR = Adjusted Odd Ratio. CI = Confidence Interval. Ref = Reference category. ** = $p < .01$. *** = $p < .001$. R_N^2 = Nagelkerke's pseudo R-squared. Δ = Delta. df = Degree of freedom.

DISCUSSION

This exploratory study aimed to investigate self-harm and suicide ideation and its predictive variables during the COVID-19 pandemic in Indonesia. Findings from this study complemented previous studies on suicidal behavior in specific age groups and limited geographical locations in Indonesia before the pandemic (18-20). It was found that nearly 40% of participants in this study reported self-harm and suicide ideation. The ideation intensity in this study was relatively low ($M = 0.5$, $SD = 0.8$), but it was higher than previous study that used the same item among female Indonesian migrant workers in Macau ($M = 0.1$, $SD = 0.3$) (34). This difference illustrated how pandemic affected psychological conditions because the data in the Macau study was not collected during a pandemic. Furthermore, the current prevalence was relatively higher compared to a longitudinal study that also used the same instrument among young people in Hong Kong, whereby 21% of the participants reported self-harm and suicide ideation in June 2020 (3); and also higher among Bangladeshi people who were only 5% of them answering "yes" on a question "Do you think about committing suicide, and are these thoughts persistent and related to COVID-19 issues?" in April 2020 (4).

The final predictive model showed that the younger age group (18-24 years old) had significantly 2.56 times higher chance of experiencing self-harm and suicide ideation than the older group (35 years or older). Our findings supported another representative population study among Indonesian but it was on anxiety, which found older age participants have lower risk of developing the disorder during the pandemic (10). We also found 50.4% of the young participants aged 18 to 24 reported having self-harm and suicide ideation, which was higher than suicidal ideation among Indonesian

adolescent students and young adults before the pandemic (19, 20, 35). The higher odds among young participants might be associated with challenges in transitioning to adulthood and adapting to online learning during the pandemic, which differed from their prior expectations (2, 11). Furthermore, previous studies found young generation accessed COVID-19 related information primarily from social media, which could lead to infodemic and triggered COVID-19 infection anxiety (10, 36).

Participants within the six provinces in Java Island, where more than 56% of total Indonesians resided (30), were predicted to have a significant 1.24 times greater chance of experiencing self-harm and suicide ideation than citizens in other areas. This might be because provinces in Java had been severely impacted by COVID-19 with the highest infection rate, more so than other provinces during the two different pandemic waves (37). Also, the infrastructure and access inequity within Java Island and disparity with other islands could be another reason. For example, 66.7% of the Class A hospitals with the best facilities and resources are located in Java Island, but they are much more accessible by people from high-income background, whereas people with low-and-middle income have limited resources (*e.g.* cannot take day off work and difficult transportation access) (21, 38). Moreover, participants in this study might be originally from outside Java and came to the island for studying or working, as evidenced by the domestic migration rate to six provinces in Java Island that reached 50.3% of total internal migration in Indonesia (39). Consequently, these domestic migrants might live alone and often do not have adequate social support, particularly young adults. This could become a risk factor as previous studies found family support could buffer the psychological problems during the crisis like the pandemic (2, 3, 40). The current study also aligned with a previous study on suicidal behavior among Indonesian people before the pandemic, whereby people living in urban area had 3.00 times higher suicide risk due to economic stress than people living in non-urban area (18).

Unemployed and freelancer groups showed greater chance of having self-harm and suicide ideation, which was up to 1.89 times higher when compared with members of civil servant, police, or military group. This finding mirrored previous nationwide studies that also found unemployed participants were 4.78 times more likely to

experience anxiety than health workers in Indonesia (10). The pressure as an economic-provider for these unemployed participants might be the reason for this psychological condition, as found in other countries with high unemployment rates (40). The self-harm and suicide ideation percentage was also found higher among participants with lower income, which is consistent with a systematic review on the association between economic inequity and poorer mental health, particularly in LMICs (7).

People with a minority-religion background significantly had 1.33 times greater odds of experiencing self-harm and suicide ideation than Muslim participants, which has the largest population in Indonesia (86.9%) (41). Movement restriction during the pandemic have made individuals unable to visit their worship places and meet their fellow believers. This could be a contributing factor to psychological problems given religious and spirituality aspects could not be detached from the sociocultural lives of Indonesians (42, 43). Moreover, individuals belonging to religious minority affiliations do not have as much support and feel more isolated from the mainstream culture (44). On the other hand, Muslim participants also could not attend mosques, however, their religious and spiritual connection could still be maintained because it is common for Indonesian mosques to use loud-speakers, which the sound (*Adzan*) is considered as an extension of a mosque in practicing the Islamic teachings (*dakwah*) and for religious purposes like a call to prayer and preaching (*khutbah*) (45). Moreover, the loudspeaker is also used for non-religious functions in daily lives like obituary news and community service (*gotong royong*) events.

Compared to men, women had 1.55 times higher likelihood for experiencing self-harm and suicide ideation. This trend was also found in an anxiety predictive model in Indonesia during the pandemic, where women had significantly 1.61 times higher odds of developing clinically significant anxiety than men (10). The reason might be because women faced more life challenges than men during the pandemic, primarily related to preconception and fertility, including unplanned pregnancy, perinatal stress, postpartum stress and depression, and miscarriage (46). Moreover, compared to men, women in the patriarchal society are expected to be more active in parenting and caregiving tasks, with additional risks of experiencing domestic violence (47).

Consequently, within these more challenging situations, women were more likely developed mental disorders such as anxiety and depression that were consistently found to be associated with suicidal behavior (3, 18).

An alternative explanation is men in the current study might be less open in expressing their feelings and mental problems because they worry about being perceived as weak and less masculine if they admit their struggles (27, 40). Previous study also found that objective living alone condition was more significant in triggering self-harm and suicidal ideation than loneliness among men (28). This condition was reflected by men's loneliness experience percentage and intensity that were relatively lower than women in the current study.

Three predictive models were developed where the R_N^2 increased from .14 in Model 1 to .40 when participant's minority conditions (sexual orientation, HIV status, and disability status) and loneliness intensity were included in Model 3, which became the novelty of this study by discovering the contribution of minority conditions and loneliness intensity to individual's self-harm and suicide ideation. It was found in the final model that non-heterosexual participants (10.4% of total participants) were significantly more likely to experience self-harm and suicide ideation, 1.72 times than heterosexual people. On the other hand, intersex/transgender/non-binary people also had higher odds in all three predictive models (2.49, 1.43, and 1.53 respectively), but the odds were statistically non-significant in the final model. This might be because this group was under-represented in the sample (0.8% of total participants).

These findings were consistent with a study among Indonesian transgender individuals and MSM that found 69.1% of the participants experienced moderate to high level of psychological distress during the pandemic that increased the risk of suicide behavior (11). Moreover, members of intersex/transgender/non-binary community lived like invisible citizens as their existence was not recognized by the government. Many of them were forced to have no citizen identity card, which means they were unable to access public services and aids. During the COVID-19 pandemic, many of them died without being surrounded by any relatives, and had to be buried without a tombstone because their identity was unknown (48).

People with positive HIV status who are also part of minority and marginalized communities had 1.90 times greater odds in the final model for experiencing self-harm and suicide ideation. However, this finding was statistically non-significant that might be due to the low representation in the overall participants (0.8%). Despite the non-significant ratio, the current finding reinforced previous studies that highlighted the very limited access to mental health services for LGBTIQ and people living with HIV worldwide, particularly in LMICs, which has worsened during the pandemic (13, 14)

People with disability were significantly predicted to have 2.18 times higher chance of experiencing self-harm and suicide ideation than people without disability. This could be because the pandemic had adversely impacted social interactions, ability to study, and opportunity to earn an income for people with disability, more severely than people without disability. For example, mask-wearing policy as part of COVID-19 health protocols meant people with deaf and hard of hearing were unable to read lip movements and communicate well with others (16). Moreover, inadequate support for people with disability may also contribute to their deteriorating mental health. For instance, sign language interpreter was not provided during the COVID-19 press conferences, so deaf people would find it extremely difficult to understand the updated situation and regulations regarding the pandemic (49). In addition, a community study in the US found that COVID 19-related stress was significantly associated with maladaptive coping strategies, including denial, substance use, and self-blame among people with disabilities and chronic conditions, which resulted in declining well-being (15).

Intensity of loneliness from social isolation among participants in this study was higher ($M = 14.0$, $SD = 4.6$) compared with the ULS-6 validation study in Indonesia, Germany, and the US, where Indonesian participants reported the lowest score ($M = 12.4$, $SD = 3.83$) (31). In the final predictive model, loneliness contributed to 18.0% of variability. The model predicted that loneliness from social isolation would increase the chance of people experiencing self-harm and suicide ideation by 1.27 times. These findings were in line with a longitudinal study during the pandemic (May-July 2020) in Brazil, where loneliness significantly increased the odds of suicidal ideation by 2.12

times fold (29).

The current study also found that prevalence of loneliness due to social isolation among younger participants was higher than the older group. These young participants might be dominated by students who studied away from their hometown and lived alone, which would amplify loneliness experience. Furthermore, chronic loneliness was consistently found to be associated with psychological disorders, particularly depression, among children, adolescents, and young adults worldwide (25). However, loneliness was statistically non-significant associated with how many people were around the individual, or the duration of conversation they had (27). In support of this, a longitudinal study in the UK found that chronic loneliness in adolescents during the COVID-19 pandemic, not how often they talked with friends, was significantly correlated with deteriorating mental health and the relationship was bidirectional (24). Therefore, more social interventions that address loneliness from social disconnectedness need to be targeted especially to the young age group.

Study strengths and limitations

The important strengths in this study were the large nationwide participants sampled from all provinces in Indonesia with diverse backgrounds and the use of validated instruments. However, there was an over-representation of participants who lived in Java Island that might have different experiences from people in other provinces outside Java Island. Also, despite the data being collected from members of minority and marginalized communities, their numbers were relatively small and might be insufficient for statistical analyses. The question of sex and gender was also mixed due to the public negative sensitivity towards gender nonconformity. Next, this online cross-sectional study does not allow for causal interpretations. The results were also limited to the data collection period and may have the risk of self-selection bias. Lastly, despite the PHQ-9 has been validated among Indonesians this instrument was originally designed to screen for depression; and used a single item (Item 9) to screen self-harm and suicide ideation in non-primary care situations might be less accurate compared with other instruments that are specially designed for this purpose. Therefore, future study should reach participants from outside of Java Island and members of minority

and marginalized communities and use a measure with higher sensitivity and specificity for screening suicide ideation in a general setting. Findings from this quantitative study should also be complemented by qualitative investigation, particularly the experiences of minority and marginalized communities that might not be suitable to be analyzed statistically.

CONCLUSION

Four in ten people in Indonesia may experience self-harm and suicide ideation during the COVID-19 pandemic. Variables that significantly contributed to predicting self-harm ideation were age, province, job status, religion, sex, sexual orientation, disability status, and intensity of loneliness from social isolation. The likelihood of experiencing self-harm and suicide ideation were higher among people aged 18-24, lived in Java Island, unemployed/student/retired and freelancer, women, members of minority and marginalized communities (non-Islam, non-heterosexual, and with disability), and those experiencing loneliness from social isolation during the pandemic. Mental health treatment and suicide prevention programs during and post-pandemic should prioritize these high-risk groups and be tailored by involving the group members in the planning processes.

DECLARATIONS

Ethics approval and consent to participate

Procedures followed were in accordance with all guidelines and regulations (institutional and national) and with the Declaration of Helsinki. This study was approved by the Research Ethics Committee of Research and Community Service Center (LPPM)-Universitas Katolik Indonesia Atma Jaya (0519A/III/LPPM-PM.10.05/05/2021). Consent to participate was collected online after an electronic informed consent page was provided to the potential participants before they started the online survey.

Consent for publication

Consent for publication was collected online before participants filled the online survey.

Availability of data and material

The Indonesian version of survey methods and results can be accessed from the Into The Light Indonesia website.² Derived data supporting the findings of this study are available from the corresponding author on request.

Competing interest

The authors declare no conflict of interest.

Funding

This study received no funding from any institution.

Authors' contributions

Study proposal: BP, SM, MJS. Study design and data collection: AL, BP, SM, MJS. Data cleaning and analysis: AL. Manuscript draft: AL. Critical feedback for the manuscript: AL, BP, SM, MJS, JH. All authors have approved the final version of the manuscript.

Acknowledgments

The authors thank Change.org Indonesia, ITL Indonesia members, and members of minority and marginalized communities for their assistance in data collection. The authors also thank Rachmadiani S. Hanifa for her contribution in the pre-study design and Rachmania P. Wardhani for providing constructive feedback on the manuscript.

² Into The Light Indonesia. (2021). *Seri Laporan ke-1: Laporan Perilaku Penggunaan Layanan Kesehatan Mental di Indonesia 2021 - Hasil Awal [Serial Report No. 1: Use of Mental Health Services in Indonesia 2021 - Preliminary Report]*. I. T. L. Indonesia. <https://www.intothelightid.org/tentang-bunuh-diri/laporan-perilaku-penggunaan-layanan-kesehatan-mental-di-indonesia-2021-hasil-awal/>

REFERENCES

1. WHO. Mental Health Atlas 2020. Geneva: World Health Organization; 2021.
2. Pramukti I, Strong C, Sitthimongkol Y, Setiawan A, Pandin MGR, Yen C-F, et al. Anxiety and suicidal thoughts during the COVID-19 pandemic: Cross-country comparative study among Indonesian, Taiwanese, and Thai University Students. *J Med Internet Res*. 2020;22:1-12. DOI: 10.2196/24487.
3. Zhu S, Zhuang Y, Lee P, Wong PWC. The changes of suicidal ideation status among young people in Hong Kong during COVID-19: A longitudinal survey. *J Affect Disord*. 2021;294:151-8. DOI: 10.1016/j.jad.2021.07.042.
4. Mamun MA, Sakib N, Gozal D, Bhuiyan AKMI, Hossain S, Bodrud-Doza M, et al. The COVID-19 pandemic and serious psychological consequences in Bangladesh: A population-based nationwide study. *J Affect Disord*. 2021;279:462-72. DOI: 10.1016/j.jad.2020.10.036.
5. Hosen I, Al-Mamun F, Mamun MA. Prevalence and risk factors of the symptoms of depression, anxiety, and stress during the COVID-19 pandemic in Bangladesh: A systematic review and meta-analysis. *Global Mental Health*. 2021;8. DOI: 10.1017/gmh.2021.49.
6. Vijayakumar L, Ray S, Fernandes TN, Pathare S. A descriptive mapping review of suicide in vulnerable populations in low and middle countries. *Asia Pac Psychiatry*. 2021;13:1-12. DOI: 10.1111/appy.12472.
7. Tibber MS, Walji F, Kirkbride JB, Huddy V. The association between income inequality and adult mental health at the subnational level—A systematic review. *Soc Psychiatry Psychiatr Epidemiol*. 2021. DOI: 10.1007/s00127-021-02159-w.
8. Pirkis J, John A, Shin S, DelPozo-Banos M, Arya V, Analuisa-Aguilar P, et al. Suicide trends in the early months of the COVID-19 pandemic: An interrupted time-series analysis of preliminary data from 21 countries. *Lancet Psychiatry*. 2021;8:579-88. DOI: 10.1016/S2215-0366(21)00091-2.
9. Oktavia MH. Indonesia Categorized as Lower Middle-Income Country: World Bank. Kompascom. 2021.
10. Anindyajati G, Wiguna T, Murtani BJ, Christian H, Wigantara NA, Putra AA, et al. Anxiety and its associated factors during the initial phase of the COVID-19 pandemic in Indonesia. *Frontiers in Psychiatry*. 2021;12(253):1-10. DOI: 10.3389/fpsy.2021.634585.
11. Septarini NW, Hendriks J, Maycock B, Burns S. Psychological distress and happiness of men who have sex with men and transgender people during the coronavirus disease-19 pandemic: Is there a need for public health policy intervention? *Frontiers in Public Health*. 2021;9:1-12. DOI: 10.3389/fpubh.2021.647548.
12. Hudiyana J, Prawira B, Kartika DA, Mahendra D, Putra IE. Gods, germs, and science: Unraveling the role of scientific literacy, germ aversion, and religious fundamentalism in predicting attitudes towards gays and lesbians. *J Community Appl Soc Psychol*. 2021. DOI: 10.1002/casp.2534.
13. Wallach S, Garner A, Howell S, Adamson T, Baral S, Beyrer C. Address exacerbated health disparities and risks to LGBTQ+ individuals during COVID-19. *Health Hum Rights*. 2020;22(2):313-6.
14. Newman PA, Prabhu SM, Akkakanjanasupar P, Tepjan S. HIV and mental health among young people in low-resource contexts in Southeast Asia: A qualitative investigation. *Global public health*. 2021;(online first):1-15. DOI: 10.1080/17441692.2021.1924822.
15. Umucu E, Lee B. Examining the impact of COVID-19 on stress and coping strategies in individuals with disabilities and chronic conditions. *Rehabil Psychol*. 2020;65:193-8. DOI: 10.1037/rep0000328.
16. The Australia-Indonesia Centre PAIR. A resilient community: Tackling COVID-19's impact on people with disability 2021 13/10/2021. Available from:

<https://pair.australiaindonesiacentre.org/research/covid-19/a-resilient-community-tackling-covid-19s-impact-on-people-with-disability/>.

17. Indriati I, Adi WS, Anggarawati T, Astuti Y, Alfriani D. Familial experiences of caring for schizophrenia patients during the COVID-19 pandemic: A qualitative study. *Jurnal Ners*. 2021;16:35-40. DOI: 10.20473/jn.v16i1.24130.

18. Mubasyiroh R, Pradono J, Nurkhotimah E, Kusumawardani N, Idaiani S. Depression as a strong prediction of suicide risk. *Global Journal of Health Science*. 2018;10:52-62. DOI: 10.5539/gjhs.v10n12p52.

19. Putra IGNE, Karin PAES, Ariastuti NLP. Suicidal ideation and suicide attempt among Indonesian adolescent students. *Int J Adolesc Med Health*. 2019;33:1-12. DOI: 10.1515/ijamh-2019-0035.

20. Ahmad SF, Turnip SS. Does family communication pattern predict suicide ideation and attempt? A longitudinal study of adolescents in Indonesia. *Advances in Social Science, Education and Humanities Research*. 2020;494:236-42. DOI: 10.2991/assehr.k.201125.020.

21. Siregar K. Why Indonesia Has The Highest Covid-19 Fatality Rate In Asia. *Channel News Asia* [Internet]. 2020 26/04/2020. Available from: <https://www.channelnewsasia.com/asia/covid-19-fatality-rate-highest-asia-indonesia-766466>.

22. Naghavi M, Postma M. Global, regional, and national burden of suicide mortality 1990 to 2016: Systematic analysis for the Global Burden of Disease Study 2016. *BMJ*. 2019;364:1-11. DOI: 10.1136/bmj.l94.

23. Aggarwal S, Borschmann R, Patton GC. Tackling stigma in self-harm and suicide in the young. *The Lancet Public Health*. 2021;6(1):e6-e7. DOI: 10.1016/S2468-2667(20)30259-0.

24. Cooper K, Hards E, Moltrecht B, Reynolds S, Shum A, McElroy E, et al. Loneliness, social relationships, and mental health in adolescents during the COVID-19 pandemic. *J Affect Disord*. 2021;289:98-104. DOI: 10.1016/j.jad.2021.04.016.

25. Loades ME, Chatburn E, Higson-Sweeney N, Reynolds S, Shafran R, Brigden A, et al. Rapid systematic review: The impact of social isolation and loneliness on the mental health of children and adolescents in the context of COVID-19. *J Am Acad Child Adolesc Psychiatry*. 2020;59:1218-39.e3. DOI: 10.1016/j.jaac.2020.05.009.

26. Lim MH, Penn DL, Thomas N, Gleeson JFM. Is loneliness a feasible treatment target in psychosis? *Soc Psychiatry Psychiatr Epidemiol*. 2020;55(7):901-6. DOI: 10.1007/s00127-019-01731-9.

27. Ernst M, Klein EM, Beutel ME, Brähler E. Gender-specific associations of loneliness and suicidal ideation in a representative population sample: Young, lonely men are particularly at risk. *J Affect Disord*. 2021;294:63-70. DOI: 10.1016/j.jad.2021.06.085.

28. Shaw RJ, Cullen B, Graham N, Lyall DM, Mackay D, Okolie C, et al. Living alone, loneliness and lack of emotional support as predictors of suicide and self-harm: A nine-year follow up of the UK Biobank cohort. *J Affect Disord*. 2021;279:316-23. DOI: 10.1016/j.jad.2020.10.026.

29. Antonelli-Salgado T, Monteiro GMC, Marcon G, Roza TH, Zimmerman A, Hoffmann MS, et al. Loneliness, but not social distancing, is associated with the incidence of suicidal ideation during the COVID-19 outbreak: A longitudinal study. *J Affect Disord*. 2021;290:52-60. DOI: 10.1016/j.jad.2021.04.044.

30. Badan Pusat Statistik-Statistics Indonesia. Hasil Sensus Penduduk 2020. Jakarta, Indonesia; 2021.

31. Hudiyana J, Lincoln TM, Hartanto S, Shadiqi MA, Milla MN, Muluk H, et al. How universal is a construct of loneliness? Measurement invariance of the UCLA Loneliness Scale in Indonesia, Germany, and the United States. *Assessment*. 2021;(online first):1-11. DOI: 10.1177/10731911211034564.

- 557 32. Kroenke K, Spitzer RL. The PHQ-9: A new depression diagnostic and severity measure.
558 Psychiatric Annals. 2002;32(9):509-15.
- 559 33. Kim S, Lee H-K, Lee K. Which PHQ-9 items can effectively screen for suicide? Machine learning
560 approaches. Int J Environ Res Public Health. 2021;18:1-10.
- 561 34. Hall BJ, Patel AR, Lao L, Liem A, Mayawati EH, Tjipto S. Structural validation of The Patient
562 Health Questionnaire-9 (PHQ-9) among Filipina and Indonesian female migrant domestic workers in
563 Macao. Psychiatry Res. 2021;295:1-7. DOI: 10.1016/j.psychres.2020.113575.
- 564 35. Prawira B, Sukmaningrum E. Suicide stigma as a predictor of help-seeking intention among
565 undergraduate students in Jakarta. Makara Human Behavior Studies in Asia. 2020;24:24-36. DOI:
566 10.7454/hubs.asia.1200120.
- 567 36. Liem A, Wang C, Dong C, Lam AIF, Latkin CA, Hall BJ. Knowledge and awareness of COVID-19
568 among Indonesian migrant workers in the Greater China Region. Public Health. 2021;197:28-35. DOI:
569 10.1016/j.puhe.2021.05.029.
- 570 37. Satuan Tugas Penanganan COVID-19. Indonesia Fights Back The Covid-19 Second Wave 2021
571 28/10/2021. Available from: [https://covid19.go.id/p/berita/indonesia-fights-back-covid-19-second-](https://covid19.go.id/p/berita/indonesia-fights-back-covid-19-second-wave)
572 [wave](https://covid19.go.id/p/berita/indonesia-fights-back-covid-19-second-wave).
- 573 38. Sambodo NP, Van Doorslaer E, Pradhan M, Sparrow R. Does geographic spending variation
574 exacerbate healthcare benefit inequality? A benefit incidence analysis for Indonesia. Health Policy Plan.
575 2021;36:1129-39. DOI: 10.1093/heapol/czab015.
- 576 39. Yuliadi I, Raharja SS. Migration and economic inequality in Indonesia: Longitudinal data
577 analysis. The Journal of Asian Finance, Economics and Business. 2020;7:541-8. DOI:
578 10.13106/jafeb.2020.vol7.no11.541.
- 579 40. Chen Y-Y, Cai Z, Chang Q, Canetto SS, Yip PSF. Caregiving as suicide-prevention: An ecological
580 20-country study of the association between men's family carework, unemployment, and suicide. Soc
581 Psychiatry Psychiatr Epidemiol. 2021. DOI: 10.1007/s00127-021-02095-9.
- 582 41. Katadata. Sebanyak 86,88% Penduduk Indonesia Beragama Islam: Persentase Pemeluk
583 Agama/Kepercayaan di Indonesia (Juni 2021) 2021 13/10/2021. Available from:
584 [https://databoks.katadata.co.id/datapublish/2021/09/30/sebanyak-8688-penduduk-indonesia-](https://databoks.katadata.co.id/datapublish/2021/09/30/sebanyak-8688-penduduk-indonesia-beragama-islam)
585 [beragama-islam](https://databoks.katadata.co.id/datapublish/2021/09/30/sebanyak-8688-penduduk-indonesia-beragama-islam).
- 586 42. Regus M. Regulating religion in a time of COVID-19 pandemic in Indonesia: context, dynamics,
587 and implication. International Journal of Sociology and Social Policy. 2021;(online first):1-19. DOI:
588 10.1108/IJSSP-07-2020-0321.
- 589 43. Liem A. "Doing my profession is also part of worship": How clinical psychologists address
590 aspects of spirituality and religion in Indonesia. Journal of Religion and Health. 2020;59:1434-57. DOI:
591 10.1007/s10943-019-00880-0.
- 592 44. Lawrence RE, Oquendo MA, Stanley B. Religion and suicide risk: A systematic review. Arch
593 Suicide Res. 2016;20:1-21. DOI: 10.1080/13811118.2015.1004494.
- 594 45. Pratisti SA. Negotiating soundscape: Practice and regulation of Adzan in Indonesia. Journal of
595 Indonesian Social Sciences and Humanities. 2020;10:75-84. DOI: 10.14203/jissh.v10i2.180.
- 596 46. Ifdil I, Fadli RP, Gusmaliza B, Putri YE. Mortality and psychological stress in pregnant and
597 postnatal women during COVID-19 outbreak in West Sumatra, Indonesia. J Psychosom Obstet Gynaecol.
598 2020;41(4):251-2. DOI: 10.1080/0167482X.2020.1779216.
- 599 47. Almeida M, Shrestha AD, Stojanac D, Miller LJ. The impact of the COVID-19 pandemic on
600 women's mental health. Arch Womens Ment Health. 2020;23:741-8. DOI: 10.1007/s00737-020-01092-
601 2.

48. Renaldi A. No ID, No Vaccines for Indonesia's Invisible People 2021 19/10/2021. Available from:
<https://restofworld.org/2021/indonesias-invisible-people-face-discrimination-and-sometimes-death-by-database/>.
49. Aulia FD, Asiah DHS, Irfan M. Peran pemerintah dalam penanganan dampak pandemi COVID-19 bagi penyandang disabilitas. Jurnal Pengabdian dan Penelitian Kepada Masyarakat. 2020;1:31-41. DOI: 10.24198/jppm.v1i1.30951.

preprint V4

SUPPLEMENT

Supplement 1. Total participants in each province

No	Province	<i>n</i>	%
1	Jawa Barat	1,172	22.49
2	DKI Jakarta	1,012	19.42
3	Jawa Timur	585	11.23
4	Jawa Tengah	509	9.77
5	Banten	431	8.27
6	DI Yogyakarta	243	4.66
7	Sumatera Utara	154	2.96
8	Bali	148	2.84
9	Sulawesi Selatan	81	1.55
10	Kalimantan Timur	77	1.48
11	Lampung	68	1.30
12	Sumatera Barat	67	1.29
13	Riau	60	1.15
14	Sumatera Selatan	59	1.13
15	Kalimantan Barat	58	1.11
16	Nusa Tenggara Timur	56	1.07
17	Kepulauan Riau	55	1.06
18	Kalimantan Selatan	53	1.02
19	DI Aceh	46	0.88
20	Sulawesi Utara	43	0.83
21	Nusa Tenggara Barat	41	0.79
22	Jambi	33	0.63
23	Sulawesi Tengah	27	0.52
24	Kalimantan Tengah	26	0.50
25	Bengkulu	16	0.31
26	Kepulauan Bangka Belitung	16	0.31
27	Sulawesi Tenggara	15	0.29
28	Gorontalo	13	0.25
29	Papua	12	0.23
30	Kalimantan Utara	11	0.21
31	Maluku	10	0.19
32	Papua Barat	8	0.15
33	Maluku Utara	4	0.08
34	Sulawesi Barat	2	0.04
Total		5,211	100.00

Note. Provinces number 1-6 are in Java Island.