Foraging Tempo: Human run patterns in multiple target search are constrained by the rate of successive responses.

Ian M. Thornton (1), Tram T. N. Nguyen (1) & Árni Kristjánsson (2, 3)

(1) Department of Cognitive Science, Faculty of Media and Knowledge Sciences, University of Malta, Malta.

(2) Faculty of Psychology, School of Health Sciences, University of Iceland, Oddi v. Sturlugötu, 101, Reykjavik, Iceland.

(3) School of Psychology, National Research University, Higher School of Economics, Moscow, Russian Federation.

Word Count: 9,235

KEYWORDS: Foraging; Foraging Tempo; Visual Search; Attention; Multiple Target Search; Temporal Constraints; Individual Differences

Corresponding author: Ian Thornton, ian.thornton@um.edu.mt

Abstract

Human foraging tasks are beginning to provide new insights into the roles of vision, attention and working memory during complex, multiple-target search. Here, we test the idea that “foraging tempo” -- the *rate* of successive target selections -- helps determine patterns of behaviour in these tasks. Previously, we established that the majority of target selections during unconstrained foraging happen at regular, rapid intervals, forming the “cruise phase” of a foraging trial. Furthermore, we noted that when the temporal interval between cruise phase responses was longer, the tendency to switch between target categories increased. To directly explore this relationship, we modified our standard iPad foraging task so that observers had to synchronize each response with an auditory metronome signal. Across trials, we increased the tempo and examined how this changed patterns of foraging when targets were defined either by a single feature or by a conjunction of features. The results were very clear. Increasing tempo systematically *decreased* the tendency for participants to switch between target categories. While this was true for both feature and conjunction trials, there was also evidence that time constraints and target complexity interacted. As in our previous work, we also observed clear individual differences in how participants responded to changes in task difficulty. Overall, our results show that foraging tempo does influence the way participants respond, and we suggest this parameter may prove be useful in further explorations of group and individual strategies during multiple target search.

Introduction

In a recent series of studies, we have used a range of 2D (Jóhannesson et al., 2016, e.g., 2017; Á. Kristjánsson et al., 2014; T. Kristjánsson et al., 2018; Thornton et al., 2019) and 3D (T. Kristjánsson, Draschkow, et al., 2020; Prpic et al., 2019) cancellation tasks to explore human foraging under conditions of varying attentional load. This work was directly inspired by the animal foraging literature (Dawkins, 1971; Heinrich et al., 1977; Jackson & Li, 2004; Pietrewicz & Kamil, 1979; Tinbergen, 1960). In our tasks, we extended the classic human visual search paradigm -- where participants are typically asked to locate a single target item embedded within a variable set size of distractor items (Hulleman & Olivers, 2017; Á. Kristjánsson & Egeth, 2020; Treisman & Gelade, 1980; Wolfe, 2010; Wolfe & Horowitz, 2004, 2017) -- to include situations where multiple targets from different categories must be located on each trial. Several other groups have also explored multiple target search (e.g., Cain et al., 2012; Fougner et al., 2015; Gilchrist et al., 2001; Hills et al., 2012, 2013; Pellicano et al., 2011; Wolfe, 2013; Wolfe et al., 2016, 2019). The common goal of such work is to better understand the relationship between vision, attention and action in complex scenarios (see Á. Kristjánsson et al., 2019 for a recent discussion).

In our original iPad task (Á. Kristjánsson et al., 2014) -- which also forms the basis of the current experimental work -- participants were required to use their finger to touch and thus cancel 40 target items on each trial. There were always 20 targets from two different categories, randomly distributed amongst 40 distractor items from two other categories. The main finding was a clear difference in patterns of foraging “runs” when the individual target items were defined by a single feature (e.g. colour; feature conditions/foraging) versus a combination of features (e.g. colour and shape; conjunction conditions/foraging). A “run” in this context simply refers to a sequence of selections from the same target category.

During feature foraging, all participants switched randomly between the two target categories, producing many short run sequences. During conjunction foraging, in contrast, the majority of participants only produced two runs, with all targets from one category being cancelled before the other. We interpreted this “exhaustive” category selection as indicating that under higher attentional load, these participants were unable or unwilling to switch rapidly between search templates, a phenomenon also discussed in the literature on animal foraging (Bond, 1983; Dawkins, 1971; Dukas & Ellner, 1993; Kamil & Bond, 2006; Tinbergen, 1960). In our original data set, we also noted that approximately 25% of participants did not show

exhaustive category selection, but rather continued to switch randomly during conjunction foraging. We termed these individuals “super-foragers” as they seemed immune to the increase in attentional load. The existence of these two basic foraging profiles – and the approximate 75/25% split -- has been replicated several times (e.g., Clarke et al., 2018; Jóhannesson et al., 2017), and we return to this interesting individual difference later in this paper.

In subsequent studies, we also identified two other aspects of foraging behaviour that form the starting point for the current study. First, examination of inter-target times (ITTs) revealed that the majority of selections happen at very regular intervals, forming what we termed the “cruise phase” of a trial (T. Kristjánsson, Thornton, et al., 2020). This regularity is interrupted only at the very start (see also Basoudan et al., 2019; Thornton & Horowitz, 2004, 2020), end and midpoint (for conjunction trials) of a trial, where response times are significantly slowed relative to the cruise phase (see T. Kristjánsson, Thornton, et al., 2020 for details).

Second, the absolute rate of selection during the cruise phase – the “foraging tempo” – appears to impose important constraints on the pattern of runs. Specifically, by analysing several previous studies from our own group and others, Thornton et al. (2019) described a clear pattern where task variants that permitted rapid ITTs (e.g., < 400 ms/item) were accompanied by a reduced tendency to switch during conjunction trials (e.g., Jóhannesson et al., 2016; Á. Kristjánsson et al., 2014; T. Kristjánsson, Thornton, et al., 2020). Conversely, those task variants where ITTs were relatively long (e.g., > 700 ms/item) gave rise to an increased tendency to switch (Clarke et al., 2018; Prpic et al., 2019; Thornton et al., 2019). These findings suggest that template complexity and temporal constraints interact to jointly determine foraging patterns. In particular, the exhaustive run behaviour found with our original iPad task may not have been solely a consequence of template complexity, but could also have relied on the fact that participants were responding rapidly.

The general idea that “search rate” can affect foraging patterns has a long and sometimes controversial history within the animal behaviour literature (Bond & Riley, 2010; Gendron & Staddon, 1983; Guilford & Dawkins, 1987, 1989a, 1989b; Hollis, 1989; Lawrence, 1989; Plaisted & Mackintosh, 1995; Reid & Shettleworth, 1992). It is generally accepted that by searching more slowly, an animal can increase the probability of detecting prey, particularly [if the prey are cryptic](#) (Gendron & Staddon, 1983). However, precisely how search rate modulates patterns of runs and how it interacts with other constraints on foraging -- such as attention (Dukas & Ellner, 1993), search images/templates (Guilford & Dawkins, 1987; Plaisted & Mackintosh, 1995) or learning (Troschianko et al., 2018) -- remains less clear. For example, while search rate is incorporated into the ‘attentive prey’ model of Dukas & Ellner (1993), their

implementation assumes ‘that there is no interaction between search rate and attention, and that the total amount of attention devoted for foraging is constant regardless of the search rate’ (p.1339).

Here, we suggest that a useful framework which may shed further light on this relationship -- and foraging tempo more generally -- is the “soft constraints” approach proposed by Wayne Gray and colleagues (Gray et al., 2006; Gray & Boehm-Davis, 2000; Gray & Fu, 2004). Before describing this approach, we should note that in the remainder of this paper we will continue to use the term “foraging tempo” to discuss temporal patterns during foraging, rather than “search rate”, as we feel the former better captures the regular response patterns in multiple target tasks, and avoids confusion with temporal estimates from traditional single target studies (Hulleman & Olivers, 2017; Á. Kristjánsson & Egeth, 2020; Treisman & Gelade, 1980; Wolfe, 2010; Wolfe & Horowitz, 2004, 2017). While the term “foraging tempo” has been used previously in the animal literature to refer to global rates of return to a nest or hive (e.g., Burkhardt, 1998; Davidson, 1997; Dyer & Seeley, 1991; Oster & Wilson, 1978), we believe the context should clearly disambiguate this usage from the current meaning.

According to Gray and colleagues then, in any interactive task a series of soft constraints govern the trade-off between low-level cognitive, perceptual and motor operations (Gray et al., 2006; Gray & Boehm-Davis, 2000; Gray & Fu, 2004). Central to this approach is the idea that ‘the control system selects sequences of interactive routines that tend to minimize performance costs measured in time while achieving expected benefits’ (Gray et al., 2006; p.463). [For any given task, the limiting factor\(s\) in terms of time could be cognitive, perceptual or motor operations.](#) If rapid motor responses are possible, for example, the limiting stages may involve perceptual or cognitive operations and constraints may be imposed on these to ensure that additional time costs are not incurred. If, on the other hand, motor responses cannot be made rapidly for some reason, the temporal bottleneck will no longer occur at the cognitive and/or perceptual stages, and constraints on the nature of the operations selected at these levels might be relaxed.

Applying these ideas to our previous findings, when using the iPad to respond, the temporal cost of cognitively switching between different conjunction templates might approach or exceed the minimum possible movement time. To avoid an increase in performance costs, constraints are applied at the cognitive level to ensure that selected operations can be completed well within the interval dictated by the motor system. Specifically, switching or activation are minimised and a single template remains active until all targets of that category have been selected. In task variants where overall response rates are slower, however, either due to the

need to navigate between target areas (Prpic et al., 2019) or because responses are being made via the mouse/keyboard rather than a touchscreen (Clarke et al., 2018; Thornton et al., 2019), conjunction template operations may not have been a limiting factor. That is, if switching between conjunction templates can be achieved within the time limit dictated by slower motor operations, then the soft constraints on the use of such a strategy may be relaxed.

The goal of the current paper was to directly test this foraging tempo hypothesis using a single task variant where operations at each level (i.e. cognitive, perceptual & motor) were held constant as we systematically varied response rate. We had participants forage through displays that were identical to those in our original iPad paper (Á. Kristjánsson et al., 2014), with the same basic success (cancel all targets from both categories) and failure (the trial terminated if a distractor was selected) criteria. However, we now also required participants to synchronize their cruise phase responses with an auditory metronome signal generated by the app. In all experimental blocks, this tempo was initially set at a moderate rate with an 800ms ITT between required responses. Across trials, the metronome tempo adapted to performance, with the gap between responses decreasing by 25 ms after 40 correct responses (i.e. a correct foraging trial), and increasing by 25 ms after two error trials. Errors were now also generated if participants deviated from the prescribed foraging tempo, by either going too fast or too slow (see Methods for details). The tempo constraints were relaxed at the start and end of the trial to account for non-cruise phase periods of responding.

As we were particularly interested in how tempo might modulate the ability to switch, we made one further important change to our usual procedure. Rather than leaving participants completely free to choose their foraging strategy, we specifically encouraged them to select targets in a “spatially efficient” manner. That is, we asked them to avoid using exhaustive runs -- where they would need to pass through the display at least twice – and to try and select all locally available targets until the increasing tempo made such switching impractical. Thus, we actively encouraged participants to switch between target categories. At very slow tempos, our review of previous studies (Thornton et al., 2019) and the soft constraint analysis described above (Gray et al., 2006) both suggest that adopting such a strategy would be possible under both feature and conjunction conditions. Our primary question was whether switching could be maintained as the tempo increased, and how this would vary as a function of condition.

We had two clear predictions. First, that the tendency to switch between target categories would generally reduce as foraging tempo increased. Second, that the influence of foraging tempo would be more pronounced during conjunction foraging than during feature foraging.

Methods

Participants

A total of 12 participants (Mean age = 25 years, SD = 5.4; 10 Female; 11 right handed) were recruited from the University of Iceland and the National Research University academic communities. Group sample size was determined prior to data collection and a detailed power analysis is provided below. All participants had normal or corrected to normal vision, were naïve as to the purpose of the research, and gave written informed consent before taking part in the experiment. All methods and procedures conformed to the Ethics and Data Protection Guidelines of the University of Malta, University of Iceland and National Research University.

Equipment

The stimuli were displayed on an iPad Air (Model A1474) with screen dimensions of 20 x 15 cm (24.6 cm diagonal) and an effective resolution of 1024 x 768 pixels at 132 ppi. The iPad was placed on a table in front of the participant in landscape mode. As viewing distance could only be approximately estimated at 50 cm, we report stimulus measures in both pixels and degrees visual angle. Stimulus presentation and response collection were carried out by a custom iPad application written in objective-C using Xcode and Cocos2d libraries.

Visual Stimuli.

The visual displays were identical to those described in our original iPad study (see Á. Kristjánsson et al., 2014 for details). Each trial presented a display containing 80 items randomly positioned within a 15 x 12 cm (approximately 17.1 x 13.7°) central viewing area. The overall layout was an invisible 10 x 8 grid, although individual items were jittered to reduce the sense of regularity. Gaps between rows and columns ensured that items never approached or occluded each other. In feature displays, the targets were red and green disks (20 pixels/0.46° visual angle) and the distractors were yellow and blue disks (counterbalanced across participants). In conjunction displays, the targets were red squares and green disks and the distractors were green squares and red disks (again counterbalanced). There were always 20 stimuli in each group, drawn on a black background. The overall spatial layout and the location of targets and distractors were generated independently on every trial.

Tempo Manipulation.

As soon as the visual display for each trial appeared, the auditory tempo cue also began. This consisted of a short (100 ms), high-frequency (fundamental freq = 2.4 kHz) mechanical tick, modelled on the sound used in Apple's "Hello Metronome" sample developer application. The sound was repeated at a regular temporal frequency, which varied across trials. Note that the selection of this particular tone was arbitrary, and we explored several sounds, selecting one that was minimally invasive, but clearly detectable at all temporal frequencies. The sound file is available on the OSF page associated with this paper at <https://osf.io/ej97c/>

During familiarization, the tempo for both conditions used an ITT of 1 second, intentionally very slow to give participants practice in synchronizing their responses. For both the feature and conjunction experimental blocks, the initial tempo used an ITT of 800 ms. After each correct trial (i.e. 40 correct responses), the ITT was reduced by 25 ms, so that the tempo gradually increased as the block of trials progressed. If participants had two consecutive error trials, the ITT was increased by 25 ms, slowing the tempo back into a range where successful trials had already been completed. Note that if participants completed all 25 trials without error, the final tempo would involve an ITT of 175 ms, which is almost certainly beyond human response rates. In reality, the fastest correct trial recorded had an ITT of 225 ms.

As our goal was to encourage participants to forage in synch with the metronome, rather than to directly measure the absolute temporal precision of their responses, we took a liberal approach when judging their adherence to the tempo. Beginning with the first correct selection, the app maintained a count of expected responses, incrementing the count with each auditory signal. This expected count was compared with the actual count of correct selection touches. If the actual count exceeded the expected count by more than 3, participants were judged to be responding too quickly. If the actual count lagged behind the expected count by more than 5, they were judged to be responding too slowly. We penalised faster responding more stringently to minimise any attempts to anticipate the auditory tone. Effectively, then, participants had the entire ITT period in which to make a response that was considered temporally correct, with this period shrinking as the tempo increased. Under this scheme, it would be possible for participants to speed up and slow down within a trial to keep the actual and expected counts synchronised. While a within trial analysis of precision might well identify such behaviour, here we looked only at the correlation between expected and actual ITTs collapsed across trials as an indicator of whether participants were attempting to adhere to the requested tempo (see Results below).

Procedure.

The task was to cancel all target items by tapping on them, while avoiding distractor items. Targets disappeared immediately following the tap. If participants selected one of the distractors the trial ended, an error message was given, and a new trial started. As already noted, we encouraged participants to switch between target categories, cancelling items in one local area, before proceeding through the display. This was suggested as a “spatially efficient” strategy, and was not enforced in any way. We also made it clear to participants that such a strategy might become impractical as tempo increased.

Participants were instructed to pay attention to the auditory tempo cue at the start of each trial, and to only begin responding once they had registered the new tempo. They were told to synchronize their responses to stay exactly in time with the auditory cue. If the cancellation rate ever exceeded the auditory tempo, the trial was terminated. Similarly, if the response rate lagged behind the tempo, the trial was also terminated. These tempo validation checks only began after the first response, allowing the participant to become familiar with the tempo, and were not implemented for the last 5 items, to take into account expected end-peaks in the response time functions (T. Kristjánsson, Thornton, et al., 2020). Each participant completed 25 trials of each condition (in counterbalanced order). One trial refers to a completed trial where all 40 targets have been cancelled.

Availability Statement.

A copy of the working app, the source code, analysis scripts and files are all available upon request. Raw data files and individual plots for all 12 participants have been uploaded as supplementary material to the Open Science Foundation (OSF) page associated with this paper at <https://osf.io/ej97c/>

Power Analysis.

As noted above, our sample size ($N = 12$) was determined prior to data collection, and was chosen to match previous studies from our group (e.g., Á. Kristjánsson et al., 2014; T. Kristjánsson, Thornton, et al., 2020; Thornton et al., 2019). Closely following the method adopted by Thornton et al., (2019), we conducted an *a priori* analysis to verify that this sample size provided sufficient power to detect within-subject differences between feature and conjunction foraging. Specifically, examination of the effect sizes in two previous studies (Jóhannesson et al., 2017; Á. Kristjánsson et al., 2014) yielded Cohen’s d_z calculations of 2.3 and 1.6, for the difference between the number of feature and conjunction runs. Using G*Power

3.1.9.4 (Faul et al., 2007), with assumed power of 0.95 and an alpha of 0.05, these effect size estimates suggested sample sizes of 5 and 8 participants, respectively.

As in Thornton et al., (2019), we opted to be conservative and run an additional 4 participants relative to the sample size suggested by the smaller effect size estimate. We did this as we believed smaller sample sizes might mask expected individual differences in other aspects of run behaviour. We were also unsure of the impact of tempo on the basic feature/conjunction manipulation. While the decision to run more participants than suggested by the *a priori* analysis could increase false discovery rates, as we had powered for an effect that has been replicated many times, we did not consider this a concern (e.g., Clarke et al., 2018; Á. Kristjánsson et al., 2014; Wolfe et al., 2019). Similarly, while using a more conservative expected power estimate of 0.8, rather than 0.95, may have been more appropriate, with the relatively large effect sizes from the previous studies, this would only have shifted sample size estimates down by one or two participants at most.

Subsequent to data collection, we also examined whether a different approach to *a priori* power analysis would have substantially altered our sample size estimates. As described by Anderson et al., (2017), the “Bias and Uncertainty Corrected Sample Size” (BUCCS) toolbox uses the reported F values and total sample size from previous factorial studies – rather than derived estimates of effect size – to generate necessary sample sizes for planned studies. As well as allowing the researcher to more easily focus on effects of interest (e.g., interactions), as the name indicates, BUCCS also corrects for other issues that can lead to underpowered studies (Anderson et al., 2017).

Here, we chose the previous study from our group (T. Kristjánsson et al., 2018) that most closely matched the current factorial design, while also involving temporal manipulations. That study included a 2 (Condition: feature/conjunction) x 4 (Time Limit: 20, 40, 60, 80) repeated measures analysis of variance and had a sample size was 17. Of most interest in the current context, was the significant Condition x Time Limit interaction, $F(2.2, 33.1) = 9.4$, $p < .001$, $\eta_p^2 = .39$.

We used this F value, along with the sample size and alpha parameter from T. Kristjánsson et al., (2018) as the basis for conducting *a priori* analysis using the BUCCS *ss.power.wa* function. We made use of the default settings of assumed alpha for the planned study = 0.05, level of assurance = 0.8, and desired power of 0.8. We specified the main factorial design of the current experiment – 2 (Condition) x 5 (Tempo) – and identified the interaction as the effect

of interest. This analysis yielded a necessary sample size of 10 participants, closely approximating our original choice.

Data Analysis.

Data from one participant (P9) was excluded from the main analysis as they were unwilling or unable to switch between target categories, even at the very slowest tempos. Their pattern of responses was thus completely different from any other participant, and we therefore treated them as an outlier. For the sake of completeness, we present their data during the evaluation of individual differences, and further discuss possible reasons for their approach to the task. The data analysis steps outlined below, were thus applied to the data for the remaining 11 participants.

The staircase method we adopted was designed to steadily increase the tempo as each block of trials progressed, up to a performance limit where successive responses could no longer be made accurately, a limit which was expected to vary from participant to participant. As an initial check to verify that participants were accurately adhering to the tempo prescribed on each trial, we first compared the expected and observed ITT as a function of Tempo and Condition. In addition to reporting the slope of these functions (i.e. correlation coefficients), we also calculated the average offset between actual and observed ITT, collapsed across Tempo. This should provide a clear indication of any consistent response delay, as function of Condition. We used separate paired t-tests to compare the slope and offset measures across Condition. We also used one-sample t-tests to assess whether offset measures were significantly different from zero.

Next, to more concisely summarise patterns across blocks we binned the 25 trials into 5 consecutive temporal bins. As each participant only completed one block per condition, dividing the tempo range into 5 sections with 5 repetitions per bin, also allowed us to better characterise average performance within a given tempo range. To confirm that these bins captured the increase in foraging speed, we conducted a 2 (Condition) x 5 (Trial Bin) repeated measures ANOVA on the average tempo values themselves.

While error rates were expected to be low, given the nature of the staircase procedure, they would also be crucial in determining the overall tempo profiles. Errors could either result from incorrect category selection, or from failing to maintain the designated tempo. We thus used a 2 (Condition) x 2 (Error Type) repeated measures ANOVA to examine the error rates.

Our main analyses concerned patterns of run behaviour. A “run” in this context simply refers to a sequence of selections from the same target category. Here, we focus on three specific

characteristics of run behaviour: i) the average number of runs per trial; ii) the average run length per trial; and iii) the proportion of trials within a block where run behaviour would be classified as non-random. We briefly discuss each of these in turn.

When participants select randomly from two target categories the total number of runs on a given trial should equal approximately the total targets divided by 2. Fewer runs per trial suggests participants are limiting selections based on category identity. As noted above, in our previous studies (Jóhannesson et al., 2016, 2017; Á. Kristjánsson et al., 2014), participants typically selected targets at random during feature foraging, but used exhaustive category selection (i.e. only two runs) during conjunction foraging trials. Here, we were specifically interested in how the number of runs changed as tempo increased.

The second characteristic of interest is the average run length on a given trial (total targets/number of runs). In our task, with a fixed number of targets on every trial, the average run length varies non-linearly as an inverse proportion of the number of runs. For equal increments in the number of runs across trials, there is thus a larger increase in run length when there are fewer runs than with many runs. This non-linearity has the potential to emphasise foraging patterns that are less obvious when only examining the number of runs. For example, previously, we have found it to be a useful indicator of individual differences (e.g., Á. Kristjánsson et al., 2014; Figure 4). We include it here as an additional probe into the nature of foraging tempo.

As an aside, the *variability* of run lengths within a trial may prove to be a particularly interesting measure to explore in future studies, in addition to or instead of the average run length. For example, with the current overall set size of 40 targets, a trial in which the two target categories are selected in 4 runs (i.e. T1, T2, T1, T2), the run lengths might be [8, 10, 12, 10] or [10, 10, 10, 10] or [5, 18, 15, 2], etc., etc. In each of these cases, the number of runs would be 4, and the average run length would be 10. However, the standard deviation would vary widely. For the sake of brevity, we do not report this measure here – there was no qualitative difference compared to average number of runs in current scenario – but we believe this observation may prove useful in future work.

Finally, while the relationship between number of runs and total targets can provide a useful heuristic to indicate whether participants select randomly from available target categories, there are additional techniques, such as the one sample runs test (Dawkins, 1971; Á. Kristjánsson et al., 2014) or random simulation (Wolfe et al., 2019), which can provide more precise quantitative measures. Here, we followed the method previously described in Á. Kristjánsson et al., (2014) to classify each individual trial. Specifically, we applied the one sample runs tests

to each individual trial, conducting 25 separate analyses per participant and using Bonferroni correction to adjust the level of alpha for multiple tests (i.e. $p < 0.002$). We then quantified the proportion of trials of that were non-random at this corrected alpha level for each participant.

For each of the above run measures, we conducted two basic analyses. First, to capture overall changes in behaviour as tempo increased, and to directly measure differences between feature and conjunction trials, we used the same 2 (Condition) x 5 (Trial Bin) repeated measures ANOVA described for the tempo data. Second, to more precisely measure the *change* in run behaviour as a function of tempo, we used linear regression to fit the data of individual participants. We report the average slopes of the run pattern x tempo functions, and compare them across condition using paired-samples t-tests. Note that as an initial examination of feature/conjunction block order showed no influence on any measures, we did not include this factor in our main analysis. Block order is recorded in the demographics information provided in the supplementary material along with the raw data.

Results

Figure 1a plots the relationship between actual and observed ITT across the whole tempo range as a function of Condition. It seems clear that participants adjusted their responses to stay in step with the required tempo. The correlation for both feature ($r = 0.99$) and conjunction ($r = 0.98$) conditions was almost perfect, and there was no difference between the slopes of these essentially parallel lines, $t(10) = 0.38$, n.s. (Figure 1b). However, both of these functions are shifted, indicating the presence of response delays. Specifically, the average observed ITTs for both the feature ($M = 9.3$ ms, $SE = 0.89$, $t(10) = 10.5$, $p < 0.001$) and conjunction ($M = 37.1$, $SE = 3.2$, $t(10) = 11.5$, $p < 0.001$) conditions were significantly longer than the requested ITTs (Figure 1c). There also appears to be fairly constant offset between them of approximately 28 ms, $t(10) = 8.7$, $p < 0.001$. Thus, while participants were able to adjust their ITTs as a function of Tempo in both conditions, they did so with varying levels of precision. The relatively constant increase in observed ITTs across tempo, and the clear difference between feature and conjunction conditions, suggests the source of these offsets may relate to the attentional demands of target detection, rather than tempo reproduction *per se*, although additional analysis -- looking at tempo precision within trials -- would be required to shed further light on this issue.

As described earlier, for the remainder of our analyses, we condensed the full tempo range by averaging the responses of each participant into 5 consecutive temporal bins. We first

wanted to verify that binning in this way still captured changes in tempo across each block of trials. Figure 2 shows how the average ITT systematically decreased across trial bin as expected, given the tempo staircase design. Note that the raw staircase data for each participant can be found in the Supplementary material. While there were main effects of both Condition, $F(1,10) = 24.2$, $p < 0.01$, $MSE = 0.007$, $\eta_p^2 = 0.71$, and Trial Bin, $F(1.8,18.0) = 1018$, $p < 0.001$, $MSE = 0.001$, $\eta_p^2 = 0.99$, these must be evaluated in the context of a significant Condition x Trial Bin interaction, $F(1.6,16.3) = 27.6$, $p < 0.001$, $MSE = 0.001$, $\eta_p^2 = 0.73$. From the same initial starting point, participants were able to reach consistently faster final (i.e. Bin 5) tempos during feature ($M = 284$ ms/item, $SE = 5$), as compared to conjunction ($M = 423$ ms/item, $SE = 20$), blocks of trials

The shallower tempo slope during the conjunction condition suggests that participants made more errors during these trials, and this is confirmed in Figure 3. There was a main effect of Condition, with far fewer total errors during feature ($M = 1.0$, $SE = 0.14$) than conjunction ($M = 4.2$, $SE = 0.58$) trials, $F(1,10) = 35.6$, $p < 0.001$, $MSE = 3.2$, $\eta_p^2 = 0.78$. There was also a main effect of Error Type, with more category selection errors ($M = 4.2$, $SE = 0.79$), than tempo related errors ($M = 1.0$, $SE = 0.25$), $F(1,10) = 10.3$, $p < 0.01$, $MSE = 10.5$, $\eta_p^2 = 0.51$. Both of these main effects need to be interpreted within the context of the significant Error Type x Condition interaction that can be clearly seen in Figure 3, where the overall pattern of errors is dominated by the selection errors during the conjunction condition, $F(1,10) = 6.2$, $p < 0.05$, $MSE = 6.2$, $\eta_p^2 = 0.38$. Note that the error rates reported in Figure 3 are for the entire block of trials, not per trial. As overall error rates were so low – a finding consistent with our previous studies – it was not feasible to examine possible speed-accuracy trade-offs as a function of tempo.

Figure 4 illustrates the relationship between the average number of runs and average tempo within each trial bin for both feature and conjunction conditions. There are three clear patterns. First, the number of runs consistently reduces as tempo increases in both feature and conjunction conditions. Second, participants appear to use substantially more runs during feature foraging than conjunction foraging, even at the slowest tempo, a difference maintained across all subsequent trial bins. Third, there is noticeably more variability in performance during conjunction foraging compared to feature foraging, both in terms of average number of runs and average tempo within a given trial bin. We return to these individual differences shortly.

The 2 (Condition) x 5 (Trial Bin) ANOVA used to examine changes in the number of runs during the course of a block confirmed these observations. There was a main effect of Condition, with more runs during feature foraging ($M = 16.1$, $SE = 0.49$) than conjunction foraging ($M = 9.8$, $SE = 0.48$), $F(1,10) = 122.6$, $p < 0.001$, $MSE = 9.1$, $\eta_p^2 = 0.93$. There was also a main effect of Trial Bin, $F(4,40) = 58.3$, $p < 0.001$, $MSE = 8.4$, $\eta_p^2 = 0.85$. The Trial Bin x Condition interaction, however, was not significant, $F(2.4, 24.4) = 1.0$, n.s., $MSE = 12.2$, $\eta_p^2 = 0.09$. Consistent with this latter finding, and the average trend line shown in Figure 4, our slope analysis indicated that there was no difference in the rate of reduction in the number of runs as a function of tempo between the feature ($M = -2.63$ runs/100 ms, $SE = 0.33$) and conjunction ($M = -3.22$ runs/100 ms, $SE = 0.38$) conditions, $t(10) = 1.5$, n.s.

Figure 5 shows the pattern of run length data. There was a main effect of Condition, $F(1,10) = 40.0$, $p < 0.001$, $MSE = 6.3$, $\eta_p^2 = 0.8$, with consistently shorter run lengths during feature foraging ($M = 3.3$, $SE = 0.25$) than conjunction foraging ($M = 6.4$, $SE = 0.45$). There was also a main effect of Trial Bin, $F(1.6,15.8) = 31.5$, $p < 0.001$, $MSE = 10.9$, $\eta_p^2 = 0.76$. In contrast to the number of runs, here there was a significant Condition x Trial Bin interaction, $F(1.9,18.7) = 4.2$, $p < 0.05$, $MSE = 8.9$, $\eta_p^2 = 0.3$. Direct comparisons of the run length x tempo slopes, confirmed that there was a more rapid increase in run length during conjunction ($M = 2.37$ items/100 ms, $SE = 0.5$) than feature ($M = 0.08$ items/100 ms, $SE = 0.02$) trial blocks, $t(10) = 3.4$, $p < 0.01$.

Figure 6 shows how the proportion of non-random run behaviour changes during a block, as a function of Condition and Tempo. For the feature condition ($M = 0.24$, $SE = 0.04$), run behaviour was exclusively random for the two earliest trial bins. Although non-random foraging becomes more prevalent in the latter stages of the block, there is considerable between-subject variability, and even at the fastest tempos proportions barely exceed 50%. In contrast, during conjunction foraging ($M = 0.69$, $SE = 0.05$), approximately 25% of trials are non-random even at the slowest tempos, rising quickly towards 100% in the latter two bins.

As with the other run measures, these patterns gave rise to main effects of both Condition, $F(1,10) = 59.7$, $p < 0.001$, $MSE = 0.07$, $\eta_p^2 = 0.86$, and Trial Bin, $F(4,40) = 43.8$, $p < 0.001$, $MSE = 0.04$, $\eta_p^2 = 0.81$. Although the Condition x Trial Bin interaction was not significant, $F(4,40) = 2.0$, n.s., $MSE = 0.06$, $\eta_p^2 = 0.17$, the slope analysis did indicate a consistently more rapid increase in the proportion of non-random trials as a function of tempo for the conjunction condition ($M = 0.23$ units/100 ms, $SE = 0.02$), compared to the feature condition ($M = 0.14$ units/100 ms, $SE = 0.02$), $t(11) = 2.4$, $p < 0.05$.

Individual Differences

As noted in the Introduction, in our previous work we have consistently found individual differences in how participants adapt their run behaviour in response to increased attentional load. Specifically, while the majority of participants change from random category selection during feature foraging to exhaustive category selection during conjunction foraging, a subset of participants – so-called super-foragers – maintain random behaviour throughout. Here, we were interested if participants varied in terms of the influence of tempo on their foraging behaviour.

The individual data plots for all observers and dependent measures are provided in the supplementary material. In this section, we illustrate what we consider to be the major individual differences, using example participant data from the analysis of average run length. In our previous work, this dependent variable had provided the most stable estimates of individual differences (see Á. Kristjánsson et al., 2014; Figure 4) and in the current work, it also appeared most sensitive to changes in tempo.

Participant 7 (upper left in Figure 7; see also participants 1, 4, 5, 10 & 12 in supplementary figures) nicely illustrates the canonical pattern (Figure 5), with increasing run length as a function of increasing tempo, with the rate of increase being steeper for the conjunction condition. Thus, the majority of participants do appear to show the expected pattern whereby attentional load and the demands of faster responding interact.

Participant 9 (upper right in Figure 7), was excluded from the main analysis, as they showed a completely unique pattern of consistent switching during the feature condition, and consistent exhaustive runs during the conjunction condition. As we had explicitly asked participants to attempt to switch between categories during the slow tempo portions of each block, this suggests that the participant was unwilling or unable to follow these instructions. Examination of the error data for this participant showed that they did not make any selection errors at all, suggesting that their performance was not mediated by difficulty in switching.

Participant 8 (see also participant 2 in supplementary files) shows little evidence of a systematic increase of run length with tempo. Rather, both profiles are relatively flat, but the performance in the conjunction condition is more variable, and perhaps more interestingly, compressed towards the lower end of the tempo range. This suggests that the participant

attempted to frequently switch during conjunction trials, but was not very successful. Consistent with this idea, both participants 8 and 2 had relatively high selection error rates.

Participant 11 on the other hand (see also participants 3 & 6 in the supplementary files) appears to have been much more successful in maintaining a pattern of conjunction switching across a wider range of tempos. The run length remains low across all trial bins, and there is a close parallel between feature and conjunction conditions. This suggests that these participants were able to cope with the dual demands of increased target complexity and increased tempo. These participants thus behaved in a manner consistent with the super-foragers from our previous studies (e.g., Jóhannesson et al., 2017; Á. Kristjánsson et al., 2014).

Discussion

In two recent papers, we suggested that foraging tempo -- the rate of successive target responses within a trial -- helps determine patterns of run behaviour in multiple target search (Prpic et al., 2019; Thornton et al., 2019). Specifically, our analysis of previous studies showed that the tendency to switch randomly between target categories was more common when the interval between responses was longer. Conversely, the tendency to use extended selection from a single category – the hallmark of attentionally demanding conjunction foraging – was more pronounced when successive responses were made rapidly. One simple explanation for these findings would be that reduced time to respond adjusts the soft constraints that are applied when selecting between available cognitive operations (Gray et al., 2006; Gray & Boehm-Davis, 2000; Gray & Fu, 2004). At faster speeds, the temporal cost of switching categories is avoided, affecting the overall pattern of foraging.

In the current paper, we directly tested this idea, asking participants to complete our standard iPad foraging task (Á. Kristjánsson et al., 2014) with the additional requirement of synchronising their responses to an auditory metronome signal, also produced by the app. Across trials, we increased the tempo of this metronome signal and examined patterns of runs as a function of both tempo and attentional load (i.e. feature versus conjunction foraging). Our main finding was very clear: changes in foraging tempo were accompanied by systematic changes in run behaviour.

In the Introduction, we made two specific predictions concerning foraging tempo. Below, we address these in turn, before examining other aspects of the data and drawing some general conclusions. First, we predicted that the tendency to switch between target categories would generally reduce as foraging tempo increased. If observers decrease their switching, the number

of runs should decrease and the run length increase with tempo. Similarly, the tendency to use non-random selection from target categories should increase as tempo increases. These are exactly the patterns we see in Figures 3-5, and in the robust main effects of Tempo in all of the related analyses.

The finding that foraging tempo does modulate patterns of runs in this way has both theoretical and practical implications. To begin with, these results provide a further example of an interactive behaviour in which temporal factors play an important role in shaping overall performance, consistent with the soft constraint approach suggested by Gray and colleagues (Gray et al., 2006; Gray & Boehm-Davis, 2000; Gray & Fu, 2004). In terms of foraging behaviour, while we had previously argued that the appearance of exhaustive run patterns in humans was solely a consequence of changing attentional demands (Á. Kristjánsson et al., 2014), this explanation appears to be incomplete. At least in humans, increases in attentional complexity—such as the standard feature/conjunction manipulation—do not appear sufficient to trigger a complete shift from freely selecting amongst all available “prey” types, towards foraging for a single category. Rather, to observe this change in run behaviour, increases in attentional demand need to be accompanied by the tendency to select/respond rapidly. While formal attempts to model the role of limited attention during foraging, for example the “attentive prey” model (Dukas, 2002, 2004; Dukas & Ellner, 1993), may be able to indirectly capture some aspects of this attention x time interaction—at least for conspicuous/cryptic prey manipulations (Gendron & Staddon, 1983)—it may be more useful to directly integrate the consequences of time constraints, along the lines suggested by the soft constraints approach (Gray et al., 2006; Gray & Boehm-Davis, 2000; Gray & Fu, 2004).

From a more practical perspective, the current results show that manipulating the time available for making selections is another useful experimental method for modulating foraging behaviour in humans. Indeed, manipulating foraging tempo would appear to give rise to graded changes in run behaviour, rather than to the random/exhaustive dichotomy we have previously observed. The virtual absence of exhaustive category selection during conjunction foraging in the current dataset is almost certainly caused by our specific task instructions—a point we return to shortly—but more importantly, tempo manipulations do seem to provide a finer level of resolution for exploring the impact of increasing task demands on foraging behaviour, particularly in the context of individually determined thresholds of performance.

Furthermore, when designing new tasks, factors that can affect the rate of responding—such as response modality (e.g., touch versus mouse) or the spatial separation between items—should clearly be borne in mind, as these may constrain run behaviour via the influence of

foraging tempo (see Thornton et al., 2019 for further discussion). Similarly, when examining run behaviour in a given context, examination of response rates may shed additional light on general task demands (Gray et al., 2006; Gray & Boehm-Davis, 2000; Gray & Fu, 2004).

Our second prediction was that the influence of foraging tempo would be more pronounced during conjunction foraging than during feature foraging. Here, the pattern of results was less clear. Examination of Figure 4 suggests that the change in the number of runs as a function of tempo was very similar between the two conditions. Even at the slowest tempo, participants consistently use fewer runs during conjunction foraging, and this gap is maintained across all time bins, giving rise to very similar tempo slopes in the two conditions. In contrast, the pattern of run length data in Figure 5 and non-random behaviour in Figure 6, both suggest that changes in foraging tempo have an earlier impact on patterns of responding during conjunction than feature foraging.

It thus appears that the interaction between the feature/conjunction manipulation and foraging tempo is not as simple we had predicted. However, this in itself raises several interesting points. First, the lack of an interaction when considering the number of runs appears to reflect the fact that foraging tempo had a stronger influence on feature foraging than we had anticipated. Increasing time pressure may thus be a simple way to change patterns of foraging without the need to vary target type. Second, while the number of runs is a simple and intuitive dependent variable, it may not always capture more subtle changes in behaviour. Here, for example, the very clear interaction evident in the run length data (Figure 5), is absent when only considering the number of runs (Figure 4). This reiterates the need to consider a range of dependent measures when exploring foraging behaviour.

Another aspect of performance in the current experiment were clear individual differences in response to variations in foraging tempo. Specifically, we found that some observers seemed to be less affected than others by the decreasing time limit, and these individuals seem to show performance patterns reminiscent of the participants that we called super-foragers in our previous work (Jóhannesson et al., 2017; Á. Kristjánsson et al., 2014). Taking participant 11 as an example (Figure 7), there was little difference between feature and conjunction run patterns, the overall tempo range was similar, and the impact of tempo on run patterns was very modest.

We should note that in our original study, the feature and conjunction behaviour of super-foragers was completely indistinguishable (see Á. Kristjánsson et al., 2014; Figure 4, participants 4, 8, 11 & 14). Here, this is not the case. Thus, although the impact of tempo for some individuals is modest, it may still add diagnostic power over and above the basic feature

and conjunction manipulation. This may prove useful in terms of identifying underlying mechanisms. For example, while previous studies with children have found clear links between foraging behaviour and other, more general cognitive abilities (Ólafsdóttir et al., 2016, 2019), this has not been the case with studies of adults (Clarke et al., 2018; Jóhannesson et al., 2017). We suggest that exploring links with tasks that emphasize temporal constraints may thus be a useful avenue for future studies of individual differences in foraging behaviour.

Remaining with the topic of individual differences, it is important to remember that in the current task we explicitly asked participants to avoid using extended runs of selections from a single category. That is, we directly encouraged switching behaviour. We did this as we were interested in monitoring at what tempo participants would no longer be able to switch between target categories and having ceiling levels of performance at the outset in one of the primary measures (i.e. number of runs) was clearly not desirable.

Our subsequent data (see Figure 4) indicates that participants were able to switch quite effectively during conjunction conditions, when encouraged to do so. So much so, in fact, that there was only very limited evidence of fully exhaustive run behaviour, even at the very fastest tempos (see individual participants plots, supplementary figures). This strongly suggests that the use of exhaustive runs in the conjunction condition of our original study was not due to participants being unable to switch. Rather it suggests that they either strategically chose not to switch or some other aspect of their behaviour constrained their ability to switch. What might have caused this pattern of behaviour?

The most obvious explanation, based on the soft constraints framework already discussed above (Gray et al., 2006; Gray & Boehm-Davis, 2000; Gray & Fu, 2004), is that participants choosing an exhaustive strategy were prioritising response speed. That is, if these participants were attempting to respond close to the upper limit imposed by their perceptual-motor system, then this may have required cognitive operations that were as simple (i.e. fast) as possible, essentially precluding the use of template switching. Consistent with this idea, examination of response time from our original study (Á. Kristjánsson et al., 2014; Figure 3) shows that the six fastest participants all used exhaustive runs during conjunction foraging.

A second possibility is that these observers were more averse to making errors. During conjunction foraging, selection errors are much more likely to occur than during feature foraging (see Figure 3). Participants who want to minimize errors – either because they have prioritized accuracy or possibly because they become frustrated with having to repeat trials -- would clearly do better by adopting an exhaustive strategy. Consistent with this idea, Tagu & Kristjánsson (2020) recently noted that during eye-gaze foraging (Jóhannesson et al., 2016)

error rates were lower for participants who adopted exhaustive search than those that would typically be classified as super-foragers. We should note, however, that in a recent study that compared performance with and without terminal error feedback, we found little impact of this factor on overall patterns of foraging (Thornton et al., 2019).

Approaching this issue from the other side, we can also ask why some participants in each of our studies – the super-foragers – do appear so comfortable with consistently switching during conjunction conditions? There is now strong evidence the participants can routinely maintain more than one target template in working memory, and are able to selectively activate or deactivate items depending on temporal expectations of task relevance (Grubert & Eimer, 2018, 2020; see also T. Kristjánsson & Kristjánsson, 2018). It seems safe to assume that during the course of a multiple-target trial, repeatedly switching activation between two conjunction templates in working memory would require more cognitive resources than maintaining activation of just one of those templates. Thus, participants who switch during conjunction conditions are choosing to place more demands on the cognitive system, than those who do not.

As noted in the Introduction, an important feature of the soft constraints approach to interactive behaviours is that individuals are able to adjust the balance between cognitive, perceptual and motor operations in order to minimize performance costs. It seems likely that such adjustments could take into account individual differences in each of these systems (e.g., working memory capacity, visual acuity, motor fluency) as well as allowing constraints to be ‘overridden by factors such as training or by deliberately adopted top-down strategies’ (Gray et al., 2006; p.463). Our original idea was that super-foragers differed from typical foragers in respect to basic cognitive abilities, such as working memory capacity or selective attention. As already mentioned, however, we have thus far not been able to identify such differences, at least in adults (Jóhannesson et al., 2017). It remains possible that differences can be found in other cognitive abilities that might more directly relate to switching, such as flexibility, or that the source of individual differences are in the perceptual or motor domains. If super-foragers have better motor fluency or more precise hand-eye coordination, for example, these could conceivably change the balance of resources across domains, freeing up cognition to handle more complex operations within a similar time frame.

In terms of top-down strategies, those participants who continue to switch during conjunction conditions may be placing more emphasis on reducing their overall movement across the display, rather than reducing time. Although we chose not to focus on inter-target distance (ITD) or overall movement in the current study, simply for the sake of brevity, our

previous work has shown that there is a very strong correlation between run behaviour and spatial efficiency. That is, the use of fewer, longer runs requires travelling further distances. In our original study, for example, the correlation between the number of runs and the distance travelled was 0.87 and 0.92 for feature and conjunction conditions, respectively (Á. Kristjánsson et al., 2014; Figure 3). Here, we've clearly shown that run behaviour changes as expected as a function of tempo, with fewer runs being used as tempo increases (Figure 4). Consistent with this pattern, there were also strong correlations between tempo and distance travelled, both for feature (0.80) and conjunction (0.83) conditions. Returning to the more general issue of individual differences, it does thus seem possible that switching could be used as a strategy by those individuals who chose to prioritize spatial efficiency over minimizing time.

Of course, it is also possible that those participants who spontaneously choose to switch during conjunction conditions are simply more engaged with the task, thus affecting their levels of arousal, vigilance or alertness (Fernandez-Duque & Posner, 1997; Kahneman, 1973; Mackworth, 1970). For example, in another variant of our iPad task, we allowed participants to respond freely, but varied the overall duration of trials (T. Kristjánsson et al., 2018). When observers had only 5, 10 or 15 s to select as many targets as they could, they tended to switch far more between target types than when they had unlimited time. We suggested that adding time pressure in this way changed their levels of concentration, facilitating switching, which in turn enabled them to select more targets than they otherwise could have with extended runs, given the spatial layout of the display. In other ongoing work in our lab, we have started to explore how the risk of simulated predation affects patterns of human foraging (for an online example of these tasks, see <https://maltacogsci.org/thePredationGame/>). Consistent with the idea of a link between levels of alertness/arousal and foraging patterns, participants who were actively “hunted” by wolf objects as they foraged were significantly more likely to switch during conjunction conditions than those for whom the wolves just served as visual distractions (Thornton et al., 2020).

Finally, another approach that might yield useful insights into the nature of individual differences would be to examine whether patterns of behaviour remain stable across different forms of foraging. Several lines of evidence now support the existence of “domain general” search strategies (Hills et al., 2008) that apply not only to movement through an environment to obtain resources – as we have modelled in our task – but also to both the internal and external search for information (Fu & Gray, 2006; Hills & Dukas, 2012; Pirolli & Card, 1999; Van den Driessche et al., 2019; see Todd & Hills, 2020 for a recent overview). Examples of internal

search for information would be recalling concepts from semantic memory (Hills et al., 2012), finding anagram solutions (Hills et al., 2010) or finding mathematical solutions (Hills, 2013). Examples of external search for information would include scanning the environment for signs of predators or prey (Hills & Dukas, 2012), navigating the internet (Pirolli & Card, 1999) or making use of interactive visualization techniques (Rensink, 2014). Behaviour across these search domains has been shown to be linked via priming studies (e.g., Hills et al., 2008, 2010) and there is also initial evidence that individual differences may be preserved across internal and external tasks (Van den Driessche et al., 2019). It would clearly be an interesting future direction to examine how the task demands of attention and tempo we have studied here impact the search for information, and whether similar individual differences in response to such demands, if they exist, transfer across domains.

Conclusion

In the current study, we have demonstrated that there is a direct link between foraging tempo and run-based patterns seen during multiple target search. Foraging tasks are increasingly being used to measure visual attention and working memory and have proven useful in identifying stable individual differences. Here, even though we explicitly encouraged all participants to switch between target categories under the more demanding conjunction condition, we still found substantial individual differences in performance. Further research is required to fully characterize these differences and to identify the strategies or mechanisms that give rise to them. More generally, our results suggest that future studies should consider how the temporal demands of a given task impact expected outcomes, and that manipulating foraging tempo could serve as a simple but effective way to manipulate group and individual strategies during human foraging.

Acknowledgments

The authors would like to thank Jérôme Tagu, Sunčica Zdravković, Ron Rensink and an anonymous reviewer for helpful comments and suggestions. We are particularly grateful to Ron Rensink for the idea of using “soft constraints” to help interpret these findings. IMT’s research is supported by the Academic Work Resource Fund and the University Research Fund of the University of Malta. AK was supported by the Icelandic Research Fund (grant #152427) as well as a grant from the Research Fund of the University of Iceland.

REFERENCES

- Anderson, S. F., Kelley, K., & Maxwell, S. E. (2017). Sample-Size Planning for More Accurate Statistical Power: A Method Adjusting Sample Effect Sizes for Publication Bias and Uncertainty. *Psychological Science*, 28(11), 1547–1562.
<https://doi.org/10.1177/0956797617723724>
- Basoudan, N., Torrens-Burton, A., Jenkins, A., Thornton, I. M., Hanley, C., Tree, J. J., Thomas, S., & Tales, A. (2019). Sequential Information Processing: The “Elevated First Response Effect” Can Contribute to Exaggerated Intra-Individual Variability in Older Adults. *The Yale Journal of Biology and Medicine*, 92(1), 13–20.
- Bond, A. B. (1983). Visual search and selection of natural stimuli in the pigeon: the attention threshold hypothesis. *Journal of Experimental Psychology: Animal Behavior Processes*, 9(3), 292.
- Bond, A. B., & Riley, D. A. (2010). Searching Image in the Pigeon: A Test of Three Hypothetical Mechanisms. *Ethology*, 87(3–4), 203–224.
<https://doi.org/10.1111/j.1439-0310.1991.tb00247.x>
- Burkhardt, J. F. (1998). Individual flexibility and tempo in the ant, *Pheidole dentata*, the influence of group size. *Journal of Insect Behavior*, 11(4), 493–505.
- Cain, M. S., Vul, E., Clark, K., & Mitroff, S. R. (2012). A Bayesian Optimal Foraging Model of Human Visual Search. *Psychological Science*, 23(9), 1047–1054.
<https://doi.org/10.1177/0956797612440460>
- Clarke, A. D. F., Irons, J., James, W., Leber, A. B., & Hunt, A. R. (2018). *Stable individual differences in strategies within, but not between, visual search tasks*.
<https://doi.org/10.31234/osf.io/bqa5v>

- Davidson, D. W. (1997). The role of resource imbalances in the evolutionary ecology of tropical arboreal ants. *Biological Journal of the Linnean Society*, 61(2), 153–181. <https://doi.org/10.1111/j.1095-8312.1997.tb01785.x>
- Dawkins, M. (1971). Shifts of “attention” in chicks during feeding. *Animal Behaviour*, 19(3), 575–582. [https://doi.org/10.1016/S0003-3472\(71\)80114-8](https://doi.org/10.1016/S0003-3472(71)80114-8)
- Dukas, R. (2002). Behavioural and ecological consequences of limited attention. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, 357(1427), 1539–1547.
- Dukas, R. (2004). Causes and Consequences of Limited Attention. *Brain, Behavior and Evolution*, 63(4), 197–210. <https://doi.org/10.1159/000076781>
- Dukas, R., & Ellner, S. (1993). Information processing and prey detection. *Ecology*, 74(5), 1337–1346.
- Dyer, F. C., & Seeley, T. D. (1991). Nesting Behavior and the Evolution of Worker Tempo in Four Honey Bee Species. *Ecology*, 72(1), 156–170. <https://doi.org/10.2307/1938911>
- Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G*Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior Research Methods*, 39(2), 175–191. <https://doi.org/10.3758/BF03193146>
- Fernandez-Duque, D., & Posner, M. I. (1997). Relating the mechanisms of orienting and alerting. *Neuropsychologia*, 35(4), 477–486. [https://doi.org/10.1016/S0028-3932\(96\)00103-0](https://doi.org/10.1016/S0028-3932(96)00103-0)
- Fougnie, D., Cormiea, S. M., Zhang, J., Alvarez, G. A., & Wolfe, J. M. (2015). Winter is coming: How humans forage in a temporally structured environment. *Journal of Vision*, 15(11), 1–1.
- Fu, W., & Gray, W. (2006). Suboptimal tradeoffs in information seeking. *Cognitive Psychology*, 52(3), 195–242. <https://doi.org/10.1016/j.cogpsych.2005.08.002>

- Gendron, R. P., & Staddon, J. E. R. (1983). Searching for Cryptic Prey: The Effect of Search Rate. *The American Naturalist*, 121(2), 172–186. <https://doi.org/10.1086/284049>
- Gilchrist, I. D., North, A., & Hood, B. (2001). Is visual search really like foraging? *Perception*, 30(12), 1459–1464.
- Gray, W. D., & Boehm-Davis, D. A. (2000). Milliseconds matter: An introduction to microstrategies and to their use in describing and predicting interactive behavior. *Journal of Experimental Psychology: Applied*, 6(4), 322–335. <https://doi.org/10.1037/1076-898X.6.4.322>
- Gray, W. D., & Fu, W.-T. (2004). Soft constraints in interactive behavior: the case of ignoring perfect knowledge in-the-world for imperfect knowledge in-the-head. *Cognitive Science*, 28(3), 359–382. https://doi.org/10.1207/s15516709cog2803_3
- Gray, W. D., Sims, C. R., Fu, W.-T., & Schoelles, M. J. (2006). The soft constraints hypothesis: A rational analysis approach to resource allocation for interactive behavior. *Psychological Review*, 113(3), 461–482. <https://doi.org/10.1037/0033-295X.113.3.461>
- Grubert, A., & Eimer, M. (2018). The Time Course of Target Template Activation Processes during Preparation for Visual Search. *The Journal of Neuroscience*, 38(44), 9527–9538. <https://doi.org/10.1523/JNEUROSCI.0409-18.2018>
- Grubert, A., & Eimer, M. (2020). Preparatory Template Activation during Search for Alternating Targets. *Journal of Cognitive Neuroscience*, 1–11. https://doi.org/10.1162/jocn_a_01565
- Guilford, T., & Dawkins, M. S. (1987). Search images not proven: A reappraisal of recent evidence. *Animal Behaviour*, 35(6), 1838–1845. [https://doi.org/10.1016/S0003-3472\(87\)80076-3](https://doi.org/10.1016/S0003-3472(87)80076-3)

- Guilford, T., & Dawkins, M. S. (1989a). Search image versus search rate: a reply to Lawrence. *Animal Behaviour*, 37, 160–162. [https://doi.org/10.1016/0003-3472\(89\)90019-5](https://doi.org/10.1016/0003-3472(89)90019-5)
- Guilford, T., & Dawkins, M. S. (1989b). Search image versus search rate: Two different ways to enhance prey capture. *Animal Behaviour*, 37, 163–165. [https://doi.org/10.1016/0003-3472\(89\)90021-3](https://doi.org/10.1016/0003-3472(89)90021-3)
- Heinrich, B., Mudge, P. R., & Deringis, P. G. (1977). Laboratory analysis of flower constancy in foraging bumblebees: *Bombus ternarius* and *B. terricola*. *Behavioral Ecology and Sociobiology*, 2(3), 247–265. <https://doi.org/10.1007/BF00299738>
- Hills, T. T. (2013). Investigating mathematical search behavior using network analysis. In R. Lesh, P. L. Galbraith, C. R. Haines, & A. Hurford (Eds.), *Modeling Students' Mathematical Modeling Competencies* (pp. 571–581). Springer Netherlands. <https://doi.org/10.1007/978-94-007-6271-8>
- Hills, T. T., & Dukas, R. (2012). The evolution of cognitive search. In P. M. Todd, T. T. Hills, & T. W. Robbins (Eds.), *Cognitive search: evolution, algorithms, and the brain* (pp. 11–24). MIT Press.
- Hills, T. T., Jones, M. N., & Todd, P. M. (2012). Optimal foraging in semantic memory. *Psychological Review*, 119(2), 431.
- Hills, T. T., Kalff, C., & Wiener, J. M. (2013). Adaptive Lévy processes and area-restricted search in human foraging. *PLoS One*, 8(4), e60488.
- Hills, T. T., Todd, P. M., & Goldstone, R. L. (2008). Search in External and Internal Spaces: Evidence for Generalized Cognitive Search Processes. *Psychological Science*, 19(8), 802–808. <https://doi.org/10.1111/j.1467-9280.2008.02160.x>
- Hills, T. T., Todd, P. M., & Goldstone, R. L. (2010). The central executive as a search process: Priming exploration and exploitation across domains. *Journal of*

- Experimental Psychology: General*, 139(4), 590–609.
<https://doi.org/10.1037/a0020666>
- Hollis, K. L. (1989). In search of a hypothetical construct: A reply to Guilford & Dawkins. *Animal Behaviour*, 37, 162–163. [https://doi.org/10.1016/0003-3472\(89\)90020-1](https://doi.org/10.1016/0003-3472(89)90020-1)
- Hulleman, J., & Olivers, C. N. (2017). On the brink: The demise of the item in visual search moves closer. *Behavioral and Brain Sciences*, 40.
- Jackson, R. R., & Li, D. (2004). One-encounter search-image formation by araneophagic spiders. *Animal Cognition*, 7(4), 247–254. <https://doi.org/10.1007/s10071-004-0219-x>
- Jóhannesson, Ó. I., Kristjánsson, Á., & Thornton, I. M. (2017). Are foraging patterns in humans related to working memory and inhibitory control? *Japanese Psychological Research*, 59(2), 152–166.
- Jóhannesson, Ó. I., Thornton, I. M., Smith, I. J., Chetverikov, A., & Kristjánsson, Á. (2016). Visual Foraging With Fingers and Eye Gaze. *I-Perception*, 7(2).
<https://doi.org/10.1177/2041669516637279>
- Kahneman, D. (1973). *Attention and effort*. Prentice-Hall.
- Kamil, A. C., & Bond, A. B. (2006). Selective attention, priming, and foraging behavior. *Comparative Cognition: Experimental Explorations of Animal Intelligence*, 106–126.
- Kristjánsson, Á., & Egeth, H. (2020). How feature integration theory integrated cognitive psychology, neurophysiology, and psychophysics. *Attention, Perception, & Psychophysics*, 82(1), 7–23. <https://doi.org/10.3758/s13414-019-01803-7>
- Kristjánsson, Á., Jóhannesson, Ó. I., & Thornton, I. M. (2014). Common Attentional Constraints in Visual Foraging. *PLoS ONE*, 9(6), e100752.
<https://doi.org/10.1371/journal.pone.0100752>

- Kristjánsson, Á., Ólafsdóttir, I. M., & Kristjánsson, T. (2019). *Visual Foraging Tasks Provide New Insights into the Orienting of Visual Attention: Methodological Considerations*. Humana Press. https://doi.org/10.1007/7657_2019_21
- Kristjánsson, T., Draschkow, D., Pálsson, Á., Haraldsson, D., Jónsson, P. Ö., & Kristjánsson, Á. (2020). Moving foraging into 3D: Feature versus conjunction-based foraging in virtual reality. *Quarterly Journal of Experimental Psychology*, 174702182093702. <https://doi.org/10.1177/1747021820937020>
- Kristjánsson, T., & Kristjánsson, Á. (2018). Foraging through multiple target categories reveals the flexibility of visual working memory. *Acta Psychologica*, 183, 108–115.
- Kristjánsson, T., Thornton, I. M., Chetverikov, A., & Kristjánsson, Á. (2020). Dynamics of visual attention revealed in foraging tasks. *Cognition*, 194, 104032. <https://doi.org/10.1016/j.cognition.2019.104032>
- Kristjánsson, T., Thornton, I. M., & Kristjánsson, Á. (2018). Time limits during visual foraging reveal flexible working memory templates. *Journal of Experimental Psychology: Human Perception and Performance*, 44(6), 827.
- Lawrence, E. S. (1989). Why blackbirds overlook cryptic prey: Search rate or search image? *Animal Behaviour*, 37, 157–160. [https://doi.org/10.1016/0003-3472\(89\)90018-3](https://doi.org/10.1016/0003-3472(89)90018-3)
- Mackworth, J. F. (1970). *Vigilance and attention: a signal detection approach*. Penguin.
- Ólafsdóttir, I. M., Gestsdóttir, S., & Kristjánsson, Á. (2019). The role of executive functions in foraging throughout development. *Journal of Vision*, 19(10), 234b–234b.
- Ólafsdóttir, I. M., Kristjánsson, T., Gestsdóttir, S., Jóhannesson, Ó. I., & Kristjánsson, Á. (2016). Understanding visual attention in childhood: Insights from a new visual foraging task. *Cognitive Research: Principles and Implications*, 1(1). <https://doi.org/10.1186/s41235-016-0016-5>

- Oster, G. F., & Wilson, E. O. (1978). *Caste and ecology in the social insects*. Princeton University Press.
- Pellicano, E., Smith, A. D., Cristino, F., Hood, B. M., Briscoe, J., & Gilchrist, I. D. (2011). Children with autism are neither systematic nor optimal foragers. *Proceedings of the National Academy of Sciences*, 108(1), 421–426.
- Pietrewicz, A. T., & Kamil, A. C. (1979). Search Image Formation in the Blue Jay (*Cyanocitta cristata*). *Science*, 204(4399), 1332–1333.
<https://doi.org/10.1126/science.204.4399.1332>
- Pirolli, P., & Card, S. (1999). Information foraging. *Psychological Review*, 106(4), 643–675.
<https://doi.org/10.1037/0033-295X.106.4.643>
- Plaisted, K. C., & Mackintosh, N. J. (1995). Visual search for cryptic stimuli in pigeons: implications for the search image and search rate hypotheses. *Animal Behaviour*, 50(5), 1219–1232. [https://doi.org/10.1016/0003-3472\(95\)80039-5](https://doi.org/10.1016/0003-3472(95)80039-5)
- Prpic, V., Kniestedt, I., Camilleri, E., Maureira, M. G., Kristjánsson, Á., & Thornton, I. M. (2019). A serious game to explore human foraging in a 3D environment. *PLOS ONE*, 14(7), e0219827. <https://doi.org/10.1371/journal.pone.0219827>
- Reid, P. J., & Shettleworth, S. J. (1992). Detection of cryptic prey: Search image or search rate? *Journal of Experimental Psychology: Animal Behavior Processes*, 18(3), 273–286. <https://doi.org/10.1037/0097-7403.18.3.273>
- Rensink, R. A. (2014). On the prospects for a science of visualization. In W. Huang (Ed.), *Handbook of human centric visualization* (pp. 147–175). Springer.
- Tagu, J., & Kristjánsson, Á. (2020). Dynamics of attentional and oculomotor orienting in visual foraging tasks. *Quarterly Journal of Experimental Psychology*, 174702182091935. <https://doi.org/10.1177/1747021820919351>

- Thornton, I. M., de'Sperati, C., & Kristjánsson, Á. (2019). The influence of selection modality, display dynamics and error feedback on patterns of human foraging. *Visual Cognition*, 1–23. <https://doi.org/10.1080/13506285.2019.1658001>
- Thornton, I. M., & Horowitz, T. S. (2004). The multi-item localization (MILO) task: Measuring the spatiotemporal context of vision for action. *Perception & Psychophysics*, 66(1), 38–50. <https://doi.org/10.3758/BF03194859>
- Thornton, I. M., & Horowitz, T. S. (2020). MILO Mobile: An iPad app to measure search performance in multi-target sequences. *I-Perception*, 11(3), 1–17. <https://doi.org/https://doi.org/10.1177/2041669520932587>
- Thornton, I. M., Tagu, J., Zdravković, S., & Kristjánsson, Á. (2020). Hunting while hunted: Simulated risk of predation affects patterns of human foraging. *Manuscript in Preparation*.
- Tinbergen, L. (1960). The Natural Control of Insects in Pinewoods I. Factors influencing the intensity of predation by songbirds. *Archives Néerlandaises de Zoologie*, 13(3), 265–343. <https://doi.org/10.1163/036551660X00053>
- Todd, P. M., & Hills, T. T. (2020). Foraging in Mind. *Current Directions in Psychological Science*, 29(3), 309–315. <https://doi.org/10.1177/0963721420915861>
- Treisman, A. M., & Gelade, G. (1980). A feature-integration theory of attention. *Cognitive Psychology*, 12(1), 97–136.
- Troscianko, J., Skelhorn, J., & Stevens, M. (2018). Camouflage strategies interfere differently with observer search images. *Proceedings of the Royal Society B: Biological Sciences*, 285(1886), 20181386. <https://doi.org/10.1098/rspb.2018.1386>
- Van den Driessche, C., Chevrier, F., Cleeremans, A., & Sackur, J. (2019). Lower Attentional Skills predict increased exploratory foraging patterns. *Scientific Reports*, 9(1). <https://doi.org/10.1038/s41598-019-46761-0>

- Wolfe, J. M. (2010). Visual search. *Current Biology*, 20(8), R346–R349.
<https://doi.org/10.1016/j.cub.2010.02.016>
- Wolfe, J. M. (2013). When is it time to move to the next raspberry bush? Foraging rules in human visual search. *Journal of Vision*, 13(3), 10–10.
- Wolfe, J. M., Aizenman, A. M., Boettcher, S. E. P., & Cain, M. S. (2016). Hybrid foraging search: Searching for multiple instances of multiple types of target. *Vision Research*, 119, 50–59. <https://doi.org/10.1016/j.visres.2015.12.006>
- Wolfe, J. M., Cain, M. S., & Aizenman, A. M. (2019). Guidance and selection history in hybrid foraging visual search. *Attention, Perception, & Psychophysics*.
<https://doi.org/10.3758/s13414-018-01649-5>
- Wolfe, J. M., & Horowitz, T. S. (2004). Opinion: What attributes guide the deployment of visual attention and how do they do it? *Nature Reviews Neuroscience*, 5(6), 495–501.
<https://doi.org/10.1038/nrn1411>
- Wolfe, J. M., & Horowitz, T. S. (2017). Five factors that guide attention in visual search. *Nature Human Behaviour*, 1(3), 58. <https://doi.org/10.1038/s41562-017-0058>

FIGURE CAPTIONS

Figure 1. a) Relationship between requested and observed ITT as a function of Tempo and Condition, averaged across participants; b) Average slope of the requested x observed ITT function; c) Average offset of the requested x observed ITT function. Error bars represent standard error of the mean.

Figure 2. Foraging tempo as a function of trial bin, averaged across participants. Error bars represent standard error of the mean.

Figure 3. Average number of errors per block, classified by error type and experimental condition. Error bars represent standard error of the mean.

Figure 4. Scatterplot showing the relationship between average number of runs and average tempo in each of the trial bins. Data are plotted for both the feature (open symbols) and conjunction (closed symbols) conditions. Dashed lines are best fitting linear slopes. Error bars represent standard error of the mean in both dimensions.

Figure 5. Scatterplot showing the relationship between average run length and average tempo in each of the trial bins. Data are plotted for both the feature (open symbols) and conjunction (closed symbols) conditions. Dashed lines are best fitting linear slopes. Error bars represent standard error of the mean in both dimensions.

Figure 6. Scatterplot showing the relationship between the proportion of non-random trials and average tempo in each of the trial bins. Data are plotted for both the feature (open symbols) and conjunction (closed symbols) conditions. Dashed lines are best fitting linear slopes. Error bars represent standard error of the mean in both dimensions.

Figure 7. Example scatterplots for individual participants, showing the relationship between average run length and average tempo in each of the trial bins. Data are plotted for both the feature (open symbols) and conjunction (closed symbols) conditions. Dashed lines are best fitting linear slopes. Error bars represent standard error of the mean in both dimensions. See supplementary files for the remaining 8 participants.

Figure 1 Relationship between actual and observed ITT

Figure 2 Tempo x Trial Bin

Figure 3

Figure 4 Number of Runs

Figure 5 Run Length

Figure 6 Proportion of non-random trials

Figure 7 – Individual differences