**Coping With COVID: Risk and Resilience Factors for Mental Health
in a German Representative Panel Study**

Antje Riepenhausen^{1,2}, Ilya M. Veer¹, Carolin Wackerhagen¹, Zala C. Reppmann¹,
Göran Köber^{3,4}, José Luis Ayuso-Mateos^{5,6,7}, Sophie A. Bögemann⁸, Giovanni
Corrao^{9,10}, Mireia Felez-Nobrega^{6,11}, Josep Maria Haro Abad^{6,11}, Erno Hermans⁸, Judith
van Leeuwen⁸, Klaus Lieb^{12,13}, Vincent Lorant¹⁴, Murielle Mary-Krause¹⁵, Roberto
Mediavilla^{5,6}, Maria Melchior¹⁵, Ellenor Mittendorfer-Rutz¹⁶, Matteo Monzio
Compagnoni^{9,10}, Kuan-Yu Pan¹⁷, Lara Puhlmann^{12,18}, Karin Roelofs^{8,19}, Marit
Sijbrandij²⁰, Pierre Smith^{14,21}, Oliver Tüscher^{12,13}, Anke Witteveen²⁰, Matthias Zerban²²,
Raffael Kalisch^{12,22*}, Hannes Kröger^{23,24*}, Henrik Walter^{1,2*}

¹Charité – Universitätsmedizin Berlin, Corporate Member of Freie Universität Berlin
and Humboldt-Universität zu Berlin, Department of Psychiatry and Neurosciences |
CCM, Research Division of Mind and Brain, Charitéplatz 1, 10117 Berlin, Germany.

²Berlin School of Mind and Brain, Faculty of Philosophy, Humboldt-Universität zu
Berlin, Berlin, Germany.

³Institute of Medical Biometry and Statistics, Faculty of Medicine and Medical Center –
University of Freiburg, Freiburg, Germany.

⁴Freiburg Center for Data Analysis and Modelling, University of Freiburg, Freiburg,
Germany.

⁵Department of Psychiatry, Universidad Autónoma de Madrid (UAM), Madrid, Spain.

Abstract

Background: The COVID-19 pandemic might affect mental health. Data from population-representative panel surveys with multiple waves including pre-COVID data investigating risk and protective factors are still rare.

Methods: In a stratified random sample of the German household population (n=6,684), we conducted survey-weighted multiple linear regressions to determine the association of various psychological risk and protective factors with changes in psychological distress (PD; measured via PHQ-4) from pre-pandemic (average of 2016 and 2019) to peri-pandemic (both 2020 and 2021) time points. Control analyses on PD change between two pre-pandemic time points (2016 and 2019) were conducted. Regularized regressions were computed to inform on which factors were statistically most influential in the multicollinear setting.

Results: PHQ-4 in 2020 ($M=2.45$) and 2021 ($M=2.21$) was elevated compared to 2019 ($M=1.79$). Several risk factors (catastrophizing, neuroticism, asking for instrumental support) and protective factors (perceived stress recovery, positive reappraisal, optimism) were identified for the peri-pandemic outcomes. Control analyses revealed that in pre-pandemic times, neuroticism and optimism were predominantly related to PD changes. Regularized regression mostly confirmed the results and highlighted perceived stress recovery as most consistent influential protective factor across peri-pandemic outcomes.

Conclusions: We identified several psychological risk and protective factors related to PD outcomes during the COVID-19 pandemic. Comparison to pre-pandemic data stress the relevance of longitudinal assessments to potentially reconcile

- 1 contradictory findings. Implications and suggestions for targeted prevention and
- 2 intervention programs during highly stressful times such as pandemics are discussed.

- 3 *Keywords:* COVID-19, psychological distress, mental health, resilience, risk
- 4 factors

5

Coping With Covid: Risk and Resilience Factors for Mental Health in a German Representative Panel Study

The spread of the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) globally affects people in various aspects of their life. Not only does the virus impose a physical threat of infection and the associated possibility of a severe course with its long-term consequences; being exposed to such threat constantly, as well as to changes in social life and the economic situation can harm mental well-being. Indeed, several studies have investigated mental health consequences of the coronavirus disease 2019 (COVID-19) pandemic in nationally representative probability samples, most of them referring to the first lockdown in spring 2020 (see *Table S1*). With some exceptions, most of these studies found higher average levels of self-reported depression and anxiety symptoms during the first months of the pandemic compared to pre-pandemic symptom levels (Daly, Sutin, & Robinson, 2021; Dawel et al., 2020; Ettman et al., 2020; Peters, Rospleszcz, Greiser, Dallavalle, & Berger, 2020; Pieh, Budimir, & Probst, 2020; Pierce et al., 2020; Sibley et al., 2020; Twenge & Joiner, 2020; Winkler et al., 2020). Meta-analytic evidence from not exclusively representative studies suggests that these increases in psychological distress were relatively small and recovered over time (Prati & Mancini, 2021; Robinson, Sutin, Daly, & Jones, 2021).

Previous research on mental health during the COVID-19 pandemic has moreover identified several relevant demographic and socio-economic risk and protective factors. Higher psychological distress during the COVID-19 pandemic has been consistently found to be associated with female gender (Daly & Robinson, 2020; Daly, Sutin, & Robinson, 2020; Gijzen et al., 2020; Holingue et al., 2020; Hyland et al.,

1 2020; Li & Wang, 2020; Niedzwiedz et al., 2020; Peters et al., 2020; Pieh et al., 2020;
2 Pierce et al., 2020; Zajacova et al., 2020), younger age (Daly & Robinson, 2020; Daly
3 et al., 2020, 2021; Every-Palmer et al., 2020; Holingue et al., 2020; Hyland et al., 2020;
4 Li & Wang, 2020; Niedzwiedz et al., 2020; Peters et al., 2020; Pieh et al., 2020; Pierce
5 et al., 2020; Zajacova et al., 2020), pre-existing mental conditions (Daly & Robinson,
6 2020; Every-Palmer et al., 2020; Holman, Thompson, Garfin, & Silver, 2020), poor
7 physical health status (Every-Palmer et al., 2020; Holman et al., 2020), and living with
8 young children (Pierce et al., 2020). The results for level of education, income, and
9 employment status are more heterogenous, with studies finding evidence of these
10 being both potential protective as well as risk factors (Daly & Robinson, 2020; Daly et
11 al., 2020; Ettman et al., 2020; Li & Wang, 2020; Niedzwiedz et al., 2020; Pieh et al.,
12 2020; Pierce et al., 2020).

13 The impact of psychological factors on mental health during the COVID-19
14 pandemic, however, has received less attention particularly in representative studies.
15 Identifying such – possibly malleable – psychological factors in the general population
16 will be of great value for informing tailored prevention and intervention efforts to
17 reduce mental health problems and improve well-being during crises (Kunzler et al.,
18 2021).

19 Insights from studies using non-random convenience sampling suggest that
20 several psychological factors are protective factors associated with lower psychological
21 distress or resilience (operationalized as lower psychological distress than expected
22 given a certain exposure to stressors) during the COVID-19 pandemic: These studies
23 found lower psychological distress to be predicted by cognitive flexibility (Dawson &

1 Golijani-Moghaddam, 2020; McCracken, Badinlou, Buhrman, & Brocki, 2020), grit
2 (McCracken et al., 2020), meaning in life (Schnell & Krampe, 2020), dispositional
3 mindfulness (Conversano et al., 2020), secure and avoidant attachment styles (Moccia
4 et al., 2020), optimism (Płomecka et al., 2020; Veer et al., 2021), emotional stability
5 (i.e., low neuroticism; Fernández, Crivelli, Guimet, Allegri, & Pedreira, 2020; Flesia et
6 al., 2020; Veer et al., 2021), self-control (Flesia et al., 2020; Schnell & Krampe, 2020),
7 perceived stress recovery (Veer et al., 2021), positive appraisal style and positive
8 appraisal specific to the COVID-19 pandemic (Veer et al., 2021), both positive (Flesia
9 et al., 2020; Zhu et al., 2020) and behavioral (Veer et al., 2021) coping skills as well as
10 coping skills specific for the COVID-19 pandemic (Fernández et al., 2020), making
11 meaning in negative experiences (Yang et al., 2021), general self-efficacy (Bendau et
12 al., 2020; Veer et al., 2021), internal locus of control (Flesia et al., 2020), and self-
13 esteem (Arima et al., 2020).

14 Due to the non-random sampling strategy of most of the studies until now, it
15 is however difficult to assess to what degree these results generalize to the general
16 population or to what degree they might be driven by (self-) selection of the
17 respondents into the sample (Fink, 2003). A further problem that prohibits reliable
18 conclusions from previous studies on psychological factors and mental health during
19 the COVID-19 pandemic is the systematic lack of pre-pandemic baseline
20 measurements. Whereas these studies can thus describe psychological distress during
21 the pandemic or make claims on average changes by referring to average pre-
22 pandemic health in other samples, they cannot draw inferences regarding measures of
23 intra-individual change.

1 In the current study, we addressed both shortcomings in the literature and
2 investigated the relationship between psychological factors (selected based on cross-
3 sectional findings in a large convenience sample (Veer et al., 2021)) and changes in
4 depression and anxiety symptoms (psychological distress; PD) during the COVID-19
5 pandemic in a sample that is both representative of the German household population
6 and has pre-pandemic baseline measures of the same individuals. Moreover, the long-
7 running panel study allowed us to compare these associations to those with changes
8 from 2016–2019, a ‘normal’ period without a singular and ubiquitous stressor like the
9 pandemic.

10 In accordance with most studies on depression and anxiety in the pandemic,
11 we assumed that the pandemic influenced PD, expecting an increase in PD in 2020 and
12 2021 compared to 2019 and 2016. We moreover hypothesized neuroticism and
13 catastrophizing to be risk factors, expecting higher scores to be associated with larger
14 increases or smaller decreases in PD. We finally expected the following psychological
15 factors to be associated with smaller increases or larger decreases in PD, as protective
16 factors: positive reappraisal, putting into perspective, acceptance, use of instrumental
17 support, positive appraisal specific to the COVID-19 pandemic, perceived stress
18 recovery, optimism, and locus of control.

Methods

Participants

The present sample is a subset of the German nationally representative panel study ‘Socio-economic Panel’ (SOEP; Goebel et al., 2019; Liebig et al., 2019). The SOEP annually surveys over 30,000 participants in more than 20,000 households which come from a stratified random sample of the German household population. For the current study, a random subset of 12,000 households (one participant per household) were contacted via telephone interviews between April 1, 2020 (67,366 confirmed cases of COVID-19 and 732 confirmed deaths related to COVID-19 in Germany so far) and July 4, 2020 (196,096 confirmed cases and 9,010 confirmed deaths) in the context of the SOEP-CoV study (Entringer et al., 2020; Kühne, Kroh, Liebig, & Zinn, 2020). Data collection was split into nine tranches. Overall, n=6,684 individuals participated in the survey in 2020. All n=6,684 participants were recontacted between January 18, 2021 (2,040,659 confirmed cases and 46,633 confirmed deaths) and February 15, 2021 (2,338,987 confirmed cases and 65,076 confirmed deaths). Altogether, n=6,006 individuals participated in this follow-up survey. Participants were also surveyed in previous years and provided information on PD in 2016 (January–September, N=5127) and 2019 (January–September; N=6399); missing values for pre-pandemic PD were imputed. Information on exact timing and size of the individual tranches in 2020 and follow-up assessment in 2021 can be found in *Table S2*.

Measures

PD were assessed using the Patient Health Questionnaire for Depression and Anxiety (PHQ-4), a four-item questionnaire screening for depressive and anxiety

1 symptoms that has already been used in pre-pandemic waves in this sample (Kroenke,
2 Spitzer, Williams, & Löwe, 2009; Löwe et al., 2010). The PHQ-4 is a validated mental
3 health screening instrument and measures general anxiety and depressive symptoms
4 using a 4-point Likert scale ranging from 0 (*'not at all'*) to 3 (*'nearly every day'*). Overall
5 sum scores range from 0 to 12 with classifications into no (0–2), mild (3–5), moderate
6 (6–8), and severe (9–12) symptoms of general anxiety and depression. The PHQ-4 was
7 assessed in 2016, 2019, 2020 and 2021.

8 The coping dimensions of positive reappraisal, putting into perspective and
9 acceptance were assessed using three single items from the Cognitive Emotion
10 Regulation Questionnaire (CERQ; Garnefski & Kraaij, 2007; Loch, Hiller, & Witthöft,
11 2011), adapted in wording to assess emotion regulation during the previous two
12 weeks. Likewise, catastrophizing was measured using a reformulated item from the
13 CERQ scale 'catastrophizing'. The rationale for reformulating the CERQ items to reflect
14 state- rather than trait-like coping was to capture emotion regulation strategies
15 specifically used during the COVID-19 pandemic. Finally, instrumental support-
16 seeking was measured using the first item of the 'using instrumental support' scale of
17 the brief COPE (Carver, 1997). These coping items were selected because they were
18 identified to load most strongly on three factors that were identified using principle
19 component analysis in yet unpublished research (for details, see Supplement 2). The
20 items for positive reappraisal, putting into perspective and acceptance loaded most
21 strongly on a factor representing positive appraisal style, using instrumental support
22 best reflected a behavioral coping style factor, whereas catastrophizing best
23 represented maladaptive coping.

1 Additionally, positive appraisal specific to the COVID-19 pandemic was
2 assessed with two self-formulated items. Perceived stress recovery was measured
3 using one item from the Brief Resilience Scale (Chmitorz et al., 2018; Smith et al.,
4 2008). All coping, COVID-19 appraisal, and recovery items were answered on a Likert
5 scale from 0 (*'don't agree at all'*) to 4 (*'fully agree'*) and were collected during the 2020
6 survey period. Optimism was assessed in 2019 using one item asking about the
7 attitude towards the future, ranging from 1 (*'pessimistic'*) to 4 (*'optimistic'*). Locus of
8 Control was assessed in 2015 and measured using a 10-item instrument with a Likert
9 scale ranging from 1 (*'disagree completely'*) to 4 (*'agree completely'*). Higher values
10 indicate an internal locus of control. Neuroticism was assessed in 2017 using the Big
11 Five Inventory – short version (BFI-S; Hahn, Gottschling, & Spinath, 2012). Answers on
12 the 7-point Likert scale range from 1 (*'does not apply'*) to 7 (*'applies fully'*).

13 Example items for all measures and information on included covariates can be
14 found in *Table 1*, which additionally summarizes the hypothesized relation between
15 psychological factors and outcomes. An overview of the timing of data assessment for
16 the different variables can be found in *Figure 1*.

17 **Statistical Analyses**

18 ***Data Preprocessing***

19 Data cleaning and analyses were performed in R v4.0.0 (R Core Team, 2020).
20 The code used for preprocessing and analyses is available at <https://osf.io/znwjt/>.

21 Missing values (4.5%) were imputed by means of the MICE R package (Buuren &
22 Groothuis-Oudshoorn, 2011) using classification and regression trees with $m=5$
23 imputations and 50 iterations. Statistical outliers were all within the range of the used

scales, therefore considered meaningful and not removed. Predictor variables were z-standardized; outcome variables were not z-standardized. This enabled a) comparison between different PD outcomes irrespective of their variance and b) clinically interpretable evaluation of the relation between psychological factors and absolute change in PD.

As shown in *Table 1* and *Figure 1*, pre-pandemic PD was calculated by averaging PHQ-4 scores from 2016 and 2019 to create a more robust baseline. Change in PD was then calculated by taking difference scores between pre-pandemic PD to 2020 (Δ PHQ 2020) and pre-pandemic PD to 2021 (Δ PHQ 2021); To better understand whether the examined psychological factors predicted change specifically during the COVID-19 pandemic or were generally related with changes in PD over time, we additionally investigated the relation of the predictors with the change in PD from 2016 to 2019 (Δ PHQ 2019).

Descriptive Statistics

We conducted survey-weighted linear models to compare levels in PD between pre- and peri-pandemic survey waves.

Testing of Main Hypotheses

The above-mentioned hypotheses were tested using separate multiple linear regression analyses for each psychological factor/outcome pair, including all covariates in each model. This resulted in 10 regressions per outcome. Results were Bonferroni-corrected and hence considered significant at $p < .005$. Baseline PD levels were added to the models as an additional covariate to control for regression to the mean; we however refrain from reporting their associations with changes in PD. To

1 counteract possible biases in sample selection and due to selective response rates,
2 population survey weights were used (Kroh, 2009; Siegers, Belcheva, & Silbermann,
3 2020). Because the survey weight was zero for 27 participants, final sample size was
4 $n=6657$ ($n=5981$ at follow-up).

5 In order to determine which of the significant predictors found were most
6 strongly associated with the outcomes in the multivariate setting with partly correlated
7 variables, and at the same time avoid overfitting in a model with many predictors, we
8 subsequently conducted LASSO (Least Absolute Shrinkage and Selection Operator)
9 regularized regression analyses (Hastie, Tibshirani, & Wainwright, 2015) using the
10 *miselect* R package (Rix & Du, 2020) and calculated inclusion frequencies. Details on
11 this analysis can be found in *Supplement 3*. Note that it was not possible to include
12 survey weights into the LASSO analysis. As background information for interpretation
13 of the unweighted LASSO results, a comparison of results from the unweighted linear
14 regressions and weighted linear regressions can therefore be found in *Table S3*.

15 ***Additional Analyses***

16 As robustness analyses, we ran multiverse or specification curve analyses
17 (Simonsohn, Simmons, & Nelson, 2020; Steegen, Tuerlinckx, Gelman, & Vanpaemel,
18 2016). Here, slightly different model specifications (linear vs. robust regression, cube-
19 root-transformation of non-normally distributed variables vs. no transformation) were
20 used to ensure that small arbitrary changes did not have major influences on the results
21 of the study (see *Supplement 4*).

1 To investigate how the psychological factors are associated with PHQ-4 in the
2 individual years (vs. the change between years), we ran linear mixed models and
3 estimated margins (mean $\pm$ 1SD) for all predictors (see *Supplement 5*).
4

Results

Sample Description

Between April and June 2020, 31% (vs. 20% in 2019 and 28% in 2016) of the population reported mild, 5% (vs. 4% in 2019 and 6% in 2016) moderate, and 2% (vs. 2% in 2019 and 2% in 2016) severe PD. Peri-pandemic PHQ-4 in 2020 (weighted $M=2.45/12$, $SEM=0.049$) was significantly elevated compared to pre-pandemic levels in 2019 (weighted $M=1.79/12$, $SEM=0.048$; $t(6655)=9.73$, $p<2.2e-16$) and 2016 (weighted $M=2.17/12$, $SEM=0.061$; $t(6655)=3.34$, $p=.002$). In January and February 2021, 29% reported mild, 5% moderate, and 2% severe symptoms. Peri-pandemic PHQ-4 in 2021 ($M=2.21/12$, $SEM=0.048$) was significantly elevated compared to 2019 ($t(6655)=6.07$, $p=1.345e-8$), but not compared to 2016 ($t(6655)=0.455$, $p=.653$), and significantly lower than in 2020 ($t(6655)=-3.41$, $p=7.31e-4$). *Figure 2* displays weighted means and 95% confidence interval of the mean for PHQ-4 across the different years (panel A) and the nine individual tranches assessed in 2020 (panel B).

Socio-Demographic Variables

With respect to socio-demographic factors, history of depression was positively related with ΔPHQ 2020 ($\beta=0.697$), ΔPHQ 2021 ($\beta=1.063$), and ΔPHQ 2019 ($\beta=2.060$). Age group 18–24 ($\beta=1.075$) and female gender ($\beta=0.419$) were positively related with ΔPHQ 2021. All other socio-demographic variables were not significantly related with the outcomes. Exact relations of all covariates with the outcomes can be found in *Tables S4, S5 and S6*.

Multiple Linear Regressions

1 As hypothesized, perceived recovery ($\beta=-0.473$) and reappraisal ($\beta=-0.192$)
2 were negatively, whereas catastrophizing ($\beta=0.553$) and neuroticism ($\beta=0.214$) were
3 positively related with Δ PHQ 2020. Contrary to our hypotheses, instrumental support-
4 seeking ($\beta=0.282$) was also positively related. All other predictors were not
5 significantly associated with Δ PHQ 2020 (see *Table S4*).

6 As expected, perceived recovery ($\beta=-0.332$) and optimism ($\beta=-0.139$) were
7 negatively, whereas catastrophizing ($\beta=0.259$) and neuroticism ($\beta=0.355$) were
8 positively associated with Δ PHQ 2021. Instrumental support-seeking ($\beta=0.170$) was
9 again positively related. All other predictors were not significantly associated with
10 Δ PHQ 2021 (see *Table S5*).

11 To see if these factors were specifically relevant during the COVID-19 pandemic
12 or also relevant before, we repeated the analyses with the change in PD during a control
13 period (2016–2019) as the outcome (see *Table S6*). Optimism ($\beta=-0.175$) was
14 negatively, whereas neuroticism ($\beta=0.421$) was positively associated with Δ PHQ 2019.
15 All other psychological factors were not related. Beta coefficients for all psychological
16 factors and all outcomes are shown in *Figure 3*.

17 **LASSO Regularized Regressions**

18 LASSO regularized regression analysis highlighted the roles of
19 catastrophizing, perceived recovery, neuroticism, and asking for instrumental
20 support for Δ PHQ 2020, of neuroticism, perceived recovery and catastrophizing for
21 Δ PHQ 2021, and of neuroticism as well as optimism for Δ PHQ 2019 (see *Table S7*).

22 **Specification Curve Analyses**

1 The performed specification curve analyses indicate that results remained
2 stable across model specifications (see *Supplement 4*).

3 **Linear Mixed Models**

4 Linear mixed models revealed similar patterns of predictors for pre- vs. peri-
5 pandemic PHQ-4 compared to the multiple linear regressions on Δ PHQ outcomes (see
6 Supplement 5).

7

Discussion

The goal of this study was to investigate if mean psychological distress (PD) increased compared to pre-pandemic times and which risk and resilience factors are associated with the change in PD in a sample representative of the German household population.

First, as expected, we found that PD was on average significantly higher in both 2020 and 2021 compared to 2019. It however must be mentioned that pre-pandemic PD in 2019 was lower than in 2016. Due to these PD fluctuations at baseline, averaged baseline scores were used in all subsequent analyses.

Second, in line with our hypotheses, we found catastrophizing and neuroticism to be risk factors for PD. Unexpectedly, asking for instrumental support also was positively associated with PD across peri-pandemic outcomes.

Third, the most consistent protective factor across all analyses was self-perceived recovery from stress, whereas other factors like optimism and positive reappraisal were only partially supported as protective factors. Contrary to our expectations, putting things into perspective, acceptance, and positive appraisal specific to the COVID-19 pandemic did not emerge as protective factors. We will discuss these results in more detail below.

PDIncrease in the General Population

In our analyses there was an increase in PD in 2020 (2.45/12) and 2021 (2.21/12) compared to 2019 (1.79/12). PD in 2020, but not 2021, was also higher than in 2016 (2.17/12). PD in 2021 was again significantly lower than in 2020.

1 These average numbers are clearly not in the pathological range, as scores of
2 6 and higher reflect moderate to severe PD. However, systematic increases of average
3 PD into the pathological range can hardly be expected in a sample consisting of over
4 6000 participants. Especially the proportion of participants reporting mild (vs. no)
5 symptoms was elevated compared to pre-pandemic times. Our findings of only small
6 but significant increases during the pandemic are in accordance with many other
7 findings in population-based studies (Daly et al., 2020; Niedzwiedz et al., 2020; Peters
8 et al., 2020; Pierce et al., 2020; Twenge & Joiner, 2020). Intriguingly, this small effect
9 may be caused by (vulnerable) subpopulations as indicated by longitudinal samples
10 (e.g., Ahrens et al., 2021).

11 Existing meta-analytic evidence suggests a recovery of PD over time (Robinson
12 et al., 2021). In our sample, PD in 2021 was still elevated, which we attribute to the
13 fact that unlike the studies included in the meta-analysis, we covered a later time point
14 in the middle of another wave of COVID-19 infections. PD in 2021 was however lower
15 than in 2020. This might on the one hand be explained by a habituation effect to the
16 pandemic consequences, including an adjustment to the changes in daily life and social
17 distancing measures. On the other hand, the existence of more precise knowledge
18 about the virus and the prospect of starting vaccination campaigns in Germany in the
19 beginning of 2021 might have led to lower uncertainty compared to 2020 and therefore
20 a different appraisal of the situation, which in turn differentially influenced mental
21 well-being.

22 **Risk Factors for PD**

1 Female gender and younger age were socio-demographic risk factors for peri-
2 pandemic PD in 2021 but not in 2020, adding to the mixed picture that although many
3 studies reported these to be risk factors (see introduction), meta-analytic evidence did
4 not find this relationship (Robinson et al., 2021). The most important psychological
5 risk factor was catastrophizing as it showed positive associations with PD changes
6 across peri-pandemic analyses (but not in the control analyses for the pre-pandemic
7 change from 2016–2019). Catastrophizing is the tendency to think that things are
8 worse than they are or will have a far worse outcome than is realistic. Confirming
9 previous research that highlights catastrophizing as one of the most prominent
10 emotion regulation strategies predicting PD (Garnefski & Kraaij, 2007; Martin & Dahlen,
11 2005), our results indicate that this type of coping is the most maladaptive of those
12 included as predictor. Neuroticism also showed a positive association with PD
13 outcomes in almost all analyses, also for the pre-pandemic control analyses, an
14 association that is well known from the literature (Lahey, 2009). Unexpectedly, asking
15 for instrumental support as coping strategy also emerged as quite consistently
16 positively associated with PD, contrary to what we hypothesized. However, it is
17 conceivable that this predictor was confounded with having negative experiences or
18 symptoms in the first place. The specific formulation of this item was: *‘I’ve been trying*
19 *to get advice or help from other people about what to do’*. People might have only
20 reached out to other people for help if they already experienced significant burden,
21 whereas individuals with less burden might not have sought to do so, especially under
22 the given pandemic circumstances.

23 **Protective Factors for PD**

1 Overall, we found perceived recovery from stress to be the most consistent
2 protective factor across peri-pandemic analyses. Optimism and positive reappraisal
3 were at least partially found to be protective factors, consistent with previous research
4 on their association with mental health (Martin & Dahlen, 2005; Plomin et al., 1992).
5 Contrary to our expectations we did not find support for putting things into
6 perspective, acceptance, and positive appraisal specific to COVID-19 to be protective
7 factors.

8 Optimism and perceived recovery were the only protective factors associated
9 also with PD change in the pre-pandemic control period.

10 These findings could thus indicate that the results regarding other
11 psychological protective factors such as positive reappraisal are specific to the COVID-
12 19 pandemic and that they are not related with changes in PD under normal
13 circumstances. However, an additional, and more likely, explanation is that the
14 temporal distance between the assessment of the pre-pandemic PD change score on
15 the one hand and psychological factors assessed in 2020 on the other hand is too large
16 to find associations. The fact that coping items such as positive reappraisal were
17 reformulated to reflect state-like coping in 2020 substantiates this possible
18 explanation, especially since perceived stress recovery, a trait-like measure assessed
19 in the same wave, does show relation with Δ PHQ 2019.

20 Limitations

21 Despite the strengths of our study such as the representativeness of the sample
22 and existence of individual pre-pandemic baseline PD, which have been considered
23 important specifically in the context of the COVID-19 pandemic (Kunzler et al., 2021;

1 Nieto, Navas, & Vázquez, 2020), as well as the comparison with change in PD during a
2 pre-pandemic period and the use of LASSO regularized regression that selects the
3 most promising variables in a model with many potential variables, there also are
4 several limitations: Most importantly, psychological factors were not assessed at all
5 survey waves (see *Figure 1*). Given that many psychological factors such as coping may
6 be variable and malleable (Compas, Forsythe, & Wagner, 1988), this impedes
7 disentangling directionality of causation between psychological factors, PD, and
8 pandemic context. We are also aware of the second major limitation that results from
9 the uneven sampling of psychological factors: We were forced to include variables that
10 were assessed in 2020 to the model predicting change from 2016 to 2019. Our
11 rationale to nevertheless include the factors into the model was to keep the models as
12 similar as possible to set the peri-pandemic results into perspective. We moreover
13 included variables that were assessed during previous survey waves, such as locus of
14 control in 2015, neuroticism in 2017, and optimism in 2019. Although it would have
15 been preferred to have more recent data, based on the literature we expect a relative
16 stability of these constructs (Cobb-Clark & Schurer, 2011; Cobb-Clark & Schurer, 2013;
17 Scheier, Carver, & Bridges, 1994). Other limitations are the self-report nature of
18 assessments and that the use of single items instead of entire validated questionnaires
19 to assess the coping dimensions, although necessary for pragmatic reasons, might
20 have led to a reduced statistical power. Lastly, we do not have knowledge of specific
21 stressors that might have occurred between the two measurements; changes in PD
22 from baseline to 2020 and baseline to 2021 can therefore only be partly, and only on
23 average, attributed to experiencing the COVID-19 pandemic. However, although there

undoubtedly are other influences on changes in PD that we did not assess and that therefore cannot be controlled for, these are expected to occur at random, whereas every participant experienced the COVID-19 pandemic during data collection.

Outlook

In the present research, we identified several psychological factors that are associated with changes in psychological distress during the COVID-19 pandemic in the general population of Germany. Although due to the used instruments our results can strictly only give insights regarding PD, in light of past findings (Veer et al., 2021) we also expect these psychological factors to be related to general mental health and resilience (i.e., mental health controlled for stressor exposure; see Kalisch et al., 2020 for further details). The exact pattern of predictors might certainly be different when investigating these slightly different outcomes. For example, it should be noted that the strongest predictors for PD in our study were those that are conceptually closest to PD (catastrophizing, asking for instrumental support, neuroticism, and perceived stress recovery), whereas psychological factors that are conceptually further away from symptoms such as positive reappraisal, optimism, or locus of control, show weaker relationships with PD. These latter factors however seem to be stronger predictors for resilience, as for instance shown in Veer et al. (2021). Future representative studies should investigate this in more detail.

Our results do point to some possibly malleable factors that we found to be prospectively associated with changes in PD during COVID-19 and that are therefore possible candidates for targeted prevention and intervention programs to improve general mental well-being during challenging times such as pandemics. Given the

1 pandemic situation, these prevention efforts should ideally be widely accessible and
2 allow for a remote delivery via internet and/or mobile phone. Above all, improving
3 stress recovery, e.g., via physical exercise in the nature (Wooller, Rogerson, Barton,
4 Micklewright, & Gladwell, 2018) or smartphone-assisted biofeedback (Hunter, Olah,
5 Williams, Parks, & Pressman, 2019), appears to be the most promising starting point.
6 Moreover, reducing catastrophizing tendencies, for example via smartphone-based
7 cognitive behavioral interventions for mental health prevention (Ebert et al., 2018;
8 Marciniak et al., 2020), increasing internal locus of control, e.g., via online
9 interventions as has been done by Nallapothula et al. (2020) in an academic context,
10 increasing optimism, e.g. using the Best Possible Self intervention (Malouff & Schutte,
11 2017), and learning to also see positive aspects in the overall challenging situation,
12 e.g., via mobile cognitive behavioral interventions (Ebert et al., 2018; Marciniak et al.,
13 2020) are promising paths to increase individual well-being. Future research should
14 corroborate these directions using interventional studies.

1 **Word Count**

2 Word count: 4469 (excluding title page, abstract, references, figures, and
3 tables). Abstract word count: 236.

4 **Acknowledgements**

5 The authors thank Anna Daniels and Sarah A. Wellan for helpful comments on
6 the manuscript.

7 **Funding Statement**

8 This project has received funding from the European Union's Horizon 2020
9 research and innovation program under Grant Agreement numbers 777084
10 (DynaMORE) and 101016127 (RESPOND). "SOEP-CoV: The Spread of the Coronavirus
11 in Germany: Socio-Economic Factors and Consequences" was funded by the German
12 Federal Ministry of Education and Research (BMBF). AR was supported by
13 Studienstiftung des deutschen Volkes. HK was supported by the Deutsche
14 Forschungsgemeinschaft (DFG, German Science Foundation) Grants – 415809395,
15 427279591,40965412. The data can be accessed via the research data center of the
16 SOEP.

17 **Conflicts of Interest**

18 None.

19 **Ethical Standards**

20 The authors assert that all procedures contributing to this work comply with
21 the ethical standards of the relevant national and institutional committees on human
22 experimentation and with the Helsinki Declaration of 1975, as revised in 2008.

23

References

- Ahrens, K. F., Neumann, R. J., Kollmann, B., Plichta, M. M., Lieb, K., Tüscher, O., & Reif, A. (2021). Differential impact of COVID-related lockdown on mental health in Germany. *World Psychiatry: Official Journal of the World Psychiatric Association (WPA)*, 20(1), 140–141. <https://doi.org/10.1002/wps.20830>
- Arima, M., Takamiya, Y., Furuta, A., Siriratsivawong, K., Tsuchiya, S., & Izumi, M. (2020). Factors associated with the mental health status of medical students during the COVID-19 pandemic: A cross-sectional study in Japan. *BMJ Open*, 10(12), e043728. <https://doi.org/10.1136/bmjopen-2020-043728>
- Bendau, A., Plag, J., Kunas, S., Wyka, S., Ströhle, A., & Petzold, M. B. (2020). Longitudinal changes in anxiety and psychological distress, and associated risk and protective factors during the first three months of the COVID-19 pandemic in Germany. *Brain and Behavior*. <https://doi.org/10.1002/brb3.1964>
- Buuren, S. van, & Groothuis-Oudshoorn, K. (2011). mice: Multivariate Imputation by Chained Equations in R. *Journal of Statistical Software*, 45(1), 1–67. <https://doi.org/10.18637/jss.v045.i03>
- Carver, C. S. (1997). You want to measure coping but your protocol' too long: Consider the brief cope. *International Journal of Behavioral Medicine*, 4(1), 92. https://doi.org/10.1207/s15327558ijbm0401_6
- Chmitorz, A., Wenzel, M., Stieglitz, R.-D., Kunzler, A., Bagusat, C., Helmreich, I., ... Tüscher, O. (2018). Population-based validation of a German version of the Brief Resilience Scale. *PloS One*, 13(2), e0192761.

- 1 <https://doi.org/10.1371/journal.pone.0192761>
- 2 Cobb-Clark, D. A., & Schurer, S. (2011). *The Stability of Big-Five Personality Traits*
- 3 (SSRN Scholarly Paper No. ID 1919414). Rochester, NY: Social Science
- 4 Research Network. <https://doi.org/10.2139/ssrn.1919414>
- 5 Cobb-Clark, D. A., & Schurer, S. (2013). Two Economists' Musings on the Stability of
- 6 Locus of Control. *The Economic Journal*, 123(570), F358-F400.
- 7 <https://doi.org/10.1111/eoj.12069>
- 8 Compas, B. E., Forsythe, C. J., & Wagner, B. M. (1988). Consistency and variability in
- 9 causal attributions and coping with stress. *Cognitive Therapy and Research*,
- 10 12(3), 305-320. <https://doi.org/10.1007/BF01176192>
- 11 Conversano, C., Di Giuseppe, M., Miccoli, M., Ciacchini, R., Gemignani, A., & Orrù, G.
- 12 (2020). Mindfulness, Age and Gender as Protective Factors Against
- 13 Psychological Distress During COVID-19 Pandemic. *Frontiers in Psychology*,
- 14 11, 1900. <https://doi.org/10.3389/fpsyg.2020.01900>
- 15 Daly, M., & Robinson, E. (2020). Psychological distress and adaptation to the COVID-
- 16 19 crisis in the United States. *Journal of Psychiatric Research*.
- 17 <https://doi.org/10.1016/j.jpsychires.2020.10.035>
- 18 Daly, M., Sutin, A. R., & Robinson, E. (2020). Longitudinal changes in mental health
- 19 and the COVID-19 pandemic: Evidence from the UK Household Longitudinal
- 20 Study. *Psychological Medicine*, 1-10.
- 21 <https://doi.org/10.1017/S0033291720004432>
- 22 Daly, M., Sutin, A. R., & Robinson, E. (2021). Depression reported by US adults in
- 23 2017-2018 and March and April 2020. *Journal of Affective Disorders*, 278,

- 1 131–135. <https://doi.org/10.1016/j.jad.2020.09.065>
- 2 Dawel, A., Shou, Y., Smithson, M., Cherbuin, N., Banfield, M., Caelear, A. L., ...
- 3 Batterham, P. J. (2020). The Effect of COVID–19 on Mental Health and
- 4 Wellbeing in a Representative Sample of Australian Adults. *Frontiers in*
- 5 *Psychiatry*, *11*, 579985. <https://doi.org/10.3389/fpsyt.2020.579985>
- 6 Dawson, D. L., & Golijani–Moghaddam, N. (2020). COVID–19: Psychological flexibility,
- 7 coping, mental health, and wellbeing in the UK during the pandemic. *Journal*
- 8 *of Contextual Behavioral Science*, *17*, 126–134.
- 9 <https://doi.org/10.1016/j.jcbs.2020.07.010>
- 10 Ebert, D. D., Van Daele, T., Nordgreen, T., Karekla, M., Compare, A., Zarbo, C., ...
- 11 Baumeister, H. (2018). Internet– and Mobile–Based Psychological
- 12 Interventions: Applications, Efficacy, and Potential for Improving Mental
- 13 Health. *European Psychologist*, *23*(2), 167–187.
- 14 <https://doi.org/10.1027/1016-9040/a000318>
- 15 Entringer, T. M., Kröger, H., Schupp, J., Kühne, S., Liebig, S., Goebel, J., ... Zinn, S.
- 16 (2020). *Psychische Krise durch Covid–19? Sorgen sinken, Einsamkeit steigt,*
- 17 *Lebenszufriedenheit bleibt stabil* (Working Paper No. 1087). SOEPpapers on
- 18 Multidisciplinary Panel Data Research. Retrieved from SOEPpapers on
- 19 Multidisciplinary Panel Data Research website:
- 20 <https://www.econstor.eu/handle/10419/222647>
- 21 Ettman, C. K., Abdalla, S. M., Cohen, G. H., Sampson, L., Vivier, P. M., & Galea, S.
- 22 (2020). Prevalence of Depression Symptoms in US Adults Before and During
- 23 the COVID–19 Pandemic. *JAMA Network Open*, *3*(9), e2019686.

- 1 <https://doi.org/10.1001/jamanetworkopen.2020.19686>
- 2 Every-Palmer, S., Jenkins, M., Gendall, P., Hoek, J., Beaglehole, B., Bell, C., ... Stanley,
3 J. (2020). Psychological distress, anxiety, family violence, suicidality, and
4 wellbeing in New Zealand during the COVID-19 lockdown: A cross-sectional
5 study. *PloS One*, 15(11), e0241658.
6 <https://doi.org/10.1371/journal.pone.0241658>
- 7 Fernández, R. S., Crivelli, L., Guimet, N. M., Allegri, R. F., & Pedreira, M. E. (2020).
8 Psychological distress associated with COVID-19 quarantine: Latent profile
9 analysis, outcome prediction and mediation analysis. *Journal of Affective*
10 *Disorders*, 277, 75–84. <https://doi.org/10.1016/j.jad.2020.07.133>
- 11 Fink, A. (2003). *How to Sample in Surveys*. SAGE.
- 12 Flesia, L., Monaro, M., Mazza, C., Fietta, V., Colicino, E., Segatto, B., & Roma, P.
13 (2020). Predicting Perceived Stress Related to the Covid-19 Outbreak through
14 Stable Psychological Traits and Machine Learning Models. *Journal of Clinical*
15 *Medicine*, 9(10), 3350. <https://doi.org/10.3390/jcm9103350>
- 16 Garnefski, N., & Kraaij, V. (2007). The Cognitive Emotion Regulation Questionnaire:
17 Psychometric features and prospective relationships with depression and
18 anxiety in adults. *European Journal of Psychological Assessment*, 23(3), 141–
19 149. *psych* (2007–12103–002). <https://doi.org/10.1027/1015-5759.23.3.141>
- 20 Gijzen, M., Shields-Zeeman, L., Kleinjan, M., Kroon, H., van der Roest, H., Bolier, L.,
21 ... de Beurs, D. (2020). The Bittersweet Effects of COVID-19 on Mental Health:
22 Results of an Online Survey among a Sample of the Dutch Population Five
23 Weeks after Relaxation of Lockdown Restrictions. *International Journal of*

- 1 *Environmental Research and Public Health*, 17(23).
- 2 <https://doi.org/10.3390/ijerph17239073>
- 3 Goebel, J., Grabka, M. M., Liebig, S., Kroh, M., Richter, D., Schröder, C., & Schupp, J.
- 4 (2019). The German Socio-Economic Panel (SOEP). *Jahrbücher Für*
- 5 *Nationalökonomie Und Statistik*, 239(2), 345–360.
- 6 <https://doi.org/10.1515/jbnst-2018-0022>
- 7 Hahn, E., Gottschling, J., & Spinath, F. M. (2012). Short measurements of personality –
- 8 Validity and reliability of the GSOEP Big Five Inventory (BFI-S). *Journal of*
- 9 *Research in Personality*, 46(3), 355–359.
- 10 <https://doi.org/10.1016/j.jrp.2012.03.008>
- 11 Hastie, T., Tibshirani, R., & Wainwright, M. (2015). *Statistical Learning with Sparsity:*
- 12 *The Lasso and Generalizations*. CRC Press.
- 13 Holingue, C., Badillo-Goicoechea, E., Riehm, K. E., Veldhuis, C. B., Thrul, J., Johnson,
- 14 R. M., ... Kalb, L. G. (2020). Mental distress during the COVID-19 pandemic
- 15 among US adults without a pre-existing mental health condition: Findings
- 16 from American trend panel survey. *Preventive Medicine*, 139, 106231.
- 17 <https://doi.org/10.1016/j.ypmed.2020.106231>
- 18 Holman, E. A., Thompson, R. R., Garfin, D. R., & Silver, R. C. (2020). The unfolding
- 19 COVID-19 pandemic: A probability-based, nationally representative study of
- 20 mental health in the United States. *Science Advances*, 6(42).
- 21 <https://doi.org/10.1126/sciadv.abd5390>
- 22 Hunter, J. F., Olah, M. S., Williams, A. L., Parks, A. C., & Pressman, S. D. (2019). Effect
- 23 of Brief Biofeedback via a Smartphone App on Stress Recovery: Randomized

- 1 Experimental Study. *JMIR Serious Games*, 7(4), e15974.
- 2 <https://doi.org/10.2196/15974>
- 3 Hyland, P., Shevlin, M., McBride, O., Murphy, J., Karatzias, T., Bentall, R. P., ...
- 4 Vallières, F. (2020). Anxiety and depression in the Republic of Ireland during
- 5 the COVID-19 pandemic. *Acta Psychiatrica Scandinavica*, 142(3), 249–256.
- 6 <https://doi.org/10.1111/acps.13219>
- 7 Kalisch, R., Köber, G., Binder, H., Ahrens, K. F., Basten, U., Chmitorz, A., ... Engen, H.
- 8 (2020). *A generic solution for the operationalization and measurement of*
- 9 *resilience and resilience processes in longitudinal observations: Rationale and*
- 10 *basic design of the MARP and LORA studies* [Preprint]. PsyArXiv.
- 11 <https://doi.org/10.31234/osf.io/jg238>
- 12 Kroenke, K., Spitzer, R. L., Williams, J. B. W., & Löwe, B. (2009). An Ultra-Brief
- 13 Screening Scale for Anxiety and Depression: The PHQ-4. *Psychosomatics*,
- 14 50(6), 613–621. [https://doi.org/10.1016/S0033-3182\(09\)70864-3](https://doi.org/10.1016/S0033-3182(09)70864-3)
- 15 Kroh, M. (2009). *Short-Documentation of the Update of the SOEP-Weights, 1984–*
- 16 *2008*. DIW Berlin.
- 17 Kühne, S., Kroh, M., Liebig, S., & Zinn, S. (2020). The Need for Household Panel
- 18 Surveys in Times of Crisis: The Case of SOEP-CoV. *Survey Research Methods*,
- 19 14(2), 195–203. <https://doi.org/10.18148/srm/2020.v14i2.7748>
- 20 Kunzler, A. M., Röthke, N., Günthner, L., Stoffers-Winterling, J., Tüscher, O., Coenen,
- 21 M., ... Lieb, K. (2021). Mental burden and its risk and protective factors during
- 22 the early phase of the SARS-CoV-2 pandemic: Systematic review and meta-
- 23 analyses. *Globalization and Health*, 17(1), 34.

- 1 <https://doi.org/10.1186/s12992-021-00670-y>
- 2 Lahey, B. B. (2009). Public health significance of neuroticism. *The American*
- 3 *Psychologist*, 64(4), 241–256. <https://doi.org/10.1037/a0015309>
- 4 Li, L. Z., & Wang, S. (2020). Prevalence and predictors of general psychiatric disorders
- 5 and loneliness during COVID–19 in the United Kingdom. *Psychiatry Research*,
- 6 297, 113267. <https://doi.org/10.1016/j.psychres.2020.113267>
- 7 Liebig, S., Goebel, J., Kroh, M., Schröder, C., Grabka, M. M., Schupp, J., ...
- 8 Zimmermann, S. (2019). *Socio–Economic Panel (SOEP), data from 1984–2018*,
- 9 version 35. Retrieved from <https://doi.org/10.5684/soep-core.v35>
- 10 Loch, N., Hiller, W., & Witthöft, M. (2011). Der Cognitive Emotion Regulation
- 11 Questionnaire (CERQ) Erste teststatistische Überprüfung einer deutschen
- 12 Adaption = The Cognitive Emotion Regulation Questionnaire (CERQ)
- 13 Psychometric evaluation of a German adaptation. *Zeitschrift Für Klinische*
- 14 *Psychologie Und Psychotherapie: Forschung Und Praxis*, 40(2), 94–106. [psych](https://doi.org/10.1026/1616-3443/a000079)
- 15 (2012–04949–003). <https://doi.org/10.1026/1616-3443/a000079>
- 16 Löwe, B., Wahl, I., Rose, M., Spitzer, C., Glaesmer, H., Wingenfeld, K., ... Brähler, E.
- 17 (2010). A 4–item measure of depression and anxiety: Validation and
- 18 standardization of the Patient Health Questionnaire–4 (PHQ–4) in the general
- 19 population. *Journal of Affective Disorders*, 122(1), 86–95.
- 20 <https://doi.org/10.1016/j.jad.2009.06.019>
- 21 Malouff, J. M., & Schutte, N. S. (2017). Can psychological interventions increase
- 22 optimism? A meta–analysis. *The Journal of Positive Psychology*, 12(6), 594–
- 23 604. <https://doi.org/10.1080/17439760.2016.1221122>

- 1 Marciniak, M. A., Shanahan, L., Rohde, J., Schulz, A., Wackerhagen, C., Kobylińska, D.,
2 ... Kleim, B. (2020). Standalone Smartphone Cognitive Behavioral Therapy–
3 Based Ecological Momentary Interventions to Increase Mental Health: Narrative
4 Review. *JMIR MHealth and UHealth*, 8(11), e19836.
5 <https://doi.org/10.2196/19836>
- 6 Martin, R. C., & Dahlen, E. R. (2005). Cognitive emotion regulation in the prediction of
7 depression, anxiety, stress, and anger. *Personality and Individual Differences*,
8 39(7), 1249–1260. <https://doi.org/10.1016/j.paid.2005.06.004>
- 9 McCracken, L. M., Badinlou, F., Buhrman, M., & Brocki, K. C. (2020). The Role of
10 Psychological Flexibility in the Context of COVID–19: Associations with
11 Depression, Anxiety, and Insomnia. *Journal of Contextual Behavioral Science*.
12 <https://doi.org/10.1016/j.jcbs.2020.11.003>
- 13 Moccia, L., Janiri, D., Pepe, M., Dattoli, L., Molinaro, M., De Martin, V., ... Di Nicola, M.
14 (2020). Affective temperament, attachment style, and the psychological
15 impact of the COVID–19 outbreak: An early report on the Italian general
16 population. *Brain, Behavior, and Immunity*, 87, 75–79.
17 <https://doi.org/10.1016/j.bbi.2020.04.048>
- 18 Nallapothula, D., Lozano, J. B., Han, S., Herrera, C., Sayson, H. W., Levis–Fitzgerald,
19 M., & Maloy, J. (2020). M–LoCUS: A Scalable Intervention Enhances Growth
20 Mindset and Internal Locus of Control in Undergraduate Students in STEM.
21 *Journal of Microbiology & Biology Education*, 21(2).
22 <https://doi.org/10.1128/jmbe.v21i2.1987>
- 23 Niedzwiedz, C. L., Green, M. J., Benzeval, M., Campbell, D., Craig, P., Demou, E., ...

- 1 Katikireddi, S. V. (2020). Mental health and health behaviours before and
2 during the initial phase of the COVID-19 lockdown: Longitudinal analyses of
3 the UK Household Longitudinal Study. *Journal of Epidemiology and*
4 *Community Health*. <https://doi.org/10.1136/jech-2020-215060>
- 5 Nieto, I., Navas, J. F., & Vázquez, C. (2020). The quality of research on mental health
6 related to the COVID-19 pandemic: A note of caution after a systematic
7 review. *Brain, Behavior, & Immunity – Health*, 7, 100123.
8 <https://doi.org/10.1016/j.bbih.2020.100123>
- 9 Peters, A., Rospleszcz, S., Greiser, K. H., Dallavalle, M., & Berger, K. (2020). The
10 Impact of the COVID-19 Pandemic on Self-Reported Health. *Deutsches*
11 *Aerzteblatt Online*. <https://doi.org/10.3238/arztebl.2020.0861>
- 12 Pieh, C., Budimir, S., & Probst, T. (2020). The effect of age, gender, income, work,
13 and physical activity on mental health during coronavirus disease (COVID-19)
14 lockdown in Austria. *Journal of Psychosomatic Research*, 136, 110186.
15 <https://doi.org/10.1016/j.jpsychores.2020.110186>
- 16 Pierce, M., Hope, H., Ford, T., Hatch, S., Hotopf, M., John, A., ... Abel, K. M. (2020).
17 Mental health before and during the COVID-19 pandemic: A longitudinal
18 probability sample survey of the UK population. *The Lancet Psychiatry*, 7(10),
19 883–892. [https://doi.org/10.1016/S2215-0366\(20\)30308-4](https://doi.org/10.1016/S2215-0366(20)30308-4)
- 20 Płomecka, M. B., Gobbi, S., Neckels, R., Radziński, P., Skórko, B., Lazzeri, S., ...
21 Jawaid, A. (2020). Mental Health Impact of COVID-19: A global study of risk
22 and resilience factors. *MedRxiv*, 2020.05.05.20092023.
23 <https://doi.org/10.1101/2020.05.05.20092023>

- 1 Plomin, R., Scheier, M. F., Bergeman, C. S., Pedersen, N. L., Nesselroade, J. R., &
2 McClearn, G. E. (1992). Optimism, pessimism and mental health: A
3 twin/adoption analysis. *Personality and Individual Differences*, 13(8), 921–
4 930. [https://doi.org/10.1016/0191-8869\(92\)90009-E](https://doi.org/10.1016/0191-8869(92)90009-E)
- 5 Prati, G., & Mancini, A. D. (2021). The psychological impact of COVID-19 pandemic
6 lockdowns: A review and meta-analysis of longitudinal studies and natural
7 experiments. *Psychological Medicine*, 51(2), 201–211.
8 <https://doi.org/10.1017/S0033291721000015>
- 9 R Core Team. (2020). *R: A Language and Environment for Statistical Computing*. R
10 Foundation for Statistical Computing, Vienna, Austria. Retrieved from
11 <http://www.R-project.org/>
- 12 Rix, A., & Du, J. (2020). miselect: Variable Selection for Multiply Imputed Data
13 (Version 0.9.0). Retrieved from <https://CRAN.R-project.org/package=miselect>
- 14 Robinson, E., Sutin, A. R., Daly, M., & Jones, A. (2021). *A systematic review and meta-*
15 *analysis of longitudinal cohort studies comparing mental health before versus*
16 *during the COVID-19 pandemic* (p. 2021.03.04.21252921).
17 <https://doi.org/10.1101/2021.03.04.21252921>
- 18 Scheier, M. F., Carver, C. S., & Bridges, M. W. (1994). Distinguishing optimism from
19 neuroticism (and trait anxiety, self-mastery, and self-esteem): A reevaluation
20 of the Life Orientation Test. *Journal of Personality and Social Psychology*,
21 67(6), 1063–1078. (1995-07978-001). [https://doi.org/10.1037/0022-](https://doi.org/10.1037/0022-3514.67.6.1063)
22 3514.67.6.1063
- 23 Schnell, T., & Krampe, H. (2020). Meaning in Life and Self-Control Buffer Stress in

- 1 Times of COVID-19: Moderating and Mediating Effects With Regard to Mental
2 Distress. *Frontiers in Psychiatry*, 11, 582352.
3 <https://doi.org/10.3389/fpsy.2020.582352>
- 4 Sibley, C. G., Greaves, L. M., Satherley, N., Wilson, M. S., Overall, N. C., Lee, C. H. J., ...
5 Barlow, F. K. (2020). Effects of the COVID-19 pandemic and nationwide
6 lockdown on trust, attitudes toward government, and well-being. *The*
7 *American Psychologist*, 75(5), 618-630.
8 <https://doi.org/10.1037/amp0000662>
- 9 Siegers, R., Belcheva, V., & Silbermann, T. (2020). *SOEP-Core v35 - Documentation of*
10 *Sample Sizes and Panel Attrition in the German Socio-Economic Panel (SOEP)*
11 *(1984 until 2018)*.
- 12 Simonsohn, U., Simmons, J. P., & Nelson, L. D. (2020). Specification curve analysis.
13 *Nature Human Behaviour*, 4(11), 1208-1214.
14 <https://doi.org/10.1038/s41562-020-0912-z>
- 15 Smith, B. W., Dalen, J., Wiggins, K., Tooley, E., Christopher, P., & Bernard, J. (2008).
16 The brief resilience scale: Assessing the ability to bounce back. *International*
17 *Journal of Behavioral Medicine*, 15(3), 194-200.
18 <https://doi.org/10.1080/10705500802222972>
- 19 Steegen, S., Tuerlinckx, F., Gelman, A., & Vanpaemel, W. (2016). Increasing
20 Transparency Through a Multiverse Analysis. *Perspectives on Psychological*
21 *Science*, 11(5), 702-712. <https://doi.org/10.1177/1745691616658637>
- 22 Twenge, J. M., & Joiner, T. E. (2020). Mental distress among U.S. adults during the
23 COVID-19 pandemic. *Journal of Clinical Psychology*, 76(12), 2170-2182.

- 1 <https://doi.org/10.1002/jclp.23064>
- 2 Veer, I. M., Riepenhausen, A., Zerban, M., Wackerhagen, C., Puhlmann, L. M. C.,
- 3 Engen, H., ... Kalisch, R. (2021). Psycho-social factors associated with mental
- 4 resilience in the Corona lockdown. *Translational Psychiatry*, 11(1), 1–11.
- 5 <https://doi.org/10.1038/s41398-020-01150-4>
- 6 Winkler, P., Formanek, T., Mlada, K., Kagstrom, A., Mohrova, Z., Mohr, P., & Csemy, L.
- 7 (2020). Increase in prevalence of current mental disorders in the context of
- 8 COVID-19: Analysis of repeated nationwide cross-sectional surveys.
- 9 *Epidemiology and Psychiatric Sciences*, 29, e173.
- 10 <https://doi.org/10.1017/S2045796020000888>
- 11 Wooller, J. J., Rogerson, M., Barton, J., Micklewright, D., & Gladwell, V. (2018). Can
- 12 Simulated Green Exercise Improve Recovery From Acute Mental Stress?
- 13 *Frontiers in Psychology*, 9. <https://doi.org/10.3389/fpsyg.2018.02167>
- 14 Yang, Z., Ji, L.-J., Yang, Y., Wang, Y., Zhu, L., & Cai, H. (2021). Meaning Making Helps
- 15 Cope with COVID-19: A Longitudinal Study. *Personality and Individual*
- 16 *Differences*, 110670. <https://doi.org/10.1016/j.paid.2021.110670>
- 17 Zajacova, A., Jehn, A., Stackhouse, M., Choi, K. H., Denice, P., Haan, M., & Ramos, H.
- 18 (2020). Mental health and economic concerns from March to May during the
- 19 COVID-19 pandemic in Canada: Insights from an analysis of repeated cross-
- 20 sectional surveys. *SSM – Population Health*, 12, 100704.
- 21 <https://doi.org/10.1016/j.ssmph.2020.100704>
- 22 Zhu, J., Sun, L., Zhang, L., Wang, H., Fan, A., Yang, B., ... Xiao, S. (2020). Prevalence
- 23 and Influencing Factors of Anxiety and Depression Symptoms in the First-Line

- 1 Medical Staff Fighting Against COVID-19 in Gansu. *Frontiers in Psychiatry*, 11,
- 2 386. <https://doi.org/10.3389/fpsyt.2020.00386>
- 3

Table 1

Overview of Variables and Instruments Used

<i>Variable</i>	<i>Instrument</i>	<i>Type</i>	<i>Expected relation to PD</i>
<i>Dependent Variables</i>			
Change in PD from pre-pandemic levels (mean of 2016 and 2019) to 2020	$\Delta\text{PHQ } 2020 = \text{PHQ-4 } 2020 - (\text{PHQ-4 } 2016 + \text{PHQ-4 } 2019)/2$		
Change in PD from pre-pandemic levels (mean of 2016 and 2019) to 2021	$\Delta\text{PHQ } 2021 = \text{PHQ-4 } 2021 - (\text{PHQ-4 } 2016 + \text{PHQ-4 } 2019)/2$		
Change in PD from 2016 to 2019	$\Delta\text{PHQ } 2019 = \text{PHQ-4 } 2019 - \text{PHQ-4 } 2016$		
<i>Independent Variables</i>			
Coping: positive reappraisal (PREAP)	CERQ positive reappraisal scale (1 item); <i>'I thought that the situation also has its positive sides'</i>	State	⊖
Coping: putting into perspective (PERSP)	CERQ putting into perspective scale (1 item); <i>'I thought that it hasn't been too bad compared to other things'</i>	State	⊖
Coping: acceptance (ACC)	CERQ acceptance scale (1 item); <i>'I thought that I have to accept the situation'</i>	State	⊖
Positive appraisal specific to the COVID-19 pandemic (PAC)	Self-formulated; positive appraisal of COVID-19 situation on a personal & societal level (2 items); <i>'I expect that I will learn something positive from the corona pandemic for my</i>	State	⊖

	<i>own life' and 'In the long run, I think that society will change for the better because of the corona pandemic'</i>		
Coping: using instrumental support (SUPP)	Brief COPE: using instrumental support scale (1 item); <i>'I've been trying to get advice or help from other people about what to do'</i>	State	⊖
Coping: catastrophizing (CATA)	CERQ: catastrophizing scale (1 item); <i>'I kept thinking about how terrible it is what I have experienced'</i>	State	⊕
Perceived stress recovery (REC)	Brief Resilience Scale (1 item); <i>'I tend to bounce back quickly after hard times'</i>	Trait	⊖
Optimism (OPT)	SOEP-specific item (1 item); <i>'If you think about the future, are you...' (1, pessimistic – 4, optimistic)</i>	Trait	⊖
Locus of Control (LOC)	SOEP-specific questionnaire (10 items); e.g., <i>'My life's course depends on me'</i> and <i>'Success is a matter of fate and luck' (-)</i>	Trait	⊖
Neuroticism (NEU)	BFI-S (3 items); <i>'I am...' 'nervous' 'a worrier', 'relaxed, able to deal with stress' (-)</i>	Trait	⊕
<i>Covariates</i>			
Age (in years)	18–24, 25–34, 35–44, 45–54, 55–64, 65–74, 75–84, 85+		
Gender	m/f		
Education	1: no degree, still in school or lower degree; 2: middle or high school degree; 3: high school degree with subsequent vocational training or university degree		
Household income	Lower/middle/upper tercile		

Risk group status for severe course in case of infection with SARS-CoV-2	Yes/no (determined based on age and BMI as well as self-report of at least one of the following diagnoses: asthma, cardiovascular disease, diabetes, cancer, stroke, high blood pressure, dementia, rheumatism, handicap)
History of diagnosed depression	Yes/no (self-report)
Lockdown status	Yes/no (participation in 2020 up to/after May 5)

Note. PD= psychological distress; PHQ=Patient Health Questionnaire; CERQ=Cognitive Emotion Regulation Questionnaire; SOEP=Socio-economic

Panel; BFI-S=Big Five Inventory, short version; m=male, f=female; SARS-CoV-2= severe acute respiratory syndrome coronavirus 2; BMI=body

mass index. Expected relation to PHQ indicates the hypothesized relationship between the respective independent variable and Δ PHQ 2020 as well as Δ PHQ 2021.

Table 2*Sample Characteristics (N=6,684)*

	%
Gender	
Male	39.32
Female	60.68
Age	
18–24	2.5
25–34	9.38
35–44	16.37
45–54	23.21
55–64	21.27
65–74	15.61
75–84	9.63
85+	2.02
Education	
No degree, still in school or lower degree	24.29
Middle or high school degree	43.88
High school degree with subsequent vocational training or university degree	31.83
Income	
Lower tertile	32.96
Medium tertile	33.49
Higher tertile	33.55
Risk group status for severe course in case of infection with SARS-CoV-2	
Yes	51.26
No	48.74
History of depression	
Yes	9.78
No	90.22

Figure 1*Timing of Data Collection for Predictors and Outcome Variables*

Note. OPT = optimism; LOC = locus of control; NEU = neuroticism; PREAP = positive reappraisal; PERSP = putting into perspective; ACC = acceptance; SUPP = asking for instrumental support; CATA = catastrophizing; PAC = positive appraisal specific to the COVID-19 pandemic; REC=perceived stress recovery; PHQ-4 = Patient Health Questionnaire, 4 item version; ΔPHQ 2019=change in PHQ-4 from 2016 to 2019; ΔPHQ 2020=change in PHQ-4 from 2019 to 2020, ΔPHQ 2021=change in PHQ-4 from 2019 to 2021.

Figure 2

Psychological Distress (PHQ-4) Across Years (A) and Across the 9 Tranches Ranging From April 1 to June 28, 2020 (B)

Note. Error bars depict the 95% confidence interval. PHQ-4 values range from 0 to 12, higher values indicating higher psychological distress. As weighted means are used, means of each individual tranche are representative for the German population. In B, weighted mean PHQ-4 values of the entire sample in 2016 and 2019 are displayed as dotted and dashed horizontal lines, respectively.

Figure 3

Beta Coefficients of Multiple Linear Regressions for Δ PHQ 2020 (A), Δ PHQ 2021 (B), and Δ PHQ 2019 (C).

Note. This figure shows beta coefficients of the psychological factors for the three outcomes.

Complete output tables of the respective linear regressions can be found in Tables S4–S6.

Predictors are z-standardized, outcomes are not standardized. Error bars depict the 95% confidence interval.