Transdiagnostic Processes as Mediators of Change in an Internet-delivered Intervention based on the Unified Protocol

Carmen Schaeuffele¹, Jonathan Baer¹, Inken Buengener¹, Raphaela Grafiadeli², Eva Heuthaler¹, Judith Strieder¹, Patricia Ziehn¹, Christine Knaevelsrud¹, Babette Renneberg¹, and Johanna Boettcher^{1, 2}

¹ Department of Education and Psychology, Freie Universität Berlin, Berlin, Germany

² Clinical Psychology and Psychotherapy, Psychologische Hochschule Berlin, Berlin, Germany

Address for Correspondence: Carmen Schaeuffele, Freie Universität Berlin, Department of Education and Psychology, Clinical Psychology and Psychotherapy, Habelschwerdter Allee 45, 14195 Berlin, Germany, email: carmen.schaeuffele@fu-berlin.de

Funding: This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Conflict of interest: The authors do not have any conflict of interest to report.

Abstract

Background: Transdiagnostic treatments target shared mechanisms between disorders to facilitate change across diagnoses. The Unified Protocol (UP) aims at changing dysfunctional reactions towards emotions by increasing mindful emotion awareness and cognitive flexibility, as well as decreasing anxiety sensitivity and emotion avoidance.

Method: We investigated whether these transdiagnostic processes were malleable by treatment and mediated the relationship between treatment and outcome in an internet-delivered adaptation of the UP. $N=129$ participants with mixed anxiety, depressive, and somatic symptom disorders were randomized to treatment or waitlist.

Results: The treatment yielded significant changes in all transdiagnostic processes over time in comparison to a waitlist condition. In separate mediator models, significant mediating effects were found for mindfulness, cognitive flexibility, behavioral activation, and experiential avoidance. When all mediators were combined in a multiple mediator model, the indirect effects through mindfulness and cognitive flexibility emerged as significant.

Conclusion: These findings add to the growing body of research on transdiagnostic processes as mediators of change and emphasize mindfulness and cognitive flexibility as a transdiagnostic treatment target. However, these results should be interpreted cautiously, as temporal precedence could not be established.

Keywords: transdiagnostic, internet-based cognitive behavioral therapy, Unified Protocol, mediators, mechanisms of change

Transdiagnostic Processes as Mediators of Change in an Internet-Based Adaptation of the Unified Protocol

Numerous studies show that Cognitive Behavioral Therapy (CBT) works (Cuijpers, 2019; Hofmann et al., 2012). In recent years, the field has seen a rise of novel transdiagnostic treatments that have the potential to provide easier access to evidence-based mental health care by addressing a wider range of disorders and lowering barriers to treatment (e.g., Dalgleish et al., 2020; Sauer-Zavala, Gutner, et al., 2017; Schaeuffele et al., 2020). Transdiagnostic treatments show promising effects when delivered face-to-face and over the internet (Newby et al., 2015; Păsărelu et al., 2017; Pearl & Norton, 2017), with equivalence to gold-standard single disorder protocols (e.g., Barlow et al., 2017; Pearl & Norton, 2017). However, overall – across single disorder and transdiagnostic treatments – there remains a need to improve effectiveness and response rates. A better understanding of the active ingredients of psychotherapy can help to create more effective treatments and lower non-response rates (Kazdin, 2007). With the emergence of personalized and modularized treatments, knowledge on change mechanisms can also help to create a better fit between therapy and patient (Hofmann & Hayes, 2019). However, after years of research and advancements, the factors underlying change are still poorly understood (Cuijpers et al., 2019; Kazdin, 2007). This may be due to the fact that investigating mechanisms of change is a complicated endeavor. Determining whether a factor constitutes a (causal) mechanism of change requires a theoretical foundation of putative mechanisms to guide the investigation, the establishment of temporal precedence (change occurs first in the mediator followed by change in symptoms), the inclusion of several mediators, and ideally a manipulation of the mediating variable (Cuijpers et al., 2019; Kazdin, 2007).

A common distinction in mechanisms of change is between common and specific factors. Common factors are active across virtually all therapies and settings, whereas specific factors, as the name suggests, are more treatment-specific (Cuijpers et al., 2019; Wampold,

2015). The most robust evidence on common factors has been found for therapeutic alliance and treatment expectancy (Baier et al., 2020; Constantino et al., 2018; Flückiger et al., 2018). Therapeutic alliance and treatment expectancy have also been shown to predict and mediate outcomes in the internet-based setting (Berger, 2017; J. Boettcher et al., 2013; El Alaoui et al., 2015, 2016; Flückiger et al., 2018; Probst et al., 2019). In comparison, the body of research on relevant specific factors is less clear (e.g., Domhardt et al., 2020; Kazantzis et al., 2018). The number of potential candidate mechanisms impedes the study of specific factors, with every disorder-specific model encompassing specific processes that cause or maintain the disorder. Evidence on how these specific processes are related to outcome is difficult to aggregate and summarize across disorders. In contrast, the transdiagnostic approach, with its emphasis on shared mechanisms that should be targeted in treatment, provides an attractive framework for investigating mediating variables (Sauer-Zavala, Gutner, et al., 2017).

The Unified Protocol (UP) is one example of a theory-driven transdiagnostic treatment for emotional disorders (Barlow et al., 2016; Ellard et al., 2010). The authors hypothesize that individuals with emotional disorders share a tendency to frequently and intensely experience negative affect and exhibit maladaptive reactions to these high negative emotions (Barlow et al., 2014; Bullis et al., 2019). The UP aims at changing these dysfunctional reactions towards emotions by increasing mindful emotion awareness as well as cognitive flexibility and decreasing avoidance (Barlow et al., 2004). These processes are increasingly recognized as transdiagnostic mediators of change across CBT treatments (e.g., Eustis et al., 2016; Goldin et al., 2016; Kocovski et al., 2015).

First studies started to investigate transdiagnostic processes in relation to outcome within the UP framework. As an important prerequisite, the modules of psychoeducation, mindful emotion awareness, cognitive flexibility, and countering emotional behaviors led to changes in these putative mechanisms (Sauer-Zavala, Cassiello-Robbins, et al., 2017). Single case experimental design studies found that mindfulness, cognitive flexibility, and anxiety

sensitivity were associated with outcome (Boswell, 2013; Boswell et al., 2014; Brake et al., 2016). Findings on avoidance are mixed. In single case experimental design studies, experiential avoidance did not exhibit substantial changes following treatment (Boswell et al., 2014; Brake et al., 2016) and did not predict or precede changes in anxiety or depression in an individual with comorbid depression and generalized anxiety (Boswell et al., 2014). In a larger RCT, Eustis et al. (2019) investigated the mediating effect of experiential avoidance across the UP and single-disorder treatment protocols for mixed anxiety disorders. In contrast to these single case studies, experiential avoidance mediated and preceded changes across the UP and single-disorder protocols, suggesting that experiential avoidance is a general transdiagnostic mechanism in CBT (Eustis et al., 2019). One meta-analysis aggregated the UP's effects on emotion regulation (Sakiris & Berle, 2019). They differentiated between adaptive (e.g., mindfulness, reappraisal) and maladaptive (e.g. experiential avoidance, thought suppression) emotion regulation skills (Sakiris & Berle, 2019). Overall, they found preliminary evidence that adaptive strategies increased moderately over the course of treatment and that there were moderate to large effects on experiential avoidance and negligible effects on thought suppression.

Taken together, while there is preliminary evidence for the putative transdiagnostic mechanisms in the UP, the majority of findings are correlational, limited to anxiety disorders, and stem from small sample sizes. Only Eustis et al. (2019) investigated mediating effects in a larger sample, but their investigation was limited to one transdiagnostic process.

Our primary goal for this study was to extend previous findings on the relevance of transdiagnostic processes, specifically mindfulness, cognitive flexibility, anxiety sensitivity, and avoidance, in the UP in an internet-based setting for emotional disorders beyond anxiety disorders. We were interested whether these transdiagnostic processes were malleable by treatment and to which extent they mediated treatment effects – both in isolation and in a multiple mediator model taking into account their interrelation.

Method

We collected data within a randomized controlled trial examining the efficacy of an internet-based version of the UP for anxiety, depressive, and somatic symptom disorders (Schaeuffele, Homeyer, et al., 2020). For this purpose, treatment was compared to waitlist in a 1:1 ratio ($N=129$). The treatment yielded large effects on symptom distress in comparison to waitlist and 69% of participants were considered responders. The proportion of responders would shift to a more conservative estimate of 55% if we considered all participants that did not report post-treatment data as non-responders. Following the waiting period of 10 weeks, participants of the waitlist received access to the intervention.

Participants

We recruited participants by self-selection in online mental health forums, on social media platforms, as well as over targeted ads on a popular search engine. We included participants with an anxiety, depressive, or somatic symptom disorder as a primary diagnosis in the trial. While the UP has primarily been applied in the treatment of anxiety disorders (e.g., Cassiello-Robbins et al., 2020), the rationale should be expandable to other disorders with strong emotional components like depression and somatic symptom disorders (Farchione et al., 2012). Participants were diagnosed with a structured clinical interview via telephone prior to randomization (Margraf et al., 2017). We excluded participants if they (1) currently underwent psychotherapy, (3) were under 18, (4) changed their medication dosage in the preceeding three months, (5) showed indications of acute suicidality, (6) were diagnosed with a substance use disorder, (7) experienced symptoms of mania, or (8) psychosis. See Figure 1 for a detailed flow of participants through the inclusion process.

Intervention

The intervention is a 10-week adaptation of the UP for internet-based use. For this purpose, the UP workbook was translated one-to-one from English into German. In a next step, the content was divided into ten modules. In several editing loops, the paragraphs were

shortened, simplified, and summarized, while retaining the core content and exercises. Figure 2 shows core concepts of the 10 final modules. Participants had to complete a module in order to start with a new module. To complete a module, participants had to complete a reflection on how this module helped them work towards their goals. We recommended that participants work on one module per week, but participants were free to complete the modules at their own pace. In theory, participants could skip exercises and leave the reflection blank to start a new module, but this only affected a negligible proportion of modules and participants. Online therapists were the first and last author as well as 16 advanced students of clinical psychology who were trained in a 1-day workshop and supervised. Online therapists provided written feedback once per week on a fixed day over the course of 10 weeks. The therapists focused on increasing engagement and motivation but also provided corrective feedback. Online therapists, for example, provided guidance when they felt that participants did not engage with the exercises in the intended manner or struggled with adaptive solutions (e.g. therapists would guide participants to come up with alternative adaptive behaviors or offer personalized examples of possible alternative thoughts).

Ethics and trial registration

The ethics committee of the Department of Education and Psychology at Freie Universitaet Berlin, Germany, approved the trial (186/2018). The trial was registered at the German Clinical Trial Registry DRKS (DRKS00014820).

Measures

All assessments were self-rated on the online platform. Figure 2 shows an overview of measurement time points. Outcome and mediators were assessed at baseline (t_0). All putative mediators were assessed immediately following the module that targeted the construct to ensure participants had received the dosage of treatment (t_1). For mechanisms targeted early in treatment (mindfulness and cognitive flexibility), we additionally administered the measures at post-treatment to explore further changes (t_2). We did not assess anxiety

sensitivity and avoidance at post-treatment due to temporal proximity to the module9/week9 assessment. We assumed that one module was completed per week. Accordingly, we assessed the transdiagnostic processes in the waitlist group in the week that corresponded to the number of the module (e.g., we assessed mindfulness after module 3 in the treatment and after 3 weeks in the waitlist group). The percentage of missing values was comparatively high (see Table 2). Technical problems at the beginning of the study led to low response rates in the waitlist control group. As many participants in the intervention group did not complete all modules, missingness rates were higher for measures administered in later modules.

Symptom Distress

We assessed symptom distress with the 18-item Brief Symptom Inventory (BSI-18) (Derogatis, 2000; German version: Franke et al., 2011). The BSI-18 is internally reliable and demonstrates satisfactory psychometric properties (Derogatis & Fitzpatrick, 2004; Franke et al., 2011; Prinz et al., 2013).

Mindfulness

We assessed mindfulness with the 16-item Southampton Mindfulness Questionnaire (SMQ) (Chadwick et al., 2008; German version: Böge et al., 2020). The SMQ assesses mindfulness as the reaction to distressing thoughts and images. Example items include “Usually when I experience distressing thoughts and images, I judge the thought/image as good or bad” and “Usually when I experience distressing thoughts and images, I just notice them and let them go”. The SMQ is unidimensional and internally reliable (Böge et al., 2020; Chadwick et al., 2008).

Cognitive Flexibility

We assessed cognitive flexibility with the 6-item reappraisal subscale of the Emotion Regulation Questionnaire (ERQ) (Gross & John, 2003; German version: Abler & Kessler, 2009). Example items include “When I want to feel less negative emotion, I change the way I’m thinking about the situation.” and “I control my emotions by changing the way I think

about the situation I'm in". The ERQ has satisfactory psychometric properties (Abler & Kessler, 2009; Gross & John, 2003).

Anxiety Sensitivity

We assessed anxiety sensitivity with the 18-item Anxiety Sensitivity Index-3 (ASI-3) (Taylor et al., 2007; German version: Kemper et al., 2009). The scale assesses three facets of anxiety sensitivity: Physical, cognitive, and social concerns. Respective example items include "When my chest feels tight, I get scared that I won't be able to breathe properly", "When my mind goes blank, I worry there is something terribly wrong with me.", and "When I begin to sweat in a social situation, I fear people will think negatively of me.". Psychometric properties are satisfactory (Kemper et al., 2009; Taylor et al., 2007).

Avoidance

The UP targets avoidance by countering avoidance tendencies and engaging in emotional exposures. For depressed patients specifically, targeting avoidance relates to overcoming tendencies of withdrawal and engaging more actively with valued activities which in turn should increase activation (Boswell et al., 2018). In order to capture different aspects of avoidance and be more reflective of the proportion of depressed participants in our sample, we assessed both experiential avoidance, which has been described as the unwillingness to remain in contact with distressing emotions, thoughts, memories, and physical sensations, as well as behavioral activation.

Experiential Avoidance. We assessed experiential avoidance with the 15-item Brief Experiential Avoidance Questionnaire (BEAQ) (Gámez et al., 2014). The BEAQ is internally reliable (Gámez et al., 2014). Example items include "The key to a good life is never feeling any pain" and "I go out of my way to avoid uncomfortable situations". We used our own German translation of the scale. Psychometric properties were investigated in a student and clinical population with comparable results to the English original. Our translation showed high internal reliability (.80-.81), seven- to thirteen-day test-retest-reliability (.77-.86), and a

favorable pattern of convergent and divergent validity in comparison to other measures of experiential avoidance. Our findings indicated that the BEAQ is well represented by a bifactor structure that includes a general experiential avoidance factor and specific factors of individual facets of experiential avoidance (e.g., distress aversion or distraction). Future studies should substantiate this preliminary finding. For the purpose of this study, we relied on the total scale.

Behavioral Activation. We assessed behavioral avoidance and activation with the 9-item Behavioral Activation for Depression Scale Short Form (BADS) (Manos et al., 2011; German version: Fuhr et al., 2016). The BADS has two subscales, an activation and an avoidance subscale. While originally developed to measure changes during behavioral activation for depression, items are phrased transdiagnostically. Example items include “I engaged in a wide and diverse array of activities” or “Most of what I did was to escape from or avoid something unpleasant”. The instrument has shown good psychometric properties (Fuhr et al., 2016; Manos et al., 2011).

Statistical Analyses

The accompanying de-identified data set is available at https://osf.io/hr5wb/?view_only=aa05542c923f41c683a6ecf61a47d25b [anonymous view-only link]. We ran all statistical analyses in RStudio, version 1.2.5033 (RStudio Team, 2019). For the transdiagnostic processes, we first investigated whether the treatment led to significant changes in the putative mediators (treatment, time, and the interaction of treatment and time as fixed and individuals as random effects, mediator as outcome) with linear mixed effects models using the lme4 package (Bates et al., 2014), obtaining p values with the lmer-test package (Kuznetsova et al., 2017). We also calculated effect sizes (Hedge’s *g*) using estimated means and pooled standard deviations.

We then tested all putative transdiagnostic processes in separate mediator models with group as predictor and symptom distress at post-treatment as outcome (see Figure 3). In a next

step, we combined all mediators that showed a significant mediating effect in the separate mediator models in a multiple mediator model (see Figure 4), where residuals of mediators were allowed to covary (Preacher & Hayes, 2008). Mediation models were fitted as path analyses with bias-corrected bootstrapping (10000 draws) and full maximum likelihood estimation to account for missing values using the lavaan package (Rosseel, 2012). If the bias-corrected bootstrapped confidence interval does not contain zero, then the effect of treatment is at least partially dependent on the mediator and the indirect effect can be considered significant (Preacher & Hayes, 2008). We controlled all mediators and outcomes for their baseline value, following findings from a simulation study that found this approach superior to other methods like difference scores in terms of error rates, bias, power, and confidence interval coverage (Valente & MacKinnon, 2017). To judge the magnitude of the mediating effect, P_M — the ratio of the indirect effect to the total effect — was calculated (Preacher & Kelley, 2011; Wen & Fan, 2015).

Results

Demographics

Demographic variables are displayed in Table 1. Groups did not differ at baseline.

Transdiagnostic Processes over Time

Means, Hedges' g effect sizes, as well as percentages of missing values at each measurement point are reported in Table 2. We found significant changes over time in comparison to waitlist for all transdiagnostic processes: mindfulness, cognitive flexibility, anxiety sensitivity, behavioral activation, and experiential avoidance ($F = 5.30-16.40$, $df = 1-2$, $p < .05$). The transdiagnostic processes changed in the hypothesized direction: mindfulness, cognitive flexibility, and behavioral activation increased ($g = 0.73 - 0.99$), while anxiety sensitivity and experiential avoidance decreased ($g = 0.4 - 0.48$). Between group effect sizes were large for behavioral activation and cognitive flexibility ($g = 0.86 - 0.88$), and medium for mindfulness, anxiety sensitivity, and experiential avoidance ($g = 0.59 - 0.76$).

Transdiagnostic Processes as Mediators

Unstandardized and standardized parameter estimates as well as fit indices of the full models can be found in the supplementary material.

Single Mediator Models

Single mediator models with transdiagnostic processes as the mediator between group and outcome are depicted in Figure 3. For mindfulness and cognitive flexibility, two measurement points were available for the comparison between treatment and waitlist and the indirect effects through both time points were significant, as shown in Figure 3. For mindfulness, the indirect effect through the first time point was 0.93, 90%CI [0.18, 2.19] and for the second -2.57, 90%CI [-4.86, -0.67]. For cognitive flexibility, the indirect effect through the first time point was -0.59, 90%CI [-1.71, 0.33] and for the second -1.06, 90%CI [-2.74, -0.06]. The indirect effect through anxiety sensitivity was not significant, while the indirect path through experiential avoidance and behavioral activation were. The treatment effect, thus, was partially mediated by mindfulness, cognitive flexibility, experiential avoidance, and behavioral activation, but not anxiety sensitivity. The effect size of the mediating effect was $P_M=.13$ for mindfulness (indirect effect through t1 and t2), $P_M=.06$ for cognitive flexibility (indirect effect through t1 and t2), $P_M=.09$ for experiential avoidance, and $P_M=.15$ for behavioral avoidance.

Multiple Mediator Models

Figure 4 depicts the multiple mediator model that includes all significant mediators from the single mediator models (all except anxiety sensitivity). In the combined multiple mediator model, the bootstrapped 95% confidence interval of the indirect path through mindfulness and cognitive flexibility did not contain zero: The treatment effect was partially mediated by mindfulness and cognitive flexibility, but not by the other putative processes. The indirect effect through mindfulness showed an effect size of $P_M=.09$ and through

cognitive flexibility $P_M=.06$. The total indirect effect (all indirect effects combined) was -4.58, corresponding to an effect size of $P_M=.48$.

Discussion

This study investigated transdiagnostic processes in an internet-based intervention based on the Unified Protocol for anxiety, depressive, and somatic symptom disorders. All transdiagnostic processes were malleable by treatment with medium to large effect sizes. In single mediator models, we found that cognitive flexibility, mindfulness, experiential avoidance, and behavioral activation partially mediated the relationship between treatment and symptom distress. When all significant mediators were collapsed into one model, the multiple mediation model revealed the indirect paths through mindfulness and cognitive flexibility as significant.

Results from the single mediator models are in line with previous research on the UP that found mindfulness, cognitive flexibility, and experiential avoidance predictive and associated with symptom change (Boswell et al., 2014; Brake et al., 2016; Eustis et al., 2019). Anxiety sensitivity – while increasingly conceptualized as a transdiagnostic process and found to be associated with symptom change in the UP (Boettcher et al., 2016; Boswell et al., 2013) – might play a lesser role in depressive disorders and thus, may not be related to outcome to the same extent in our fairly depressed sample. Future studies should investigate whether these findings on transdiagnostic processes are dependent on diagnosis, as suggested by a recent study on mindfulness as a predictor of outcome in the UP (Woods et al., 2020). The non-significant finding on anxiety sensitivity also raises the question of whether all participants benefit from the interoceptive exposure module that targets anxiety sensitivity. This highlights the potential of a flexible, modular application of the UP, where an individualized sequence of treatment components could be provided based on transdiagnostic processes (Fisher et al., 2019; Sauer-Zavala et al., 2018).

To our knowledge, transdiagnostic processes have not been investigated jointly in one mediation model for the UP as of yet. Mindfulness and cognitive flexibility, as measured by the SMQ and the reappraisal subscale of the ERQ, might capture the overarching aim of the UP best with its focus on how individuals react to distressing thoughts and images (and by that probably also emotions). These findings also tap into the debate whether it is more beneficial to decrease maladaptive or increase adaptive emotion regulation strategies in treatment (e.g., Southward et al., 2021). In our study, the adaptive emotion regulation processes – mindfulness, cognitive flexibility, and behavioral activation – seemed to showed more stable indirect effects than experiential avoidance (and anxiety sensitivity).

The disentanglement of transdiagnostic processes seems quite complex, given they are all related, have fuzzy boundaries, and may share a common underlying basis (e.g., Mansell & McEvoy, 2017; Spinhoven et al., 2017). For instance, mindfulness training seems to facilitate exposure and decrease avoidance in some participants treated with the UP, highlighting the interplay and carryover effects between processes (Curreri et al., 2020). CBT treatments have also shown to be mediated by transdiagnostic processes without explicitly targeting them, for example by mindfulness (e.g., Goldin et al., 2016; Kocovski et al., 2015) or experiential avoidance (e.g., Eustis et al., 2016). These findings also challenge our theory-driven decision to measure mechanisms following the module that target them. Instead, additional research on the structure and hierarchy of transdiagnostic processes as well as dismantling component studies are needed to get a better understanding of the interplay of the processes in therapy and isolate effects.

Our findings need to be interpreted in the light of several limitations. (a) The trial suffered from a high rate of missing values. Besides participants in the treatment group failing to fill out post-treatment assessments, dropout from the intervention affected the assessment of the transdiagnostic processes in the later modules. In order to ensure that participants received the proper dosage of treatment, we measured the transdiagnostic processes following

the module that targeted them. But since the treatment was self-paced and the mean number of modules completed was $M=7.22$ (Schaeuffele, Homeyer, et al., 2020), this naturally meant that the assessments after completion of module 7 and 9 were limited to participants who progressed to this stage of treatment. Thus, our findings are preliminary and should be replicated with larger samples. Our sample size also did not permit us to analyze mediation in different subsamples, e.g. responders versus non-responders. However, we believe that examining mechanisms of change particularly in those patients experiencing change is a promising route of future research (DeRubeis et al., 2014). The dropout also limits the generalizability of our results. We have several hypotheses as to why participants might have not completed the intervention. The self-paced setup and “one size fits all” transdiagnostic approach of the intervention may have been overwhelming and lacked personalization for participants, especially in a guided self-help context. In addition, the lack of adherence-fostering features like personalized reminders and progress reinforcement, as they are implemented in other guided internet-based interventions, may have been detrimental to adherence. Future applications could address these concerns by delivering the intervention in a personalized, modular fashion and including adherence-fostering features. (b) When comparing treatment and waitlist, temporal precedence, that has been described as an important prerequisite for causation (Kazdin, 2007), could not be established with how measurements were set up. While several other of the quality criteria of mediation studies outlined by Lemmens et al. (2016) could be established (RCT with control group, $n>40$ per group, assessment of several mediators), the lack of temporal precedence does not allow to draw any conclusions on causation. However, internet-based treatments are shorter in comparison to face-to-face and it seems to take time for the effect on transdiagnostic processes to unfold as the delayed changes in mindfulness and cognitive flexibility indicate. These circumstances complicate the establishment of temporality. Our results were obtained within an internet-based delivery of the UP. While the face-to-face and internet-based setting

generally seem to share findings on relevant mechanisms of change (e.g., for depression: Domhardt et al., 2020; Lemmens et al., 2016), we cannot rule out that setting-specific influences affected our findings and they, thus, may not translate to the face-to-face setting to the full extent. Direct comparisons between face-to-face and internet-based deliveries and their accompanying change mechanisms are sparse and should be subject of future studies. (c) Linear mediation models, as employed in our study, may not be suitable to capture the complexity of therapeutic processes, since the relationship between outcome and mediator are dynamic, bidirectional, and non-linear (Hofmann et al., 2020). More frequent and parallel assessments of mediators and outcomes are needed to understand the dynamic and agents of change in a more fine-grained manner. To facilitate this, shorter and change sensitive measures should be utilized. (d) The study of common factors, like expectation and working alliance, is constrained to the group receiving treatment (Cuijpers et al., 2019) and thus could not be investigated in comparison with a waitlist control group in our design. To what extent common and specific factors contribute to outcome and how they interact to drive change is still unclear (for meta-analyses from the face-to-face setting see Kazantzis et al., 2018; Lemmens et al., 2016; for a study with a large depressed sample in an internet-based setting see Zagorscak et al., 2020). Future studies should include a control group where alliance and expectation can be measured to unravel effects between common and specific factors.

As Cuijpers (2019) points out, even if temporal precedence and a dose-response relationship is established, possible third variables might still be responsible for changes in mediator and outcome. To put it simply: Only what is measured can be found. In this regard, the use of qualitative studies that openly explore changes and facilitators of changes from the client's perspective might offer valuable insights beyond the constraints of quantitative explorations of mechanisms (Elliott, 2010).

Taken together, we found preliminary evidence that mindfulness, cognitive flexibility, experiential avoidance, and behavioral activation are partially mediating treatment effects in

an internet-based adaptation of the UP. When combined into one model including all five mediators, only the indirect paths through mindfulness and cognitive flexibility were significant. While preliminary, these results emphasize mindfulness and cognitive flexibility as a transdiagnostic treatment target. Future studies should substantiate these findings with larger sample sizes and more frequent and parallel assessments.

References

- Abler, B., & Kessler, H. (2009). Emotion Regulation Questionnaire – Eine deutschsprachige Fassung des ERQ von Gross und John. *Diagnostica*, 55(3), 144–152.
<https://doi.org/10.1026/0012-1924.55.3.144>
- Baier, A. L., Kline, A. C., & Feeny, N. C. (2020). Therapeutic alliance as a mediator of change: A systematic review and evaluation of research. *Clinical Psychology Review*, 82, 101921. <https://doi.org/10.1016/j.cpr.2020.101921>
- Barlow, D. H., Allen, L. B., & Choate, M. L. (2004). Toward a unified treatment for emotional disorders. *Behavior Therapy*, 35(2), 205–230.
[https://doi.org/10.1016/S0005-7894\(04\)80036-4](https://doi.org/10.1016/S0005-7894(04)80036-4)
- Barlow, D. H., Allen, L. B., & Choate, M. L. (2016). Toward a unified treatment for emotional disorders—Republished article. *Behavior Therapy*, 47(6), 838–853.
<https://doi.org/10.1016/j.beth.2016.11.005>
- Barlow, D. H., Ellard, K. K., Sauer-Zavala, S., Bullis, J. R., & Carl, J. R. (2014). The origins of neuroticism. *Perspectives on Psychological Science : A Journal of the Association for Psychological Science*, 9(5), 481–496. <https://doi.org/10.1177/1745691614544528>
- Barlow, D. H., Farchione, T. J., Bullis, J. R., Gallagher, M. W., Murray-Latin, H., Sauer-Zavala, S., Bentley, K. H., Thompson-Hollands, J., Conklin, L. R., Boswell, J. F., Ametaj, A., Carl, J. R., Boettcher, H. T., & Cassiello-Robbins, C. (2017). The Unified Protocol for transdiagnostic treatment of emotional disorders compared with diagnosis-specific protocols for anxiety disorders: A randomized clinical trial. *JAMA Psychiatry*. <https://doi.org/10.1001/jamapsychiatry.2017.2164>
- Bates, D., Mächler, M., Bolker, B., & Walker, S. (2014). Fitting linear mixed-effects models using lme4. *ArXiv:1406.5823 [Stat]*. <http://arxiv.org/abs/1406.5823>
- Berger, T. (2017). The therapeutic alliance in internet interventions: A narrative review and suggestions for future research. *Psychotherapy Research : Journal of the Society for*

- Psychotherapy Research*, 27(5), 511–524.
<https://doi.org/10.1080/10503307.2015.1119908>
- Boettcher, H., Brake, C. A., & Barlow, D. H. (2016). Origins and outlook of interoceptive exposure. *Journal of Behavior Therapy and Experimental Psychiatry*, 53, 41–51.
<https://doi.org/10.1016/j.jbtep.2015.10.009>
- Boettcher, J., Renneberg, B., & Berger, T. (2013). Patient expectations in internet-based self-help for social anxiety. *Cognitive Behaviour Therapy*, 42(3), 203–214.
<https://doi.org/10.1080/16506073.2012.759615>
- Böge, K., Schaeuffele, C., Jacobsen, P., Chadwick, P., Ergen, E., Hahne, I., Bergmann, N., Boettcher, J., Wingefeld, K., Bajbouj, M., & Hahn, E. (2020). Validation of the German version of the Southampton Mindfulness Questionnaire (SMQ). *Mindfulness*, 11(9), 2219–2234. <https://doi.org/10.1007/s12671-020-01447-x>
- Boswell, J. F. (2013). Intervention strategies and clinical process in transdiagnostic cognitive-behavioral therapy. *Psychotherapy (Chicago, Ill.)*, 50(3), 381–386.
<https://doi.org/10.1037/a0032157>
- Boswell, J. F., Anderson, L. M., & Barlow, D. H. (2014). An idiographic analysis of change processes in the unified transdiagnostic treatment of depression. *Journal of Consulting and Clinical Psychology*, 82(6), 1060–1071. <https://doi.org/10.1037/a0037403>
- Boswell, J. F., Conklin, L. R., Oswald, J. M., & Bugatti, M. (2018). The unified protocol for major depressive disorder. In *Applications of the Unified Protocol for Transdiagnostic Treatment of Emotional Disorders*. Oxford University Press.
- Boswell, J. F., Farchione, T. J., Sauer-Zavala, S., Murray, H. W., Fortune, M. R., & Barlow, D. H. (2013). Anxiety sensitivity and interoceptive exposure: A transdiagnostic construct and change strategy. *Behavior Therapy*, 44(3), 417–431.
<https://doi.org/10.1016/j.beth.2013.03.006>

- Brake, C. A., Sauer-Zavala, S., Boswell, J. F., Gallagher, M. W., Farchione, T. J., & Barlow, D. H. (2016). Mindfulness-based exposure strategies as a transdiagnostic mechanism of change: An exploratory alternating treatment design. *Behavior Therapy, 47*(2), 225–238. <https://doi.org/10.1016/j.beth.2015.10.008>
- Bullis, J. R., Boettcher, H., Sauer-Zavala, S., Farchione, T. J., & Barlow, D. H. (2019). What is an emotional disorder? A transdiagnostic mechanistic definition with implications for assessment, treatment, and prevention. *Clinical Psychology: Science and Practice, 26*(2), e12278. <https://doi.org/10.1111/cpsp.12278>
- Cassidello-Robbins, C., Southward, M. W., Tirpak, J. W., & Sauer-Zavala, S. (2020). A systematic review of unified protocol applications with adult populations: Facilitating widespread dissemination via adaptability. *Clinical Psychology Review, 78*, 101852. <https://doi.org/10.1016/j.cpr.2020.101852>
- Chadwick, P., Hember, M., Symes, J., Peters, E., Kuipers, E., & Dagnan, D. (2008). Responding mindfully to unpleasant thoughts and images: Reliability and validity of the Southampton mindfulness questionnaire (SMQ). *The British Journal of Clinical Psychology, 47*(Pt 4), 451–455. <https://doi.org/10.1348/014466508X314891>
- Constantino, M. J., Vîslă, A., Coyne, A. E., & Boswell, J. F. (2018). A meta-analysis of the association between patients' early treatment outcome expectation and their posttreatment outcomes. *Psychotherapy, 55*(4), 473–485. <https://doi.org/10.1037/pst0000169>
- Cuijpers, P. (2019). Targets and outcomes of psychotherapies for mental disorders: An overview. *World Psychiatry, 18*(3), 276–285. <https://doi.org/10.1002/wps.20661>
- Cuijpers, P., Reijnders, M., & Huibers, M. J. H. (2019). The role of common factors in psychotherapy outcomes. *Annual Review of Clinical Psychology, 15*(1), 207–231. <https://doi.org/10.1146/annurev-clinpsy-050718-095424>

- Curreri, A. J., Farchione, T. J., Sauer-Zavala, S., & Barlow, D. H. (2020). Mindful emotion awareness facilitates engagement with exposure therapy: An idiographic exploration using single case experimental design. *Behavior Modification*, 014544552094766. <https://doi.org/10.1177/0145445520947662>
- Dalgleish, T., Black, M., Johnston, D., & Bevan, A. (2020). Transdiagnostic approaches to mental health problems: Current status and future directions. *Journal of Consulting and Clinical Psychology*, 88(3), 179–195. <https://doi.org/10.1037/ccp0000482>
- Derogatis, L. R. (2000). *BSI-18: Brief Symptom Inventory 18—Administration, scoring, and procedures manual*. NCS Pearson.
- Derogatis, L. R., & Fitzpatrick, M. (2004). The SCL-90-R, the Brief Symptom Inventory (BSI), and the BSI-18. In *The use of psychological testing for treatment planning and outcomes assessment: Instruments for adults*. Lawrence Erlbaum Associates Publishers.
- DeRubeis, R. J., Gelfand, L. A., German, R. E., Fournier, J. C., & Forand, N. R. (2014). Understanding processes of change: How some patients reveal more than others—and some groups of therapists less—about what matters in psychotherapy. *Psychotherapy Research*, 24(3), 419–428. <https://doi.org/10.1080/10503307.2013.838654>
- Domhardt, M., Steubl, L., Boettcher, J., Buntrock, C., Karyotaki, E., Ebert, D. D., Cuijpers, P., & Baumeister, H. (2020). Mediators and mechanisms of change in internet- and mobile-based interventions for depression: A systematic review. *Clinical Psychology Review*, 101953. <https://doi.org/10.1016/j.cpr.2020.101953>
- El Alaoui, S., Ljótsson, B., Hedman, E., Kaldø, V., Andersson, E., Rück, C., Andersson, G., & Lindefors, N. (2015). Predictors of symptomatic change and adherence in internet-based cognitive behaviour therapy for social anxiety disorder in routine psychiatric care. *PLOS ONE*, 10(4), e0124258. <https://doi.org/10.1371/journal.pone.0124258>

El Alaoui, S., Ljótsson, B., Hedman, E., Svanborg, C., Kaldø, V., & Lindefors, N. (2016).

Predicting outcome in internet-based cognitive behaviour therapy for major depression: A large cohort study of adult patients in routine psychiatric care. *PLOS ONE*, *11*(9), e0161191. <https://doi.org/10.1371/journal.pone.0161191>

Ellard, K. K., Fairholme, C. P., Boisseau, C. L., Farchione, T. J., & Barlow, D. H. (2010).

Unified protocol for the transdiagnostic treatment of emotional disorders: Protocol development and initial outcome data. *Cognitive and Behavioral Practice*, *1*, 88–101. <https://doi.org/10.1016/j.cbpra.2009.06.002>

Elliott, R. (2010). Psychotherapy change process research: Realizing the promise.

Psychotherapy Research : Journal of the Society for Psychotherapy Research, *20*(2), 123–135. <https://doi.org/10.1080/10503300903470743>

Eustis, E. H., Cardona, N., Nauphal, M., Sauer-Zavala, S., Rosellini, A. J., Farchione, T. J., &

Barlow, D. H. (2019). Experiential avoidance as a mechanism of change across cognitive-behavioral therapy in a sample of participants with heterogeneous anxiety disorders. *Cognitive Therapy and Research*, *135*(4), 166.

<https://doi.org/10.1007/s10608-019-10063-6>

Eustis, E. H., Hayes-Skelton, S. A., Roemer, L., & Orsillo, S. M. (2016). Reductions in

experiential avoidance as a mediator of change in symptom outcome and quality of life in acceptance-based behavior therapy and applied relaxation for generalized anxiety disorder. *Behaviour Research and Therapy*, *87*, 188–195.

<https://doi.org/10.1016/j.brat.2016.09.012>

Farchione, T. J., Fairholme, C. P., Ellard, K. K., Boisseau, C. L., Thompson-Hollands, J.,

Carl, J. R., Gallagher, M. W., & Barlow, D. H. (2012). Unified protocol for transdiagnostic treatment of emotional disorders: A randomized controlled trial.

Behavior Therapy, *43*(3), 666–678. <https://doi.org/10.1016/j.beth.2012.01.001>

- Fisher, A. J., Bosley, H. G., Fernandez, K. C., Reeves, J. W., Soyster, P. D., Diamond, A. E., & Barkin, J. (2019). Open trial of a personalized modular treatment for mood and anxiety. *Behaviour research and therapy. Behaviour Research and Therapy*, 116, 69–79. <https://doi.org/10.1016/j.brat.2019.01.010>
- Flückiger, C., Del Re, A. C., Wampold, B. E., & Horvath, A. O. (2018). The alliance in adult psychotherapy: A meta-analytic synthesis. *Psychotherapy*, 55(4), 316. <https://doi.org/10.1037/pst0000172>
- Franke, G. H., Ankerhold, A., Haase, M., Jäger, S., Tögel, C., Ulrich, C., & Frommer, J. (2011). Der Einsatz des Brief Symptom Inventory 18 (BSI-18) bei Psychotherapiepatienten. *Psychotherapie, Psychosomatik, Medizinische Psychologie*, 61(2), 82–86. <https://doi.org/10.1055/s-0030-1270518>
- Fuhr, K., Hautzinger, M., Krisch, K., Berking, M., & Ebert, D. D. (2016). Validation of the Behavioral Activation for Depression Scale (BADs)-Psychometric properties of the long and short form. *Comprehensive Psychiatry*, 66, 209–218. <https://doi.org/10.1016/j.comppsy.2016.02.004>
- Gámez, W., Chmielewski, M., Kotov, R., Ruggero, C., Suzuki, N., & Watson, D. (2014). The brief experiential avoidance questionnaire: Development and initial validation. *Psychological Assessment*, 26(1), 35–45. <https://doi.org/10.1037/a0034473>
- Goldin, P. R., Morrison, A., Jazaieri, H., Brozovich, F., Heimberg, R., & Gross, J. J. (2016). Group CBT versus MBSR for social anxiety disorder: A randomized controlled trial. *Journal of Consulting and Clinical Psychology*, 84(5), 427–437. <https://doi.org/10.1037/ccp0000092>
- Gross, J. J., & John, O. P. (2003). Individual differences in two emotion regulation processes: Implications for affect, relationships, and well-being. *Journal of Personality and Social Psychology*, 85(2), 348–362. <https://doi.org/10.1037/0022-3514.85.2.348>

- Hofmann, S. G., Asnaani, A., Vonk, I. J. J., Sawyer, A. T., & Fang, A. (2012). The efficacy of cognitive behavioral therapy: A review of meta-analyses. *Cognitive Therapy and Research*, 36(5), 427–440. <https://doi.org/10.1007/s10608-012-9476-1>
- Hofmann, S. G., Curtiss, J. E., & Hayes, S. C. (2020). Beyond linear mediation: Toward a dynamic network approach to study treatment processes. *Clinical Psychology Review*, 76, 101824. <https://doi.org/10.1016/j.cpr.2020.101824>
- Hofmann, S. G., & Hayes, S. C. (2019). The future of intervention science: Process-based therapy. *Clinical Psychological Science*, 7(1), 37–50. <https://doi.org/10.1177/2167702618772296>
- Kazantzis, N., Luong, H. K., Usatoff, A. S., Impala, T., Yew, R. Y., & Hofmann, S. G. (2018). The processes of cognitive behavioral therapy: A review of meta-analyses. *Cognitive Therapy and Research*, 42(4), 349–357. <https://doi.org/10.1007/s10608-018-9920-y>
- Kazdin, A. E. (2007). Mediators and mechanisms of change in psychotherapy research. *Annual Review of Clinical Psychology*, 3, 1–27. <https://doi.org/10.1146/annurev.clinpsy.3.022806.091432>
- Kemper, C. J., Ziegler, M., & Taylor, S. (2009). Überprüfung der psychometrischen Qualität der deutschen Version des Angstsensitivitätsindex-3. *Diagnostica*, 55(4), 223–233. <https://doi.org/10.1026/0012-1924.55.4.223>
- Kocovski, N. L., Fleming, J. E., Hawley, L. L., Ho, M.-H. R., & Antony, M. M. (2015). Mindfulness and acceptance-based group therapy and traditional cognitive behavioral group therapy for social anxiety disorder: Mechanisms of change. *Behaviour Research and Therapy*, 70, 11–22. <https://doi.org/10.1016/j.brat.2015.04.005>
- Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest package: Tests in linear mixed effects models. *Journal of Statistical Software*, 82(13). <https://doi.org/10.18637/jss.v082.i13>

Lemmens, L. H. J. M., Müller, V. N. L. S., Arntz, A., & Huibers, M. J. H. (2016).

Mechanisms of change in psychotherapy for depression: An empirical update and evaluation of research aimed at identifying psychological mediators. *Clinical Psychology Review*, 50, 95–107. <https://doi.org/10.1016/j.cpr.2016.09.004>

MacKinnon, D. P. (2008). *Introduction to statistical mediation analysis*. Lawrence Erlbaum Associates.

Manos, R. C., Kanter, J. W., & Luo, W. (2011). The behavioral activation for depression scale-short form: Development and validation. *Behavior Therapy*, 42(4), 726–739. <https://doi.org/10.1016/j.beth.2011.04.004>

Mansell, W., & McEvoy, P. M. (2017). A test of the core process account of psychopathology in a heterogenous clinical sample of anxiety and depression. A case of the blind men and the elephant? *Journal of Anxiety Disorders*, 46, 4–10.

Margraf, J., Cwik, J. C., Suppiger, A., & Schneider, S. (2017). *DIPS open access: Diagnostic interview for mental disorders. [DIPS Open Access: Diagnostisches Interview bei psychischen Störungen.]*. Mental Health Research and Treatment Center, Ruhr-Universität Bochum.

Newby, J. M., McKinnon, A., Kuyken, W., Gilbody, S., & Dalgleish, T. (2015). Systematic review and meta-analysis of transdiagnostic psychological treatments for anxiety and depressive disorders in adulthood. *Clinical Psychology Review*, 40, 91–110. <https://doi.org/10.1016/j.cpr.2015.06.002>

Păsărelu, C. R., Andersson, G., Bergman Nordgren, L., & Dobrea, A. (2017). Internet-delivered transdiagnostic and tailored cognitive behavioral therapy for anxiety and depression: A systematic review and meta-analysis of randomized controlled trials. *Cognitive Behaviour Therapy*, 46(1), 1–28. <https://doi.org/10.1080/16506073.2016.1231219>

- Pearl, S. B., & Norton, P. J. (2017). Transdiagnostic versus diagnosis specific cognitive behavioural therapies for anxiety: A meta-analysis. *Journal of Anxiety Disorders*, 46, 11–24. <https://doi.org/10.1016/j.janxdis.2016.07.004>
- Preacher, K. J., & Hayes, A. F. (2008). Asymptotic and resampling strategies for assessing and comparing indirect effects in multiple mediator models. *Behavior Research Methods*, 40(3), 879–891. <https://doi.org/10.3758/BRM.40.3.879>
- Preacher, K. J., & Kelley, K. (2011). Effect size measures for mediation models: Quantitative strategies for communicating indirect effects. *Psychological Methods*, 16(2), 93–115. <https://doi.org/10.1037/a0022658>
- Prinz, U., Nutzinger, D. O., Schulz, H., Petermann, F., Braukhaus, C., & Andreas, S. (2013). Comparative psychometric analyses of the SCL-90-R and its short versions in patients with affective disorders. *BMC Psychiatry*, 13, 104. <https://doi.org/10.1186/1471-244X-13-104>
- Probst, G. H., Berger, T., & Flückiger, C. (2019). The alliance-outcome relation in internet-based interventions for psychological disorders: A correlational meta-analysis. *Verhaltenstherapie*, 1–12. <https://doi.org/10.1159/000503432>
- Rosseel, Y. (2012). lavaan: An R package for structural equation modeling. *Journal of Statistical Software*, 48(2). <https://doi.org/10.18637/jss.v048.i02>
- RStudio Team. (2019). *RStudio: Integrated Development Environment for R*. RStudio, Inc. <http://www.rstudio.com/>
- Sakiris, N., & Berle, D. (2019). A systematic review and meta-analysis of the Unified Protocol as a transdiagnostic emotion regulation based intervention. *Clinical Psychology Review*, 72, 101751. <https://doi.org/10.1016/j.cpr.2019.101751>
- Sauer-Zavala, S., Cassiello-Robbins, C., Ametaj, A. A., Wilner, J. G., & Pagan, D. (2018). Transdiagnostic treatment personalization: The feasibility of ordering unified protocol

- modules according to patient strengths and weaknesses. *Behavior Modification*, 518–543. <https://doi.org/10.1177/0145445518774914>
- Sauer-Zavala, S., Cassiello-Robbins, C., Conklin, L. R., Bullis, J. R., Thompson-Hollands, J., & Kennedy, K. A. (2017). Isolating the unique effects of the unified protocol treatment modules using single case experimental design. *Behavior Modification*, 41(2), 286–307. <https://doi.org/10.1177/0145445516673827>
- Sauer-Zavala, S., Gutner, C. A., Farchione, T. J., Boettcher, H. T., Bullis, J. R., & Barlow, D. H. (2017). Current definitions of “transdiagnostic” in treatment development: A search for consensus. *Behavior Therapy*, 48, 128–138. <https://doi.org/10.1016/j.beth.2016.09.004>
- Schaeuffele, C., Homeyer, S. L., Perea, L., Scharf, L., Schulz, A., Knaevelsrud, C., Renneberg, B., & Boettcher, J. (2020). *The unified protocol as an internet-based intervention for emotional disorders: Randomized controlled trial* [Preprint]. PsyArXiv. <https://doi.org/10.31234/osf.io/528tw>
- Schaeuffele, C., Schulz, A., Knaevelsrud, C., Renneberg, B., & Boettcher, J. (2020). CBT at the crossroads: The rise of transdiagnostic treatments. *International Journal of Cognitive Therapy*. <https://doi.org/10.1007/s41811-020-00095-2>
- Southward, M. W., Sauer-Zavala, S., & Cheavens, J. S. (2021). Specifying the mechanisms and targets of emotion regulation: A translational framework from affective science to psychological treatment. *Clinical Psychology: Science and Practice*. <https://doi.org/10.1037/cps0000003>
- Spinhoven, P., van Hemert, A. M., & Penninx, B. W. J. H. (2017). Experiential avoidance and bordering psychological constructs as predictors of the onset, relapse and maintenance of anxiety disorders: One or Many? *Cognitive Therapy and Research*, 41(6), 867–880. <https://doi.org/10.1007/s10608-017-9856-7>

- Taylor, S., Zvolensky, M. J., Cox, B. J., Deacon, B., Heimberg, R. G., Ledley, D. R., Abramowitz, J. S., Holaway, R. M., Sandin, B., Stewart, S. H., Coles, M., Eng, W., Daly, E. S., Arrindell, W. A., Bouvard, M., & Cardenas, S. J. (2007). Robust dimensions of anxiety sensitivity: Development and initial validation of the Anxiety Sensitivity Index-3. *Psychological Assessment, 19*(2), 176–188.
<https://doi.org/10.1037/1040-3590.19.2.176>
- Valente, M. J., & MacKinnon, D. P. (2017). Comparing models of change to estimate the mediated effect in the pretest-posttest control group design. *Structural Equation Modeling : A Multidisciplinary Journal, 24*(3), 428–450.
<https://doi.org/10.1080/10705511.2016.1274657>
- Wampold, B. E. (2015). How important are the common factors in psychotherapy? An update. *World Psychiatry, 14*(3), 270–277. <https://doi.org/10.1002/wps.20238>
- Wen, Z., & Fan, X. (2015). Monotonicity of effect sizes: Questioning kappa-squared as mediation effect size measure. *Psychological Methods, 20*(2), 193–203.
<https://doi.org/10.1037/met0000029>
- Woods, B. K., Sauer-Zavala, S., Farchione, T. J., & Barlow, D. H. (2020). Isolating the effects of mindfulness training across anxiety disorder diagnoses in the unified protocol. *Behavior Therapy, 51*(6), 972–983.
<https://doi.org/10.1016/j.beth.2020.01.001>
- Zagorscak, P., Heinrich, M., Schulze, J., Böttcher, J., & Knaevelsrud, C. (2020). Factors contributing to symptom change in standardized and individualized Internet-based interventions for depression: A randomized-controlled trial. *Psychotherapy, 57*(2), 237–251. <https://doi.org/10.1037/pst0000276>

Table 1*Demographic Characteristics of Participants at Baseline*

Demographic Variable	Treatment (n=65)	Waitlist (n=64)	Total	Statistical test of group difference		
				df	χ^2 / t	p
Gender				1	.48	.49
Male	23 (35.4%)	18 (28.1%)	41 (31.8%)			
Female	42 (64.6%)	46 (71.9%)	88 (68.2%)			
Age				127	.18	.86
Mean (SD)	37.51 (11.99)	37.11 (13.04)	37.31 (12.47)			
Range	18-67	18-66	18-67			
Relationship				1	1.99	.16
in a relationship	40 (61.5%)	30 (46.9%)	70 (54.3%)			
single	25 (38.5%)	34 (53.1%)	59 (45.7%)			
Highest Education				4	4.7	.32
up to 9 years of school education	4 (6.2%)	5 (7.8%)	9 (7%)			
Secondary school	10 (15.4%)	11 (17.2%)	21 (16.3%)			
College entrance qualification	16 (24.6%)	25 (39.1%)	41 (31.8%)			
College / university degree	34 (52.3%)	22 (34.4%)	56 (43.3%)			
other	1 (1.5%)	1 (1.6%)	2 (1.6%)			
Employment status				5	2.99	.7
employed	33 (50.8%)	29 (45.3%)	62 (48.1%)			
self-employed	4 (6.2%)	5 (7.8%)	9 (7%)			
in education	12 (18.5%)	17 (26.6%)	29 (22.5%)			
pensioned	3 (4.6%)	5 (7.8%)	8 (6.2%)			
unemployed	9 (13.8%)	6 (9.4%)	15 (11.6%)			
other	4 (6.2%)	2 (3.1%)	6 (4.7%)			
Currently on medication				1	0	.99
yes	16 (24.6%)	16 (25%)	32 (24.8%)			
no	49 (75.4%)	48 (75%)	97 (75.2%)			
Prior Psychotherapy				1	1.95	.16
yes	44 (67.7%)	51 (79.7%)	95 (73.6%)			
no	21 (32.3%)	13 (20.3%)	34 (26.4%)			
Primary Diagnosis				9	5.66	.77
Agoraphobia	3 (4.6%)	4 (6.3%)	7 (5.4%)			
Generalized Anxiety Disorder	9 (13.8%)	5 (7.8%)	14 (10.8%)			
Panic Disorder	5 (7.7%)	5 (7.8%)	10 (7.8%)			
Social Anxiety Disorder	13 (20%)	16 (25%)	29 (22.5%)			
Major Depressive Disorder	14 (21.5%)	17 (26.6%)	31 (24%)			
Persistent Depressive Disorder	15 (23.1%)	12 (18.8%)	27 (20.9%)			
Somatic Symptom Disorder	2 (3.1%)	3 (6.3%)	5 (3.9%)			
Illness Anxiety Disorder	4 (6.2%)	2 (3.1%)	6 (4.7%)			
Comorbidity						
1 comorbid diagnosis	51 (78.5%)	48 (75%)	99 (76.7%)			
2 comorbid diagnoses	27 (41.5%)	28 (43.7%)	55 (42.6%)			
3 comorbid diagnoses	18 (27.7%)	18 (28.1%)	36 (27.9%)			
4 comorbid diagnoses	12 (18.5%)	10 (15.6%)	22 (17.1%)			
5 comorbid diagnoses	5 (7.7%)	4 (6.2%)	9 (6.9%)			

Table 2*Means, standard deviations/errors, percentage of missing values and effect sizes*

		Treatment			Waitlist			Treatment		
		M	SD/SE	% Missings	M	SD/SE	% Missings	Within Hedge's g baseline to post- module	Within Hedge's g Baseline to post	Between group effect size
Mindfulness (SMQ)	baseline	34.88	15.25	0	34.8	11.83	0			
	module 3	35.6	1.69	15.38	30.95	1.97	46.88	0.05 [-.29, .4]	0.73 [.38, 1.09]	0.76 [0.4, 1.11]
	post	45.7	1.85	36.92	35.7	1.72	17.19			
Cognitive Flexibility (ERQ)	baseline	21.97	7.43	1.53	22.17	7.18	0			
	module 5	26.77	0.95	20.0	23.06	0.95	18.75	0.73 [.38, 1.09]	0.99 [.63, 1.35]	0.86 [0.5, 1.22]
	post	28.77	1.05	38.46	22.96	0.94	17.19			
Anxiety Sensitivity (ASI-3)	baseline	36.12	14.12	1.53	37.97	11.56	1.56	-0.4		-0.65
	module 7	30.58	1.8	35.38	38.96	1.72	20.31	[-.75, -0.05]	-	[-1.0, -0.29]
Experiential Avoidance (BEAQ)	baseline	52.34	9.28	0	53.75	9.53	0	-0.48		-0.59
	module 9	47.77	1.63	53.85	53.83	1.32	20.31	[-0.83, -0.13]	-	[-0.94, -0.24]
Behavioral Activation (BADS)	baseline	23.65	8.21	0	23.52	8.7	0	0.91		0.88
	module 9	31.38	1.47	53.85	23.55	1.17	18.75	[0.55, 1.27]	-	[0.52, 1.24]

Note. SMQ: Southampton Mindfulness Questionnaire, ERQ: Emotion Regulation Questionnaire – Reappraisal Subscale, ASI-3: Anxiety Sensitivity Schedule 3, BEAQ: Brief Experiential Avoidance Questionnaire, BADS: Behavioral Activation for Depression Scale. Between group effect sizes reported have been obtained for the comparison at post for SMQ and ERQ and for module/week for ASI-3, BEAQ, and BADS.

Figure 1*Consort Flow Chart*

Figure 2*Core Concepts of the Intervention and Measurement Points of Mediators and Outcome*

Note. SMQ: Southampton Mindfulness Questionnaire, BSI: Brief Symptom Distress Inventory, ERQ: Emotion Regulation Questionnaire, ASI: Anxiety Sensitivity Index, BEAQ: Brief Experiential Avoidance Questionnaire, BADs: Behavioral Activation for Depression Scale

Figure 3*Single Mediator Models: Indirect Effect of Intervention on Symptom Distress through Transdiagnostic Processes***(1) Mediation through Mindfulness** **(2) Mediation through Cognitive Flexibility** **(3) Mediation through Anxiety Sensitivity** **(4) Mediation through Experiential Avoidance** **(5) Mediation through Behavioral Activation**

Note. Graphical depiction of the mediator models of (1) mindfulness, (2) cognitive flexibility, (3) anxiety sensitivity, (4) experiential avoidance, and (5) behavioral activation as mediators between intervention and symptom distress. Mindfulness and cognitive flexibility were measured twice, after the module that targeted the mechanism (module/week 3 and module/week 5 respectively; t1) and at post-treatment (t2). Anxiety sensitivity, experiential avoidance, and behavioral activation were solely assessed after the module that targeted them (module/week 7 or module/week 9). Symptom distress was measured at post-treatment. Mediators and outcome were controlled by their respective baseline value. These paths were omitted from the figure for clarity. CIs represents upper and lower limit of 95% Bootstrap-corrected confidence intervals. * $p < .05$

Figure 4*Multiple Mediator Model*

Note. Graphical depiction of the multiple mediation model including mindfulness, cognitive flexibility, experiential avoidance, and behavioral activation as mediators between intervention and symptom distress. Mindfulness and cognitive flexibility were measured twice, after the module that targeted the mechanism (module/week 3 and module/week 5 respectively; t1) and at post-treatment (t2). Experiential avoidance and behavioral activation were solely assessed after module/week 9. Symptom distress was measured at post-treatment. Mediators and outcome were controlled by their respective baseline value and mediators were allowed to covary. These paths were omitted from the figure for clarity. Indirect effect 1 and 2 refer to the indirect effects through t1 and t2. CIs represents upper and lower limit of 95% Bootstrap-corrected confidence intervals. * $p < .05$