**Limits of cross-modal plasticity? Short-term visual deprivation does not
enhance cardiac interoception, thermosensation, or tactile spatial acuity**

Dominika Radziun^{1*}, Laura Crucianelli¹, H. Henrik Ehrsson¹

¹ Department of Neuroscience, Karolinska Institutet, Stockholm, Sweden

E-mail addresses: dominika.radziun@ki.se; laura.crucianelli@ki.se; henrik.ehrsson@ki.se

* Corresponding author:

Dominika Radziun

Department of Neuroscience

Karolinska Institutet

Solnavägen 9, 171 65 Solna

Stockholm, Sweden

Telephone: (+46) 8 524 87 989

Email: dominika.radziun@ki.se

Abstract

In the present study, we investigated the effect of short-term visual deprivation on discriminative touch, cardiac interoception, and thermosensation by asking 64 healthy volunteers to perform four behavioral tasks. The experimental group contained 32 subjects who were blindfolded and kept in complete darkness for 110 minutes, while the control group consisted of 32 volunteers who were not blindfolded but were otherwise kept under identical experimental conditions. Both groups performed the required tasks three times: before and directly after deprivation (or control) and after an additional washout period of 40 minutes, in which all participants were exposed to normal light conditions. Our results showed that short-term visual deprivation had no effect on any of the senses tested. This finding suggests that short-term visual deprivation does not modulate basic bodily senses and extends this principle beyond tactile processing to the interoceptive modalities of cardiac and thermal sensations.

Keywords: cross-modal plasticity, blindfolding, interoception, thermosensation, touch

1. Introduction

Neuroplasticity is the brain's capacity to adapt and change in response to phenomena such as learning, developmental factors, and aging as well as injury or a loss of peripheral input (see Pascual-Leone, Amedi, Fregni & Merabet, 2005). Cross-modal plasticity, a type of neuroplasticity, occurs after sensory deprivation, which could be a result of disease, brain damage, or other factors and can lead to the strengthening of one or more sensory systems to compensate for the lack of another, reflecting an adaptive strategy (see Merabet & Pascual-Leone, 2009). Among other hypotheses (see Singh, Phillips, Merabet & Sinha, 2018), it has been proposed that the source of these cross-modal changes could be a process of "unmasking" and subsequent strengthening of weak cross-modal connections that are suppressed under normal conditions (see Pascual-Leone & Hamilton, 2001; Merabet et al., 2007; Merabet et al., 2008; Striem-Amit, Cohen, Dehaene & Amedi, 2012; Qin & Yu, 2013; Lazzouni & Lepore, 2014). Indeed, such pre-existing cortico-cortical connections between, for example, visual areas and other (preserved) modality areas, which are suppressed under normal circumstances, could facilitate information transfer to the visual cortex (Schroeder et al., 2003; Ptito & Kupers, 2005; Masuda, Dumoulin, Nakadomari & Wandell, 2008; Cappe, Rouiller & Barone, 2009; Masuda et al., 2010). Given the compelling results from both animal and human studies (Convento, Vallar, Galantini & Bolognini, 2013; Humanes-Valera, Aguilar & Foffani, 2013; Makin & Bensmaia, 2017), the "unmasking" hypothesis is thought to explain at least some of the general processes observed in the reorganization of the adult cortex (see Singh et al., 2018). Although the specific mechanism underlying the rerouting of non-visual information to the visual cortex has not been fully understood, sensory deprivation studies remain an attractive method of exploring one of the most fascinating properties of the human brain, namely, plasticity.

A well-studied example of massive cross-modal plasticity is the neural changes that follow blindness. Those changes reportedly lead to enhancements in the following senses: touch (e.g., Goldreich & Kanics, 2006; Chebat, Rainville, Kupers & Ptito, 2007; Bauer, Yazzolino, Hirsch, Cattaneo, Vecchi & Merabet, 2015), hearing (e.g., Voss et al., 2004; Gougoux, Zatorre, Lassonde, Voss & Lepore, 2005; Collignon, Voss, Lassonde & Lepore, 2008), and smell (e.g., Rosenbluth, Grossman & Kaitz, 2000; Cuevas, Plaza, Rombaux, De Volder & Renier, 2009; Beaulieu-Lefebvre, Schneider, Kupers & Ptito, 2011; Kupers et al., 2011). Interestingly, a considerable number of studies have suggested that brain plasticity can also be triggered in healthy individuals by short-term visual deprivation for periods as short as 90 minutes – an observation that can shed light on the mechanisms of neuroplasticity (e.g., Facchini & Aglioti, 2003; Weisser et al., 2005; Lewald, 2007; Lazzouni, Voss & Lepore, 2012; Landry, Shiller & Champoux, 2013; Fengler, Nava & Röder, 2015; Pagé, Sharp, Landry & Champoux, 2016; Schwenk, Van Rullen & Bremmer, 2020; but see also: Wong, Hackeman, Hurd & Goldreich, 2011; Crabtree & Norman, 2014; Cambieri et al., 2017). Furthermore, it has been shown that blindfolding in sighted individuals leads to increased excitability of the visual cortex (Boroojerdi et al., 2000; Fierro et al., 2005), which may become engaged in processing non-visual stimuli (Weisser et al., 2005; Merabet et al., 2007; 2008). Taken together, there is evidence that visual deprivation through blindfolding reversibly affects several perceptual abilities, which indicates that short-term deprivation, to some degree, can produce similar perceptual effects as blindness (Merabet et al., 2008).

Recently, sensory abilities related to the body were shown to be altered following visual impairment; for example, studies have suggested that blind people discriminate heat better than sighted individuals (Slimani, Ptito & Kupers, 2015) and present lower pain thresholds for both cold and heat (Slimani et al., 2013). In addition to being somatosensory submodalities, these processes of temperature perception and heat pain have been reclassified as interoception based

on anatomical considerations and the fact that they provide information about the physiological condition of the body, which is a key function of interoception (Craig, 2003a; see also: Khalsa et al., 2018). Classic definitions of interoception were originally focused on visceral sensations only (see Sherrington, 1948), whereas more recent accounts frame interoceptive signals more broadly and include stimuli mediated by the skin and transmitted through lamina I of the spinal cord, e.g., sharp and burning pain, innocuous warmth and cold, itch, or affective touch (see Purves et al., 2019). Such signals help the organism maintain an optimal internal state via the activation of homeostatic mechanisms (see von Mohr & Fotopoulou, 2018). Therefore, interoception, in its broader definition used in this paper, refers to signals originating from the internal body and visceral organs, such as cardiac or gastric sensations, as well as to skin-mediated signals that facilitate homeostasis, such as pain, thermal sensations or affective touch because of their motivational relevance in physiological regulation (see Craig, 2003b; Hua, Strigo, Baxter, Johnson & Craig, 2005; Björnsdotter, Morrison & Olausson, 2010; Fealey, 2013; Ceunen, Vlaeyen & Van Diest, 2016; Gentsch, Crucianelli, Jenkinson & Fotopoulou, 2016; Crucianelli, Krahé, Jenkinson & Fotopoulou, 2018; Gilam, Gross, Wager, Keefe & Mackey, 2020; Wei & Van Someren, 2020).

In a study by Noel et al. (2018) on audiovisual deprivation and cardiac interoceptive accuracy, a very short (15 minutes) deprivation through blindfolding and testing in an anechoic room did not alter interoceptive accuracy on a group level (approximately half of the participants showed improved interoceptive accuracy while the other half showed worse accuracy). It has also been shown that short-term deprivation of exteroceptive senses, including vision, through Reduced Environmental Stimulation Therapy (REST), leads to heightened interoceptive awareness in patients with high levels of anxiety sensitivity (Feinstein et al., 2018). Similarly, REST seems to decrease pain intensity ratings and pain widespreadness in patients with chronic pain (Loose, Manuel, Karst, Schmidt & Beissner, 2021). Interestingly,

Zubek and colleagues (1964) showed that sighted subjects who were visually deprived for a week showed an increase in sensitivity to heat and pain. However, except for these studies, the influence of short-term purely visual deprivation has not yet been examined using a battery of interoceptive tasks, instead focusing on one interoceptive modality. Based on a number of behavioral studies of interoception in blind individuals showing hypersensitivity to heat and cold pain (Slimani et al., 2013), enhanced innocuous heat discrimination (Slimani et al., 2015) and faster central processing of C-fiber input (Slimani, Plaghki, Ptito & Kupers, 2016) following congenital blindness, it could be hypothesized that short-term visual deprivation can also have an effect on interoceptive modalities in sighted individuals.

One of the measures most often employed in interoception research is the heartbeat counting task (Dale & Anderson, 1978; Schandry, 1981), in which participants count their heartbeats for a given amount of time without touching their body and then their estimation is compared with the number of their real recorded heartbeats. The measurement is supposed to determine the participant's access to sensory information from the heart. The task is short and easy to implement, and it should offer a relatively direct measure of cardiac interoception. Compared with the classic cardioceptive heartbeat discrimination task (Katkin, Reed & Deroo, 1983), where participants need to judge whether a sequence of stimuli is presented in synchrony with their heartbeat or not, the heartbeat counting task also offers the advantage of not having additional potentially confounding factors and task demands related to tones or flashes (see Garfinkel et al., 2016a). However, the heartbeat counting task has been criticized in recent years because the results may be influenced by several factors, such as beliefs about or knowledge of the resting heart rate as well as the heart rate itself (Ring, Brener, Knapp, & Mailloux, 2015; Murphy et al., 2018; Ring & Brener, 2018; Zamariola, Maurage, Luminet & Corneille, 2018). Therefore, to identify additional measures of interoceptive submodalities, our group has recently developed a task focused on thermosensation, namely, the thermal matching task

(Crucianelli, Enmalm & Ehrsson, 2021), in which participants are asked to identify a previously perceived thermal stimulus (i.e., a stroke on the skin) in a sequence of colder or warmer stimuli presented in increasing or decreasing order. The results obtained in two separate samples suggest that it is possible to broaden the testable interoceptive modalities beyond cardiac signals to include temperature perception and other skin-based modalities that supposedly also rely on input from C-fibers (Crucianelli et al., 2018, 2021).

In this experiment, we aimed to explore the role of short-term visual deprivation on tactile, thermosensory, and cardiac perception by asking 64 healthy sex-balanced volunteers to perform four behavioral tasks. Three different tasks focusing on two separate interoceptive submodalities, cardiac and thermosensory, were chosen to provide a multifaceted overview of the effects of short-term deprivation on interoception. Cardiac interoceptive perception was operationalized here as the degree of accuracy in the heartbeat counting task (Dale & Anderson, 1978; Schandry, 1981), while thermosensory perception was operationalized as the degree of accuracy in the newly established thermal matching task (Crucianelli et al., 2021) as well as the sensitivity and consistency in detecting temperature changes in the temperature detection task, a widely used task in clinical settings (Fruhstorfer, Lindblom & Schmidt, 1976; see also Heldestad, Linder, Sellersjö & Nordh, 2010). As a measure of tactile acuity, we implemented a commonly used test of passive tactile spatial acuity, the tactile grating orientation task (see Van Boven, Hamilton, Kauffman, Keenan & Pascual-Leone, 2000; Facchini & Aglioti, 2003; Wong et al., 2011a). Thirty-two test subjects were blindfolded and kept in complete darkness for 110 minutes, while the 32 control-group volunteers were not blindfolded and kept under the same experimental settings. Both groups performed the tasks three times: before and directly after deprivation (or control) and after an additional washout period of 40 minutes, in which all participants were exposed to normal light conditions. We tested the hypothesis that a short period of blindfolding would lead to reversible improvement of cardiac and thermal sensations.

We thus predicted that in the thermal matching task, temperature detection task, and heartbeat counting task, participants in the deprived group would show significant improvement after blindfolding and that the effect would disappear when the blindfold was removed. We also re-examined the hypothesis (Facchini & Aglioti, 2003) that blindfolding would improve tactile acuity and predicted that in the tactile grating orientation task, the blindfolded group should show significantly better performance (higher acuity) than the control group.

2. Methods

2.1 Participants

The experiment was completed by a total of 64 healthy right-handed volunteers: 32 in the deprived group (mean age = 26.4, range = 18-39, 16 females, 16 males) and 32 in the non-deprived group (mean age = 26.5, range = 19-46, 16 females, 16 males). The sample size was determined before the experiment started and mirrored the previous study by Crucianelli et al. (2021), who included the same thermal matching and thermal detection tasks as used in the present study as well as the heartbeat counting task; moreover, this sample size is similar to those used before for blindfolding experiments on the tactile acuity task (e.g., Wong et al., 2011). Both groups were sex-balanced due to reports suggesting that women presented higher interoceptive sensibility (tendency to notice bodily sensations more often) but lower accuracy (Grabauskaitė, Baranauskas & Griškova-Bulanova, 2017) as well as a higher performance in the grating orientation task (Wong et al., 2011a). There was no significant difference in age between the groups ($t(62) = -.077$, $p = .939$). Body mass index (BMI) data were collected for the subjects since this parameter has been shown to influence cardiac interoceptive accuracy (Murphy, Geary, Millgate, Catmur & Bird, 2017). The BMI was 22.5 (SD = 3.3) for participants

in the deprived group and 23.2 (SD = 5) in the non-deprived group, with no significant difference between groups ($t(62) = -.68, p = .499$). The average age and BMI were similar to values from other studies of interoception in healthy samples (e.g., Pollatos, Gramann & Schandry, 2006; Garfinkel, Seth, Barrett, Suzuki & Critchley, 2015).

Each subject was assigned to one of the two groups (deprived or non-deprived) according to a randomization schedule. The participants were recruited through advertisements on the campus of Karolinska Institutet and social media and tested within an 8-month period between July and February. All participants reported that they had normal or corrected-to-normal vision. Exclusion criteria included a history of neurological or psychiatric disorders, skin conditions (e.g., psoriasis) or alterations (scars, tattoos), and finger pad calluses or injuries because these conditions can affect the perception of touch, warmth and cold.

The study was approved by the Ethics Review Authority in Sweden (2016/2398-31/4 and 2019-03823), and the experiment was carried out in accordance with the approved guidelines. All participants provided written informed consent before the study. The participants were compensated for participation with 1000 SEK (an equivalent of €100).

2.2 Tasks and procedures

Figure 1. Timeline of the experiment. *Prep* represents the time allotted for the questionnaires and instructions. *S1*, *S2* and *S3* represent session I, session II and session III, respectively.

225 The three testing sessions (I, II, III) consisted of four tasks (see below) and were
226 separated by fixed intervals (Figure 1). The order of tasks was kept constant across the sessions
227 and groups: (1) heartbeat counting task, (2) thermal matching task, (3) tactile grating orientation
228 task, and (4) temperature detection task (see Table 1). We decided on such order to avoid
229 potential effect of thermal and tactile tasks demands on cardiac reactivity, and to separate two
230 thermal tasks with a procedure focusing on another sensory modality. Both groups performed
231 the tasks before and directly after deprivation (or control) and after an additional washout period
232 of 40 minutes, in which all participants were exposed to normal room light conditions and could
233 see the entirety of the testing room. The duration of blindfolding and washout was based on
234 previous studies on short-term visual deprivation (e.g., Fierro et al., 2005; Weisser, Stilla,
235 Peltier, Hu & Sathian, 2005; Lewald, 2007; Merabet et al., 2008; Landry et al., 2013). All of
236 the subjects were blindfolded while performing the tasks to make the experimental conditions
237 identical for both of the groups. The blindfold, which also covered the nose area, prevented all
238 light from reaching the eyes; in the deprived group, medical tape was placed around the mask
239 to avoid accidental displacement. Blinking or eye movements were not prevented by the
240 blindfold. Only the subjects from the deprived group were blindfolded during the interval
241 between sessions I and II, with the blindfold remaining in place from the end of session I to the
242 end of session II, while the subjects from the control group removed the mask and were re-
243 exposed to light after every task during session II. This design allows for a straightforward
244 comparison of baseline performance (no procedural differences between the groups in session
245 I) without potentially confounding factors (i.e., possible effect of light deprivation already within
246 session I due to the relatively long duration of the session). Participants were alert and listening
247 to a previously prepared playlist of music or a podcast of their choice for the whole duration of
248 the blindfolding/control. They were accompanied by the experimenter, who informed
249 participants about the time left every 15 minutes and made sure that the participants did not

show any signs of drowsiness or discomfort. All participants remained alert for the whole duration of the experiment and verbally reported to the experimenter by confirming they understood the information. In the deprived group, during session II, the lights in the room were turned off to ensure that the participants were indeed kept in complete darkness, and the experimenter conducted the task with the minimal light needed to apply the stimuli and record the answers. However, to further make sure that the blindfold indeed covered all the visual input, at the beginning of the deprivation period, participants were asked about their light perception under normal room illumination. Each volunteer confirmed that no light was noticed. Also, by the end of the experiment all participants confirmed that no light was noticed during the entire procedure involving the blindfold. Participants in the non-deprived group spent the remainder of the period in a room with normal light conditions. The same conditions of no view restrictions and normal light were administered for the period between sessions II and III (washout) for both groups. Upon experiment completion, all participants were debriefed about the purpose of the study.

Task order	Task	Task description	Outcome measures
1	Heartbeat counting task	6 trials (25 s, 30 s, 35 s, 40 s, 45 s, 50 s)	Values from 0–1
2	Thermal matching task	3 temperatures (30 °C, 32 °C, 34 °C) 2 body locations (palm and forearm) 2 orders of stimulation (warming and cooling)	Values from 0–1
3	Tactile grating orientation task	20 trials of up to 8 gratings	Grating of 70% accuracy
4	Temperature detection task	Method of limits 5 trials for warming, 5 trials for cooling	Temperature detection and standard deviations

Table 1. Overview of the structure of tasks. The basic tasks described in the order they were completed during the experimental procedure.

Participants were asked to fill out two questionnaires regarding their bodily experiences and psychological functioning (see *Questionnaires*) at the very beginning of the experiment prior to the behavioral tasks, which allowed for any potentially elevated heart rates due to walking/cycling fast pace to the building, etc. to return to a normal level in all participants since increased physiological arousal has been shown to provide an advantage for heartbeat perception (Pollatos, Herbert, Kaufmann, Auer & Schandry, 2007). For the same reason, participants were also asked not to consume any caffeinated drinks on the day of the experiment (see Hartley, Lovallo & Whitsett, 2004; McMullen, Whitehouse, Shine, Whitton, & Towell, 2012). Before the start of the first session, all participants were informed about the experimental setup and received a short description of the experiment. Then, the participants sat on a chair in a comfortable position. The temperature of the testing room was inspected before, during and after the experiment, and it was kept at approximately a neutral temperature of 22.5 °C. The subjects were well adjusted to the room temperature before starting the first behavioral task. All thermal tasks were conducted on the left nondominant palm or forearm, which is consistent with the procedures used in Crucianelli et al. (2021). The grating orientation task, however, was conducted on the right dominant index finger, in accordance with previous experiments on tactile acuity (Facchini & Aglioti, 2003; Wong et al., 2011a). All tasks were administered by the same experimenter (D.R.) in all participants.

2.2.1 Questionnaires

Participants were asked to complete two self-report questionnaires. The Body Awareness Questionnaire (BAQ; Shields, Mallory & Simon, 1989) is an 18-item scale measuring attentiveness to normal bodily processes. The Depression, Anxiety and Stress Scale (DASS-21; Lovibond & Lovibond, 1995) is a self-report questionnaire consisting of 21 items, with 7 items per subscale on depression, anxiety and stress. It was implemented to serve as a control measure for possible subclinical manifestations of depression, anxiety and stress, which have all been suggested to be associated with altered interoception (e.g., Dunn, Dalgleish, Ogilvie & Lawrence, 2007, Domschke, Stevens, Pfleiderer & Gerlach, 2010; Paulus & Stein, 2010, respectively). No significant differences emerged between the groups in any of the measurements (see Table 2).

	BAQ	Depression	Anxiety	Stress
Total sample	84.39 (12.84)	2.28 (2.52)	3.08 (3.19)	4.45 (3.33)
Deprived group	83.81 (11.34)	1.69 (1.96)	2.75 (2.90)	3.84 (2.92)
Non-deprived group	84.97 (14.35)	2.88 (2.88)	3.41 (3.48)	5.06 (3.64)
t (p) values	-.358 (.722)	-1.928 (.058)	-.820 (.415)	-1.478 (.144)

Table 2. Mean and standard deviations for the Body Awareness Questionnaire (BAQ) and each subscale of the Depression, Anxiety and Stress Scale (DASS-21).

2.2.2 Heartbeat counting task

A heart rate baseline reading was obtained over a 5-minute period before the beginning of the heartbeat counting task. The participants' heart rate was recorded using a Biopac MP150 BN-PPGED (Goleta, CA, United States) pulse oximeter attached to their nondominant (left)

308 index finger and connected to a laptop with AcqKnowledge software (version 5.0), which
309 recorded the number of heartbeats after preset time. The number of heartbeats was then
310 quantified using the embedded ‘count peaks’ function. To reduce the possibility that
311 participants would perceive the pulsation in fingers due to the grip of the pulse oximeter, which
312 has been shown to facilitate the performance and inflate the confidence ratings (Murphy et al.,
313 2019), special attention was focused on ensuring a comfortable and not overtight fit of the finger
314 cuff. The resting heart rates were 74.38 BPM (SD = 7.38) in the deprived group and 75.85 BPM
315 (SD = 12.37) in the non-deprived group in S1; 66.75 BPM (SD = 10.03) in the deprived group
316 and 66.13 BPM (SD = 9.69) in the non-deprived group in S2; and 66.06 (SD = 10.80) in the
317 deprived group and 66.17 (SD = 9.68) in the non-deprived group in S3. This finding is
318 consistent with other experiments showing resting heart rates of 68-76 beats per minute for
319 healthy people aged 20-39 years (Hart, 2015), which corresponded to the vast majority of our
320 sample.

321 Participants were asked to breathe normally and given the following instructions:
322 *Without manual checking, can you silently count each heartbeat you feel in your body from the*
323 *time you hear “start” to when you hear “stop”. Do not take your pulse. You are only allowed*
324 *to feel the sensation of your heart beating.* A cue from the experimenter signaled when to start
325 and stop counting. After the trial, the participants verbally reported the number of heartbeats
326 counted, and they did not receive any feedback regarding their performance. Immediately after
327 reporting the number of counted heartbeats, the participants were asked to rate their confidence
328 in perceived accuracy of response (see Garfinkel et al., 2015). This confidence judgment was
329 reported on a scale from 0 (total guess/no heartbeat awareness) to 10 (complete confidence/full
330 perception of heartbeat). To produce a global measure of mean confidence in perceived
331 accuracy of response, the mean confidence during the heartbeat counting task was calculated
332 by averaging the confidence judgments over all experimental trials. The task was repeated six

times to form six trials, using intervals of 25, 30, 35, 40, 45 and 50 seconds (as in the original procedure of Schandry, 1981), with a break of 30 seconds between intervals. Participants received no information about the interval length. The interval order was randomized between participants and sessions.

For each trial, an accuracy score was derived using the formula based on Schandry (1981):

$$\frac{1}{6} \sum \left(1 - \frac{|recorded\ heartbeats - counted\ heartbeats|}{recorded\ heartbeats} \right)$$

The resulting scores were averaged over 6 trials. The interoceptive accuracy scores obtained following this transformation usually vary between 0 and 1, with higher scores indicating a better discrimination of the heartbeats (i.e., smaller differences between estimated and actual heartbeats).

2.2.3 Thermosensory tasks

Before each thermal task, the temperature of the dorsal surface of the left hand and ventral surface of the left hand was measured using an infrared thermometer (Microlife NC 150, Taipei, Taiwan) at three different locations at each site to control for any significant individual differences in skin temperature that could potentially influence the performance in the task (for skin temperature values, see Supplementary Table 1). The thermal stimuli were delivered through a 25 x 50 mm thermode attached to a thermal stimulator (Somedic SenseLab AB, Hörby, Sweden), with a precision of ± 0.1 - 0.2 °C.

2.2.3.1 Thermal matching task

358

359 A range of non-noxious temperatures from 22 °C (cool) to neutral (32 °C; typical human
360 arm skin temperature; Arens and Zhang, 2006) to 42 °C (warm) was applied. The temperatures
361 were presented in a systematically increasing (from cool to warm) or decreasing (from warm
362 to cool) order in trials consisting of gradual changes in temperature (2 °C at a time), with up to
363 9 increments in total. The participant's task was to verbally indicate the temperature that was
364 presented at the beginning of the trial ("reference temperature"). The task was repeated six
365 times to form six trials in an increasing/decreasing manner presented in a randomized order,
366 with 30 °C, 32 °C and 34 °C (within the range of neutral/innocuous temperatures) used as
367 reference temperatures. The temperature at the start of the trial was $\pm 8^{\circ}\text{C}$ of the reference
368 temperature (range from 22-38 °C for 30 °C; range from 24-40 °C for 32 °C; and range from
369 26-42 °C for 34 °C). The same procedure was introduced for the forearm (hairy skin) and palm
370 (non-hairy skin). The starting the task from palm/forearm was counterbalanced across
371 participants and sessions. The duration of each stroke was kept constant at 3 seconds, and the
372 velocity of the touch was approximately 3 cm/s. For a full description of the task, see Crucianelli
373 et al. (2021).

374 Immediately after reporting their perception of the reference temperature, the
375 participants were asked to rate their confidence in the accuracy of the response (see Garfinkel
376 et al., 2015). This confidence judgment was reported using a scale from 0 (total guess) to 10
377 (complete confidence).

378 To calculate the accuracy, the following formula was applied (from Crucianelli et al.,
379 2021):

380

381
$$1 - \left(\sum \frac{(|\text{reported temperature} - \text{reference temperature}|)/2}{12} \right)$$

382

where 12 represents the total number of options presented to the participants (regardless of direction – overestimation or underestimation of temperature) across the three trials. The formula provides a value between 0 and 1, with 0 suggesting poor performance and 1 suggesting optimal performance in the task.

Two variables were introduced: location (palm/forearm) and order of temperature change (increasing/decreasing). For each subject, one increasing value and one decreasing accuracy value for the forearm and for the palm were obtained. To ensure that our analysis was consistent with previous studies showing different skin and central activations for warming and cooling (Hua et al., 2005), we considered the increasing and decreasing trials separately for each skin location.

2.2.3.2 Temperature detection task

The detection of cold and warm static thermal stimuli was measured using the well-established Martsock methods of the limits (Fruhstorfer et al., 1976). We used the same protocol adopted by Heldestad et al. (2010) and Crucianelli et al. (2021). The experimenter kept the thermode on the forearm or palm of the participant without applying additional pressure. Participants were asked to hold a response button in their right hand and to press it as soon as they perceived a change in the temperature in any direction (i.e., warmer or colder than the previously perceived temperature; see Heldestad et al., 2010). The starting point of the stimulation was 32 °C. As soon as the participants pressed the button, the temperature automatically changed in the opposite direction and returned to the baseline temperature, where it stayed for 5 seconds before moving to the next trial. The temperature changed at a rate of 1 °C/s and returned to baseline at a speed of 4 °C/s. Participants completed 5 trials of the task per

order (increasing/decreasing temperature), both on the palm and the forearm. Starting the task from the palm or forearm was counterbalanced across participants and sessions.

We did not ask the participants to rate their confidence in the accuracy of response since the task followed a standardized method, which would be disrupted by applying additional measures.

Optimal performance was operationalized as (1) sensitivity to temperature change, i.e., average difference from the target temperature (32 °C), and (2) consistency in perceiving changes in temperature, i.e., standard deviation of the trials within an increasing (warmth perception)/decreasing (cold perception) block for both palm and forearm (for a similar approach, see Crucianelli et al., 2021).

2.2.4 Tactile grating orientation task

The experimental stimuli consisted of eight hemispheric plastic domes with stamped parallel bars and grooves of equal width (JVP [Johnson-Van Boven-Phillips] Spatial Discrimination Domes, Stoelting, Inc. Wood Dale, IL). The different grating widths were as follows: 0.35, 0.5, 0.75, 1, 1.2, 1.5, 2 and 3 mm. The same exact set of gratings was used in several other studies examining the effect of visual deprivation on tactile acuity (e.g., Van Boven et al., 2000; Merabet et al., 2008; Norman & Bartholomew, 2011; Crabtree & Norman, 2014). The right index finger of the subject was fixated on a table in a palm-up position and immobilized using adhesive tape applied to the nail. Gratings were manually applied to the distal pad of the right index finger for ~1.5 seconds, with moderate force. Previous reports demonstrated that these stimuli can be delivered manually because performance in this task has been shown to be independent of subtle changes in time and pressure of application (Van Boven & Johnson, 1994; Johnson & Phillips, 1981; Vega-Bermudez & Johnson, 1999). The

orientation of the gratings was placed either horizontally or vertically relative to the long axis of the finger. In each trial, a two-alternative forced-choice procedure was used in which subjects had to verbally report whether the grating was oriented horizontally or vertically. The task was terminated when the subject reached the chance level, i.e., 50% or less correct responses. Every experimental session consisted of up to eight blocks, with one for each grating width. Each block consisted of 20 randomized trials, half with horizontal gratings and half with vertical gratings. The sequence of the blocks corresponded to a decreasing width order of the gratings. Care was taken by a trained experimenter to avoid any movement of the finger during contact with the grating. No feedback about the accuracy of the response was given to the subjects at any time.

Immediately after reporting the orientation of the grating, the participants were asked to rate their confidence in the accuracy of the response (see Garfinkel et al., 2015). This confidence judgment was made using a scale from 0 (total guess) to 10 (complete confidence).

The percentage of correct responses was computed for each block, and the grating orientation threshold was calculated by linear interpolation between grating widths spanning 70% correct responses (see Van Boven & Johnson, 1994; Merabet et al., 2008; Wong et al., 2011a; Garfinkel et al., 2016b). Six participants from the deprived group and 9 participants from the non-deprived group were excluded from the data analysis because they could not complete the majority of the test blocks or could not perform the task beyond the expected level (70% accuracy).

2.3 Data analysis

The data were tested for normality using the Shapiro-Wilk test and found to be not distributed normally ($p < .05$). However, we decided to use parametric tests for all analyses

because of their utility in factorial designs and ANOVAs, such as in the current study (see Guterstam, Larsson, Zeberg & Ehrsson, 2019 for a similar approach). The use of non-parametric tests yielded the same results as our parametric approach, which is consistent with the notion that t statistics are reasonably robust to non-normality (Sawilowsky & Blair, 1992; see Supplementary Results). Bonferroni correction was used as a follow-up for significant effects and interactions. All p values were two-tailed. For the Bayesian analyses, the default Cauchy prior was used. For data visualization, raincloud plots in R were used (Allen, Poggiali, Whitaker, Marshall & Kievit, 2019).

3. Results

3.1 Heartbeat counting task

We predicted that in the heartbeat counting task, participants from the deprived group would show significant improvement after blindfolding but that their accuracy would return to the baseline level after the wash-out (light re-exposure) period, while the non-deprived group would not show this pattern of accuracy changes across sessions. However, our results did not support this hypothesis (no main effect of *group*, $F = .039$, $p = .844$, no interaction between *group* and *session*, $F = 1.333$, $p = .267$). Rather, our results revealed that the interoceptive accuracy improved over time, independent of whether the participant had experienced visual deprivation (main effect of *session*, $F = 12.981$, $p < .001$; Figure 2). Thus, no effect of specific blindfolding per se could be established. This conclusion was confirmed by further exploratory analyses: in the blindfolded group, the Bonferroni-corrected post hoc comparisons ($\alpha = 0.017$) revealed a significant difference between sessions I and II ($t(31) = -2.838$, $p = .008$; $M_1 = .747$, $SD_1 = .151$; $M_2 = .800$, $SD_2 = .122$) and no significant difference between sessions II and III

482 $(t(31) = .641, p = .526; M_3 = .789, SD_3 = .127)$. Similarly, in the non-blindfolded group, we
483 found a significant difference between session I and session II ($t(31) = -3.147, p = .004; M_1 =$
484 $.720, SD_1 = .125; M_2 = .790, SD_2 = .146$) and no significant difference between session II and
485 session III ($t(31) = -.932, p = .358; M_3 = .809, SD_3 = .121$). Further exploratory analysis of
486 performance between the groups during each session did not reveal statistically significant
487 differences (session I, session II and session III: $t(62) = .783, p = .437; t(62) = .283, p = .778;$
488 $t(62) = -.633, p = .529$, respectively).

489 To test whether our data provided evidence for the absence of an interaction of group
490 and session, which would support the null hypothesis, we performed a 2 x 3 Bayesian ANOVA.
491 The Bayesian analysis revealed a Bayes factor of 7.072 in favor of the null hypothesis ($BF_{01} =$
492 7.072) of no interaction between *group* and *session*, indicating that the data were 7.072 times
493 more likely under the null hypothesis than under the alternative hypothesis. Furthermore, a
494 Bayesian paired t-test run for a direct comparison of S2 between the blindfolded and control
495 groups revealed a Bayes factor of 3.784 ($BF_{01} = 3.784$) in support of the null hypothesis.

496 Given that beliefs about or knowledge of the resting heart rate as well as the heart rate
497 itself have been shown to influence the performance of the heartbeat counting task (see the
498 *Introduction*), we decided to run an exploratory analysis examining potential fluctuations in
499 heart rate across sessions. We found no main effect of *group* ($F = .020, p = .889$) but a
500 significant main effect of *session* ($F = 60.238, p < .001$), with no *group* x *session* interaction (F
501 $= .654, p = .522$), suggesting that the heart rate of all participants changed (decreased)
502 significantly throughout the course of the experiment. Then, we ran another exploratory
503 analysis examining whether the average number of reported heartbeats changed across sessions
504 in any of the groups. We found no main effect of *group* ($F = .001, p = .982$), no main effect of
505 *session* ($F = 1.151, p = .320$), and no significant interaction between *group* and *session* ($F =$
506 2.993, $p = .054$). Taken together, the improvement in accuracy might be driven by natural

fluctuations in heart rate and not by task demands, which is consistent with previous studies (see Introduction).

The pattern observed in the accuracy measurements was mirrored by the confidence ratings: we did not observe a main effect of *group* ($F = .172, p = .680$) or an interaction between *group* and *session* ($F = .759, p = .470$) but did observe a main effect of *session* ($F = 5.634, p = .005$; Supplementary Figure 1). These findings show that not only the accuracy but also confidence increased across sessions, regardless of the group, although without any notable effect from the blindfolding procedure.

Figure 2. Accuracy in the heartbeat counting task. Individual data points, boxplots, density plots and group means with standard errors are shown. All the following figures are formatted in the same fashion.

The baseline performance in the heartbeat counting task in both groups was comparable with the results obtained in other studies using this paradigm (e.g., Borhani, Ládavas,

Fotopoulou & Haggard, 2017), which highlights that the task was successfully implemented in the present study.

3.2 Thermal matching task

We predicted that in the thermal matching task, participants in the deprived group would show significant reversible improvement after blindfolding while participants in the non-deprived group would not show these changes in accuracy across sessions. The analysis of the effect of visual deprivation revealed no main effect of *group* ($F = .686$, $p = .411$), although a significant main effect of *session* ($F = 3.551$, $p = .032$) and a significant interaction between *group* and *session* ($F = 3.502$, $p = .033$; Figure 3) were observed. Bonferroni-corrected post hoc tests ($\alpha = 0.017$) revealed that there was no significant difference in the average accuracy between the groups in any of the sessions ($t(62) = 1.886$, $p = .064$; $t(62) = -.285$, $p = .777$; $t(62) = .542$, $p = .590$ for sessions I, II and III; $M_1 = .839$, $SD_1 = .051$; $M_2 = .836$, $SD_2 = .064$; $M_3 = .844$, $SD_3 = .072$ in the deprived group and $M_1 = .798$, $SD_1 = .110$; $M_2 = .842$, $SD_2 = .097$; $M_3 = .833$, $SD_3 = .099$ in the non-deprived group). However, further Bonferroni-corrected post hoc comparisons ($\alpha = 0.017$) revealed a significant difference between sessions I and II ($t(31) = -3.223$, $p = .003$) and no significant difference between sessions II and III ($t(31) = .947$, $p = .351$) in the control group. In turn, in the blindfolded group, we did not find a significant difference between session I and session II ($t(31) = .226$, $p = .823$) or between session II and session III ($t(31) = -.757$, $p = .455$). Taken together, these results suggest that visual deprivation did not significantly influence thermosensation as measured by the thermal matching task.

Figure 3. Average accuracy across conditions in the thermal matching task.

In line with Crucianelli et al. (2021), we found that both groups showed higher accuracy in the baseline session when stimulated on the forearm compared to the palm (main effect of *location [arm/palm]*, $F = 27.697$, $p < .001$; Supplementary Figure 2). Therefore, we reproduced the basic effect of the thermal matching task on a group with the same number of participants who had similar demographic backgrounds as reported in the original paper of Crucianelli et al. (2021; see *Participants*), which was examined to highlight that the task was successfully implemented in the present study. The analysis of all sessions revealed no interaction between *group* and *location* ($F = .730$, $p = .396$) or between *group* and *temperature* ($F = .323$, $p = .572$), suggesting that the blindfolding procedure did not influence the *forearm* vs. *palm* difference (Supplementary Figure 3).

In the case of the confidence ratings, we did not observe a main effect of *group* ($F = .020$, $p = .887$) or an interaction between *group* and *session* ($F = .058$, $p = .943$), and we also did not observe a main effect of *session* ($F = .820$, $p = .443$; Supplementary Figure 4), thus

showing that confidence did not increase across sessions, regardless of the group. As in Crucianelli et al. (2021), we observed a significant main effect of *location* ($F = 21.776$, $p < .001$), with participants being more confident with their answers for the forearm than the palm. We did not observe a main effect of *temperature* ($F = 2.266$, $p = .137$) or an interaction between *location* and *temperature* ($F = .352$, $p = .555$). Our analyses revealed no interaction between *group* and *location* ($F = .700$, $p = .406$) or between *group* and *temperature* ($F = .082$, $p = .776$).

3.3 Temperature detection task

As in the thermal matching task (Section 3.2), we predicted that participants in the deprived group would show significant reversible improvement after blindfolding while participants in the non-deprived group would not show these changes in accuracy across sessions. The analysis of the effect of visual deprivation on sensitivity to temperature change revealed no main effect of *group* ($F = 1.274$, $p = .263$), no main effect of *session* ($F = 1.532$, $p = .220$), and no interaction between *group* and *session* ($F = .686$, $p = .505$; Figure 4A).

Figure 4. Sensitivity to temperature change (A) and consistency in perceiving temperature change (B) in the temperature detection task.

Similarly, we observed no main effect of *group* ($F = .919$, $p = .341$), no main effect of *session* ($F = .041$, $p = .960$), and no interaction between *group* and *session* ($F = .126$, $p = .882$; Figure 4B) in terms of consistency in perceiving temperature change (Figure 4B).

To test whether our data provided evidence for the absence of an interaction of session and group, which would support the null hypothesis, we performed a 2 x 3 Bayesian ANOVA, separately for sensitivity and for consistency. For sensitivity, the Bayesian analysis revealed a Bayes factor of 8.889 in favor of the null hypothesis ($BF_{01} = 8.889$) of no interaction between *group* and *session*. Similarly, for consistency, the Bayesian analysis revealed a Bayes factor of 9.495 in favor of the null hypothesis ($BF_{01} = 9.495$) of no interaction between *group* and *session*. Therefore, our results suggest that visual deprivation does not influence thermosensation as measured by the temperature detection task.

Additionally, we found that both groups showed higher sensitivity in the baseline session when detecting cooling stimuli compared to warming stimuli (main effect of *temperature*, $F = 204.040$, $p < .001$), thereby replicating the effect observed in Crucianelli et al. (2021). The analysis of all sessions revealed no interaction between *group* and *temperature* ($F = .007$, $p = .933$) or between *group* and *location* ($F = 1.844$, $p = .179$; Supplementary Figure 5).

3.4 Tactile grating orientation task

We predicted that in the grating orientation task, the blindfolded group would show significantly better performance (higher acuity) than the control group in the second session, which will return to the baseline level after light re-exposure, and the control group would not show such changes in accuracy across sessions. However, these predictions were not met. We found no main effect of *group* ($F = .110$, $p = .742$), no main effect of *session* ($F = 1.628$, $p = .202$) and no interaction between *group* and *session* ($F = .120$, $p = .887$; Figure 5). An exploratory analysis of the differences in performance between the groups during each session revealed no statistical significance (session I, session II and session III: $t(47) = .001$, $p = .999$;

609 $t(47) = -.288, p = .774$; $t(47) = -.497, p = .621$, respectively; $M_1 = 1.463, SD_1 = .514$; $M_2 =$
610 $1.279, SD_2 = .459$; $M_3 = 1.308, SD_3 = .554$ in the deprived group, respectively, and $M_1 = 1.463,$
611 $SD_1 = .721$; $M_2 = 1.324, SD_2 = .642$; $M_3 = 1.397, SD_3 = .703$ in the non-deprived group,
612 respectively). Overall, visual deprivation did not have a significant effect on tactile acuity, and
613 tactile acuity did not improve (or change) over time in either of the two groups.

614 To test whether our data provide evidence for the absence of an interaction of session
615 and group, which would support the null hypothesis, we performed a Bayesian 2 x 3 ANOVA.
616 Bayesian analysis revealed a Bayes factor of 8.19 ($BF_{01} = 8.19$) in favor of the null hypothesis
617 of no interaction of session and group, indicating that the data were 8.19 times more likely
618 under the null hypothesis than under the alternative hypothesis. Furthermore, a Bayesian paired
619 t-test run for a direct comparison of S2 between the blindfolded and control groups revealed a
620 Bayes factor of 3.386 ($BF_{01} = 3.386$) in support of the null hypothesis. Therefore, our results
621 suggest that visual deprivation does not influence tactile acuity as measured by the grating
622 orientation task.

623

624 **Figure 5.** Tactile grating orientation threshold.

The baseline performance in the task in both groups was comparable with the results obtained in other studies using this paradigm (Merabet et al., 2008; Wong et al., 2011; Bola et al., 2016), which highlights that the task was successfully implemented in the present study.

3.5 Relationship across tasks

No significant relationship was found between performance in any of the tasks (Supplementary Table 2). The lack of significant correlations between tasks further supports the idea of independent processing across interoceptive submodalities (in line with Crucianelli et al., 2021; see also Ferentzi et al., 2018).

4. Discussion

In this study, we tested the influence of short-term (110 minutes) visual deprivation in healthy adults on cardiac interoception, thermosensation, and tactile spatial acuity to systematically address the potential influence of visual deprivation on bodily senses through cross-modal plasticity in sighted individuals. Both the deprived and non-deprived (control) groups performed a battery of tasks three times: before and directly after deprivation (or control) and after an additional washout period of 40 minutes, in which all participants were exposed to normal light conditions. We found that both cardiac interoception and skin-based interoception (thermosensation) were resistant to the effects of short-term deprivation, which was confirmed by the observation of the exact same pattern of results in both the classic (static temperature detection task) and newly established (dynamic thermal matching task) thermosensation tests as well as the lack of effect of blindfolding on performance in the heartbeat counting task. We

also did not observe a blindfolding-driven change in tactile spatial acuity. Taken together, our results showed no effect on any of the senses tested, suggesting that basic bodily senses are resistant to cross-modal plastic changes induced by short-term visual deprivation.

In a study by Noel et al. (2018) on audiovisual deprivation and cardiac interoceptive accuracy, a very short deprivation period of 15 minutes did not alter interoceptive accuracy in a significant way when considering the whole sample. However, the result of Noel and colleagues might simply be explained by the brief duration of blindfolding because such short periods of visual deprivation have not been shown to lead to any changes on behavioral or neural levels, with 30 minutes being the shortest known deprivation period to produce a reliable effect (Leon-Sarmiento, Bara-Jimenez & Wassermann, 2005). However, why was a period of 110 minutes insufficient to increase the cardiac interoceptive accuracy? One reason could be that the potential sensory enhancements within interoceptive senses are not driven by ‘pure’ cross-modal compensatory plasticity processes but require training periods to be reinforced, which is similar to the tactile improvement observed after very prolonged visual deprivation in blindness (see Alary et al., 2009; Sathian & Stilla, 2010; Voss, 2011; Wong, Gnanakumaran & Goldreich, 2011; for a cardiac interoceptive training example, see Quadt, Mulcahy, Critchley & Garfinkel, 2020). Moreover, Slimani and colleagues (2013) did not find a difference in thresholds for non-painful thermal stimulation between blind and sighted groups, which might suggest that thermosensation is overall resistant to the effects of both short- and long-term visual deprivation, including congenital blindness. In a follow-up experiment, Slimani and colleagues (2014) showed no pain hypersensitivity in late blind individuals, which suggests that enhanced sensitivity to pain following blindness is potentially a result of brain plasticity changes related to early but not late vision loss. Taken together, it could be speculated that some processes ascribed to interoception (e.g., heat pain perception) might be altered only as a result of congenital visual deprivation; moreover, some of these processes (cardiac and thermal

interoception) might remain unchanged even under such circumstances, which was further confirmed by the lack of influence of short-term deprivation on the thermal abilities of sighted individuals found in our study.

In keeping with recent research (Wong, Hackeman, Hurd & Goldreich, 2011; Crabtree & Norman, 2014) but inconsistent with the original findings of Facchini and Aglioti (2003), we did not see an improvement in tactile spatial acuity after short-term visual deprivation. Our results follow a number of studies in blind individuals in which improved tactile acuity in blindness were suggested to be mostly driven by experience-dependent mechanisms – for example, increased training of fingertips due to braille reading, and not necessarily by the lack of vision itself (Alary et al., 2009; Sathian & Stilla, 2010; Voss, 2011; see also Wong, Gnanakumaran & Goldreich, 2011).

Interestingly, our negative results in the four bodily tasks examining single modalities were also consistent with recent studies showing multisensory but not unisensory enhancement following short-term deprivation. Fengler and colleagues (2015) did not find any changes in the basic perceptual tasks implemented in their procedure (two unisensory perceptual threshold measures, auditory and visual) but showed a reduced interference effect on multisensory affective prosody judgments. It is worthy to note, however, that multisensory audio-visual discrimination task was not influenced by the blindfolding procedure. Furthermore, in the study of Radziun and Ehrsson (2018), which used a non-visual version of the well-known paradigm of rubber-hand illusion (Botvinick & Cohen, 1998; Ehrsson, Holmes & Passingham, 2005) to probe the dynamic plasticity of body representation, participants from the blindfolded group showed a significantly larger recalibration of hand position sense towards the location of the rubber hand than the control group (“proprioceptive drift”), which is a commonly used behavior index of the illusion. However, the blindfolded group’s accuracy in localizing their finger before the illusion, i.e., a unisensory proprioceptive task, showed no significant difference from

700 the control non-deprived group. Similarly, Petkova and colleagues (2012) did not find a
701 difference between blind and sighted participants in a proprioceptive task testing the basic
702 proprioceptive ability to localize their hand in space without vision, although they did observe
703 an altered (abolished) somatic rubber hand illusion in the blind group (Petkova et al., 2012).
704 The spatial recalibration associated with the somatic rubber hand illusion depends on the
705 integration of congruent tactile and proprioceptive signals from the two upper limbs (Ehrsson
706 et al., 2005; Petkova, Zetterberg & Ehrsson 2012), which is a process that can be implemented
707 by sensory integration mechanisms in the frontal and parietal association cortices and the
708 cerebellum (Ehrsson et al 2005). This more complex integration process of bodily signals was
709 specifically affected in both blindfolded (Radziun & Ehrsson, 2018) and blind (Petkova et al.,
710 2012) participants, in contrast to basic proprioception, tactile acuity, or interoception, that
711 presumably rely predominantly on the primary and secondary somatosensory cortex (e.g.,
712 Eickhoff et al., 2006; Khalsa, Rudrauf, Feinstein & Tranel, 2009; Haag et al., 2015; Lutz &
713 Bensmaia, 2021) and the insula (e.g., Livneh et al., 2017; Evrard, 2019). The present negative
714 findings also suggest that the previously observed effects of blindfolding on the recalibration
715 of the felt hand position (in the somatic rubber hand illusion; Radziun & Ehrsson, 2018) were
716 due to the altered multisensory integration rather than changes in tactile acuity (i.e., increased
717 sensitivity to tactile incongruence), thermosensation (increased sensitivity to thermal
718 incongruence), or cardiac interoception (Tsakiris, Tajadura- Jiménez & Costantini, 2011, but
719 see also Crucianelli et al., 2018; Horváth et al., 2020). Moreover, multisensory integration
720 within the bodily senses has been shown to be altered in blind individuals, suggesting that visual
721 experiences shape both behavioral and neural responses to tactile–proprioceptive stimulation
722 (Crollen et al., 2017), which again points to the role of vision in multimodal interactions, even
723 when visual input is not directly involved.

Blindfolding in sighted individuals has been shown to modify the excitation/inhibition balance in the visual cortex and lead to increased activation of the visual areas (Boroojerdi et al., 2000; Fierro et al., 2005), which have been reported to become engaged in processing non-visual stimuli (Weisser et al., 2005; Merabet et al., 2007; 2008). Moreover, short-term visual deprivation was demonstrated to be associated with increased excitability of the motor cortex (Leon-Sarmiento et al., 2005); however, the evidence is mixed (Cambieri et al., 2017). Among the investigations of the effects of blindfolding on the brain, electrophysiological studies have shown signatures of improvement of haptic recognition memory (Santaniello, Sebastián, Carretié, Fernández-Folgueiras & Hinojosa, 2018), plasticity of the auditory steady-state response (Lazzouni et al., 2012), and slow-wave changes in cortical visual areas (Bernardi et al., 2019). However, to the best of our knowledge, there are no neuroimaging or electrophysiological studies examining the effects of visual deprivation on active areas and electrophysiological signatures related to cardiac interoception or thermosensation, such as by using the insula as a region of interest. Further studies might throw light on the potential links between various forms of visual deprivation and bodily senses on a neural level.

Importantly, the results of the heartbeat counting task, in which the performance of both groups was compared across three sessions, are consistent with studies highlighting the effect of repeated performance on participants' accuracy (e.g., Ring, Brener, Knapp & Mailloux, 2015). This finding may indicate that the heartbeat counting task is not optimally suited to quantifying cardiac interoception in repeated-measures designs (for a recent debate on the validity of the heartbeat counting task, see: Zamariola et al., 2018; Ainley, Tsakiris, Pollatos, Schulz & Herbert, 2020; Zimirich, Nusser & Pollatos, 2020; Corneille, Desmedt, Zamariola, Luminet & Maurage, 2020). In contrast, none of the thermal tasks showed an effect of practice on the performance. Thus, thermosensation might provide more stable and robust results regarding the consistency and reliability of participants' performance.

Blindfolding paradigms provide a useful method of inducing and measuring behavioral proxies of neuroplasticity, with the aim of better understanding the rapid plastic changes in the brain. Our work suggests that in cases of cardiac interoception, thermosensation, and discriminative touch, 110 minutes of visual deprivation is not enough to produce any changes on a behavioral level. Further studies might help elucidate why improvement of only some perceptual processes and abilities can be observed after short-term visual deprivation and why basic bodily sensations, such as cardiac interoception, thermosensation, tactile acuity, or proprioception, do not seem to be affected.

Funding

This work was supported by the Swedish Research Council (grant no: 2017-03135). Laura Crucianelli was supported by the Marie Skłodowska-Curie Intra-European Individual Fellowship (grant no: 891175). The funding sources had no involvement in study design, collection, analyses and interpretation of the data or in writing this paper.

Competing Interests

The authors declare no competing interests.

References

Ainley, V., Tsakiris, M., Pollatos, O., Schulz, A., & Herbert, B. M. (2020). Comment on “Zamariola et al. (2018), Interoceptive Accuracy Scores are Problematic: Evidence from Simple Bivariate Correlations”—The empirical data base, the conceptual reasoning and the analysis behind this statement are misconceived and do not support the authors’ conclusions. *Biological Psychology*, 152, 107870.

Alary, F., Duquette, M., Goldstein, R., Chapman, C. E., Voss, P., Buissonnière-Ariza, V. L., & Lepore, F. (2009). Tactile acuity in the blind: A closer look reveals superiority over the sighted in some but not all cutaneous tasks. *Neuropsychologia*, 47(10), 2037-2043.

Allen, M., Poggiali, D., Whitaker, K., Marshall, T. R., & Kievit, R. A. (2019). Raincloud Plots: A multi-platform tool for robust data visualization. *Wellcome Open Research*, 4, 63.

Arens, E., & Zhang, H. (2006). The skin's role in human thermoregulation and comfort. In: *Thermal and moisture transport in fibrous materials* (Pan N., Gibson P., eds), pp 560–602. Cambridge, UK: Woodhead Publishing Ltd.

Bauer, C., Yazzolino, L., Hirsch, G., Cattaneo, Z., Vecchi, T., & Merabet, L. B. (2015). Neural correlates associated with superior tactile symmetry perception in the early blind. *Cortex*, 63, 104-117.

Beaulieu-Lefebvre, M., Schneider, F. C., Kupers, R., & Ptito, M. (2011). Odor perception and odor awareness in congenital blindness. *Brain Research Bulletin*, 84(3), 206-209.

Bernardi, G., Betta, M., Cataldi, J., Leo, A., Haba-Rubio, J., Heinzer, J., Cirelli, C., Tononi, G., Pietrini, P., Ricciardi, E., Siclari, F. (2019). Visual imagery and visual perception induce similar changes in occipital slow waves of sleep. *Journal of Neurophysiology*, 121, 2140-2152.

Björnsdotter, M., Morrison, I., & Olausson, H. (2010). Feeling good: On the role of C fiber mediated touch in interoception. *Experimental Brain Research*, 207(3-4), 149-155.

Bola, Ł., Siuda-Krzywicka, K., Paplińska, M., Sumera, E., Hańczur, P., & Szwed, M. (2016). Braille in the Sighted: Teaching Tactile Reading to Sighted Adults. *Plos One*, 11(5).

Boroojerdi, B. (2000). Enhanced Excitability of the Human Visual Cortex Induced by Short-term Light Deprivation. *Cerebral Cortex*, 10(5), 529-534.

Borhani, K., Lâdavas, E., Fotopoulou, A. & Haggard, P. (2017). “Lacking warmth”: Alexithymia trait is related to warm-specific thermal somatosensory processing. *Biological Psychology*, 128, 132-140.

Botvinick, M., & Cohen, J. (1998). Rubber hands ‘feel’ touch that eyes see. *Nature*, 391(6669), 756-756.

Brener, J., & Ring, C. (2016). Towards a psychophysics of interoceptive processes: The measurement of heartbeat detection. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 371(1708).

Cambieri, C., Iacovelli, E., Gori, M. C., Onesti, E., Ceccanti, M., Frasca, V., & Inghilleri, M. (2017). Effects of visual deprivation on primary motor cortex excitability: A study on healthy subjects based on repetitive transcranial magnetic stimulation. *Experimental Brain Research*, 235(7), 2059-2067.

Cappe, C., Rouiller, E., & Barone, P. (2009). Multisensory anatomical pathways. *Hearing Research*, 258(1-2), 28-36.

Ceunen, E., Vlaeyen, J. W., & Diest, I. V. (2016). On the Origin of Interoception. *Frontiers in Psychology*, 7.

Chebat, D., Rainville, C., Kupers, R., & Ptito, M. (2007). Tactile–‘visual’ acuity of the tongue in early blind individuals. *NeuroReport*, 18(18), 1901-1904.

Collignon, O., Voss, P., Lassonde, M., & Lepore, F. (2008). Cross-modal plasticity for the spatial processing of sounds in visually deprived subjects. *Experimental Brain Research*, 192(3), 343-358.

Convento, S., Vallar, G., Galantini, C., & Bolognini, N. (2013). Neuromodulation of Early Multisensory Interactions in the Visual Cortex. *Journal of Cognitive Neuroscience*, 25(5), 685-696.

Couto, B., Salles, A., Sedeño, L., Peradejordi, M., Barttfeld, P., Canales-Johnson, A., . . . Ibanez, A. (2013). The man who feels two hearts: The different pathways of interoception. *Social Cognitive and Affective Neuroscience*, 9(9), 1253-1260.

Corneille, O., Desmedt, O., Zamariola, G., Luminet, O., & Maurage, P. (2020). A heartfelt response to Zimprich et al. (2020), and Ainley et al. (2020)'s commentaries: Acknowledging issues with the HCT would benefit interoception research. *Biological Psychology*, 152, 107869.

Crabtree, C. E., & Norman, J. F. (2014). Short-Term Visual Deprivation, Tactile Acuity, and Haptic Solid Shape Discrimination. *PLoS ONE*, 9(11).

Craig, A. D., Chen, K., Bandy, D., & Reiman, E. M. (2000). Thermosensory activation of insular cortex. *Nature Neuroscience*, 3(2), 184-190.

Craig, A. D. (2003a). How do you feel? Interoception: The sense of the physiological condition of the body. *Nature Reviews Neuroscience*, 3(8), 655-666.

Craig, A. D. (2003b). Pain Mechanisms: Labeled Lines Versus Convergence in Central Processing. *Annual Review of Neuroscience*, 26(1), 1-30.

Crollen, V., Lazzouni, L., Rezk, M., Bellemare, A., Lepore, F., & Collignon, O. (2017). Visual experience shapes the neural networks remapping touch into external space. *Journal of Neuroscience*, 37(42), 10097-10103.

Crucianelli, L., Krahé, C., Jenkinson, P. M., & Fotopoulou, A. (2018). Interoceptive ingredients of body ownership: Affective touch and cardiac awareness in the rubber hand illusion. *Cortex*, 104, 180-192.

Crucianelli, L., Enmalm, A., & Ehrsson, H. H. (2021). Thermosensation as a novel method to probe interoception and its relationship with other interoceptive modalities. *bioRxiv*.

Cuevas, I., Plaza, P., Rombaux, P., Volder, A. G., & Renier, L. (2009). Odour discrimination and identification are improved in early blindness. *Neuropsychologia*, 47(14), 3079-3083.

Dale, A., & Anderson, D. (1978). Information Variables in Voluntary Control and Classical Conditioning of Heart Rate: Field Dependence and Heart-Rate Perception. *Perceptual and Motor Skills*, 47(1), 79-85.

Domschke, K., Stevens, S., Pfleiderer, B., & Gerlach, A. L. (2010). Interoceptive sensitivity in anxiety and anxiety disorders: An overview and integration of neurobiological findings. *Clinical Psychology Review*, 30(1), 1-11.

Dunn, B. D., Dalgleish, T., Ogilvie, A. D., & Lawrence, A. D. (2007). Heartbeat perception in depression. *Behaviour Research and Therapy*, 45(8), 1921-1930.

Ehrsson, H. H., Holmes, N. P., & Passingham, R. E. (2005). Touching a Rubber Hand: Feeling of Body Ownership Is Associated with Activity in Multisensory Brain Areas. *Journal of Neuroscience*, 25(45), 10564-10573.

- Eickhoff, S. B., Lotze, M., Wietek, B., Amunts, K., Enck, P., & Zilles, K. (2006). Segregation of visceral and somatosensory afferents: An fMRI and cytoarchitectonic mapping study. *NeuroImage*, 31(3), 1004-1014.
- Ernst, M. O., & Banks, M. S. (2002). Humans integrate visual and haptic information in a statistically optimal fashion. *Nature*, 415(6870), 429-433.
- Evrard, H. C. (2019). The organization of the primate insular cortex. *Frontiers in Neuroanatomy*, 13.
- Facchini, S., & Aglioti, S. M. (2003). Short term light deprivation increases tactile spatial acuity in humans. *Neurology*, 60(12), 1998-1999.
- Fealey, R. D. (2013). Interoception and autonomic nervous system reflexes thermoregulation. *Autonomic Nervous System Handbook of Clinical Neurology*, 79-88.
- Feinstein, J. S., Khalsa, S. S., Yeh, H., Al Zoubi, O., Arevian, A. C., Wohlrab, C., Pantino, M. K., Cartmell, L. J., Simmons, W. K., Stein, M. B., & Paulus, M. P. (2018). The elicitation of relaxation and interoceptive awareness using floatation therapy in individuals with high anxiety sensitivity. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, 3(6), 555-562.
- Fengler, I., Nava, E., & Röder, B. (2015). Short-term visual deprivation reduces interference effects of task-irrelevant facial expressions on affective prosody judgments. *Frontiers in Integrative Neuroscience*, 9.
- Ferentzi, E., Bogdány, T., Szabolcs, Z., Csala, B., Horváth, Á., & Köteles, F. (2018). Multichannel investigation of interoception: Sensitivity is not a generalizable feature. *Frontiers in Human Neuroscience*, 12.
- Fierro, B., Brighina, F., Vitello, G., Piazza, A., Scalia, S., Giglia, G., Daniele, O. & Pascual-Leone, A. (2005). Modulatory effects of low- and high-frequency repetitive transcranial magnetic stimulation on visual cortex of healthy subjects undergoing light deprivation. *The Journal of Physiology*, 565(2), 659-665.
- Fruhstorfer, H., Lindblom, U., & Schmidt, W. C. (1976). Method for quantitative estimation of thermal thresholds in patients. *Journal of Neurology, Neurosurgery & Psychiatry*, 39(11), 1071-1075.
- Garfinkel, S. N., Seth, A. K., Barrett, A. B., Suzuki, K., & Critchley, H. D. (2015). Knowing your own heart: Distinguishing interoceptive accuracy from interoceptive awareness. *Biological Psychology*, 104, 65-74.
- Garfinkel, S. N., Tiley, C., O'Keeffe, S., Harrison, N. A., Seth, A. K., & Critchley, H. D. (2016a). Discrepancies between dimensions of interoception in autism: Implications for emotion and anxiety. *Biological Psychology*, 114, 117-126.
- Garfinkel, S. N., Manassi, M. F., Hamilton-Fletcher, G., In den Bosch, Y., Critchley, H. D., & Engels, M. (2016b). Interoceptive dimensions across Cardiac and respiratory axes. *Philosophical Transactions of the Royal Society B*, 371 (1708).
- Gentsch, A., Crucianelli, L., Jenkinson, P., & Fotopoulou, A. (2016). The Touched Self: Affective Touch and Body Awareness in Health and Disease. In *Affective Touch and the Neurophysiology of CT Afferents* (pp. 355-384). Springer: New York.
- Gilam, G., Gross, J. J., Wager, T. D., Keefe, F. J., & Mackey, S. C. (2020). What Is the Relationship between Pain and Emotion? Bridging Constructs and Communities. *Neuron*, 107(1), 17-21.

Goldreich, D., & Kanics, I. M. (2006). Performance of blind and sighted humans on a tactile grating detection task. *Perception & Psychophysics*, 68(8), 1363-1371.

Gougoux, F., Zatorre, R. J., Lassonde, M., Voss, P., & Lepore, F. (2005). A Functional Neuroimaging Study of Sound Localization: Visual Cortex Activity Predicts Performance in Early-Blind Individuals. *PLoS Biology*, 3(2).

Goldreich, D., & Kanics, I. M. (2006). Performance of blind and sighted humans on a tactile grating detection task. *Perception & Psychophysics*, 68(8), 1363-1371.

Grabauskaitė, A., Baranauskas, M., & Griškova-Bulanova, I. (2017). Interoception and gender: What aspects should we pay attention to? *Consciousness and Cognition*, 48, 129-137.

Green, B. G., & Akirav, C. (2007). Individual differences in temperature perception: Evidence of common processing of sensation intensity of warmth and cold. *Somatosensory & Motor Research*, 24(1-2), 71-84.

Guterstam, A., Larsson, D. E., Zeberg, H., & Ehrsson, H. H. (2019). Multisensory correlations—Not tactile expectations—Determine the sense of body ownership. *Plos One*, 14(2).

Haag, L. M., Heba, S., Lenz, M., Glaubitz, B., Höffken, O., Kalisch, T., . . . Schmidt-Wilcke, T. (2015). Resting bold fluctuations in the primary somatosensory cortex correlate with tactile acuity. *Cortex*, 64, 20-28.

Hart, J. (2015). Normal resting pulse rate ranges. *Journal of Nursing Education and Practice*, 5(8), 95–98.

Hartley, T. R., Lovallo, W. R., & Whitsett, T. L. (2004). Cardiovascular effects of caffeine in men and women. *The American Journal of Cardiology*, 93(8), 1022–1026.

Heldestad, V., Linder, J., Sellersjö, L., & Nordh, E. (2010). Reproducibility and influence of test modality order on thermal perception and thermal pain thresholds in quantitative sensory testing. *Clinical Neurophysiology*, 121(11), 1878-1885.

Horváth, Á., Ferentzi, E., Bogdány, T., Szolcsányi, T., Witthöft, M., & Köteles, F. (2020). Proprioception but not cardiac interoception is related to the rubber hand illusion. *Cortex*, 132, 361–373.

Hua, L. H., Strigo, I. A., Baxter, L. C., Johnson, S. C., & Craig, A. D. (Bud). (2005). Anteroposterior somatotopy of innocuous cooling activation focus in human dorsal posterior insular cortex. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 289(2), R319–R325.

Humanes-Valera, D., Aguilar, J., & Foffani, G. (2013). Reorganization of the Intact Somatosensory Cortex Immediately after Spinal Cord Injury. *PLoS ONE*, 8(7).

Johnson, K. O., & Phillips, J. R. (1981). Tactile spatial resolution. I. Two-point discrimination, gap detection, grating resolution, and letter recognition. *Journal of Neurophysiology*, 46(6), 1177-1192.

Katkin, E. S., Reed, S. D., & Deroo, C. (1983). A methodological analysis of 3 techniques for the assessment of individual-differences in heartbeat detection. *Psychophysiology*, 20(4), 452-452.

Khalsa, S. S., Rudrauf, D., Feinstein, J. S., & Tranel, D. (2009). The pathways of interoceptive awareness. *Nature Neuroscience*, 12(12), 1494-1496.

Khalsa, S. et al. (2017). Interoception and Mental Health: A Roadmap. *Biological Psychiatry: CNI*, 3, 501-513.

Kupers, R., Beaulieu-Lefebvre, M., Schneider, F., Kassuba, T., Paulson, O., Siebner, H., & Ptito, M. (2011). Neural correlates of olfactory processing in congenital blindness. *Neuropsychologia*, 49(7), 2037-2044.

Landry, S. P., Shiller, D. M., & Champoux, F. (2013). Short-term visual deprivation improves the perception of harmonicity. *Journal of Experimental Psychology: Human Perception and Performance*, 39(6), 1503-1507.

Lazzouni, L., Voss, P., & Lepore, F. (2012). Short-term crossmodal plasticity of the auditory steady-state response in blindfolded sighted individuals. *European Journal of Neuroscience*, 35(10), 1630-1636.

Lazzouni, L., & Lepore, F. (2014). Compensatory plasticity: Time matters. *Frontiers in Human Neuroscience*, 8.

Lewald, J. (2007). More accurate sound localization induced by short-term light deprivation. *Neuropsychologia*, 45(6), 1215-1222.

Leon-Sarmiento, F. E., Bara-Jimenez, W., & Wassermann, E. M. (2005). Visual deprivation effects on human motor cortex excitability. *Neuroscience Letters*, 389(1), 17-20.

Livneh, Y., Ramesh, R. N., Burgess, C. R., Levandowski, K. M., Madara, J. C., Fenselau, H., . . . Andermann, M. L. (2017). Homeostatic circuits selectively gate food cue responses in insular cortex. *Nature*, 546(7660), 611-616.

Loose, L. F., Manuel, J., Karst, M., Schmidt, L. K., & Beissner, F. (2021). Flotation restricted environmental stimulation therapy for chronic pain. *JAMA Network Open*, 4(5).

Lovibond, S.H. & Lovibond, P.F. (1995). Manual for the Depression Anxiety & Stress Scales. (2nd Ed.) Sydney: Psychology Foundation.

Lutz, O. J., & Bensmaia, S. J. (2021). Proprioceptive representations of the hand in somatosensory cortex. *Current Opinion in Physiology*, 21, 9-16.

Makin, T. R., & Bensmaia, S. J. (2017). Stability of Sensory Topographies in Adult Cortex. *Trends in Cognitive Sciences*, 21(3), 195-204.

Masuda, Y., Dumoulin, S. O., Nakadomari, S., & Wandell, B. A. (2008). V1 Projection Zone Signals in Human Macular Degeneration Depend on Task, not Stimulus. *Cerebral Cortex*, 18(11), 2483-2493.

Masuda, Y., Horiguchi, H., Dumoulin, S. O., Furuta, A., Miyauchi, S., Nakadomari, S., & Wandell, B. A. (2010). Task-Dependent V1 Responses in Human Retinitis Pigmentosa. *Investigative Ophthalmology & Visual Science*, 51(10), 5356.

McMullen, M. K., Whitehouse, J. M., Shine, G., Whitton, P. A., & Towell, A. (2012). Caffeine in hot drinks elicits cephalic phase responses involving cardiac activity. *Food & Function*, 9.

Merabet, L. B., Swisher, J. D., McMains, S. A., Halko, M. A., Amedi, A., Pascual-Leone, A., & Somers, D. C. (2007). Combined Activation and Deactivation of Visual Cortex During Tactile Sensory Processing. *Journal of Neurophysiology*, 97(2), 1633-1641.

Merabet, L. B., Hamilton, R., Schlaug, G., Swisher, J. D., Kiriakopoulos, E. T., Pitskel, N. B., Kauffman, T. & Pascual-Leone, A. (2008). Rapid and Reversible Recruitment of Early Visual Cortex for Touch. *PLoS ONE*, 3(8).

Merabet, L. B., & Pascual-Leone, A. (2009). Neural reorganization following sensory loss: The opportunity of change. *Nature Reviews Neuroscience*, 11(1), 44-52.

- Murphy, J., Geary, H., Millgate, E., Catmur, C., & Bird, G. (2017). Direct and indirect effects of age on interoceptive accuracy and awareness across the adult lifespan. *Psychonomic Bulletin & Review*, 25(3), 1193-1202.
- Murphy, J., Millgate, E., Geary, H., Ichijo, E., Coll, M., Brewer, R., Catmur, C. & Bird, G. (2018). Knowledge of resting heart rate mediates the relationship between intelligence and the heartbeat counting task. *Biological Psychology*, 133, 1-3.
- Murphy, J., Brewer, R., Coll, M.-P., Plans, D., Hall, M., Shiu, S. S., Catmur, C. & Bird, G. (2019). I feel it in my finger: Measurement device affects cardiac interoceptive accuracy. *Biological Psychology*, 148, 107765.
- Norman, J. F., & Bartholomew, A. N. (2011). Blindness enhances tactile acuity and haptic 3-D shape discrimination. *Attention, Perception, & Psychophysics*, 73(7), 2323-2331.
- Noel, J.-P., Park, H.-D., Pasqualini, I., Lissek, H., Wallace, M., Blanke, O., & Serino, A. (2018). Audio-visual sensory deprivation degrades visuo-tactile peri-personal space. *Consciousness and Cognition*, 61, 61–75.
- Pagé, S., Sharp, A., Landry, S. P., & Champoux, F. (2016). Short-term visual deprivation can enhance spatial release from masking. *Neuroscience Letters*, 628, 167-170.
- Pascual-Leone, A., & Hamilton, R. (2001). The metamodal organization of the brain. *Progress in Brain Research Vision: From Neurons to Cognition*, 427-445.
- Pascual-Leone, A., Amedi, A., Fregni, F., & Merabet, L. B. (2005). The plastic human brain cortex. *Annual Review of Neuroscience*, 28(1), 377-401.
- Paulus, M. P., & Stein, M. B. (2010). Interoception in anxiety and depression. *Brain Structure and Function*, 214(5-6), 451-463.
- Petkova, V. I., Zetterberg, H., & Ehrsson, H. H. (2012). Rubber Hands Feel Touch, but Not in Blind Individuals. *PLoS ONE*, 7(4).
- Pollatos, O., Gramann, K., & Schandry, R. (2006). Neural systems connecting interoceptive awareness and feelings. *Human Brain Mapping*, 28(1), 9-18.
- Pollatos, O., Schandry, R., Auer, D. P., & Kaufmann, C. (2007). Brain structures mediating cardiovascular arousal and interoceptive awareness. *Brain Research*, 1141, 178-187.
- Pollatos, O., Gramann, K., & Schandry, R. (2006). Neural systems connecting interoceptive awareness and feelings. *Human Brain Mapping*, 28(1), 9-18.
- Pollatos, O., Herbert, B. M., Kaufmann, C., Auer, D. P. & Schandry, R. (2007). Interoceptive awareness, anxiety and cardiovascular reactivity to isometric exercise. *International Journal of Psychophysiology*, 65(2), 167–173.
- Ptito, M., & Kupers, R. (2005). Cross-Modal Plasticity In Early Blindness. *Journal of Integrative Neuroscience*, 4(4), 479-488.
- Qin, W., & Yu, C. (2013). Neural Pathways Conveying Novisual Information to the Visual Cortex. *Neural Plasticity*, 1-14.
- Quadt, L., Mulcahy, J., Critchley, H. D. & Garfinkel, S. N. (2020). Impact of interoceptive training on anxiety in autistic adults: the ‘Aligning Dimensions of Interoceptive Experience’ (ADIE) trial. *Psychosomatic Medicine*, 82(6), A208-A209.
- Radziun, D., & Ehrsson, H. H. (2018). Short-term visual deprivation boosts the flexibility of body representation. *Scientific Reports*, 8(1).

Ring, C., Brener, J., Knapp, K., & Mailloux, J. (2015). Effects of heartbeat feedback on beliefs about heart rate and heartbeat counting: A cautionary tale about interoceptive awareness. *Biological Psychology*, 104, 193-198.

Ring, C., & Brener, J. (2018). Heartbeat counting is unrelated to heartbeat detection: A comparison of methods to quantify interoception. *Psychophysiology*, 55(9).

Rosenbluth, R., Grossman, E. S., & Kaitz, M. (2000). Performance of Early-Blind and Sighted Children on Olfactory Tasks. *Perception*, 29(1), 101-110.

Santaniello, G., Sebastián, M., Carretié, L., Fernández-Folgueiras, U., & Hinojosa, J. A. (2018). Haptic recognition memory following short-term visual deprivation: Behavioral and neural correlates from ERPs and alpha band oscillations. *Biological Psychology*, 133, 18-29.

Sathian, K., & Stilla, R. (2010). Cross-modal plasticity of tactile perception in blindness. *Restorative Neurology and Neuroscience*, 28(2), 271-281.

Sawilowsky, S. S., & Blair, R. C. (1992). A more realistic look at the robustness and Type II error properties of the t test to departures from population normality. *Psychological Bulletin*, 111(2), 352-360.

Schandry, R. (1981). Heart Beat Perception and Emotional Experience. *Psychophysiology*, 18(4), 483-488.

Schroeder, C. E., Smiley, J., Fu, K. G., McGinnis, T., O'Connell, M. N., & Hackett, T. A. (2003). Anatomical mechanisms and functional implications of multisensory convergence in early cortical processing. *International Journal of Psychophysiology*, 50(1-2), 5-17.

Sherrington, C.S. (1948). *The Integrative Action of the Nervous System*. Cambridge, UK: Cambridge University Press.

Schwenk, J. C. B., Van Rullen, R. & Bremmer, F. (2020). Dynamics of Visual Perceptual Echoes Following Short-Term Visual Deprivation. *Cerebral Cortex Communications*, 1, 1-11.

Striem-Amit, E., Cohen, L., Dehaene, S., & Amedi, A. (2012). Reading with Sounds: Sensory Substitution Selectively Activates the Visual Word Form Area in the Blind. *Neuron*, 76(3), 640-652.

Shields, S. A., Mallory, M. E., & Simon, A. (1989). The Body Awareness Questionnaire: Reliability and Validity. *Journal of Personality Assessment*, 53(4), 802-815.

Singh, A. K., Phillips, F., Merabet, L. B., & Sinha, P. (2018). Why Does the Cortex Reorganize after Sensory Loss? *Trends in Cognitive Sciences*, 22(7), 569-582.

Slimani, H., Danti, S., Ricciardi, E., Pietrini, P., Ptito, M., & Kupers, R. (2013). Hypersensitivity to pain in congenital blindness. *Pain*, 154(10), 1973-1978.

Slimani, H., Danti, S., Ptito, M., & Kupers, R. (2014). Pain Perception Is Increased in Congenital but Not Late Onset Blindness. *PLoS ONE*, 9(9).

Slimani, H., Ptito, M., & Kupers, R. (2015). Enhanced heat discrimination in congenital blindness. *Behavioural Brain Research*, 283, 233-237.

Slimani, H., Plaghki, L., Ptito, M., & Kupers, R. (2016). Pain hypersensitivity in congenital blindness is associated with faster central processing of C-fibre input. *European Journal of Pain*, 20(9), 1519-1529.

Templeton, G. F. (2011). A two-step approach for transforming continuous variables to normal: implications and recommendations for IS research. *Communications of the Association for Information Systems*, 28(1), 4.

- 1078 Tsakiris, M., Jiménez, A. T., & Costantini, M. (2011). Just a heartbeat away from one's
1079 body: Interoceptive sensitivity predicts malleability of body-representations. *Proceedings of*
1080 *the Royal Society B: Biological Sciences*, 278(1717), 2470-2476.
- 1081 Van Boven, R. W., & Johnson, K. O. (1994). A psychophysical study of the mechanisms
1082 of sensory recovery following nerve injury in humans. *Brain*, 117(1), 149-167.
- 1083 Van Boven, R. W., Hamilton, R. H., Kauffman, T., Keenan, J. P., & Pascual-Leone, A.
1084 (2000). Tactile spatial resolution in blind Braille readers. *Neurology*, 54(12), 2230-2236.
- 1085 von Mohr, M., & Fotopoulou, A. (2018). The Cutaneous borders of interoception:
1086 Active and social inference of pain and pleasure on the skin. In M. Tsakiris & H. de Preester
1087 (Eds.), *Interoceptive Mind: From Homeostasis to Awareness* (1st ed., pp. 102–120). Oxford,
1088 United Kingdom: Oxford University Press.
- 1089 Vega-Bermudez, F., & Johnson, K. O. (1999). SA1 and RA Receptive Fields, Response
1090 Variability, and Population Responses Mapped with a Probe Array. *Journal of*
1091 *Neurophysiology*, 81(6), 2701-2710.
- 1092 Voss, P., Lassonde, M., Gougoux, F., Fortin, M., Guillemot, J., & Lepore, F. (2004).
1093 Early- and Late-Onset Blind Individuals Show Supra-Normal Auditory Abilities in Far-Space.
1094 *Current Biology*, 14(19), 1734-1738.
- 1095 Voss, P. (2011). Superior Tactile Abilities in the Blind: Is Blindness Required? *Journal*
1096 *of Neuroscience*, 31(33), 11745-11747.
- 1097 Voss, P. (2018). Brain (re)organization following visual loss. *Wiley Interdisciplinary*
1098 *Reviews: Cognitive Science*, 10(1).
- 1099 Weisser, V., Stilla, R., Peltier, S., Hu, X., & Sathian, K. (2005). Short-term visual
1100 deprivation alters neural processing of tactile form. *Experimental Brain Research*, 166(3-4),
1101 572-582.
- 1102 Wei, Y., & Someren, E. J. (2020). Interoception relates to sleep and sleep disorders.
1103 *Current Opinion in Behavioral Sciences*, 33, 1-7.
- 1104 Wong, M., Hackman, E., Hurd, C., & Goldreich, D. (2011a). Short-Term Visual
1105 Deprivation Does Not Enhance Passive Tactile Spatial Acuity. *PLoS ONE*, 6(9).
- 1106 Wong, M., Gnanakumaran, V., & Goldreich, D. (2011b). Tactile Spatial Acuity
1107 Enhancement in Blindness: Evidence for Experience-Dependent Mechanisms. *Journal of*
1108 *Neuroscience*, 31(19), 7028-7037.
- 1109 Zamariola, G., Maurice, P., Luminet, O., & Corneille, O. (2018). Interoceptive accuracy
1110 scores from the heartbeat counting task are problematic: Evidence from simple bivariate
1111 correlations. *Biological Psychology*, 137, 12–17.
- 1112 Zimprich, D., Nusser, L., & Pollatos, O. (2020). Are interoceptive accuracy scores from
1113 the heartbeat counting task problematic? A comment on Zamariola et al. (2018). *Biological*
1114 *Psychology*, 152, 107868.
- 1115 Zubek, J. P., Flye, J., & Aftanas, M. (1964). Cutaneous Sensitivity after Prolonged
1116 Visual Deprivation. *Science*, 144(3626), 1591-1593.