Title:

The self and its nature: A psychopathological perspective on the risk-reducing effects of environmental green space for psychosis

Author:

Sjoerd J.H. Ebisch
Department of Neuroscience, Imaging and Clinical Sciences
Institute of Advanced Biomedical Technologies (ITAB)
G. d'Annunzio University of Chieti-Pescara
Chieti (CH), Italy

Correspondence to:

Sjoerd J. H. Ebisch
s.ebisch@unich.it

Keywords:

intrinsic self, extrinsic self, environment, green space, natural surroundings, psychosis, schizophrenia, urban, self-disturbance

ABSTRACT

Epidemiological studies have shown that environmental green space contributes to the reduction of psychosis incidence in the population. Clarifying the psychological and neuro-functional mechanisms underlying the risk-decreasing effects of green surroundings could help optimize preventive environmental interventions. This perspective article specifically aims to open a new window on the link between environmental green space and psychosis by considering its core psychopathological features. Psychotic disorders, such as schizophrenia, are essentially characterized by self-disturbances. The psychological structure of the self has been described as a multidimensional phenomenon that emerges from the reciprocal interaction with the environment through intrinsic and extrinsic self-processes. The intrinsic self refers to the experience of mental activity and environmental information as inherently related to one's own person, which involves self-referential processing, self-reflection, memory, interoception, and emotional evaluation. The extrinsic self refers to sensorimotor interactions with the environment and the sense of agency, that is, the experience of being the source of one's own actions and the multisensory consequences thereof. In psychosis, anomalous self-processing has been related to a functional fragmentation of intrinsic and extrinsic self-processes and related brain networks. Moreover, evidence from cognitive neuroscience suggests that green space could have beneficial effects on self-related processing. Based on the literature, it could be hypothesized that self-processing is involved in mediating the beneficial effects of green space for psychosis. Considering the multidimensionality of the self, it is proposed that urban green space design aimed at improving mental health ideally impacts the complexity of self-facets and thus restores the individual's self.

INTRODUCTION

Psychotic disorders are severe mental illnesses that involve a “loss of vital contact with reality” (Parnas and Henriksen, 2014; Minkowski, 1927). The most studied and pervasive psychotic disorder is likely schizophrenia, but psychotic episodes can happen in various psychiatric, neurological, and neurodevelopmental conditions and can also be induced by the use of medications or psychoactive drugs. Socioeconomically, psychosis has been associated with premature mortality, morbidity, financial and social burdens, and poor outcomes (Fazel et al., 2014; van Os and Linscott, 2012; Rössler et al., 2005). Psychoses, particularly schizophrenia, place large demands on public healthcare and its budgets (Rössler et al., 2005).

The high heritability of psychosis indicates the existence of genetic risk factors (Smigielski et al., 2020), although genetic components cannot fully explain the development of psychosis (van Os et al., 2010; Pries et al., 2018; Torrey and Yolken, 2019). The risk of developing a psychosis during one’s life is also determined by biological, psychological, and social factors (McGrath et al., 2004; van Os et al., 2010; Fusar-Poli et al., 2017). This knowledge provides important information for intervention programs aimed at detecting high-risk individuals and preventing them from becoming psychotic (Ruhrman et al., 2010; van Os et al., 2017). However, the implementation of large-scale intervention strategies is still limited. In addition to therapeutic interventions (Fusar-Poli et al., 2019; McGorry et al., 2009), other possibilities for preventive intervention have been implied. Epidemiological studies suggest that environmental factors are involved in the development of psychosis as well (van Os et al., 2010; Dragt et al., 2011; Fusar-Poli et al., 2017). Moreover, longitudinal studies suggest that the modulation of psychosis incidence through environmental factors likely involves causal relationships that cannot be explained by other mediating epidemiological variables (van Os et al., 2010).

Some of these factors may be attractive targets for global risk-reducing environmental interventions (Sussman and Hollander, 2014). Among the various environmental factors that have been related to psychosis risk, urbanicity has been consistently associated with an elevated risk of psychosis as compared to living in rural environments (Lewis et al., 1992; Pedersen and Mortensen, 2001; van Os et al., 2004; Newbury et al., 2016; Sundquist et al., 2004; see for reviews, Krabbendam and van Os, 2005; Kelly et al., 2010). Given that the majority of the world population currently lives in urban areas and continues to increase, this is an issue of concern for mental healthcare (United Nations, 2014). One notable difference between urban and rural environments that has recently received increasing attention in the scientific literature concerning environmental effects on mental health is the quantity and kind of natural or green space (Verheij et al., 2008), which could be hypothesized as constituting a risk-modulating factor for psychosis. Although it should be emphasized that the diminished availability of green space is by no means the only characteristic of urban environments and that psychosis incidence in the population is likely modulated by a complexity of interacting environmental and genetic factors (Schmitt et al., 2014; Tsuang, 2000), in the present article, I will specifically examine the utilization potential of green space to reduce the incidence of psychosis.

In particular, this perspective article aims at developing a new hypothesis regarding the psychological and neuronal mechanisms that could explain the risk-decreasing effects of natural surroundings for psychosis. For the purpose of focus, an investigation of how the effects of green space on psychosis incidence can be embedded in the full complexity of genetic and environmental factors and events remains beyond the scope of the present article, although this will be a mandatory topic for ensuing dialogues. A better understanding of the underlying risk-reducing mechanisms of concrete environmental features could help in optimally exploiting these features in urban design. Given the fact that structural and direct investigations in compromising mental disorders like psychosis are still sparse, one approach to exploring these mechanisms would be to link the core psychopathological features of psychosis to green space effects.

GREEN SPACE EXPOSURE IS RELATED TO REDUCED PSYCHOSIS RISK

Natural environments usually refer to surroundings that include green space (essentially typified by the presence of vegetation) but also can include blue space (water bodies), which could be developed with or without human intervention (Smith et al., 2017; Völker and Kistemann, 2011; Bratman et al., 2012). The definition of green space, as used in studies that investigate its psychological effects through experience, covers a broad range of environments, from pristine nature to human cultivated gardens, and more specifically includes urban greens, forests/woodlands, countryside/farmland, and the wilderness (Bratman et al., 2012). Urban green space, in turn, can be characterized by the more formal or informal design of gardens and parks (Twedt et al., 2016). The efficiency of green space exposure in modulating health and wellbeing has been studied more generally in terms of the presence of green space in people's living environments and also more specifically in terms of visual exposure (e.g., window views and virtual environments) and physically spending time in or interacting with green space (Bratman et al., 2012).

The beneficial effects of natural surroundings on daily life have been shown for physical and mental health, social well-being, academic and job performance, and happiness (Maas et al., 2009; Hartig, 1993; Hartig et al., 2014; James et al., 2015; van den Berg et al., 2015; Cohen-Cline et al., 2015; Sarkar et al., 2018; White et al., 2013; Bratman et al., 2012). Although the nature of such effects remains to be clarified, natural surroundings, as compared to urban areas, are generally thought to decrease air and noise pollution, improve social cohesion and stress regulation, facilitate the restoration of attention and fatigue, improve the mood, and reduce depressive symptoms (Markevych et al., 2017; Twohig-Bennett et al., 2018; Bratman et al., 2012; Kaplan and Kaplan, 1989; Ullrich et al., 1991; Berman et al., 2008). A recent study highlighted the relevance of environmental interventions by showing that individuals frequenting urban environments with few green resources and an increased risk for mental illness are particularly receptive of the presence of urban green space (Tost et al., 2019). Some further evidence suggests that being exposed to real green and blue space induces psychophysiological and behavioral responses that can be differentiated from those induced by the same environments when reproduced by visual media (Huang et al., 2009, 2019). Similarly, a recent meta-analysis showed that real natural settings benefit mood to a greater degree than simulated settings (Browning et al., 2020).

To date, the relationship between the presence of residential natural spaces and a lower risk for psychosis has been investigated by few studies. Boers and colleagues (2018) showed that psychotic patients had a significantly lower amounts of available green space (agricultural areas, natural areas, and artificially installed greenery) but not blue space near their residence, while controlling for age, gender, urbanicity, and socioeconomic status. The same study did not find any significant relationship between green space and the length of stay in a psychiatric ward as a measure of illness severity. However, because no other measures of illness severity were included, further studies may need to consider more direct measures as well, including illness impact and symptomatology. A nation-wide, population-based study including more than 900,000 people showed that a decreased presence of green space near one's residence during childhood (calculated as the normalized difference vegetation index, NDVI, based on remote sensing satellite images) was associated with higher incidence rate ratios for many psychiatric disorders, including schizophrenia and related psychotic disorders (Engemann et al., 2019a). A dose-response relationship over time was also observed in this study, and the lowest levels of green space were associated with up to a 55% higher risk of psychiatric disorders. Additionally, when adjusting for urbanization, parents' socioeconomic status, family history, parental age, and municipal socioeconomic factors, these effects remained significant for most disorders. Another nationwide population-based study by the same group focusing on schizophrenia further showed a dose-response relationship between the amount of residential green space present during childhood (NDVI), but not green space heterogeneity, and the risk of developing

schizophrenia later in life (Engemann et al., 2018). Also, in this case, the results remained stable after adjustment for urbanization, age, sex, and socioeconomic status. Finally, in a series of follow-up studies, Engemann and colleagues more specifically associated agricultural areas and near-natural green and blue space with lower schizophrenia rates as compared to urban areas, whereas vegetation density (NVDI) was negatively associated with schizophrenia rates in a dose-response manner independently for urban and agricultural areas (Engemann et al., 2019b). Moreover, based on hazard ratios, additive effects on schizophrenia risk were found for childhood green space exposure (NVDI close to residence based on remote sensing satellite images) and genetic liability, while an interaction between these factors could not be detected (Engemann et al., 2020). Taken together, these epidemiological results suggest that the presence of and access to natural surroundings in rural and urban areas may reflect a modulatory environmental factor that contributes to the prevention of psychosis and that green space effects may be disentangled from urbanicity and genetic effects.

DISRUPTED SELF-PROCESSING IS A CORE FEATURE OF PSYCHOSIS

One of the questions that is derived from these studies is “What are the critical aspects of psychosis that might be influenced by green space?” From a historical, psychopathological perspective, schizophrenia and related psychotic disorders have been described as self-disorders (Raballo et al., 2011). Building on these early insights, more recently, psychosis has been characterized as a complexity of self-disturbances that is generally described by an exaggerated self-consciousness and a disrupted sense of owning one’s personal perceptions and thoughts (Sass and Parnas, 2003; Nelson et al., 2014). Because self-disturbances are often reported as subclinical phenomena that have already occurred prior to the onset of psychosis, remain stable after the attenuation of frank psychotic symptoms, and are highly predictive of conversion to psychosis, self-disturbances are commonly considered to reflect the core features of psychosis (Nelson et al., 2012; Schultze-Lutter, 2009). These properties of self-disturbances make them interesting targets for preventive intervention programs. Indeed, as reviewed above, green space may positively act on the mechanisms underlying psychosis onset during childhood.

To develop a more concrete understanding of the exact nature of self-disturbances in psychosis, it is useful to take a closer look at the psychological structure of the self-concept. The self and its disturbance have been increasingly portrayed from multiple perspectives in psychology and cognitive neuroscience. A major distinction is that between intrinsic and extrinsic self-processes (Vanhaudenhuyse et al., 2011; Gallagher, 2000), which can be traced to the self-aspects of the “Me,” as the conscious person who is known by himself or herself (e.g., “this is me”), and the “I,” who experiences and interacts with the environment (e.g., “I did that”), as described by William James (James, 1890). Specifically, intrinsic self-processing refers to the perception of information as belonging to oneself or as personally relevant, which allows self-reflection and experiencing a sense of identity (van der Meer et al., 2010; Damasio, 1999; Northoff and Bermpohl, 2004). This is mainly an internally directed process that integrates stimuli and thoughts with inner information from memory, personal narrative, and interoception and is related to emotional evaluation through the link with transient bodily states (van der Meer et al., 2010; Northoff and Panksepp, 2008; Qin and Northoff, 2011). In contrast, extrinsic self-processing regards the experience of oneself as the source of one’s own actions and their consequences, that is, the perception of oneself as an entity with a sense of agency (Gallagher, 2000; Haggard, 2017). It can be conceived of as an externally directed process based on the experience of one’s intentional control over the environment through action (Jeannerod, 2003; Gallese and Sinigaglia, 2010). Action-related processes that could be involved include multisensory integration, self-monitoring, and sensorimotor predictions (Blanke, 2012). For instance, the integration of predicted somatosensory, motor, visual, and proprioceptive perceptions with the actual multisensory action consequences is crucial for a coherent self-experience during intentional behavior (Frith et al., 2000; van Kemenade et al., 2019).

This literature suggests that the self is composed of distinguishable but complementary processes. Such processes have been associated with largely distinct brain networks (Vanhaudenhuyse et al., 2011). However, in order to permit self-awareness, the networks involved in intrinsic and extrinsic self-processing (e.g., default mode, sensorimotor, attention, and executive networks) likely interact (Molnar-Szakacs and Uddin, 2013), possibly with an additional modulatory role for the salience network in the insula (Menon and Uddin, 2010). Indeed, recent empirical findings from neuroimaging (Di Plinio et al., 2020) suggest that intrinsic and extrinsic self-processes constitute a multidimensional phenomenon that emerges from a bidirectional interaction between an internal self and its external environment, where such interactions could be facilitated by efficient information exchange and integration across intrinsic and extrinsic self-networks in the brain.

Considering psychosis from this perspective, it can be proposed that anomalous intrinsic and extrinsic self-processes, as well as their functional imbalance, could explain self-disturbances. In particular, it has been proposed that the experience of a disrupted sense of self in its reciprocal interaction with the environment in psychosis could be explained by the fragmented functioning of intrinsic and extrinsic self-networks (Ebisch and Aleman, 2016). Furthermore, a biopsychological neural model has been proposed to describe how psychotic symptoms can be explained via a disequilibrium of network states and interference between network activities due to impaired auto-excitation and collateral inhibition (Looijestijn et al., 2015). As a consequence, at the phenomenological level, the distinction between internally and externally generated information may blur, leading to impaired self-recognition, depersonalization, and the tendency to experience one's thoughts, internal speech, or actions as belonging to an external agent or force, rather than oneself (Waters et al., 2012; Nelson et al., 2009; Kircher and Leube, 2003). Because whether and how the beneficial effects of natural surroundings on psychosis can be related to these self-disturbances remain poorly understood, attempts to clarify this issue would require examining the way in which environmental factors associated with natural spaces affect the cognitive and neural mechanisms underlying self-related processing.

CAN GREEN SPACE IMPROVE SELF-PROCESSING?

To address this particular question in more detail, a literature search was performed for articles empirically investigating the relationship between green space and self-processing. PubMed and Google Scholar were screened for peer-reviewed articles published from January 1990 to June 2020, using the terms “self” AND “nature experience” OR “nature exposure” OR “green space” OR “natural space.” Similar searches were performed for studies considering self-disturbances, using the terms “psychosis” OR “schizophrenia” AND “self” AND “nature experience” OR “nature exposure” OR “green space” OR “natural space.” Articles that resulted from these searches and relevant references cited in those articles were reviewed, and this selection was further completed by searches in the author's personal files, where articles published in English were included. The literature search results showed that this specific issue received limited attention. After close inspection of the articles' content, seven empirical studies were found to be relevant to the present perspective article (i.e., Hayhurst et al., 2015; Hunter et al., 2010; Lederbogen et al., 2010; Margalit and Ben-Ari, 2014; Bratman et al., 2015; O'Brien and Lomas, 2017; Fuller et al., 2017), which are discussed below.

A recent systematic review (Mygind et al., 2019) concluded that there is conditional evidence in children and adolescents for a beneficial effect on the part of diversified nature experiences on self-efficacy, a concept closely related to the sense of agency, though this was based on studies with a note regarding the risk of biased results (mountain and lake activities in O'Brien and Lomas, 2017; wilderness therapy intervention in Margalit and Ben-Ari, 2014; outdoor residential experiences in countryside with woodland in Fuller et al., 2017; and developmental sail voyage experiences in Hayhurst et al., 2015, as well as in Hunter et al., 2010). Similarly positive effects were reported for self-esteem, resilience, and academic and cognitive performance, whereas evidence regarding self-concept, problem solving, and mood were inconclusive (see Mygind et al., 2019). In addition, some

cognitive neuroscience studies began to provide relevant insights into the cognitive and neural mechanisms underlying the association between green space and self-processing. In a cross-sectional, functional magnetic resonance imaging (fMRI) study involving non-clinical, healthy individuals, Lederbogen and colleagues (2011) dissociated the impact of urban upbringing and city living on brain activity when participants were exposed to social evaluative stress. Their results showed that current city living was linked to increased amygdala activity, whereas urban upbringing affected the perigenual anterior cingulate cortex. In another fMRI study, Bratman and colleagues (2015) investigated the link between nature experiences (a 90-minute walk in green space consisting of grassland with scattered trees, shrubs, and fauna, as well as views of hills and a bay, versus an urban walk in a busy city thoroughfare) and rumination in a sample of non-clinical, healthy participants. Relevantly, rumination can be considered a dysfunctional self-process defined as a maladaptive pattern of self-referential thought and self-relational emotions. Behaviorally, the results showed that the nature experiences, as compared to the urban experiences, led to decreased rumination. Neuro-functionally, nature experiences, as compared to urban experiences, led to decreased neural activity (cerebral blood flow) in the subgenual prefrontal cortex and perigenual anterior cingulate cortex during rumination.

Notably, both fMRI studies (Lederbogen et al, 2011; Bratman et al., 2015) linked the ventromedial frontal cortex, specifically the perigenual anterior cingulate cortex, to urbanicity or green experiences. Of interest, the perigenual anterior cingulate cortex has been identified as a crucial brain structure for many self-related processes that comprise self-continuity, self-consciousness, self-reflection, self-regulation, and self-other similarity (Northoff, 2017; van der Meer et al., 2010; Scalabrini et al., 2019). The perigenual anterior cingulate cortex is located in the default mode network, which is characterized by high levels of activity when individuals are involved in free thought, such as mind-wandering (Northoff et al., 2010). Moreover, the perigenual anterior cingulate cortex is a key node within a network that integrates information from the memory and interoception and is affected by its interactions with the anterior insula, medial temporal regions, and posterior cingulate cortex, (Northoff and Panksepp, 2008). Regarding its involvement in psychosis, fMRI experiments applying explicit tasks of self-evaluation reported deficient intrinsic self-processing in the perigenual anterior cingulate cortex and adjacent ventromedial prefrontal cortices in patients with schizophrenia as compared to healthy controls (van der Meer et al., 2013; Tan et al., 2015; Kuhn and Gallinat, 2013; Modinos et al., 2011).

DISCUSSION

The evidence reviewed above begins to provide new insights into the mechanisms that may explain how green space exposure can reduce psychosis risk. On the one hand, epidemiological studies consistently show that green space exposure decreases the incidence of psychosis, whereas urbanicity is associated with an increased risk for psychosis. On the other hand, cognitive and neuroscience findings preliminarily suggest that exposure to and experiences of green space, as well as blue space, could have beneficial effects on self-related processing. Based on these findings, it is proposed that self-processing and related brain network interactions may constitute a relevant mechanism that mediates between green space exposure and psychosis risk (Figure 1).

Because this hypothesis has received very little or no attention in published empirical investigations using clinical samples and specific reports of positive and null results regarding the link between green space and psychosis-related self-disturbances are lacking, novel studies are encouraged to directly and systematically investigate how the environmental planning of green space could comprehensively facilitate intrinsic and extrinsic self-experiences in non-clinical samples, as well as in psychotic disorders or high-risk populations. The literature reviewed in this article provides concrete perspectives for further investigation. For example, psychological and neuroscientific studies with high levels of control over experimental conditions and modulating factors should

reproduce the epidemiological findings. Moreover, definite issues for future studies of the health benefits of green space include the modulating effects of particular types and features of green space and the kind of interactions individuals have with green space on intrinsic/extrinsic self-processing and psychosis risk.

Further limitations of the present review must be mentioned. Because it focuses specifically on the risk-reducing effects of green space for psychosis, it is important to remember that the onset of psychosis depends on a multiplicity of factors and events. The lack of green space exposure is one out of many characteristics of urbanicity that contribute to psychosis risk, in addition to other physical, social, familiar, occupational, and relationship variables (van Os et al., 2010; Brown, 2011; Schmitt et al., 2014). Green space is a piece of an intricate puzzle of interactions between numerous environmental variables, as well as genetic factors, that explains the variance in psychosis onset. It will be crucial for future studies to increase our understanding of how the effects of green space can be embedded in this full complexity of factors, both to explain and prevent psychosis onset (Tsuang, 2000; Howes et al., 2004; Meyer-Lindenberg and Tost, 2012). Finally, in this article, the effect of green space on psychosis prevention is examined in the context of self-disturbances, which are the core features of psychosis and sensitive predictors of its onset, while the relationships with positive symptoms, social impairments, and reduced insight into illness are not considered. It would be relevant to expand this investigating by including these domains too.

In conclusion, self-awareness is a continuous, integrative phenomenon that emerges from a stream of sensory perceptions, actions, bodily states, memories, motivations, thoughts, and imaginations. These processes support a coherent sense of self with a past (autobiographical memories), a present (actual experiences), and a future (prospective thoughts). Green space may be an environment with auspicious qualities that can modulate mental and bodily self-experiences over time and space on various scales, transforming itself continuously as it evolves and changes over time, while at the same time remaining a familiar environment (Northoff and Huang, 2017; Kuo, 2015). It is therefore hypothesized that green space potentially reduces psychosis risk by offering a dynamic environment that within a complex architecture of environmental and genetic factors, supports a stable, multidimensional self-experience over time. Enhancing our knowledge about the exact relationships between green space experiences and multidimensional self-processing would help in environmental planning, especially in urban areas, to optimize the positive impact of green space on mental health and illness prevention.

REFERENCES

- Boers, S., Hagoort, K., Scheepers, F., Helbich, M. (2018). Does residential green and blue space promote recovery in psychotic disorders? A cross-sectional study in the province of Utrecht, the Netherlands. *Int. J. Environ. Res. Public Health* 15(10), 2195. doi: 10.3390/ijerph15102195
- Bratman, G.N., Hamilton, J.P., Daily, G.C. (2012). The impacts of nature experience on human cognitive function and mental health. *Ann. N. Y. Acad. Sci.* 1249(1), 118-136. doi: 10.1111/j.1749-6632.2011.06400.x
- Bratman, G.N., Hamilton, J.P., Hahn, K.S., Daily, G.C., Gross, J.J. (2015). Nature experience reduces rumination and subgenual prefrontal cortex activation. *Proc. Natl. Acad. Sci. U.S.A.* 112(28), 8567-8572. doi: 10.1073/pnas.1510459112
- Brown, A.S. (2011). The environment and susceptibility to schizophrenia. *Progress in Neurobiology*, 93(1), 23-58. Doi: 10.1016/j.pneurobio.2010.09.003

- Browning, M.H., Shipley, N., McAnirlin, O., Becker, D., Yu, C.P., Hartig, T., & Dzhambov, A. M. (2020). An actual natural setting improves mood better than its virtual counterpart: A meta-analysis of experimental data. *Frontiers in Psychology* 11, 2200. Doi: 10.3389/fpsyg.2020.02200
- Berman, M.G., Jonides, J., Kaplan, S. (2008). The cognitive benefits of interacting with nature. *Psychol. Sci.* 19(12), 1207-1212. doi: 10.1111/j.1467-9280.2008.02225.x
- Blanke, O. (2012). Multisensory brain mechanisms of bodily self-consciousness. *Nat. Rev. Neurosci.* 13(8), 556-571. doi: 10.1038/nrn3292
- Cohen-Cline, H., Turkheimer, E., Duncan, G.E. (2015). Access to green space, physical activity and mental health: a twin study. *J. Epidemiol. Com. Health*, 69(6), 523-529. doi: 10.1136/jech-2014-204667
- Ćurčić-Blake, B., van der Meer, L., Pijnenborg, G.H., David, A.S., Aleman, A. (2015). Insight and psychosis: Functional and anatomical brain connectivity and self-reflection in schizophrenia. *Hum. Brain Mapp.* 36(12), 4859-4868. doi: 10.1002/hbm.22955
- Ćurčić-Blake, B., Ford, J.M., Hubl, D., Orlov, N.D., Sommer, I.E., Waters, F., Allen, P., Jardri, R., Woodruff, P.W., David, O. Mulert, C. (2017). Interaction of language, auditory and memory brain networks in auditory verbal hallucinations. *Prog. Neurobiol.* 148, 1-20. doi: 10.1016/j.pneurobio.2016.11.002
- Damasio, A.R. (1999). *The feeling of what happens: Body and emotion in the making of consciousness.* Houghton Mifflin Harcourt.
- Di Plinio, S., Perrucci, M.G., Aleman, A., Ebisch, S.J. (2020). I am Me: Brain systems integrate and segregate to establish a multidimensional sense of self. *NeuroImage* 205:116284. doi: 10.1016/j.neuroimage.2019.116284
- Di Plinio, S., Perrucci, M.G., Ebisch, S.J.H. (2020). The prospective sense of agency is rooted in local and global properties of intrinsic functional brain networks. Submitted.
- Dragt, S., Nieman, D.H., Veltman, D., Becker, H.E., van de Fliert, R., de Haan, L., Linszen, D.H. (2011). Environmental factors and social adjustment as predictors of a first psychosis in subjects at ultra high risk. *Schizophr. Res.* 125(1), 69-76. doi: 10.1016/j.schres.2010.09.007
- Ebisch, S.J., Mantini, D., Northoff, G., Salone, A., De Berardis, D., Ferri, F., Ferro, F.M., Di Giannantonio, M., Romani, G.L. and Gallese, V. (2014). Altered brain long-range functional interactions underlying the link between aberrant self-experience and self-other relationship in first-episode schizophrenia. *Schizophr. Bull.* 40(5), 1072-1082. doi: 10.1093/schbul/sbt153.
- Ebisch, S.J., and Gallese, V. (2015). A neuroscientific perspective on the nature of altered self-other relationships in schizophrenia. *J. Consc. Stud.* 22(1-2), 220-240.
- Ebisch, S.J., and Aleman, A. (2016). The fragmented self: imbalance between intrinsic and extrinsic self-networks in psychotic disorders. *Lancet Psychiatry* 3(8), 784-790. doi: 10.1016/S2215-0366(16)00045-6

- 399 Engemann, K., Pedersen, C.B., Arge, L., Tsirogiannis, C., Mortensen, P.B., Svenning, J.C. (2018).
400 Childhood exposure to green space – A novel risk-decreasing mechanism for
401 schizophrenia? *Schizophr. Res.* 199, 142-148. doi: 10.1016/j.schres.2018.03.026
402
- 403 Engemann, K., Pedersen, C.B., Arge, L., Tsirogiannis, C., Mortensen, P.B., Svenning, J.C. (2019a).
404 Residential green space in childhood is associated with lower risk of psychiatric disorders from
405 adolescence into adulthood. *Proc. Natl. Acad. Sci. U.S.A.* 116(11), 5188-5193. doi:
406 10.1073/pnas.1807504116
407
- 408 Engemann, K., Svenning, J.C., Arge, L., Brandt, J., Geels, C., Mortensen, P.B., Plana-Ripoll, O.,
409 Tsirogiannis, C. Pedersen, C.B. (2019b). Natural surroundings in childhood are associated with
410 lower schizophrenia rates. *Schizophr. Res.* in press. doi: 10.1016/j.schres.2019.10.012
411
- 412 Fazel, S., Wolf, A., Palm, C., Lichtenstein, P. (2014). Violent crime, suicide, and premature
413 mortality in patients with schizophrenia and related disorders: a 38-year total population study in
414 Sweden. *Lancet Psychiatry* 1(1), 44-54. doi: 10.1016/S2215-0366(14)70223-8
415
- 416 Frith, C.D., Blakemore, S.J., Wolpert, D.M. (2000). Explaining the symptoms of schizophrenia:
417 abnormalities in the awareness of action. *Brain Res. Rev.* 31(2-3), 357-363. doi: 10.1016/s0165-
418 0173(99)00052-1
419
- 420 Fuller, C., Powell, D., Fox, S. (2017). Making gains: the impact of outdoor residential experiences
421 on students' examination grades and self-efficacy. *Educational Review* 69, 232-247. Doi:
422 10.1080/00131911.2016.1199538
423
- 424 Fusar-Poli, P., Tantardini, M., De Simone, S., Ramella-Cravaro, V., Oliver, D., Kingdon, J.,
425 Kotlicka-Antczak, M., Valmaggia, L., Lee, J., Millan, M.J., Galderisi, S. (2017). Deconstructing
426 vulnerability for psychosis: Meta-analysis of environmental risk factors for psychosis in subjects at
427 ultra high-risk. *Eur. Psychiatry* 40, 65-75. doi: 10.1016/j.eurpsy.2016.09.003
428
- 429 Fusar-Poli, P., Davies, C., Solmi, M., Brondino, N., De Micheli, A., Kotlicka-Antczak, M., Shin,
430 J.I., Radua, J. (2019). Preventive Treatments for Psychosis: Umbrella Review (Just the
431 Evidence). *Front. Psychiatry* 10:764 doi: 10.3389/fpsy.2019.00764
432
- 433 Gallagher, S. (2000). Philosophical conceptions of the self: implications for cognitive
434 science. *Trends Cogn. Sci.* 4(1), 14-21. doi: 10.1016/S1364-6613(99)01417-5
435
- 436 Gallese, V., and Sinigaglia, C. (2010). The bodily self as power for action. *Neuropsychologia* 48(3),
437 746-755. doi: 10.1016/j.neuropsychologia.2009.09.038
438
- 439 Haggard, P. (2017). Sense of agency in the human brain. *Nat. Rev. Neurosci.* 18(4), 196-207. doi:
440 10.1038/nrn.2017.14
441
- 442 Hartig, T. (1993). Nature experience in transactional perspective. *Landsc. Urban Plan.* 25(1-2), 17-
443 36. doi: 10.1016/0169-2046(93)90120-3
444
- 445 Hartig, T., Mitchell, R., De Vries, S., Frumkin, H. (2014). Nature and health. *Annu. Rev. Public*
446 *Health* 35, 207-228. doi: 10.1146/annurev-publhealth-032013-182443
447
- 448 Hayhurst, J., Hunter, J.A., Kafka, S., Boyes, M. (2015). Enhancing resilience in youth through a 10-
449 day developmental voyage. *Journal of Adventure Education and Outdoor Learning* 15, 40–

52. doi: 10.1080/14729679.2013.843143

Howes, O.D., McDonald, C., Cannon, M., Arseneault, L., Boydell, J., Murray, R.M. (2004). Pathways to schizophrenia: the impact of environmental factors. *International Journal of Neuropsychopharmacology*, 7, S7-S13. Doi: 10.1017/S1461145704004122

Huang, S. C. L. (2009). The validity of visual surrogates for representing waterscapes. *Landscape Res.* 34(3), 323-335. doi: 10.1080/01426390902867984

Huang, S.C.L., Chiang, N.C., Kuo, N.F., & Chen, Y.J. (2019). An exploratory approach for using EEG to examine person-environment interaction. *Landscape Res.* 44(6), 702-715. doi: 10.1080/01426397.2018.1548586

Hunter, J.A., Kafka, S., Hayhurst, J., Clark, H., Dickerson, D., Harold, G., Boyes, M., O'Brien, K., Stringer, M. (2010). Increased self-efficacy following a ten-day developmental voyage. *Journal of Child & Adolescent Mental Health* 22, 63–65. doi: 10.2989/17280583.2010.496943

James, P., Banay, R.F., Hart, J.E., & Laden, F. (2015). A review of the health benefits of greenness. *Curr. Epidemi. Rep.* 2(2), 131-142. doi: 10.1007/s40471-015-0043-7

James, W. (1890). The perception of reality. *Principles of Psychology*, 2, 283-324.

Jeannerod, M. (2003). The mechanism of self-recognition in humans. *Behav. Brain Res.* 142(1-2), 1-15. doi: 10.1016/s0166-4328(02)00384-4

Kaplan, R., and Kaplan, S. (1989). The experience of nature: A psychological perspective. CUP Archive.

Kelly, B.D., O'Callaghan, E., Waddington, J.L., Feeney, L., Browne, S., Scully, P.J., Clarke, M., Quinn, J.F., McTigue, O., Morgan, M.G., Kinsella, A. (2010). Schizophrenia and the city: A review of literature and prospective study of psychosis and urbanicity in Ireland. *Schizophr. Res.*, 116(1), 75-89. doi: 10.1016/j.schres.2009.10.015

Kircher, T.T., and Leube, D.T. (2003). Self-consciousness, self-agency, and schizophrenia. *Consc. Cogn.* 12(4), 656-669. doi: 10.1016/s1053-8100(03)00071-0

Kuo, M. (2015). How might contact with nature promote human health? Promising mechanisms and a possible central pathway. *Front. Psychology*, 6, 1093. doi: 10.3389/fpsyg.2015.01093

Krabbendam, L., and Van Os, J. (2005). Schizophrenia and urbanicity: a major environmental influence—conditional on genetic risk. *Schizophr. Bull.* 31(4), 795-799. doi: 10.1093/schbul/sbi060

Kühn, S., and Gallinat, J. (2013). Resting-state brain activity in schizophrenia and major depression: a quantitative meta-analysis. *Schizophr. Bull.* 39(2), 358-365. doi: 10.1093/schbul/sbr151

Lederbogen, F., Kirsch, P., Haddad, L., Streit, F., Tost, H., Schuch, P., Wüst, S., Pruessner, J.C., Rietschel, M., Deuschle, M., Meyer-Lindenberg, A. (2011). City living and urban upbringing affect neural social stress processing in humans. *Nature* 474(7352), 498-501. doi: 10.1038/nature10190

- 501 Lewis, G., David, A., Andréasson, S., Allebeck, P. (1992). Schizophrenia and city
502 life. *Lancet* 340(8812), 137-140. doi: 10.1016/0140-6736(92)93213-7
503
- 504 Looijestijn, J., Blom, J. D., Aleman, A., Hoek, H. W., Goekoop, R. (2015). An integrated network
505 model of psychotic symptoms. *Neurosci. Biobeh. Rev.* 59, 238-250. doi:
506 10.1016/j.neubiorev.2015.09.016
507
- 508 Liemburg, E. J., van der Meer, L., Swart, M., Curcic-Blake, B., Bruggeman, R., Knegtering, H.,
509 Aleman, A. (2012). Reduced connectivity in the self-processing network of schizophrenia patients
510 with poor insight. *PLoS One* 7(8):e42707 doi: 10.1371/journal.pone.0042707
511
- 512 Maas, J., Verheij, R.A., de Vries, S., Spreeuwenberg, P., Schellevis, F.G., Groenewegen, P.P.
513 (2009). Morbidity is related to a green living environment. *J. Epidem. Com. Health* 63(12), 967-
514 973. doi: 10.1136/jech.2008.079038
515
- 516 Margalit, D., and Ben-Ari, A. (2014). The effect of wilderness therapy on adolescents' cognitive
517 autonomy and self-efficacy: Results of a non-randomized trial. In *Child & Youth Care Forum* (Vol.
518 43, No. 2, pp. 181-194). Springer US.
519
- 520 Markevych, I., Schoierer, J., Hartig, T., Chudnovsky, A., Hystad, P., Dzhambov, A.M., De Vries,
521 S., Triguero-Mas, M., Brauer, M., Nieuwenhuijsen, M.J. Lupp, G. (2017). Exploring pathways
522 linking greenspace to health: Theoretical and methodological guidance. *Environ. Res.* 158, 301-
523 317. doi: 10.1016/j.envres.2017.06.028
524
- 525 McGrath, J., Saha, S., Welham, J., El Saadi, O., MacCauley, C. Chant, D. (2004). A systematic
526 review of the incidence of schizophrenia: the distribution of rates and the influence of sex,
527 urbanicity, migrant status and methodology. *B.M.C. Med.* 2, 13. doi: 10.1186/1741-7015-2-13
528
- 529 McGorry, P.D., Nelson, B., Amminger, G.P., Bechdolf, A., Francey, S.M., Berger, G., Riecher-
530 Rössler, A., Klosterkötter, J., Ruhrmann, S., Schultze-Lutter, F. Nordentoft, M. (2009). Intervention
531 in individuals at ultra-high risk for psychosis: a review and future directions. *J. Clin. Psychiatry*
532 70(9):1206-1212. doi: 10.4088/JCP.08r04472
533
- 534 Menon, V., and Uddin, L. Q. (2010). Saliency, switching, attention and control: a network model of
535 insula function. *Brain Struct. Funct.* 214(5-6), 655-667. doi: 10.1007/s00429-010-0262-0
536
- 537 Meyer-Lindenberg, A. and Tost, H. (2012). Neural mechanisms of social risk for psychiatric
538 disorders. *Nat. Neurosci.*, 15(5), 663. doi: 10.1038/nn.3083
539
- 540 Minkowski, E. (1927). *La schizophrénie. Psychopathologie des schizoïdes et des schizophrènes.*
541 Payot.
542
- 543 Modinos, G., Renken, R., Ormel, J., Aleman, A. (2011). Self-reflection and the psychosis-prone
544 brain: an fMRI study. *Neuropsychology* 25(3), 295. doi: 10.1037/a0021747
545
- 546 Molnar-Szakacs, I., and Uddin, L. Q. (2013). Self-processing and the default mode network:
547 interactions with the mirror neuron system. *Front. Hum. Neurosci.* 7:571. doi:
548 10.3389/fnhum.2013.00571.
549
- 550 Mygind, L., Kjeldsted, E., Hartmeyer, R., Mygind, E., Bølling, M., & Bentsen, P. (2019). Mental,
551 physical and social health benefits of immersive nature-experience for children and adolescents: A

- 552 systematic review and quality assessment of the evidence. *Health & Place* 58, 102136. Doi:
553 10.1016/j.healthplace.2019.05.014
554
- 555 Nelson, B., Thompson, A., Yung, A.R. (2012). Basic self-disturbance predicts psychosis onset in
556 the ultra high risk for psychosis “prodromal” population. *Schizophr. Bull.* 38(6), 1277-1287. doi:
557 10.1093/schbul/sbs007
558
- 559 Nelson, B., Fornito, A., Harrison, B.J., Yücel, M., Sass, L.A., Yung, A.R., Thompson, A., Wood,
560 S.J., Pantelis, C. and McGorry, P.D. (2009). A disturbed sense of self in the psychosis prodrome:
561 linking phenomenology and neurobiology. *Neurosci. Biobehav. Rev.* 33(6), 807-817. doi:
562 10.1016/j.neubiorev.2009.01.002
563
- 564 Nelson, B., Parnas, J., Sass, L.A. (2014). Disturbance of minimal self (ipseity) in schizophrenia:
565 clarification and current status. *Schizophr. Bull.* 40(3), 479–482. doi: 10.1093/schbul/sbu034
566
- 567 Newbury, J., Arseneault, L., Caspi, A., Moffitt, T.E., Odgers, C.L., Fisher, H.L. (2016). Why are
568 children in urban neighborhoods at increased risk for psychotic symptoms? Findings from a UK
569 longitudinal cohort study. *Schizophr. Bull.* 42(6), 1372-1383. doi: 10.1093/schbul/sbw052
570
- 571 Northoff, G., and Bermpohl, F. (2004). Cortical midline structures and the self. *Trends Cogn.*
572 *Sci.* 8(3), 102-107. doi: 10.1016/j.tics.2004.01.004
573
- 574 Northoff, G., and Panksepp, J. (2008). The trans-species concept of self and the subcortical–cortical
575 midline system. *Trends Cogn. Sci.* 12(7), 259-264. doi: 10.1016/j.tics.2008.04.007
576
- 577 Northoff, G., Qin, P., Nakao, T. (2010). Rest-stimulus interaction in the brain: a review. *Trends*
578 *Neurosci.* 33(6), 277-284. doi: 10.1016/j.tins.2010.02.006
579
- 580• Northoff, G. (2017). Personal identity and cortical midline structure (CMS): do temporal features of
581 CMS neural activity transform into “Self-Continuity”? *Psychological Inquiry*, 28(2-3), 122-131.
582 doi: 10.1080/1047840X.2017.1337396
583
- 584 Northoff, G., and Huang, Z. (2017). How do the brain’s time and space mediate consciousness and
585 its different dimensions? Temporo-spatial theory of consciousness (TTC). *Neurosci. Biobehav.*
586 *Rev.* 80, 630-645. doi: 10.1016/j.neubiorev.2017.07.013
587
- 588 O’Brien, K., and Lomas, T. (2017). Developing a Growth Mindset through outdoor personal
589 development: can an intervention underpinned by psychology increase the impact of an outdoor
590 learning course for young people? *Journal of Adventure Education and Outdoor Learning* 17(2),
591 133-147. Doi: 10.1080/14729679.2016.1232199
592
- 593 Parnas, J. and Henriksen, M.G. (2014). Disordered self in the schizophrenia spectrum: a clinical and
594 research perspective. *Harvard Review of Psychiatry*, 22(5), 251. doi:
595 10.1097/HRP.0000000000000040
596
- 597 Pedersen, C.B., and Mortensen, P.B. (2001). Evidence of a dose-response relationship between
598 urbanicity during upbringing and schizophrenia risk. *Arch. Gen. Psychiatry* 58(11), 1039-1046.
599 doi: 10.1001/archpsyc.58.11.1039
600
- 601 Pries, L.K., Guloksuz, S., Ten Have, M., De Graaf, R., Van Dorsselaer, S., Gunther, N.,
602 Rauschenberg, C., Reininghaus, U., Radhakrishnan, R., Bak, M., Rutten, B.P. (2018). Evidence that

- environmental and familial risks for psychosis additively impact a multidimensional subthreshold psychosis syndrome. *Schizophr. Bull.* 44(4), 710-719. doi: 10.1093/schbul/sby051
- Qin, P., and Northoff, G. (2011). How is our self related to midline regions and the default-mode network? *Neuroimage*, 57(3), 1221-1233. doi: 10.1016/j.neuroimage.2011.05.028
- Raballo, A., Sæbye, D., Parnas, J. (2011). Looking at the Schizophrenia Spectrum Through the Prism of Self-disorders: An Empirical Study, *Schizophrenia Bulletin*, 37(2), 344–351. doi: 10.1093/schbul/sbp056
- Raichle, M.E., MacLeod, A.M., Snyder, A.Z., Powers, W.J., Gusnard, D.A., Shulman, G.L. (2001). A default mode of brain function. *Proc. Natl. Acad. Sci. U.S.A.* 98(2), 676-682. doi: 10.1073/pnas.98.2.676
- Rössler, W., Salize, H.J., van Os, J., Riecher-Rössler, A. (2005). Size of burden of schizophrenia and psychotic disorders. *Eur. Neuropsychopharmacol.* 15(4), 399-409. doi: 10.1016/j.euroneuro.2005.04.009
- Ruhrmann, S., Schultze-Lutter, F., Salokangas, R.K., Heinimaa, M., Linszen, D., Dingemans, P., Birchwood, M., Patterson, P., Juckel, G., Heinz, A. and Morrison, A. (2010). Prediction of psychosis in adolescents and young adults at high risk: results from the prospective European prediction of psychosis study. *Arch. Gen. Psychiatry* 67(3), 241-251. doi: 10.1001/archgenpsychiatry.2009.206
- Sarkar, C., Webster, C., Gallacher, J. (2018). Residential greenness and prevalence of major depressive disorders: a cross-sectional, observational, associational study of 94 879 adult UK Biobank participants. *Lancet Planet. Health* 2(4):e162-e173. doi: 10.1016/S2542-5196(18)30051-2
- Sass, L. A., and Parnas, J. (2003). Schizophrenia, consciousness, and the self. *Schizophr. Bull.* 29(3), 427-444. doi: 10.1093/oxfordjournals.schbul.a007017
- Scalabrini, A., Ebisch, S.J., Huang, Z., Di Plinio, S., Perrucci, M.G., Romani, G.L., Mucci, C. Northoff, G. (2019). Spontaneous Brain Activity Predicts Task-Evoked Activity During Animate Versus Inanimate Touch. *Cereb. Cortex* 29(11), 4628-4645. doi: 10.1093/cercor/bhy340
- Schmitt, A., Malchow, B., Hasan, A. Fallkai, P. (2014). The impact of environmental factors in severe psychiatric disorders. *Frontiers Neurosci.*, 8, 19. doi: 10.3389/fnins.2014.00019
- Schultze-Lutter, F. (2009). Subjective symptoms of schizophrenia in research and the clinic: the basic symptom concept. *Schizophr. Bull.* 35, 5-8. doi: 10.1093/schbul/sbn139
- Shad, M.U., and Keshavan, M.S. (2015). Neurobiology of insight deficits in schizophrenia: an fMRI study. *Schizophr. Res.*, 165(2-3), 220-226. doi: 10.1016/j.schres.2015.04.021
- Smigielski, L., Jagannath, V., Rössler, W., Walitza, S., Grünblatt, E. (2020). Epigenetic mechanisms in schizophrenia and other psychotic disorders: a systematic review of empirical human findings. *Mol. Psychiatry*, 1-31. doi: 10.1038/s41380-019-0601-3
- Smith, G., Cirach, M., Swart, W., Dèdelè, A., Gidlow, C., van Kempen, E., Kruize, H., Gražulevičienė, R., Nieuwenhuijsen, M.J. (2017). Characterisation of the natural environment:

- quantitative indicators across Europe. *Int. J. Health Geograph.* 16(1), 16. doi: 10.1186/s12942-017-0090-z
- Spreng, R.N., Mar, R.A., Kim, A.S. (2009). The common neural basis of autobiographical memory, prospection, navigation, theory of mind, and the default mode: a quantitative meta-analysis. *J. Cogn. Neurosci.* 21(3), 489-510. doi: 10.1162/jocn.2008.21029
- Sundquist, K., Frank, G., Sundquist, J.A.N. (2004). Urbanisation and incidence of psychosis and depression: follow-up study of 4.4 million women and men in Sweden. *Br. J. Psychiatry* 184(4), 293-298. doi: 10.1192/bjp.184.4.293
- Sussman, A., and Hollander, J.B. (2014). *Cognitive architecture: Designing for how we respond to the built environment.* Routledge.
- Tan, S., Zhao, Y., Fan, F., Zou, Y., Jin, Z., Zen, Y., Zhu, X., Yang, F., Tan, Y. and Zhou, D. (2015). Brain correlates of self-evaluation deficits in schizophrenia: a combined functional and structural MRI study. *PloS One* 10(9):e0138737. doi: 10.1371/journal.pone.0138737
- Torrey, E.F., Yolken, R.H. (2019). Schizophrenia as a pseudogenetic disease: A call for more gene-environmental studies. *Psychiatry Res.* 278, 146-150. doi: 10.1016/j.psychres.2019.06.006
- Tost, H., Reichert, M., Braun, U., Reinhard, I., Peters, R., Lautenbach, S., Hoell, A., Schwarz, E., Ebner-Priemer, U., Zipf, A. and Meyer-Lindenberg, A. (2019). Neural correlates of individual differences in affective benefit of real-life urban green space exposure. *Nat. Neurosci.* 22(9), 1389-1393. doi: 10.1038/s41593-019-0451-y
- Tsuang, M. (2000). Schizophrenia: genes and environment. *Biological Psychiatry*, 47(3), 210-220. Doi: 10.1016/S0006-3223(99)00289-9
- Twedt, E., Rainey, R.M., & Proffitt, D.R. (2016). Designed natural spaces: informal gardens are perceived to be more restorative than formal gardens. *Frontiers in psychology* 7, 88. Doi: 10.3389/fpsyg.2016.00088
- Twohig-Bennett, C. and Jones, A. (2018). The health benefits of the great outdoors: A systematic review and meta-analysis of greenspace exposure and health outcomes. *Environ. Res.* 166, 628-637. doi: 10.1016/j.envres.2018.06.030
- Ulrich, R.S., Simons, R.F., Losito, B.D., Fiorito, E., Miles, M.A., Zelson, M. (1991). Stress recovery during exposure to natural and urban environments. *J. Environ. Psychol.* 11(3), 201-230. doi: 10.3390/ijerph7031036
- United Nations, Department of Economic and Social Affairs, Population Division (2014) *World urbanization prospects: The 2014 revision, highlights (ST/ESA/SER.A/352).* Available at <https://esa.un.org/unpd/wup/Publications/Files/WUP2014-Highlights.pdf>.
- Van den Berg, M., Wendel-Vos, W., van Poppel, M., Kemper, H., van Mechelen, W., Maas, J. (2015). Health benefits of green spaces in the living environment: A systematic review of epidemiological studies. *Urban For. Urban Green.* 14(4), 806-816. doi: 10.1016/j.ufug.2015.07.008

- van der Meer, L., Costafreda, S., Aleman, A., David, A.S. (2010). Self-reflection and the brain: a theoretical review and meta-analysis of neuroimaging studies with implications for schizophrenia. *Neurosci. Biobehav. Rev.* 34(6), 935-946. doi: 10.1016/j.neubiorev.2009.12.004
- van der Meer, L., de Vos, A.E., Stiekema, A.P., Pijnenborg, G.H., van Tol, M.J., Nolen, W.A., David, A.S., Aleman, A. (2013). Insight in schizophrenia: involvement of self-reflection networks? *Schizophr. Bull.* 39(6), 1288. doi: 10.1093/schbul/sbs122
- Vanhaudenhuyse, A., Demertzi, A., Schabus, M., Noirhomme, Q., Bredart, S., Boly, M., Phillips, C., Soddu, A., Luxen, A., Moonen, G., Laureys, S. (2011). Two distinct neuronal networks mediate the awareness of environment and of self. *J. Cogn. Neurosci.* 23(3), 570-578. doi: 10.1162/jocn.2010.21488
- van Kemenade, B. M., Arikan, B. E., Podranski, K., Steinsträter, O., Kircher, T., Straube, B. (2019). Distinct roles for the cerebellum, angular gyrus, and middle temporal gyrus in action–feedback monitoring. *Cereb. Cortex* 29(4), 1520-1531. doi: 10.1093/cercor/bhy048
- van Os, J., Pedersen, C.B., Mortensen, P.B. (2004). Confirmation of synergy between urbanicity and familial liability in the causation of psychosis. *Am. J. Psychiatry* 161(12), 2312-2314. doi: 10.1176/appi.ajp.161.12.2312
- van Os, J., Kenis, G., Rutten, B. (2010). The environment and schizophrenia. *Nature* 468, 203–212 . doi: 10.1038/nature09563
- van Os, J., and Linscott, R.J. (2012). Introduction: the extended psychosis phenotype—relationship with schizophrenia and with ultrahigh risk status for psychosis. *Schizophr. Bull.* 38(2), 227-230. doi: 10.1093/schbul/sbr188
- van Os, J., Reininghaus, U., Meyer-Lindenberg, A. (2017). The search for environmental mechanisms underlying the expression of psychosis: introduction. *Schizophr. Bull.* 43(2):283-286. doi: 10.1093/schbul/sbw178
- Verheij, R.A., Maas, J., Groenewegen, P.P. (2008). Urban—rural health differences and the availability of green space. *European Urban and Regional Studies*, 15(4), pp.307-316. Doi: 10.1177/0969776408095107
- Völker, S., and Kistemann, T. (2011). The impact of blue space on human health and well-being—Salutogenetic health effects of inland surface waters: A review. *Int. J. Hyg. Environ. Health* 214(6), 449-460. doi: 10.1016/j.ijheh.2011.05.001
- Waters, F., Woodward, T., Allen, P., Aleman, A., Sommer, I. (2012). Self-recognition deficits in schizophrenia patients with auditory hallucinations: a meta-analysis of the literature. *Schizophr. Bull.* 38(4), 741-750. doi: 10.1093/schbul/sbq144
- White, M.P., Alcock, I., Wheeler, B.W., Depledge, M.H. (2013). Would you be happier living in a greener urban area? A fixed-effects analysis of panel data. *Psychol. Sci.* 24(6), 920-928. doi: 10.1177/0956797612464659

750

751

752

753

754

755

756

757

758

759

760

Figure 1. A hypothetical model illustrating the modulation of the integrity of self-related processing, which is impaired in psychosis, through brain network interactions by green space. The experience of natural surroundings (green space area) in one's living environment (A; increased interactions and overlap between green space and self-aspects) or being distantiated from natural surroundings during daily life (B; decreased interactions and overlap between green space and self-aspects) is associated with more or less integration of intrinsic (red; internally directed) and extrinsic (blue; externally directed) self-aspects, respectively. The green space photo depicts a view on Val d'Orcia in Tuscany, Italy, an UNESCO World Heritage landscape developed in the 14th and 15th centuries to idealize a well-balanced dialogue between humans and nature.