Does Centrality in a Cross-Sectional Network Suggest Intervention Targets for Social Anxiety Disorder?

Thomas L. Rodebaugh
Natasha A. Tonge
Marilyn L. Piccirillo
Washington University in St Louis

Eiko Fried
University of Amsterdam

Arielle Horenstein
Temple University

Amanda S. Morrison
Stanford University

Philippe Goldin
University of California Davis

James J. Gross
Stanford University

Michelle H. Lim
Katya Fernandez
Washington University in St Louis

Carlos Blanco
Franklin R. Schneier
Columbia University

Ryan Bogdan
Renee J. Thompson
Washington University in St Louis

Richard G. Heimberg
Temple University

Thomas L. Rodebaugh, Natasha A. Tonge, Marilyn L. Piccirillo, Michelle H. Lim, Katya C. Fernandez, Ryan Bogdan, Renee J. Thompson, Department of Psychological and Brain Sciences, Washington University in St. Louis. Eiko Fried, Psychological Methods Group, University of Amsterdam. Arielle Horenstein and Richard G. Heimberg, Department of

Psychology, Temple University. Amanda A. Morrison, James J Gross, Department of Psychology, Stanford University. Philippe Goldin, Betty Irene Moore School of Nursing, University of California Davis. Carlos Blanco and Franklin R. Schneier, Anxiety Disorders Clinic, New York State Psychiatric Institute.

Michelle Lim is now at the Brain and Psychological Sciences Center, Swinburne University of Technology. Katya Fernandez is now at the Department of Psychiatry and Behavioral Sciences, Stanford University. Carlos Blanco is now Director of the Division of Epidemiology, Services, and Prevention Research at the National Institute on Drug Abuse.

This research was supported in part by NIMH grant R21 MH090308 and McDonnell Center for Systems Neuroscience New Resource Grant to Thomas L. Rodebaugh; NIH grant UL1 RR024992 to Washington University St. Louis; NIMH grant R01 MH064481-01A1 and GlaxoSmithKline Pharmaceuticals grant 101618 to Richard G. Heimberg; NIH grant K02 DA023200 to Carlos Blanco; ERC Consolidator Grant no. 647209 to Eiko Fried; NIMH R01 MH092416, NCCAM Grant AT003644, and NIMH Grant R01 MH076074 to James Gross; and 1F31MH115641-01 to Marilyn L. Piccirillo. Thanks to Amit Bernstein for some helpful thoughts about this project.

Correspondence concerning this article should be addressed to Thomas L. Rodebaugh, CB 1125, Psychology Building, Washington University in St. Louis, St. Louis, MO 63130.

©American Psychological Association, 2018. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article will be available, upon publication, via its DOI: <http://dx.doi.org/10.1037/ccp0000336>

Abstract

Objective: Network analysis allows us to identify the most interconnected (i.e., *central*) symptoms, and multiple authors have suggested that these symptoms might be important treatment targets. This is because change in central symptoms (relative to others) should have greater impact on change in all other symptoms. It has been argued that networks derived from cross-sectional data may help identify such important symptoms. We tested this hypothesis in social anxiety disorder.

Method: We first estimated a state-of-the-art regularized partial correlation network based on participants with social anxiety disorder ($N = 910$) to determine which symptoms were more central. Next, we tested whether change in these central symptoms were indeed more related to overall symptom change in a separate dataset of participants with social anxiety disorder who underwent a variety of treatments ($N = 244$). We also tested whether relatively superficial item properties (infrequency of endorsement and variance of items) might account for any effects shown for central symptoms.

Results: Centrality indices successfully predicted how strongly changes in items correlated with change in the remainder of the items. Findings were limited to the measure used in the network and did not generalize to three other measures related to social anxiety severity. In contrast, infrequency of endorsement showed associations across all measures.

Conclusions: The transfer of recently published results from cross-sectional network analyses to treatment data is unlikely to be straightforward.

Key Words: network analysis; social anxiety disorder; research methods

Public Health Significance: Researchers have recently asserted that network analyses might uncover the most important symptoms to target in treatment, even when the data used were

collected at a single time point. We examined this issue in generalized social anxiety disorder, and found modest support for the notion. However, simply counting how many participants endorsed the symptom as clearly present was a superior method for identifying important symptoms.

Does Centrality in a Cross-Sectional Network Suggest Intervention Targets for Social Anxiety Disorder?

Many studies of psychopathology seem to assume what might be called a common cause perspective. This approach involves thinking of clinical symptoms largely as passive measurements of an underlying mental disorder. Thus, a person has anxiety and avoidance about a variety of social situations as a consequence of having social anxiety disorder (SAD). From this viewpoint, the important causes and consequences are those related to the underlying latent variable of SAD itself, rather than to specific symptoms of SAD. Multiple authors have recently proposed that network perspectives offer an important alternative to a common cause perspective (Borsboom & Cramer, 2013; Borsboom et al., 2016; Cramer, Waldorp, van der Maas, & Borsboom, 2010a, 2010b; Fried, Borkulo, et al., 2016; McNally et al., 2015).

In a network conception, symptoms are understood as potentially causal agents in their own right (Borsboom, 2008; Borsboom, 2017). Instead of SAD being an entity to study, it could simply be a label for a set of symptoms (or other factors) that cause each other over time. A large number of authors have suggested that both the network conception and its related analyses could help researchers uncover central, important, or key symptoms that may provide viable treatment targets (Borsboom & Cramer, 2013; Borsboom, Cramer, Schmittmann, Epskamp, & Waldorp, 2011; Bringmann et al., 2016; Bringmann et al., 2013; Cramer et al., 2010a; Fried, Epskamp, Nesse, Tuerlinckx, & Borsboom, 2016; McNally et al., 2015; Robinaugh, LeBlanc, Vuletich, & McNally, 2014; Ruzzano, Borsboom, & Geurts, 2015; van de Leemput et al., 2014; Wichers, 2014; Wichers, Groot, & Psychosystems, 2016).

One can distinguish a network theory of psychopathology from network psychometrics—the statistical techniques used to estimate network models. These network analyses, like any

other statistical technique, can be applied in a variety of ways to a variety of types of data. Many network analyses presented in the literature have focused on cross-sectional data (with some notable exceptions; see, e.g., Bringmann et al., 2013). Multiple authors have suggested that treatment of the symptoms identified as the most important in cross-sectional network analyses may result in the greatest overall treatment gains (McNally et al., 2015; Ruzzano et al., 2015). The implication is that cross-sectional network analyses might identify important treatment targets.

Figure 1 provides an example of a quantitative indicator of network importance, the centrality index *strength*. In this network, nodes (e.g., symptoms or items) are represented by circles, and the strength of the relationships between nodes is depicted by the thickness of the lines between the nodes that are called edges. In the type of graphs typically presented (e.g., in the literature cited above), the nodes are positioned based on the strength of their relationships with other nodes. In Figure 1, the nodes have varied properties, with Node 1 having the greatest strength in its edges with other nodes, where strength is defined as the sum of all absolute edge weights connected to a node. Strength is one of many centrality indices; we also investigate closeness and betweenness in this paper. As is hopefully clear from Panel B of Figure 1, centrality is not always easily determined by visually inspecting a graph: Although Node 1 has the greatest strength, it is not literally at the center of the figure. This is because centrality indices are inferences of high-dimensional network structures that cannot always be mapped in two dimensions in an ideal way and therefore may not correspond obviously to visual cues (e.g., such as how close to the center a node is).

The idea that centrality indices should identify symptoms that are important for treatment rests upon the inference that centrality indices, by identifying symptoms with strong quantitative

relationships with others, also identify those symptoms with a strong causal role during treatment. That is, consider Node 1 in Panel B of Figure 1. It has a strength index of .85, whereas all other nodes in the figure have lower strength values (Node 2 has the second highest value, at .54). If the edges shown involve causal relationships directed from Node 1 to other nodes, then a change in Node 1, compared the same amount of change in the other nodes, would be expected to produce the strongest changes in the other nodes (all other factors being equal).

The intuitive appeal of the idea that high centrality involves high causal impact is clear, but changing Node 1 may do nothing if the high strength of its associated edges are entirely produced by other nodes causing Node 1. Similarly, an edge between Nodes 1 and 2 can result from failing to include important variables in the network that covary with both, in which case changing Node 1 may not have an impact on Node 2. More generally, the question of whether and how cross-sectional relationships in complex models are related to causality over time is a contentious one. Among the authors of the current manuscript, for example, there are a wide variety of viewpoints on this issue. Some of us view models based on cross-sectional data as a first step, useful for initial testing of theories that may or may not generalize to other types of data. All of us agree at least on the idea that a cross-sectional relationship between two variables implies some shared causal path that involves those two variables (even if the shared path is that a third variable causes both). At the same time, some of us find it implausible that there will be any systematic correspondence between cross-sectional data and either experimental or longitudinal data, and point to such findings of those of Maxwell and Cole to support this pessimism (Cole & Maxwell, 2003; Gollob & Reichardt, 1987; Maxwell & Cole, 2007; Maxwell, Cole, & Mitchell, 2011).

For central symptoms to be reliably important for treatment, centrality must have a high tendency to signal cause and effect in some form: A symptom is key to treatment if changing that symptom causes an important effect on other symptoms. This situation raises an empirical question: What relation is there, if any, between central symptoms identified in a cross-sectional network and change in other symptoms across treatment? The current paper aims to answer this question in the context of SAD treatment.

Looking at the prior literature on the topic, there are two especially relevant papers that came to opposite conclusions. One study, focusing on depression symptoms, examined whether centrality (measured by strength) estimated from a cross-sectional network predicted strength in individual person-level networks across time (de Vos et al., 2017). These authors found no evidence that cross-sectional centrality indices clearly signal how much nodes predict other nodes over time. However, the sample was relatively small for a network analysis ($N = 104$), and no information was provided regarding the stability of the centrality estimates.

In a second study, Robinaugh and colleagues (Robinaugh, Millner, & McNally, 2016) demonstrated that, in a group of older adults observed naturalistically, symptoms identified as more central to the network at a single time point appeared more clearly connected to change in other symptoms over time. That is, Robinaugh and colleagues computed how much change in an item correlated with change in the remaining items, and then examined whether that item's centrality was associated with that correlation. Returning to the example in Figure 1, when Node 1 in Panel B changes, how much it changes should be highly correlated with how much the other nodes change (due to its high strength in the network), if indeed the edges radiating from Node 1 are related to causal pathways.

The results from this naturalistic study could be extended to the context of treatment. If symptoms identified as more central in cross-sectional data are indeed more important in predicting change in other symptoms across treatment as well, then change in these symptoms should be strongly associated with change in the entire network. For example, take SAD as a set of clinical symptoms involving fear and avoidance of social situations. If fear of one type of social situation, such as talking with authority figures, were found to be more central in a cross-sectional network, change in fear of talking with authority figures across treatment should relate strongly to changes in fear and avoidance of other situations. That is, if centrality in a cross-sectional network corresponds to importance for treatment, then change in the more central symptoms—whether they were targeted or not—should be a particularly good predictor of change in the rest of the network. Further, changes in these more central symptoms might also show stronger relationships with changes in other symptom measures, demonstrating their potential causal importance not only within the network, but also outside of the modeled network, in a similar realm of symptoms (and thus potentially within the same conceptual network). To the best of our knowledge, such a test has not yet been published in the literature.

We thus investigated whether symptoms identified as more central to a cross-sectional network of social anxiety symptoms (from the Liebowitz Social Anxiety Scale: LSAS; Liebowitz, 1987) showed evidence of being important for change during treatment. We used one sample to obtain centrality indices and a second sample to examine change across treatment, approximating a clinician's application of results from the literature; however, we also examined whether estimating the network based on the pre-treatment data made any difference. We examined change both within the LSAS network and outside of the modeled network (i.e., by focusing on other measures of social anxiety severity). We hypothesized that change in items

with higher centrality (versus those with lower centrality) would prove to be stronger predictors of change both within the LSAS network and in additional measures of social anxiety severity. We also expected the strongest relationships to be found within the LSAS network (due, e.g., to the common finding of stronger correlations within a measure than across measures). Finally, we tested whether item properties (i.e., infrequency of endorsement, item variance) with no obvious causal properties might account for any findings related to centrality. These latter tests are important because centrality indices can be affected by item properties such as rates of endorsement (Terluin, de Boer, & de Vet, 2016). Because we were concerned about the ability to predict *decreases* in symptoms across treatment, we were most concerned with restriction of range (i.e., variance) and floor effects (i.e., infrequency of endorsement). That is, an item, even if it measures an important causal factor, will have difficulty associating with reductions across treatment if that item lacks sufficient range or lacks sufficient numbers of participants who endorse it prior to treatment. We will refer to variance and infrequency of endorsement collectively as *relatively superficial item properties*; we mean by this phrase not that they are unimportant, but that these are properties that are relatively easy to manipulate (e.g., by changing the response scale) without changing the property we believe is being measured.

Method

Participants

Participants with generalized social anxiety disorder were pooled from several archival data. All participants provided informed consent for their data to be collected as part of a research project approved by the appropriate Institutional Review Board. Data sets were examined as two samples. The first sample (Sample A) was used to estimate the cross-sectional network, whereas in the second (Sample B), we examined change across treatment. The two

samples did not differ in regard to age or gender ($ps > .138$), but did differ by ethnicity in that Asian Americans were more well-represented in the treatment sample, $\chi^2(1, n = 605) = 126.08, p < .001$. Our intent in using the two samples was to simulate what would happen if an existing cross-sectional network analysis were used as a guide in a new treatment sample; as such, some differences between the samples are expected. However, we also examined in follow-up tests whether conclusions were different if centrality indices were drawn from the treatment sample instead.

Network analysis sample (Sample A). A total of 910 participants diagnosed with DSM-IV (American Psychiatric Association, 1994) generalized social anxiety disorder (GSAD) via structured clinical interview were included in Sample A. These data were drawn from nine separate datasets that had been collected as part of several studies conducted at metropolitan and urban research centers. Overall characteristics of the sample, including demographics, are displayed in Table 1 (full details of each subsample are available in the supplemental material). All participants completed the clinician-administered version of the LSAS. In all cases, only pre-treatment data were used, although most participants were in studies including treatment.

Treatment sample (Sample B). An additional, non-overlapping sample of 244 participants was included in analyses focused on treatment; a total of 155 participants provided at least some data at post-treatment. Participants were recruited for three treatment studies that included cognitive behavioral therapy, mindfulness-based stress reduction, aerobic exercise, and wait list conditions (see the supplemental material for full information). Participants were diagnosed with GSAD or SAD via a structured clinical interview, and the data were maintained in three separate datasets that were collected from a large, West-coast university located outside a metropolitan area. All participants completed a self-report version of the LSAS. Participant

characteristics are provided in Table 2 (additional details are provided in the supplementary material).

Measures

Liebowitz Social Anxiety Scale: Clinician-Administered Version (LSAS-CA; Liebowitz, 1987). The LSAS-CA is a 48-item clinician-administered measure that assesses social fear and avoidance across 24 separate social performance and interaction situations. Clinicians instruct individuals to report their level of fear and avoidance of the given situation during the past week using a 4-point Likert-type scale. The fear scale ranges from 0 (None) to 3 (Severe) and the avoidance scale ranges from 0 (Never) to 3 (Usually). The LSAS-CA has demonstrated excellent internal consistency, as well as strong convergent validity with other clinician-administered and self-report measures of social anxiety and divergent validity with measures of depression (Heimberg et al., 1999; Heimberg, Mueller, Holt, Hope, & Liebowitz, 1992). The LSAS-CA was used in Datasets 1-9 (i.e., Sample A). The internal consistency for the items composing the total score was excellent ($\alpha = .98$).

Liebowitz Social Anxiety Scale: Self-Report Version (LSAS-SR; Fresco et al., 2001). The self-report version of the LSAS (LSAS-SR) uses the same situations and scales as the LSAS-CA. Instructions for the LSAS-SR are adapted from the LSAS-CA and are provided at the top of the measure for the participant to review as necessary. The LSAS-SR was used in Sample B. For one of the three treatment studies, as well as at follow-up time-points for the two other treatment studies, the LSAS-SR was delivered online. Previous studies have demonstrated that the LSAS-SR and its subscales have good internal consistency and the total score from the LSAS-SR is strongly correlated with total score from the LSAS-CA, $r = .85, p < .05$ (Baker, Heinrichs, Kim, & Hofmann, 2002; Fresco et al., 2001; Oakman, Van Ameringen, Mancini, &

Farvolden, 2003). The internal consistencies for items composing the total score pre- and post-treatment were excellent ($\alpha = .91, .95$, respectively).

Social Interaction Anxiety Scale – Straightforward items (SIAS-S; Rodebaugh, Woods, & Heimberg, 2007). The SIAS-S is modified from the original SIAS that was developed by Mattick and Clarke (1998) and includes the 17 straightforward items from the original 20-item scale. The items assess social anxiety in various social interaction situations. The SIAS-S uses a 5-point Likert-type scale ranging from 0 (Not at all) to 4 (Extremely) to assess level of social anxiety in a given situation. The SIAS-S has demonstrated a unifactorial structure with high internal consistency (Rodebaugh et al., 2007; Rodebaugh, Woods, Heimberg, Liebowitz, & Schneier, 2006). Furthermore, it has displayed strong construct validity in both undergraduate and clinical samples, as well as strong convergent validity with other measures of social anxiety and divergent validity with other psychological or personality constructs (Rodebaugh et al., 2007; Rodebaugh et al., 2011).

Brief Fear of Negative Evaluation Scale – Straightforward items (BFNE-S; Rodebaugh et al., 2004). The BFNE-S is modified from the original 12-item BFNE that was developed by Leary (1983) and includes only the 8 straightforwardly-worded items. The BFNE-S uses a 5-point Likert-type scale ranging from 1 (Not at all characteristic of me) to 5 (Extremely characteristic of me). Psychometric studies have suggested that the eight straightforward items of the scale, as compared to the four reverse-scored items or the entire BFNE, demonstrate the strongest reliability and validity (Carleton, Collimore, McCabe, & Antony, 2011; Rodebaugh et al., 2011; Rodebaugh et al., 2004; Weeks et al., 2005).

Sheehan Disability Scale (SDS; Sheehan, 1983). The SDS is a 3-item measure that assesses the degree to which an individual's symptoms affect their work, social, and home life.

The items are measured using a 10-point visual analog scale that ranges from 0 (Not at all) to 10 (Extremely), and total scores range from 0, indicating no impairment, to 30, indicating significant impairment. The SDS has demonstrated a unifactorial structure with acceptable internal consistency (Leon, Shear, Portera, & Klerman, 1992). It has been demonstrated to discriminate between those who are experiencing psychiatric symptoms and those who are symptom free, suggesting good construct validity (Leon et al., 1997; Leon et al., 1992; Olfson et al., 1997). Among people with SAD, the SDS has also shown signs of good validity and modest internal consistency, although longer scales measuring disability perhaps unsurprisingly showed stronger properties (Hambrick, Turk, Heimberg, Schneier, & Liebowitz, 2004).

Data Analytic Procedure

A priori tests versus revised analyses. We originally examined all 48 of the LSAS items (24 fear and avoidance situations) in a network. However, as pointed out during the review process, the covariance matrix of the symptoms was not positive definite,¹ presumably due to high collinearity between anxiety and avoidance ratings of each situation. Thus, we investigated the anxiety and avoidance items separately. We examined three commonly-used centrality measures: Betweenness, centrality, and strength. We determined that centrality indices were moderately to highly correlated across the two items sets (betweenness: $r_s = .38-.43$, $p_s = .006-.06$; closeness: $r_s = .43-.46$, $p_s = .025-.037$; strength: $r_s = .85-.89$, $p_s < .001$; note that strength was the most stable index in each set of items). We therefore decided to add anxiety and avoidance items together for each situation. In addition to adding together anxiety and avoidance items across *situations*, we observed that two *situations* were not only very similar in concept, but also so highly correlated as to suggest that they were measuring the same construct: *talking*

¹We thank Sacha Epskamp, who provided a signed review, for uncovering this problem.

to people you don't know very well and *meeting strangers* (fear: $r = .61$; avoidance: $r = .64$); and *performing in front of an audience* and *giving a report to a group* (both fear and avoidance r s = $.77$). Treating these situations as separate could produce nonsensical estimates in the same manner that would occur if one ran a regression with two highly-correlated measures of anxiety included as separate predictors. We therefore additionally summed the two highly correlated situation pairs to generate 22 nodes for analysis. Finally, we rescaled the two nodes drawn from 4 items by using the cut function in R (R Core Team, 2017) to reproduce the same 0-6 scale of all other nodes.

The a priori analyses we originally conducted were predicated on the assumption that there would be a set of items that either clearly had higher centrality than all others, or were at least more stable in their high centrality than others. In the revised analyses, this was no longer the case. Accordingly, we adopted the method used by Robinaugh et al. (2016). Although this is not the method we had selected a priori, it is suitable for the situation in the revised data, in which the nodes all have fairly stable rankings in terms of centrality and no small set of nodes is clearly higher in centrality than the others.

Revised analyses. Here we focus on the overall plan for the analyses (more detailed information on each aspect of the data analytic procedure is available in the supplemental materials). We first created five multiple imputation data sets Sample A to handle sporadic missing data, using random forest imputation. A total of 11% of participants with GSAD in Sample A had at least some missing data. Because there is no best standard for how to deal with multiply imputed data in network analyses, we focused on consistent findings across all five data sets. Following the compositing described above (resulting in 22 nodes), network estimation and network stability tests were conducted in accordance with current standards using R packages

qgraph (version 1.4.1; Epskamp, Cramer, Waldorp, Schmittmann, & Borsboom, 2012) and bootnet (version 0.3; Epskamp, Borsboom, & Fried, 2017), respectively. See Supplementary Materials for information on all R packages utilized. An undirected regularized partial correlation network was estimated, resulting in edges that can be interpreted as partial correlation coefficients (an association between two items controlling for all other associations among items). Regularization ensures that the estimated network structure balances sensitivity with specificity, and leads to a sparse network structure that avoids obtaining spurious edges (Epskamp & Fried, in press). We determined which centrality indices to consider further based on their correlation stability coefficient estimated in bootnet. Indices were determined to be stable if at least 25% of the cases could be removed and the order of nodes maintained a correlation of 0.7 (with 95% probability) with the original sample (see Epskamp, Borsboom, & Fried, 2017 for a description of the correlation stability coefficient). To ensure that results were not due to mere item properties that can bias centrality indices, we also examined frequency of endorsement (i.e., floor effects) and standard deviation (i.e., effects for the range of the item) (cf. Terluin et al., 2016). Infrequency was defined as the number of participants who scored 0 on that node (with nodes values ranging from 0 to 6); notably, a score of 0 would indicate the participant endorsed a 0 on all of the items that ultimately comprised that node. The standard deviation was simply the *SD* for that node.

Following the method used by Robinaugh and colleagues (2016) we then examined centrality indices across nodes. The outcomes of interest were the degree to which change in a given node correlated with change, from Sample B, in (a) the remainder of the LSAS items, (b) the SIAS-S items, (c) the BFNE-S items, and (d) the SDS items. For (a), the remainder of the LSAS items were defined as the total of the items minus the investigated node. Again, the notion

is that, for each outcome, higher centrality should be associated with a higher correlation between change in that node and change in the outcome. Note that because some participants dropped out of treatment or did not provide the given measure, we obtained these correlations pairwise; each correlation was estimated using between 133 and 155 cases. To determine whether any correlations might be better explained by relatively superficial item properties, we also conducted multiple regressions in which centrality and item properties were included as predictors. When conducting multiple regressions, we examined the correlations between the centrality indices and composited them if correlations were high (e.g., $> .50$) because the intent was to assess the usefulness of centrality indices rather than pit them against each other.

We also examined the SDBeta statistic, which identifies items that are overly influential, such that their removal would have a strong impact on the regression coefficient. We considered this test essential given the small sample size for these analyses (i.e., 22 nodes). We report below when we removed nodes that had an SDBeta value in excess of 1 or -1 (Neter, Wasserman, & Kutner, 1989).

Results

Network Estimation

We first estimated the overall network structure for each of the five imputations of Sample A. Figure 2 displays the results for the first imputed dataset; the (very similar) figures for the other imputed datasets are presented in supplementary material (Figures S1 and S2). The results presented below were across all of the imputed datasets, and not simply the first one. We examined the centrality stability indices for strength, closeness, and betweenness to determine which metrics were appropriate for further analysis. Using the cut-off of 0.25 suggested by simulation studies (Epskamp, Borsboom, & Fried, 2016), we determined that strength

($CS[\text{cor}=0.7] = .59-.67$) and betweenness ($CS[\text{cor}=0.7] = .28$) were sufficiently stable across imputations to justify use in subsequent analyses. *Strength* refers to how strongly a node relates to other nodes, whereas *betweenness* refers to how important a node is in paths between other nodes (cf. Epskamp et al. 2017). Another centrality index, closeness, was less stable ($CS[\text{cor}=0.7] = .21 - .28$) and we refrain from interpreting it in the primary analyses.

Treatment Response Prediction

Within-LSAS-network prediction. In the within-network prediction, the test was whether centrality indices predicted the correlation between change in a node and change in the rest of the network. We will call the dependent variable for this set of analyses *node-LSAS change correlation*. Notably, these analyses were conducted with *nodes* as the unit of analysis and not people: Values for the nodes were obtained from Sample A (for centrality and item properties) and Sample B (for correlations). Because the data analyzed were by node, the entire dataset is represented in Table S1 in the supplementary material.

As hypothesized, both strength ($r = .48, p = .026$) and betweenness ($r = .53, p = .011$) were related to the node-LSAS change correlation. This finding indicates that the centrality indices successfully identified nodes for which their change was more strongly associated with change in the rest of the network. Because strength and betweenness were strongly correlated ($r = .66, p = .001$), they were z-scored and composited for remaining analyses (and referred to as the *centrality composite*). We did this because entering them as individual predictors would lead to a focus on their unique properties, whereas we were interested in centrality overall.

Infrequency of item endorsement ($r = -.61, p = .003$) was also strongly associated (whereas node *SD* was not: $r = .04, p = .874$) with the node-LSAS change correlation. Further, strength and betweenness showed a pattern of correlations with infrequency and variance that

might indicate that this association partially explained their relationship to change (e.g., the centrality composite's correlation with infrequency was $-.30, p = .179$; with $SD, r = .48, p = .025$). To determine the relative strength of prediction for these associations, the three node properties (centrality composite, infrequency, and SD) were entered in a regression equation predicting the node-LSAS change correlation. The coefficients for these analyses are presented in Table 2. All three predictors were statistically significant. Notably, the result for SD showed the opposite sign of what was expected (suggesting statistical suppression) and did not survive correction for nodes with excessive $SD\beta$ values (see Table 2). Thus, *both* the centrality composite and infrequency identified nodes with a stronger relationship with change in other items. When items with higher centrality changed, other LSAS items were more likely to change in comparison to when lower centrality items changed. In addition, when items that were infrequently endorsed changed, other LSAS items were less likely to change in comparison to when items that were more frequently endorsed changed.

Outside LSAS-network prediction of social anxiety severity. We next repeated the analyses conducted above for three measures not included in the LSAS network. More specifically, the question was how strongly centrality (from Sample A) predicted how change in a given node correlated with change in the social anxiety severity measure in question (from Sample B). Examining zero-order correlations, betweenness displayed at best marginal relationships (SIAS-S: $r = .18, p = .451$; BFNE-S: $r = .19, p = .388$, SDS: $r = .38, p = .079$), and neither strength nor the centrality composite showed any sign of predicting ($ps > .10$). Infrequency strongly predicted across measures (SIAS-S: $r = -.68, p = .001$; BFNE-S: $r = -.73, p < .001$, SDS: $r = -.78, p < .001$). In contrast, SD showed no relationship ($ps > .23$). Multiple

regressions are shown in Table 2 and are consistent with zero-order correlations: Only infrequency predicted strength of association in the additional social anxiety severity measures.²

Follow-up Tests

Rationale. Although there was some support for the hypothesis that highly central nodes predict more change in other nodes, infrequency of endorsement was a much more robust predictor. Three explanations for this pattern presented themselves (that are not mutually exclusive). First, the Sample A network indices may have been influenced by superficial item properties. Second, the Sample A network might not be consistent with the network structure for Sample B. Third, the fact that participants all had GSAD might produce a distorted network structure because participants were selected based in part on properties of the network (i.e., people with GSAD typically have higher LSAS scores than those without). We attempted to address each concern below to determine whether addressing these concerns (a) reduced effects for infrequency or (b) increased effects for centrality. The interested reader can re-run these analyses using the data provided in Table S1.

²Given this unexpected result, we also checked in Sample B itself to be sure that change on the LSAS, SIAS-S, BFNE-S, and SDS were correlated: They were ($r_s > .35$, $p_s < .001$, $n_s > 133$). We also checked whether changes in at least some nodes were correlated with changes in the SIAS-S, BFNE-S, and SDS; this was also true, with each measure having multiple nodes for which changes were correlated at a level of $p < .001$. This result was therefore not due to the LSAS failing to correlate with other measures, either on the level of the entire LSAS or the individual nodes. On a related note, Sacha Epskamp, who provided a signed review, pointed out that even if no casual process over time were involved in regard to the LSAS scores, centrality might predict in the manner seen here due merely to regression towards the mean. This is obviously an essential point, but one we were not able to address adequately here. We encourage further exploration (via mathematical proof, simulation, and experimental manipulation) of under what conditions centrality indices should be expected to identify nodes that are central to change due to causal processes (as opposed to statistical necessities).

Ising network. We addressed the first problem by reducing the probability that infrequency was having an effect on centrality indices. We dichotomized all nodes in Sample A using a median split of each node such that responses below the median were coded as 0. In cases where the median was 0, responses of 0 and 1 were coded as 0 instead. This procedure minimizes the effect of variance and infrequency (as well as all other item properties) on network estimation, because most had equivalent numbers of participants who endorsed 0. We then conducted the same analyses described above using the centrality indices from an Ising network. We investigated strength and closeness because these indices had interpretable levels of stability in 4 of 5 imputed datasets; none of the indices from the Ising network had acceptable stability across all 5 imputed datasets. Because the two indices were highly correlated ($r = .81, p < .001$) they were composited and combined for analysis. Substantive results for this centrality composite were identical to those reported above; it predicted regarding the LSAS (part $r = .45, p = .012$, corrected for SDBeta) but not other measures ($ps > .30$); infrequency predicted for all ($ps < .008$). Dichotomizing items and thereby addressing skewed distributions in the data did not consistently change the pattern of results, making it unlikely that skew is the main reason for the results obtained.

Correspondence of centrality indices between cross-sectional and treatment data. To address the second problem, that the results from Sample A might vary widely from the results Sample B (i.e., the treatment sample itself), we repeated the procedure we used for Sample A in Sample B. Of the centrality indices, none showed acceptable stability in these data; however, strength was the most stable metric ($CS[cor=0.7] = .21$). Strength from the pre-treatment and cross-sectional data showed reasonably good correspondence across datasets ($r = .61, p = .003$; two-way random ICC for the single measure = $.57, p = .002$, 95% CI: $.21 - .80$). The entire

correlation matrix of the centrality indices is provided in Table S2. We then repeated the multiple regressions, this time using strength, infrequency, and variance from Sample B. There were no significant predictors for the LSAS ($ps > .06$ after correction for SDBetas). For the other measures, strength did not predict ($ps > .21$), whereas infrequency predicted for the SIAS-S and BFNE-S ($ps < .03$) but not the SDS ($p = .094$; all ps correcting for SDBetas). Relying on Sample B produced more mixed results that showed no increased effects for centrality indices.

Addition of participants without GSAD. We added participants to Sample A who were diagnosed using the same procedures as the GSAD participants, but who were either recruited as normal control participants or who did not meet for GSAD diagnosis despite expectations from screening that they would. A total of 197 participants were added to Sample A. We then re-ran the original procedures and extracted centrality indices from this larger dataset. Strength ($CS[cor=0.7] = .59-.67$) and betweenness ($CS[cor=0.7] = .28-.36$) showed acceptable stability in all imputations, whereas closeness did not ($CS[cor=0.7] = .21-.28$). The correlation table for all three centrality indices from the larger versus smaller cross-sectional dataset is provided in Table S2; strength and betweenness were strongly correlated ($r = .78, p < .001$) and were therefore standardized and combined. This centrality composite, infrequency, and *SD* from the expanded sample were used as competing predictors in a multiple regression. The results from these analyses are presented in Table 2 because it presented the only instance in which infrequency demonstrated reduced effects. For the LSAS, only centrality predicted strength of association. In contrast, for the SIAS-S, BFNE-S, and SDS, only infrequency predicted strength of association and the other predictors did not.

From the above results, it was unclear whether the apparent improvement in the prediction by centrality indices regarding the LSAS was due to improved estimation of centrality

indices or reduced utility of infrequency estimates from the expanded version of Sample A. We therefore also ran this regression using the original Sample A infrequency and *SD* estimates. In this analysis, infrequency did predict regarding the LSAS at about the same level as centrality (infrequency: part $r = -.46$, $p = .006$; centrality: part $r = .44$, $p = .009$ after correction for *SDBeta*), but the evidence for centrality indices remained more convincing than in the primary analyses. There was thus some evidence that centrality performed more in keeping with hypothesis when the sample was not restricted to participants who are expected to score high on the LSAS, but this improvement was not seen for measures other than the LSAS, where infrequency continued to show the strongest associations. Infrequency, in contrast, showed stronger associations when estimated based on GSAD participants alone.

Nodes of particular interest. The nodes with highest centrality, based on the z-scored and combined centrality composite from the expanded Sample A analysis were: (a) the combined Situations 11 and 12 (talking with unfamiliar people), (b) Situation 15 (center of attention), (c) Situation 7 (going to a party). The least central were (a) Situation 17 (test-taking), (b) Situation 21 (asking someone on a date) and (c) Situation 1 (telephoning). The nodes with the lowest infrequency from the primary analyses were: (a) the combined Situations 6 and 20 (public speaking or performance), (b) Situation 16 (commenting during a meeting), (c) Situation 15 (center of attention). The nodes with the highest infrequency were: (a) Situation 13 (urinating in restroom), (b) Situation 4 (drinking in public), and (c) Situation 9 (writing while observed). Node values on specific centrality indices, as well as the different versions of each variable, can be obtained through examination of the data in Table S1 in the Supplementary Material.

Discussion

We sought to determine whether central symptoms identified via network analysis of cross-sectional data would predict the correlation between change in a given node and change in other symptoms across treatment in a second dataset. We hypothesized that items identified as highly central in the LSAS would have a stronger ability to predict change of symptoms across treatment, in accordance with other suggestions in the literature (McNally et al., 2015; Ruzzano et al., 2015). We found that centrality did predict which nodes were more strongly associated with change above and beyond other predictors. However, this prediction was restricted entirely to the LSAS itself. LSAS nodes with higher centrality indices showed no promise as useful indicators of change in other measures of social anxiety severity. In contrast, how frequently items were endorsed showed a more consistent ability to predict node importance, both within the LSAS and in extension to other measures. Nodes that were more frequently endorsed were much more likely to show signs of being influential across treatment.

What do our results suggest about the assertion that centrality from a cross-sectional network is a good guide to determining which symptoms are important to focus on in treatment? Our findings clearly run counter to the pessimistic view that centrality indices from cross-sectional data would tell us nothing about associations over time. An optimistic reading of our results might conclude that centrality indices, and particularly strength (a more stable index), might provide some information about which symptoms are more important for treatment. Higher centrality in our data was indeed associated with a stronger association with change across the entire LSAS network: Targeting the highly central symptoms might therefore promote generalization of treatment gains across the LSAS as a whole. Clinical scientists might therefore take our results as license to interpret existing centrality findings as indicating good targets for treatment. However, there are at least three caveats to this conclusion.

Caveat One: Select Items with Care

First, our results imply that simply analyzing the items of a given measure may not produce such promising results unless care is taken in determining in selecting nodes for the network analyses. Our initial results obtained by simply analyzing all of the items (see Supplemental Material) seemed to indicate that centrality indices were conflated with infrequency of endorsement, whereas our revised analyses did not indicate this was the case. Avoiding very high correlations among nodes appeared to ameliorate the effects of infrequency of endorsement on the network. Perhaps importantly, the LSAS is a frequently-used measure that is widely regarded as having great clinical utility based on a strong evidence base for its validity in measuring symptoms of SAD (see Fresco et al., 2001 for a review). Unfortunately, analyzing the individual items of this arguably gold-standard measure proved inadvisable due to very high correlations among fear and avoidance nodes that represented the same situation. Deciding in what form to include items so as to avoid nonsensical results took considerable thought and care; this may serve as a warning to researchers who may be inclined to analyze the items of a measure without consideration of item properties and intercorrelations, a method that might seem to be endorsed by early demonstrations of network analysis focused on psychopathology (cf. Fried & Cramer, 2017).

To the extent that other researchers have followed these early examples, clinicians and researchers should take care in interpreting published findings regarding centrality indices. For example, during the review process for this manuscript we became aware of two network analyses focusing on the LSAS (Heeren, Jones, & McNally, 2018; Heeren & McNally, 2018), one of which (Heeren & McNally) focused on the LSAS alone, as we did here. These researchers indeed followed the same procedure typically used in previous network analyses: All items were

included in the network in their original form. We contacted the authors (Heeren & McNally), who confirmed that their data produce the same error message we received (i.e., a warning regarding a non-positive definite matrix) when the data are analyzed using the same method we used (A. Heeren, personal communication, June 7 and June 13, 2018).³ We expect that even in cases in which the statistical error does not arise, the conceptual problem of including multiple nodes that measure the same construct may be common. The clinical scientist who is inclined to interpret centrality indices optimistically as a result of our results should be aware that many researchers in the area have only recently begun to consider that it may not be optimal to include all items on a measure as separate nodes in analysis.

Caveat Two: The Importance of Item Properties

Second, our evidence indicated a medium-to-large effect for centrality, whereas infrequency generally showed larger effects. Some readers might object that there is no reason for the “whereas” in the previous sentence: It should be no surprise that floor effects are important. However, it may be a surprise to clinicians to find that our positive results for centrality indices come with the context that it would be even better to treat common symptoms (which, arguably, existing treatments tend to focus on already). Of course, our results indicate that it would be an even better idea to use both indices to select items: That is, clinicians could treat select symptoms on the basis of both centrality and frequency of endorsement. We are of two minds on this point; some of the current authors see no contradiction between centrality indices and floor effects having an influence on results. For some of the current authors, however, this situation is unsatisfying because it indicates that relatively superficial item properties may be more important, in some instances, than centrality indices derived from

³We are grateful to the authors for their speedy and open discussion of this issue. At the time of this writing, the authors are working with the editor of the journal to publish an erratum.

sophisticated analyses that researchers and clinicians hope will uncover causal processes. Of course, it remains possible that the highly endorsed items play an important causal role. It could be the case that fear of giving presentations has a unique causal role for SAD; we simply cannot separate any such causal role from the item properties given our data. We do believe, however, that our results can serve as a warning (that echoes those of other authors, e.g., Terluin et al., 2016) to researchers to examine the issue of item properties, such as endorsement rates, when conducting any statistical analysis, including network analysis.

In network analysis, the typical current practice is to examine sets of single item scores as if items themselves measure constructs directly. This practice maximizes the chances that results could be influenced by relatively superficial item properties. We have shown in this paper, however, that careful examination of items can help reduce this possibility. It may also be useful to re-examine the latent variable models that served as the spur to move in a different direction (cf. Fried & Cramer, 2017). We say this because latent variables provide a method to combine multiple items based on their inferred relationship with the underlying variable that is being measured. This property of latent variables cannot completely eliminate the influence of factors such as floor effects, but it can at least reduce that influence. Indeed, we arguably accomplished a similar goal in a less precise way here by combining items. A rapprochement between latent variable models and network analyses may be fruitful, as has been previously suggested (Fried & Cramer, 2017). Epskamp and colleagues have recently presented an approach that allows a combination of the two methods, although multiple challenges remain (Epskamp, Rhemtulla, & Borsboom, 2017). An alternative would be to develop measures that are free of floor or ceiling effects, but we have no optimism that this will be accomplished soon for the measurement of clinical problems.

Caveat Three: The Gates of Causality

There remains a third caveat for those who might take our results as indicating that cross-sectional networks will yield informative centrality indices. Centrality indices showed no ability to detect items that showed influence outside of the LSAS, whereas frequency of item endorsement did. This puts a reader inclined to an optimistic reading in an awkward position. Centrality seems to identify important LSAS items, but only within the wall, so to speak, of the LSAS itself. Accepting this result as evidence of the importance of centrality indices would require the reader to also accept that our results imply that importance stops at the gates of the LSAS. This conclusion seems awkward to us because the other measures assess constructs that should be close neighbors to the constructs assessed by the LSAS. For example, the LSAS includes items assessing social interaction anxiety: Why, then, should centrality not also identify items showing signs of greater influence on the SIAS-S, a unifactorial measure of social interaction anxiety? Although we have thought of answers for that question, none of the answers are particularly satisfying or explain how centrality indices can index important causal processes and still provide the current result. For example, centrality might identify which items are most heavily saturated with variance that is unique to the LSAS, such as some form of method variance that remained consistent across Sample A and Sample B. For example, both versions of the LSAS used include a specific instruction to focus on the past week; the other measures we examined do not.⁴ Whether due to the focus on the past week or for some other reason related to

⁴The SDS, however, implies a focus on the past week (by including an option to check a box indicating that one was out of work for the past week for reasons other than the disorder) without specific instructions to that effect. One might therefore argue that the SDS should contain similar method variance. However, as can be seen in Table 2, the results from the SDS could be interpreted as being more positive regarding centrality than for the other measures. Thus, the possibility that this method issue is important is not contraindicated.

method variance unique to the LSAS, the result we obtained would be expected: The highly central items are important within the LSAS, but not outside of it (unless another measure shares that method component). The point is that it is difficult to rectify centrality as an important index of causality with an effect that ends at the gates of a specific measure. On the other hand, it may be possible to read these results optimistically, as an indication that networks should contain the entire set of variables involved in a causal network. From this point of view, the problem is that only the LSAS was included in the network.

Limitations and Future Directions

Our initial results were vulnerable to several concerns: (a) the network may have been affected by specific item properties (such as infrequency and variance); (b) the network may have varied meaningfully from the network that would have been estimated from the smaller pre-treatment dataset, such that the pre-treatment network would produce more useful estimates; and (c) the network may have been affected by the selection of GSAD participants. Our follow-up tests provided clarity: Only the addition of participants who did not have GSAD resulted in any sign of improved prediction for centrality indices, and even in this case infrequency remained a more robust predictor (at least outside of the LSAS network). Thus, none of these concerns appear to explain the effects observed.

We believe that the most important limitation of our work is that we did not select symptoms for specific intervention and test the resulting changes in a network when those symptoms are targeted (in comparison to when other symptoms are targeted). That scenario is clearly the desired goal of network analyses that many researchers have referred to. We focused on tests that were plausible and possible given available data; further, we tested a logical extension of the idea that centrality indices will identify key symptoms for treatment. Clearly,

however, direct tests of randomly assigned interventions at the symptom level are sorely needed. One advantage of such direct tests is that they would provide the ability for true prospective tests of centrality indices. Here, we were limited to testing whether centrality was correlated with how change in nodes correlated with change in other nodes and measures. Although these analyses involved time, they do not represent fully prospective prediction, which would be preferable. Among other reasons, fully prospective prediction with random assignment to a meaningful control condition would allow one to rule out regression to the mean as a competing explanation.

Notably, our findings, even if replicated, do not rule out the possibility that there might be some instances in which cross-sectional networks offer important information about symptoms that are key to changing not only that network, but beyond. Perhaps social anxiety symptoms, or the LSAS items in particular, are very different from other types of symptoms, and centrality indices from cross-sectional networks for another disorder would show different properties. Although the extent to which cross-sectional data have meaning for causality is complex and contentious, some authors have proposed situations under which cross-sectional data should be expected to yield causal insights (Pearl, 2000). In brief, among other conditions, the modeling strategy of directed acyclic graphs presented by Pearl requires that there cannot be feedback loops or vicious circles; more generally, only one direction of effect can be modeled between two variables in cross-sectional data (if A causes B, B cannot cause A). Pearl's approach also assumes all important variables in the causal system are included in the network. It is unclear to us whether clinical researchers generally consider these issues regarding cross-sectional data, but it is possible that giving them greater attention (cf. Morgan & Winship, 2015; Pearl, 2000) might lead to more positive results. Notably, however, one or two sets of

psychological symptoms (i.e., the typical focus of most network analysis papers thus far) do not seem like plausible candidates for a set of variables that would satisfy these conditions.

Some additional concerns are worth discussing. First, although our samples were large and reasonably diverse, greater diversity (e.g., racial diversity) would have been desirable. Second, we do not believe the kind of data examined here would ever be expected to hold to the conditions that have been suggested as necessary for cross-sectional data to comment on causality, but in theory the measurement could have been improved to increase applicability. For example, the full range of theorized causes of social anxiety could have been included in the model, rather than social anxiety symptoms alone. That said, our examination was focused on determining whether currently common network analyses would be successful in determining key symptoms, and the available networks rarely move beyond the confines of symptoms of one or two disorders (cf. Fried, 2016). Finally, there are other centrality indices available (e.g., Haslbeck & Fried, 2017), and we are aware of others that are under development. It is always possible that these indices will prove more useful, although we encourage skepticism given our current results.

Conclusions

Keeping those limitations in mind, we have several recommendations for treatment providers and treatment outcome researchers. First, we suggest caution in interpreting existing networks using cross-sectional data as indicating important symptoms that should be focused on in treatment. It remains possible that some published networks might eventually be shown to have pointed in a useful direction, but our data clearly indicate that cross-sectional networks, even in a large dataset, cannot always be taken as a clear indicator of symptoms that are important in predicting change across treatment for symptoms overall (as opposed to the specific

items used for analysis). Of note, attempting to find one or two most central symptoms to focus on gains even less support from our analyses, which (a) focused on centrality indices as *continuous* variables (i.e., not one or two most central symptoms) and (b) revealed that, unsurprisingly, the precise symptoms that were most central varied by analysis, sample, and centrality index (see Table S1).

Second, for those instances in which researchers wish to focus on cross-sectional data to inform treatment research, we urge them to carefully consider whether their data and methods are consistent with the recommendations of theorists who at least find it plausible that this is a fruitful exercise (e.g., Pearl, 2000). Third, we note that longitudinal and experimental network analysis studies are far rarer than cross-sectional studies at this time. This imbalance in the literature is unfortunate: Causal relationships might be better addressed using a combination of longitudinal (i.e., both in groups of participants and individuals) and experimental studies examining networks of symptoms (cf. Fried & Cramer, 2017) alongside carefully conducted tests in cross-sectional data. This call has been made before in regard to network analysis (e.g., Borsboom, & Cramer, 2013; van de Leemput et al., 2014, among others), but we believe the call should be repeated. Longitudinal and experimental studies are difficult and expensive, but they may be our best hope for discovering symptoms that are actually crucial for intervention: We would have to devote more attention to such studies to fully evaluate their promise.

References

- American Psychiatric Association (1994). *Diagnostic and statistical manual of mental disorders: DSM-IV* (4th ed.). Washington, DC: American Psychiatric Association.
- Baker, S. L., Heinrichs, N., Kim, H.-J., & Hofmann, S. G. (2002). The Liebowitz Social Anxiety Scale as a self-report instrument: A preliminary psychometric analysis. *Behaviour Research and Therapy*, 40, 701-715. doi:10.1016/S0005-7967(01)00060-2
- Blanco, C., Heimberg, R. G., Schneier, F. R., Fresco, D. M., Chen, H., Turk, C. L., ... & Campeas, R. (2010). A placebo-controlled trial of phenelzine, cognitive behavioral group therapy, and their combination for social anxiety disorder. *Archives of General Psychiatry*, 67, 286-295.
- Borsboom, D. (2008). Psychometric perspectives on diagnostic systems. *Journal of Clinical Psychology*, 64, 1089-1108. doi:10.1002/jclp.20503
- Borsboom, D. (2017). A network theory of mental disorders. *World Psychiatry*, 16, 5-13. doi:10.1002/wps.20375
- Borsboom, D., & Cramer, A.J. (2013). Network analysis: An integrative approach to the structure of psychopathology. *Annual Review of Clinical Psychology*, 99, 91-121. doi:10.1146/annurev-clinpsy-050212-185608
- Borsboom, D., & Cramer, A. O. (2013). Network analysis: an integrative approach to the structure of psychopathology. *Annual Review of Clinical Psychology*, 9, 91-121. doi:10.1146/annurev-clinpsy-050212-185608
- Borsboom, D., Cramer, A. O. J., Schmittmann, V. D., Epskamp, S., & Waldorp, L. J. (2011). The small world of psychopathology. *PLoS ONE*, 6, ArtID: e27407. doi:10.1371/journal.pone.0027407

- Borsboom, D., Rhemtulla, M., Cramer, A. O. J., van der Maas, H. L. J., Scheffer, M., & Dolan, C. V. (2016). Kinds versus continua: A review of psychometric approaches to uncover the structure of psychiatric constructs. *Psychological Medicine*, *46*, 1567-1579. doi:10.1017/s0033291715001944
- Bringmann, L. F., Lemmens, L. H. J. M., Huibers, M. J. H., Borsboom, D., & Tuerlinckx, F. (2015). Revealing the dynamic network structure of the Beck Depression Inventory-II. *Psychological Medicine*, *45*, 747-757. doi:10.1017/s0033291714001809
- Bringmann, L. F., Pe, M. L., Vissers, N., Ceulemans, E., Borsboom, D., Vanpaemel, W., . . . Kuppens, P. (2016). Assessing temporal emotion dynamics using networks. *Assessment*, *23*, 725-735.
- Bringmann, L. F., Vissers, N., Wichers, M., Geschwind, N., Kuppens, P., Peeters, F., . . . Tuerlinckx, F. (2013). A network approach to psychopathology: new insights into clinical longitudinal data. *PLoS ONE*, *8*, e60188. doi:10.1371/journal.pone.0060188
- Brown, T. A., & Barlow, D. H. (1994). Anxiety Disorders Interview Schedule for DSM-IV (ADIS-IV).
- Brown, T. A., & Barlow, D. H. (2014). Anxiety and Related Disorders Interview Schedule for DSM-5 (ADIS-5L): Client interview schedule.
- Carleton, R. N., Collimore, K. C., McCabe, R. E., & Antony, M. M. (2011). Addressing revisions to the Brief Fear of Negative Evaluation scale: Measuring fear of negative evaluation across anxiety and mood disorders. *Journal of Anxiety Disorders*, *25*, 822-828.
- Cole, D. A., & Maxwell, S. E. (2003). Testing mediational models with longitudinal data: Questions and tips in the use of structural equation modeling. *Journal of Abnormal Psychology*, *112*, 558-577. doi:10.1037/0021-843x.112.4.558

- Cramer, A. O. J., Waldorp, L. J., van der Maas, H. L. J., & Borsboom, D. (2010a). Comorbidity: A network perspective. *Behavioral and Brain Sciences*, 33, 137-150.
doi:10.1017/s0140525x09991567
- Cramer, A. O. J., Waldorp, L. J., van der Maas, H. L. J., & Borsboom, D. (2010b). Complex realities require complex theories: Refining and extending the network approach to mental disorders. *Behavioral and Brain Sciences*, 33, 178-193.
doi:10.1017/s0140525x10000920
- de Vos, S., Wardenaar, K. J., Bos, E. H., Snippe, E., Hartmann, J. A., Simons, C. J. P., . . . Wichers, M. (2017). Can we jump from cross-sectional to dynamic interpretations of networks? Implications for the network perspective in psychiatry. *Psychotherapy and Psychosomatics*, 86, 175-177. doi:10.1159/000453583
- DiNardo, P. A., Brown, T. A., & Barlow, D. H. (1994). *Anxiety Disorders Interview Schedule for DSM-IV: Lifetime Version (ADIS-IV-L)*. San Antonio, TX: The Psychological Corporation.
- Epskamp, S., Borsboom, D., & Fried, E. I. (2017). Estimating psychological networks and their accuracy: A tutorial paper. *Behavior Research Methods*. doi:10.3758/s13428-017-0862-1
- Epskamp, S., Cramer, A. O., Waldorp, L. J., Schmittmann, V. D., & Borsboom, D. (2012). qgraph: Network visualizations of relationships in psychometric data. *Journal of Statistical Software*, 48, 1-18.
- Epskamp, S., & Fried, E. I. (in press). A tutorial on regularized partial correlation networks. *Psychological Methods*.

- Epskamp, S., Rhemtulla, M., & Borsboom, D. (2017). Generalized network psychometrics: Combining network and latent variable models. *Psychometrika*. doi:10.1007/s11336-017-9557-x
- First, M., Williams, J., Karg, R., & Spitzer, R. (2015). Structured Clinical Interview for DSM-5—research version. *American Psychiatric Association*.
- First, M. B., Spitzer, R. L., Gibbon, M., & Williams, J. B. W. (2002). Structured Clinical Interview for DSM-IV-TR Axis I Disorders, Research Version, Patient Edition (SCID-I/P).
- Fresco, D. M., Coles, M. E., Heimberg, R. G., Liebowitz, M. R., Hami, S., Stein, M. B., & Goetz, D. (2001). The Liebowitz Social Anxiety Scale: A comparison of the psychometric properties of self-report and clinician-administered formats. *Psychological Medicine*, 31, 1025-1035. doi:10.1017/s0033291701004056
- Fried, E. I., Borkulo, C. D., Cramer, A. O. J., Boschloo, L., Schoevers, R. A., & Borsboom, D. (2016). Mental disorders as networks of problems: A review of recent insights. *Social Psychiatry and Psychiatric Epidemiology*, 52, 1-10. doi:10.1007/s00127-016-1319-z
- Fried, E. I., & Cramer, A. O. J. (2017). Moving Forward: Challenges and directions for psychopathological network theory and methodology. *Perspectives on Psychological Science*, 1745691617705892. doi:10.1177/1745691617705892
- Fried, E. I., Epskamp, S., Nesse, R. M., Tuerlinckx, F., & Borsboom, D. (2016). What are 'good' depression symptoms? Comparing the centrality of DSM and non-DSM symptoms of depression in a network analysis. *Journal of Affective Disorders*, 189, 314-320. doi:10.1016/j.jad.2015.09.005

- Goldin, P. R., Morrison, A., Jazaieri, H., Brozovich, F., Heimberg, R., & Gross, J. J. (2016). Group CBT versus MBSR for social anxiety disorder: A randomized controlled trial. *Journal of Consulting and Clinical Psychology, 84*, 427-437. doi:10.1037/ccp0000092
- Goldin, P. R., Ziv, M., Jazaieri, H., Werner, K., Kraemer, H., Heimberg, R. G., & Gross, J. J. (2012). Cognitive reappraisal self-efficacy mediates the effects of individual cognitive-behavioral therapy for social anxiety disorder. *Journal of Consulting and Clinical Psychology, 80*, 1034-1040. doi:10.1037/a0028555
- Gollob, H. F., & Reichardt, C. S. (1987). Taking account of time lags in causal models. *Child Development, 58*, 80-92.
- Hambrick, J. P., Turk, C. L., Heimberg, R. G., Schneier, F. R., & Liebowitz, M. R. (2004). Psychometric properties of disability measures among patients with social anxiety disorder. *Journal of Anxiety Disorders, 18*, 825-839.
- Harpole, J. K., Levinson, C. A., Woods, C. M., Rodebaugh, T. L., Weeks, J. W., Brown, P. J., . . . Liebowitz, M. (2015). Assessing the straightforwardly-worded Brief Fear of Negative Evaluation Scale for differential item functioning across gender and ethnicity. *Journal of Psychopathology and Behavioral Assessment, 37*, 306-317. doi:10.1007/s10862-014-9455-9
- Haslbeck, J. M. B., & Fried, E. I. (2017). How predictable are symptoms in psychopathological networks? A reanalysis of 18 published datasets. *Psychological Medicine*. doi:10.1017/S0033291717001258
- Heeren, A., Jones, P. J., McNally, R. J. (2018). Mapping network connectivity among symptoms of social anxiety and comorbid depression in people with social anxiety disorder. *Journal of Affective Disorders, 228*, 75-82. doi: <https://doi.org/10.1016/j.jad.2017.12.003>

- Heeren, A. & McNally, R. J. (2018). Social anxiety disorder as a densely interconnected network of fear and avoidance of social situations. *Cognitive Therapy and Research*, 42, 103-113. doi: 10.1007/s10608-017-9876-3
- Heimberg, R. G., Horner, K. J., Juster, H. R., Safren, S. A., Brown, E. J., Schneier, F. R., & Liebowitz, M. R. (1999). Psychometric properties of the Liebowitz Social Anxiety Scale. *Psychological Medicine*, 29, 199-212.
- Heimberg, R. G., Mueller, G. P., Holt, C. S., Hope, D. A., & Liebowitz, M. R. (1992). Assessment of anxiety in social interaction and being observed by others: The Social Interaction Anxiety Scale and the Social Phobia Scale. *Behavior Therapy*, 23, 53-73.
- Jazaieri, H., Goldin, P. R., Werner, K., Ziv, M., & Gross, J. J. (2012). A Randomized Trial of MBSR Versus Aerobic Exercise for Social Anxiety Disorder. *Journal of Clinical Psychology*, 68, 715-731. doi:10.1002/jclp.21863
- Leary, M. R. (1983). A brief version of the Fear of Negative Evaluation Scale. *Personality and Social Psychology Bulletin*, 9, 371-375.
- Ledley, D. R., Heimberg, R. G., Hope, D. A., Hayes, S. A., Zaider, T. I., Dyke, M. V., . . . Fresco, D. M. (2009). Efficacy of a manualized and workbook-driven individual treatment for social anxiety disorder. *Behavior Therapy*, 40, 414-424. doi:10.1016/j.beth.2008.12.001
- Leon, A. C., Olfson, M., Portera, L., Farber, L., & Sheehan, D. V. (1997). Assessing psychiatric impairment in primary care with the Sheehan Disability Scale. *The International Journal of Psychiatry in Medicine*, 27, 93-105. doi:10.2190/T8EM-C8YH-373N-1UWD

- Leon, A. C., Shear, M. K., Portera, L., & Klerman, G. L. (1992). Assessing impairment in patients with panic disorder: the Sheehan Disability Scale. *Social Psychiatry and Psychiatric Epidemiology*, 27, 78-82.
- Liebowitz, M. R. (1987). Social phobia. *Modern Problems of Pharmacopsychiatry*, 22, 141-173.
- Mattick, R. P., & Clarke, J. C. (1998). Development and validation of measures of social phobia scrutiny fear and social interaction anxiety. *Behaviour Research and Therapy*, 36, 455-470.
- Maxwell, S. E., & Cole, D. A. (2007). Bias in cross-sectional analyses of longitudinal mediation. *Psychological Methods*, 12, 23-44. doi:10.1037/1082-989x.12.1.23
- Maxwell, S. E., Cole, D. A., & Mitchell, M. A. (2011). Bias in cross-sectional analyses of longitudinal mediation: Partial and complete mediation under an autoregressive model. *Multivariate Behavioral Research*, 46, 816-841. doi:10.1080/00273171.2011.606716
- McNally, R. J., Robinaugh, D. J., Wu, G. W. Y., Wang, L., Deserno, M. K., & Borsboom, D. (2015). Mental disorders as causal systems: A network approach to posttraumatic stress disorder. *Clinical Psychological Science*, 3. doi:10.1177/2167702614553230
- Morgan, S. L., & Winship, C. (2015). *Counterfactuals and causal inference : methods and principles for social research* (Second Edition). New York, NY: Cambridge University Press.
- Muthén, L. K., & Muthén, B. O. (1998-2015). *Mplus user's guide*. (Seventh Edition ed.). Los Angeles, CA: Muthén & Muthén.
- Neter, J., Wasserman, W., & Kutner, M. G. (1989). *Applied linear regression analysis*. Homewood, IL: Irwin.

- Oakman, J., Van Ameringen, M., Mancini, C., & Farvolden, P. (2003). A confirmatory factor analysis of a self-report version of the Liebowitz Social Anxiety Scale. *Journal of Clinical Psychology, 59*, 149-161. doi:10.1002/jclp.10124
- Olfson, M., Fireman, B., Weissman, M. M., Leon, A. C., Sheehan, D. V., Kathol, R. G., . . . Farber, L. (1997). Mental disorders and disability among patients in a primary care group practice. *American Journal of Psychiatry, 154*, 1734-1740. doi:10.1176/ajp.154.12.1734
- Pearl, J. (2000). *Causality : models, reasoning, and inference*. Cambridge, U.K. ; New York: Cambridge University Press.
- Robinaugh, D. J., LeBlanc, N. J., Vuletich, H. A., & McNally, R. J. (2014). Network analysis of persistent complex bereavement disorder in conjugally bereaved adults. *Journal of Abnormal Psychology, 123*, 510-522. doi:10.1037/abn0000002
- Robinaugh, D. J., Millner, A. J., & McNally, R. J. (2016). Identifying highly influential nodes in the complicated grief network. *Journal of Abnormal Psychology, 125*. doi:10.1037/abn0000181
- Rodebaugh, T. L., Heimberg, R. G., Brown, P. J., Fernandez, K. C., Blanco, C., Schneier, F. R., & Liebowitz, M. R. (2011). More reasons to be straightforward: Findings and norms for two scales relevant to social anxiety. *Journal of Anxiety Disorders, 25*, 623-630. doi:10.1016/j.janxdis.2011.02.002
- Rodebaugh, T. L., Lim, M. H., Fernandez, K. C., Langer, J. K., Weisman, J. S., Tonge, N., . . . Shumaker, E. A. (2014). Self and friend's differing views of social anxiety disorder's effects on friendships. *Journal of Abnormal Psychology, 123*, 715-724. doi:10.1037/abn0000015

- Rodebaugh, T. L., Woods, C. M., & Heimberg, R. G. (2007). The reverse of social anxiety is not always the opposite: The reverse-scored items of the Social Interaction Anxiety Scale do not belong. *Behavior Therapy*, 38, 192-206.
- Rodebaugh, T. L., Woods, C. M., Heimberg, R. G., Liebowitz, M. R., & Schneier, F. R. (2006). The factor structure and screening utility of the Social Interaction Anxiety Scale. *Psychological Assessment*, 18, 231-237. doi:10.1037/1040-3590.18.2.231
- Rodebaugh, T. L., Woods, C. M., Thissen, D. M., Heimberg, R. G., Chambless, D. L., & Rapee, R. M. (2004). More information from fewer questions: The factor structure and item properties of the original and Brief Fear of Negative Evaluation scale. *Psychological Assessment*, 16, 169-181. doi:10.1037/1040-3590.16.2.169
- Ruzzano, L., Borsboom, D., & Geurts, H. M. (2015). Repetitive behaviors in autism and obsessive-compulsive disorder: New perspectives from a network analysis. *Journal of Autism and Developmental Disorders*, 45. doi:10.1007/s10803-014-2204-9
- Sheehan, D. V. (1983). *The anxiety disease*. New York: Scribner.
- Sheehan, D. V., Lecrubier, Y., Sheehan, K. H., Amorim, P., Janavs, J., Weiller, E., . . . Dunbar, G. C. (1998). The Mini-International Neuropsychiatric Interview (M.I.N.I.): The development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *Journal of Clinical Psychiatry*, 59(Suppl 20), 22-33.
- Stekhoven, D. J., & Buhlmann, P. (2012). MissForest--non-parametric missing value imputation for mixed-type data. *Bioinformatics*, 28, 112-118. doi:10.1093/bioinformatics/btr597
- Terluin, B., de Boer, M. R., & de Vet, H. C. W. (2016). Differences in Connection Strength between Mental Symptoms Might Be Explained by Differences in Variance: Reanalysis

of Network Data Did Not Confirm Staging. *PLoS ONE*, 11, e0155205.

doi:10.1371/journal.pone.0155205

van de Leemput, I. A., Wichers, M., Cramer, A. O. J., Borsboom, D., Tuerlinckx, F., Kuppens, P., . . . Scheffer, M. (2014). Critical slowing down as early warning for the onset and termination of depression. *PNAS Proceedings of the National Academy of Sciences of the United States of America*, 111, 87-92. doi:10.1073/pnas.1312114110

Weeks, J. W., Heimberg, R. G., Fresco, D. M., Hart, T. A., Turk, C. L., Schneier, F. R., & Liebowitz, M. R. (2005). Empirical validation and psychometric evaluation of the Brief Fear of Negative Evaluation Scale in patients with social anxiety disorder. *Psychological Assessment*, 17, 179-190.

Wichers, M. (2014). The dynamic nature of depression: a new micro-level perspective of mental disorder that meets current challenges. *Psychological Medicine*, 616, 1-12.

doi:10.1017/S0033291713001979

Wichers, M., Groot, P. C., & Psychosystems, E. G., EWS Group. (2016). Critical Slowing Down as a Personalized Early Warning Signal for Depression. *Psychotherapy and Psychosomatics*, 85, 114-116. doi:10.1159/000441458

Table 1

Frequencies and Descriptive Statistics from Sample A and Sample B

	Sample A (<i>N</i> = 910)	Missing <i>n</i> (%)	Sample B (<i>N</i> = 244)	Missing <i>n</i> (%)
Age (years), <i>M</i> (<i>SD</i>)	33.96 (12.10)	2 (0.22)	33.13 (8.33)	4 (1.64)
Female, <i>n</i> (%)	398 (43.74)	33 (3.63)	126 (51.64)	3 (1.23)
Race and Ethnicity, <i>n</i> (%)*		303 (33.30)		3 (1.23)
Caucasian	363 (39.89)		116 (47.54)	
African-American	124 (13.63)		2 (0.82)	
Asian or Pacific Islander	36 (3.96)		87 (35.66)	
American Indian or Alaska Native	7 (0.77)		1 (0.41)	
Multiracial**	9 (0.99)		14 (5.74)	
Unlisted racial minority	15 (1.65)		0 (0.00)	
Hispanic	53 (5.82)		21 (8.61)	
LSAS total, <i>M</i> (<i>SD</i>)***	78.31 (38.82)	38 (4.18)	88.04 (17.82)	17 (7.00)
LSAS total, post-treatment, <i>M</i> (<i>SD</i>)			51.71 (21.50)	81 (33.20)

Note. LSAS = Leibowitz Social Anxiety Scale.

*For 221 (24.30%) of the participants in Sample A and for 130 (53.30%) of the participants in Sample B, Hispanic ethnicity was assessed as an option when reporting race, rather than assessed separately as part of ethnicity. Thus, we are missing additional racial information on participants who chose to select the Hispanic option when reporting their race. The frequency for Hispanic represents to the frequency of participants who endorsed Hispanic ethnicity plus a racial category in addition to the frequency of participants who endorsed Hispanic ethnicity when it was assessed as a racial category. The frequency for Missing represents the frequency of participants who were missing all racial or ethnic data. We did not include the frequency of participants who reported Hispanic ethnicity but who were missing additional racial information in this estimate. Percentages do not add precisely to 100 due to rounding.

**Multiracial was provided as an option for only 125 (13.70%) of the participants in Sample A.

***The LSAS total score represents the pre-treatment score for Sample B.

Table 2

Multiple Regression Results (Part *rs*) for Centrality Indices

Predictor	Main Analyses (No Correction)				Main Analyses (Corrected)			Added Participants			
	LSAS	SIAS-S	BFNE-S	SDS	LSAS	SIAS-S	BFNE-S	LSAS	SIAS-S	BFNE-S	SDS
Centrality	.48**	.02	.11	.18	.40*	.25	-.16	.44*	.01	-.03	.20
Composite											
Infrequency	-.50**	-.68**	-.65**	-.72***	-.55**	-.54**	-.78***	-.26	-.57**	-.56**	-.55**
Variance	-.31*	-.14	.24	-.15	-.09	-.22	-.02	-.02	-.18	-.13	-.16
(SD)											

Note. These regressions were run using the data that can be found in Table S1 in the supplementary material. Each column heading lists the dependent variable, which in each case is the correlation between change in a node and the change in the measure listed. In each regression, all of the predictors were included. Main Analyses = Initial analysis with Sample A centrality indices. Coefficient under the (No Correction) heading is before correction by removing nodes with excessive SDBetas; coefficient Under the (Corrected) heading (if any) is after correction. The SDS analysis did not require correction. Added Participants = Sample A with participants who did not have generalized social anxiety disorder included. LSAS = Liebowitz Social Anxiety Scale; SIAS-S = Social Interaction Anxiety Scale-Straightforward; BFNE-S = Brief Fear of Negative Evaluation-Straightforward; SDS = Sheehan Disability Scale; Centrality Composite = Combined strength and closeness.

* $p < .05$, ** $p < .01$, *** $p < .001$

Figure 1. Example network demonstrating hypothetical relationships between nodes. The relationships between Nodes 1 have been arranged so that Node 1 has a higher value for strength (i.e., stronger correlations with Nodes 2-6 as indicated by thicker green lines; strength = .81) All other nodes have weaker strength in this example; Node 2 (strength = .54) is the next strongest node in the network.

Figure 2. Network model of LSAS items. The figure displays the network for the first of the five imputed data sets; network models of all five imputations can be found in the supplement. The blue (darker gray) solid lines represent positive relations, whereas the red (lighter gray and dashed) lines represent negative relations between items. Nodes of items numbered 1 through 24 refer to fear or anxiety of a given situation and avoidance of the same situation. Note that item pairs 6 and 20 and 11 and 12 have been combined into a single node due to high correlations between these items pairs. The situations represented are, in brief (i.e., not verbatim): 1 = Public

telephone use, 2 = Small groups, 3 = Eating in public, 4 = Drinking in public, 5 = Talking to authority figures, 6 = Acting, performing, giving a talk, 7 = Going to a party, 8 = Working while observed, 9 = Writing while observed, 10 = Calling a relatively unknown person, 11 = Talking to a relatively unknown person, 12 = Meeting strangers, 13 = Urinating in a public restroom, 14 = Entering a room where others are seated, 15 = Center of attention, 16 = Commenting during a meeting, 17 = Taking test, 18 = Expressing disagreement to relatively unknown person, 19 = Looking relatively unknown person in the eyes, 20 = Report to group, 21 = Asking someone on a date, 22 = Returning goods, 23 = Giving party, 24 = Resisting salesperson.

Data Transparency

Most of the data reported in this manuscript have been previously published or were collected as part of other data collections focusing on other issues. One or more primary publications for each data set are listed in the supplementary material. Although the Liebowitz Social Anxiety Scale was reported on in some of these manuscripts, none of these previous publications focuses on a network analysis in those data; further, no previous manuscript focuses on the entire set of data reported on here.

Supplemental Material

The material in this supplement serves several purposes. First, it provides detail regarding the data sets. Second, it provides more detail regarding analyses and their results. Third, it provides the original Results section from a previous version of the paper, provided as a warning regarding the pitfalls of simply analyzing all of the items on a particular measure.

Note that the data (mostly covariance matrices; raw data cannot be shared) and code is available in the associated OSF project in a zipped file. See: <https://osf.io/nphy7/>

Details on Network Estimation Sample (Sample A)

Participants in Dataset 1 ($n = 97$) and Dataset 2 ($n = 58$) were adult individuals seeking treatment for social anxiety in one of two large, urban, Northeastern cities and their surrounding metropolitan areas. Participants ranged in age from 19 – 66 years ($Mdn = 31.00$ years, $SD = 10.43$) and were primarily male ($n = 92$, 59.40%) and white ($n = 95$, 61.30%). Participants were recruited from primary health care offices and mental health clinics, or were self-referred from media advertisements. A select group of participants in this dataset were recruited as part of a nationally-funded, multi-site, treatment study that enrolled participants from 1995-2001 and examined the combined effects of phenelzine (Nardil) and cognitive-behavioral group therapy (CBGT) for SAD. Participants were assessed using the Anxiety Disorders Interview Schedule for DSM-IV, Lifetime version (ADIS-IV-L; DiNardo, Brown, & Barlow, 1994) or the Structured Clinical Interview for DSM-IV (SCID; First, Spitzer, Gibbon, & Williams, 2002). Graduate students or post-doctoral fellows affiliated with the university clinic conducted the clinical interviews, under the supervision of the clinic director. Some assessments were also conducted by staff psychologists. Most participants in this dataset went on to receive CBGT for SAD and some received treatment as part of the previously described clinical trial. Portions of this data have been published previously (e.g., Blanco et al., 2010).

Participants in Dataset 3 ($n = 149$) were adult individuals recruited as part of a nationally-funded, multi-site, treatment study examining the combined effects of paroxetine (Paxil) and cognitive-behavioral therapy (CBT) for SAD that took place from 2005-2007. Participants ranged in age from 18 - 62 years ($Mdn = 30.00$ years, $SD = 10.88$) and were primarily male ($n = 126$, 60.90%) and white ($n = 72$, 34.80%). Participants were recruited from one of two large, urban, northeastern cities and their surrounding metropolitan areas via referrals from primary health care offices and mental health clinics. Participants also were self-referred from advertisements. Participants were diagnosed with GSAD after undergoing a structured clinical interview, using either the ADIS-IV-L (DiNardo et al., 1994) or the SCID (First et al., 2002). Interviews were conducted by graduate students and post-doctoral fellows at both institutions and were supervised by the primary investigators or affiliated licensed clinical psychologists.

Participants in Dataset 4 ($n = 146$) were adult individuals seeking treatment for SAD, from the years 2002-2016. Participants ranged in age from 18 - 68 years ($Mdn = 26.00$ years, $SD = 10.20$) and were primarily female ($n = 85$, 58.20%) and white ($n = 103$, 70.50%). Participants were referred from primary healthcare clinics, mental health clinics, or via advertisements and were living in a large, urban, northeastern city and surrounding metropolitan area. Participants were

diagnosed with GSAD or SAD using either the ADIS-IV-L or the Anxiety and Related Disorders Interview Schedule for DSM-5, Lifetime version (ADIS-5L; Brown & Barlow, 2014). Graduate students or post-doctoral fellows affiliated with the university adult anxiety clinic conducted the clinical interviews. The clinic director or affiliated clinical psychologist supervised all assessments. Data from this treatment-seeking sample has been published or reported on previously (e.g., Harpole et al., 2015; Rodebaugh et al., 2011), but never in combination with the other samples presented here.

Participants in Dataset 5 ($n = 298$) were adult individuals that received CBGT for SAD at an anxiety clinic over a roughly 5-year period, ranging from 1990-1995. This sample ranged in age from 18 - 64 years ($Mdn = 34.00$ years, $SD = 9.69$) and participants were primarily male ($n = 157$, 52.70%). Participants were referred from primary healthcare clinics, mental health clinics, or via advertisements. A select group of participants were enrolled as part of a nationally-funded, multi-site, treatment study to examine combined treatment for SAD using CBGT and phenelzine. Participants enrolled in the clinical trial were excluded if they met diagnostic criteria for schizophrenia, major depression or bipolar I disorders, alcohol or substance abuse within the last six months, or reported imminent suicidality or any other serious medical condition or organic mental disorder. Participants were also excluded if they had received CBT in the past or psychiatric treatment with MAOIs for social anxiety disorder. All participants were diagnosed with social phobia after undergoing a structured clinical interview using the Anxiety Disorder Interview Schedule – Revised (ADIS-R; DiNardo & Barlow, 1988). Interviews were conducted by graduate students and post-doctoral fellows and were supervised by the primary investigator or an affiliated licensed clinical psychologist. Portions of this dataset have been reported on or published previously (Heimberg et al., 1999).

Participants in Dataset 6 ($n = 37$) were adult individuals recruited as part of a nationally-funded, multi-site, treatment study that enrolled participants beginning in 2001, and examined an individual manualized 16-session treatment protocol for SAD. Participants ranged in age from 18 - 76 years ($Mdn = 35.00$ years, $SD = 14.69$) and were primarily female ($n = 21$, 65.80%) and white ($n = 29$, 78.40%). Participants presented to one of two anxiety specialty clinics and were referred from primary healthcare clinics, mental health clinics, or via advertisements. Participants were living in either a large, urban Northeastern city and surrounding metropolitan area, or a medium-sized, urban, Midwestern city and surrounding metropolitan area. Participants were diagnosed with SAD using either the ADIS-IV-L or the Anxiety Disorders Interview Schedule for DSM-IV (ADIS-IV; Brown & Barlow, 1994) and a significant portion of the participants met criteria for GSAD. Participants were excluded if they met criteria for current substance dependence, current bipolar disorder, current or past psychotic disorder, or if they were imminently suicidal. Graduate students or post-doctoral fellows affiliated with the university anxiety clinics conducted the clinical interviews and they were supervised by clinical psychologists who specialized in the assessment and treatment of social anxiety. Participants were then randomized to an immediate treatment condition or a delayed treatment condition, with treatment beginning after a 20-week waiting period. Data from this sample has been published previously (Ledley et al., 2009).

Participants in Dataset 7 ($n = 75$) were recruited during 2011-2012 from a medium-sized, urban, Midwestern city and surrounding metropolitan area. Participants were adult individuals living in the community and ranged in age from 18 - 64 years ($Mdn = 42.00$ years, $SD = 13.56$). They were primarily female ($n = 54$, 72.00%) and white ($n = 38$, 50.70%). Participants were recruited through flyers posted in community clinics and public areas, a volunteer register, and listing of the study on the laboratory website. Participants completed a brief telephone screen and were excluded if they reported symptoms consistent with current psychosis, mania, substance abuse, or imminent suicidality. Assessments took place within an anxiety laboratory, housed within a medium-sized, urban, Midwestern university and participants were diagnosed with GSAD using the SCID-IV (M. B. First et al., 2002) in combination with the clinician-administered LSAS. Data from this study have been published in multiple studies (e.g., Rodebaugh et al., 2014, in combination with Dataset 8).

Participants in Dataset 8 ($n = 31$) were recruited between 2007 and 2011, from a medium-sized, urban, Midwestern city and surrounding metropolitan area. Participants were adult individuals living in the community and ranged in age from 18 - 63 ($Mdn = 35.00$, $SD = 12.41$). They were primarily female ($n = 17$, 54.80%) and white ($n = 20$, 64.50%). Participants were diagnosed with GSAD using the Mini International Neuropsychiatric Interview (Sheehan et al., 1998) in combination with the clinician-administered LSAS. Graduate students and a clinical psychologist affiliated with the university conducted all clinical interviews, with the graduate students conducting interviews under the supervision of a clinical psychologist.

Participants in Dataset 9 ($n = 19$) were recruited during 2015-2016 from a medium-sized, urban, Midwestern city and surrounding metropolitan area. Participants were adult individuals living in the community, ranged in age from 19 - 62 years ($Mdn = 32.00$ years, $SD = 33.33$), were primarily female ($n = 15$, 78.90%), and were primarily white ($n = 15$, 78.90%). Participants were recruited using a volunteer registry and flyers posted in community mental health clinics, as well as university and community areas. Participants completed a brief telephone screen and were excluded if they reported symptoms consistent with current psychosis, mania, substance abuse, or reported imminent suicidality. Assessments took place within an anxiety laboratory, housed within a medium-sized, urban, Midwestern university and participants were diagnosed with SAD using the SCID-V (First, Williams, Karg, & Spitzer, 2015) in combination with the clinician-administered LSAS. Graduate students completed the clinical interviews, under supervision of the laboratory director. Data from this study have not yet been published and recruitment is ongoing.

There were 197 participants across Sample A who did not receive a GSAD diagnosis. Participants were mostly women ($n = 125$; 63.50%) and the mean age was 35.10 years ($SD = 12.27$; 18-69 years). A majority of participants were White (67.50%), and 20.80% were Black, 3.60% were Asian or Pacific Islander, 3.00% identified as Multiracial, and 0.50% identified as American Indian or Alaska Native. There were 3.00% participants who identified as Hispanic.

Details for Treatment Datasets (Sample B)

Datasets 10-12. Participants in Dataset 10 ($n = 76$) were recruited over a 5-year period, 2006-2011, and ranged in age from 21 - 53 years ($Mdn = 33.00$ years, $SD = 8.96$). Participants were primarily male ($n = 39$, 51.30%) and white ($n = 43$, 56.60%), with considerable diversity as nearly one-third of the sample were Asian or Pacific Islander ($n = 21$, 27.60%). Participants were recruited from clinician referrals and community listings and completed a brief telephone screening. Participants were excluded if they reported a history of medical disorders, head trauma, or if they met criteria for select comorbid psychiatric disorders. Participants were included if they met criteria for generalized anxiety disorder, agoraphobia without a history of panic attacks, or specific phobia, in addition to GSAD. Participants were also excluded if they reported current mental health treatment, or if they reported history of receiving CBT. Assessments were conducted by clinical psychologists and took place within an anxiety laboratory, affiliated with a large, West-coast university. Participants were diagnosed with GSAD using the ADIS-IV-L (DiNardo et al., 1994). After completing the assessment, participants were randomized to either treatment with individual CBT or a waitlist control. Participants received treatment at no cost and completed assessments at pre-treatment, post-treatment, and at a 1-year follow-up. Data from this study have been published in several papers (e.g., Goldin et al., 2012).

Participants in Dataset 11 ($n = 54$) were recruited over a 3-year period, 2007-2010, and ranged in age from 23 - 53 years ($Mdn = 31.00$ years, $SD = 8.42$). They were primarily female ($n = 28$, 51.90%) and were primarily white ($n = 22$, 40.70%) or Asian or Pacific Islander ($n = 23$, 42.60%). Participants were recruited from clinician referrals and community listings and completed a brief telephone screening. Participants were excluded if they reported a history of medical disorders, head trauma, or select comorbid psychiatric disorders. Participants were allowed to participate if they met criteria for generalized anxiety disorder, major depressive disorder, obsessive-compulsive disorder, agoraphobia without a history of panic attacks, or specific phobia, in addition to GSAD. Participants were also excluded if they reported current mental health treatment, history of participation in a mindfulness-based stress reduction (MBSR) course, or reported engagement in a regular meditation or exercise regimen. Assessments were conducted by clinical psychologists and took place within an anxiety laboratory affiliated with a large West-coast university. Participants were diagnosed with GSAD using the ADIS-IV-L (DiNardo et al., 1994) and, after completing the assessment, participants were randomized to one of two treatment groups – MBSR or aerobic exercise. Participants received treatment at no cost and completed assessments at pre-treatment, post-treatment, and at a 3-month follow-up. Data from this study have been published in several papers (e.g., Jazaieri, Goldin, Werner, Ziv, & Gross, 2012).

Participants in Dataset 12 ($n = 114$) were recruited over a 3-year period, 2011 - 2014, and ranged in age from 21 - 56 years ($Mdn = 31.41$ years, $SD = 7.92$). Participants were primarily female ($n = 62$, 54.40%) and white ($n = 51$, 44.70%), although a significant number of individuals were Asian or Pacific Islander ($n = 43$, 37.70%). Participants were recruited from clinician referrals and community listings and completed a brief telephone screening. Participants were excluded if they reported a history of mental health treatment in the past year, a history of CBT treatment for an anxiety disorder in the past two years, or if they reported experience with meditation or

MBSR. Participants were also excluded if they reported a history of cardiovascular disorders or neurological disorders, or comorbid psychiatric disorders. However, participants were included if they met criteria for generalized anxiety disorder, specific phobia, obsessive compulsive disorder, panic disorder, major depressive disorder, or dysthymia, in addition to GSAD. Assessments took place within an anxiety laboratory affiliated with a large West-coast university. Participants were diagnosed with GSAD using the ADIS-IV-L (DiNardo et al., 1994) and graduate students completed the clinical interviews, under supervision of the laboratory director. After completing the assessment, participants were randomized to one of two treatment groups – MBSR or CBGT for SAD – or a waitlist control. Participants treatment at no cost and completed assessments at post-treatment and at 3-month intervals during the 1-year follow-up. Participants received monetary compensation at the 1-year follow-up. Data from this study have been published in several papers (e.g., Goldin et al., 2016).

Details for Analytic Procedure

Random forest imputation. Random forest imputation is a non-parametric imputation method that was used to impute unplanned missing values from LSAS scores for the initial network model. Random forest uses machine learning to build decision trees based on underlying patterns in data. Decision trees are used to build a predictive model that is used to estimate and impute missing values. The R package *missforest* (Stekhoven & Buhlmann, 2012), was used to create five imputed datasets. We created non-identical imputations by setting a new random seed before beginning each imputation process. A means of formally handling multiple imputation in network analyses is not yet available, so we examined findings across all five datasets to determine whether findings were consistent. Imputed values were reviewed from each dataset to ensure they did not exceed the minimum or maximum for LSAS variables in the original dataset.

Network estimation. Network estimation and network stability tests were conducted using R packages *qgraph* (version 1.4.1; Epskamp et al., 2012) and *bootnet* (version 0.3; Epskamp, Borsboom, et al., 2017), respectively. An undirected regularized partial correlation network was estimated (Epskamp & Fried, in press), resulting in edges that can be interpreted as partial correlation coefficients or conditional dependence relations (an association between two items controlling for all other associations among items). Regularization shrinks all edge weights and sets many to zero, which leads to a sparse (or parsimonious) network and avoids estimating spurious relationships. The covariance matrix was estimated using the *cor_auto* procedure, which detected and treated appropriately the ordinal items (i.e., computing polychoric correlations). Visualizations of the 5 imputations of the networks are presented in Figure S1 and Figure S2, and matrices that can be used to reproduce many of our results is provided in a supplementary attachment.

Network stability tests were conducted as recommended by Epskamp et al. (2017) to determine: (a) the stability of edge weights, by estimating 95% confidence intervals around all edge weights via a bootstrapping procedure; and (b) the accuracy of centrality coefficients using a subsetting routine, resulting in a correlation-stability (CS) coefficient that should be above .25 to indicate moderate stability, but preferably above .5 for strong stability. Visualizations of these tests are presented as Figure S3 and Figure S4, respectively.

Estimation of treatment models. LSAS item-level data in Sample B was missing for some of the sample due to an error in the data collection software ($n = 9$); in addition, many participants either dropped out of treatment or provided only partial data on some measures. Because analysis focused on properties of nodes, missing data were handled by estimating correlations pairwise based on all available data. As noted in the main text, at least 134 participants were available to estimate each correlation.

Alternative network models. We also constructed network models using the following specifications: (a) Sample B used instead of Sample A; (b) Ising model with Sample A and data dichotomized such that each item was approximately 50% endorsed; (c) Sample A with participants without GSAD included. For (a), missing data (7% of cases) was ignored and cases were deleted listwise. Otherwise, MI was used and the first imputation results are provided in the main text. Visualizations and matrices that can be used to reproduce many of our findings is provided in an additional supplemental zipped file.

Follow-up tests using statistics from these models are summarized in the main text; relevant raw data are provided in Table S1.

Table S1.

Node-Level Data used in Follow-up Analyses

Node	Close	Strength	Between	Infreq	SD	I- Close	I- Strength	I- Between	Pre Tx Closeness	Pre Tx Strength	Pre Tx Between	Pre Tx Infreq	Pre Tx SD	All Close	All Strength	All Between	All Infreq	All SD	Change Other LSAS	Change SIAS	Change BFNE	Change SDS
1	.00	.77	10.00	506.00	1.76	.01	3.14	7.00	.00	.57	22.00	61.00	1.90	.00	.75	2.00	677.00	1.70	.47	.20	.19	.25
2	.00	.92	12.00	81.00	1.64	.01	4.30	22.00	.00	.87	11.00	13.00	1.54	.00	.96	9.00	235.00	1.91	.58	.19	.21	.36
3	.00	1.10	8.00	535.00	1.83	.01	4.23	18.00	.00	1.18	23.00	128.00	1.51	.00	1.03	9.00	709.00	1.76	.57	.19	.10	.16
4	.00	.94	10.00	575.00	1.72	.01	4.39	15.00	.00	.93	16.00	109.00	1.79	.00	.94	15.00	754.00	1.64	.52	.12	.11	.18
5	.00	.77	10.00	106.00	1.73	.01	3.07	4.00	.00	.58	7.00	10.00	1.48	.00	.80	10.00	224.00	1.88	.61	.29	.26	.31
6/20	.00	1.00	10.00	9.00	2.78	.01	3.27	25.00	.00	.82	5.00	1.00	1.03	.00	1.01	10.00	80.00	2.01	.46	.34	.29	.35
7	.00	1.08	16.00	98.00	1.83	.01	4.32	30.00	.00	.88	22.00	5.00	1.49	.00	1.08	20.00	243.00	2.10	.56	.23	.22	.36
8	.00	.96	21.00	170.00	1.84	.01	3.62	13.00	.00	.96	10.00	15.00	1.68	.00	1.00	23.00	304.00	1.96	.58	.23	.24	.17
9	.00	1.03	19.00	539.00	1.85	.01	3.58	22.00	.00	.68	3.00	64.00	1.96	.00	.93	16.00	707.00	1.78	.46	.02	.01	.11
10	.00	1.01	12.00	199.00	1.88	.01	3.86	23.00	.00	.77	32.00	8.00	1.44	.00	.99	14.00	329.00	1.95	.64	.29	.29	.29
11/12	.00	1.30	30.00	72.00	3.45	.01	5.04	29.00	.00	.85	27.00	1.00	1.37	.00	1.26	32.00	191.00	2.04	.60	.21	.20	.31
13	.00	.82	2.00	645.00	1.64	.01	3.43	6.00	.00	.57	5.00	159.00	1.46	.00	.77	2.00	813.00	1.58	.36	.17	.16	.06
14	.00	1.01	26.00	176.00	1.81	.01	4.22	22.00	.00	.88	26.00	39.00	1.66	.00	1.00	21.00	322.00	1.95	.58	.32	.22	.28
15	.00	1.03	31.00	38.00	1.60	.01	3.98	13.00	.00	.70	4.00	2.00	1.11	.00	1.13	22.00	157.00	2.05	.64	.38	.40	.46
16	.00	1.00	5.00	36.00	1.57	.01	2.70	3.00	.00	1.15	15.00	3.00	1.14	.00	1.04	4.00	154.00	2.00	.62	.32	.27	.37
17	.00	.70	2.00	488.00	1.73	.01	2.02	2.00	.00	.71	8.00	106.00	1.68	.00	.64	3.00	624.00	1.71	.41	.18	.16	.15
18	.00	1.01	13.00	93.00	1.77	.01	3.14	2.00	.00	.77	8.00	2.00	1.36	.00	.99	13.00	193.00	1.95	.57	.37	.31	.32
19	.00	.85	3.00	207.00	1.96	.01	3.55	3.00	.00	.59	.00	23.00	1.68	.00	.88	6.00	349.00	2.04	.57	.18	.24	.23
21	.00	.72	1.00	83.00	1.89	.0040	4.93	.0012	.00	.29	.00	12.00	1.61	.00	.72	2.00	146.00	2.07	.48	.25	.23	.26
22	.00	.94	5.00	416.00	1.89	.0038	5.76	NA	.00	.71	6.00	64.00	1.67	.00	.93	6.00	576.00	1.86	.42	.27	.24	.23
23	.00	.90	15.00	141.00	2.05	.0033	4.91	.0013	.00	.80	24.00	11.00	1.62	.00	.94	14.00	275.00	2.27	.54	.23	.21	.23
24	.00	1.00	10.00	329.00	2.06	.0047	5.61	.0013	.00	.66	15.00	43.00	1.81	.00	.89	10.00	465.00	2.06	.52	.19	.17	.13

Note. The above are the raw data for the analyses presented in follow-up. The table is designed to be copied and pasted into another form (e.g., spreadsheet, data file); for visual parsing, please zoom in on a large monitor. Nodes are the units rather than participants.

Formation of the nodes is described in the main text; in brief, each node consists of the fear and avoidance items for the LSAS situations corresponding to the numbers given. Close = Closeness from Sample A; Strength = Strength from Sample A; Between = Betweenness from Sample A. Infreq = Number of participants with the lowest value for that node in Sample A; SD = Variance for that node in Sample A, in terms of standard deviation; I- Close = Closeness from Sample A, computed using an Ising network; I-Strength = Strength from Sample A, computed using an Ising network; I-Between = Betweenness from Sample A, computed using an Ising network; Pre Tx Closeness = Closeness computed with Sample B; Pre Tx Strength = Strength computed with Sample B; Pre Tx Betweenness = Betweenness computed with Sample B; Pre Tx Infreq = Number of participants with the lowest value for that node in Sample B; Pre Tx SD = Variance for that node in Sample B, in terms of standard deviation; All Close = Closeness from Sample A with added participants without generalized social anxiety disorder; All Strength = Strength from Sample A with added participants without generalized social anxiety disorder; All Between = Betweenness from Sample A with added participants without generalized social anxiety disorder. All Infreq = Number of participants with the lowest value for that node in Sample A with added participants without generalized social anxiety disorder; All SD = Variance for that node in Sample A, in terms of standard deviation with added participants without generalized social anxiety disorder; Other LSAS = Correlation of change in this item with change in the rest of the Liebowitz Social Anxiety Scale items, computed via the treatment dataset; Change SIAS = Correlation of change in this item with change in the Straightforward Social Interaction Anxiety Scale, computed via the treatment dataset; Change BFNE = Correlation of change in this item with change in the Straightforward Brief Fear of Negative Evaluation, computed via the treatment dataset; Change SDS = Correlation of change in this item with change in the Sheehan Disability Scale, computed via the treatment dataset.

Table S2

Correlations Between Centrality Indices

	Close (A)	Strength (A)	Between (A)	Close (B)	Strength (B)	Between (B)	Close (I)	Strength (I)	Between (I)	Close (A-All)	Strength (A-All)	Between (A-All)
Close (A)	.21											
Strength (A)	.56**	.59										
Between (A)	.89***	.66**	.28									
Close (B)	.43*	.41	.30	.00								
Strength (B)	.17	.61**	.26	.50*	.21							
Between (B)	.37	.47*	.35	.78***	.49*	.00						
Close (I)	.68**	.58**	.61**	.56**	.23	.50*	.21					
Strength (I)	.53*	.75***	.64**	.57**	.45*	.54**	.81***	.28				
Between (I)	.50*	.73***	.60**	.49*	.42	.52*	.78***	.77***	.13			
Close (A- All)	.90***	.59**	.84***	.28	.25	.38	.67**	.54*	.57**	.21		
Strength (A- All)	.65**	.94***	.72**	.37	.62**	.40	.62**	.72***	.66**	.71***	.67	

Between (A-All)	.75***	.76***	.92***	.23	.35	.40	.63**	.71***	.67**	.83***	.78***	.36
--------------------	--------	--------	--------	-----	-----	-----	-------	--------	-------	--------	--------	-----

Note. Notation in parentheses indicates the source of the centrality index. (A) = Sample A, main analysis; (B) = Sample B; (I) = Sample A, Ising network; (A-All) = Sample A, participants without generalized social anxiety disorder added. This table can be produced by running correlations based on the values given in Table S1. Centrality stability indices are provided on the diagonal for the first imputation (except for Sample B, for which imputations were not used); these are not correlations or standard reliability indices, but higher values are preferred, as described in the main text.

* $p < .05$; ** $p < .01$; *** $p < .001$

Figure S1. Network models of five imputations of the Liebowitz Social Anxiety Scale items. The figure displays the network for each of the five imputed data sets. The blue lines represent positive relations, whereas the red lines represent negative relations between items. Stronger relationships are characterized by thicker, bolder lines. Nodes of items numbered 1 through 24 refer to fear or anxiety of a given situation and avoidance of the same situation; note that two nodes are composites of two situations. Placement of nodes was conducted by the estimation algorithm; please see Figure S2 for the same figures with the nodes constrained to their average position.

Figure S2. Network models of five imputations of the Liebowitz Social Anxiety Scale items. Unlike with Figure S1, the node positions have been constrained to be an average of all five imputations. Note that, as a result, imputations are visibly very similar; however, subtle differences in imputations do exist.

Figure S3. Visualizations of the edge weights from the first imputed dataset for the primary analyses. Each point represents the value of a node pair edge and grey lines indicate the bootstrapped confidence interval. Edges are ordered by node (e.g., first point is item1-item2 edge, second point is item1-item3 edge).

Figure S3. Graphical depiction of the stability of centrality indices for Imputation 1 for the primary analyses. The percentage of the original data sampled is on the x-axis and the correlation of node centrality with the original sample is on the y-axis. Higher correlations (shallower slopes) indicate more stable indices.

Original Material Prior to Peer Review

For the purpose of transparency and completeness, we provide below the material originally included for peer review as a part of the main text. Importantly, peer review uncovered problems in estimation for the network, *so these results are not to be trusted*. Please see the main text for details. We provide it in part to illustrate that, as discussed in the main text, simply conducting a network analysis on a set of symptoms without careful consideration of whether and how items should be combined may lead to misleading results. Supplemental material for the original analyses are not included, but can be obtained from the first author.

Data Analytic Procedure

Here we focus on the overall plan for the analyses (more detailed information on each aspect of the data analytic procedure is available in the supplemental materials). We first created five multiple imputation data sets for the LSAS network model dataset to handle sporadic missing data, using random forest imputation. Because there is no best standard for how to deal with multiply imputed data in network analyses, we focused on consistent findings across all five data sets. Network estimation and network stability tests were conducted in accordance with current standards using R packages qgraph (version 1.4.1; Epskamp, Cramer, Waldorp, Schmittmann, & Borsboom, 2012) and bootnet (version 0.3; Epskamp, Borsboom, & Fried, 2017), respectively. An undirected regularized partial correlation network was estimated, resulting in edges that can be interpreted as partial correlation coefficients (an association between two items controlling for all other associations among items). Regularization ensures that the estimated network structure balances sensitivity with specificity, and leads to a sparse network structure that avoids obtaining spurious edges (Epskamp & Fried, in press).

We then used bootstrapping (via bootnet) to select the items from the pool of 48 LSAS items that were reliably more central than all others (i.e., were the most central in each dataset and remained most central in smaller subsamples according to bootnet); there happened to be four such items, which we refer to as the highly central (HC) items. We used a similar procedure to select four least central (LC) items for the purposes of comparison. To ensure that LC items were indeed less central than HC items, we used a centrality difference test in bootnet to compute bootstrapped confidence intervals around the difference between bootstrapped centrality index means for the four HC items and the four corresponding LC items. Nodes were said to be significantly different from each other if the confidence intervals resulting from the difference test did not contain 0 ($\alpha = .05$). Finally, we examined item properties, including level of endorsement and variance, to determine whether any superficial item properties might be related to centrality.

We then turned to the treatment data. We created HC and LC composites at pre- and post-treatment by summing the items identified from the network sample. The remaining LSAS items (i.e., those included neither in the HC nor the LC item sets) were summed to create an LSAS index of medium-centrality items (LSAS-MC). We used a structural equation modeling approach in the treatment data, using Mplus version 7.4 (Muthén & Muthén, 1998-2015) and the MLR estimator, in part because this allowed the handling of sporadic missing data. In addition,

this approach allowed a latent estimation of change via a latent change model. We could thus test whether change in HC items predicted change in MC items above and beyond change in LC items. Finally, we tested whether change in HC items was a better predictor of change in overall social anxiety severity as measured by the SIAS-S, BFNE-S, and SDS. The intent was to examine whether HC items showed an enhanced ability to predict change in social anxiety severity outside of the LSAS network as well as within the LSAS network. Mplus code for the primary latent change model (depicted in Figure 3) is included in supplemental material.

Results

Network Estimation

We first estimated the overall network structure for each of the five data imputations. Figures S5 and S6 display the results for the imputed data sets. The results presented below were across all of the imputed datasets, and not simply the first one. We examined the centrality indices strength, closeness, and betweenness and determined that we should not continue to examine betweenness because its stability coefficient of 0.2 (in all five imputed datasets) was below the recommended minimum cut-off of .25 (Epskamp, Borsboom, & Fried, 2016). Coefficients for strength and closeness were better, with values in all five datasets .4 or higher, well above the suggested minimum cut-off of .25 (Epskamp, Borsboom, & Fried, 2016). We thus focused on the strength and closeness of the items.

We then used bootnet to determine which of the highly central items showed stability such that they would remain the highest in centrality in a data set half the size. Four items for strength and closeness fit these criteria across all imputations and comprised the set of HC items: Fear and avoidance of acting, performing, or giving a talk in front of an audience (LSAS items 6F and 6A, respectively); and fear and avoidance of giving a report to a group (LSAS items 20F and 20A, respectively). Notably, the four items identified as having the highest closeness and strength values were still the highest at 40% of the current sample size, suggesting that similar results are reasonably likely to be found in a sample as small as $N = 364$ (i.e., 40% of the current sample).

Across imputations, four items consistently had the least centrality (LC) in the network according to strength and closeness metrics: Fear and avoidance of drinking with others in public places (items 4F and 4A, respectively) and fear and avoidance of resisting a high pressure salesperson (items 24F and 24A, respectively). All individual HC items were significantly more central on indices of strength and closeness than all individual LC items according to bootstrapped confidence intervals. Notably, however, HC items and LC items did not differ significantly from all of the other items.

Correlations between centrality and item properties. In regard to item properties, we found moderately large negative correlations between the standard deviations of the nodes and their centrality indices (for closeness, all r s across imputations rounded to $-.36$; for strength, r s ranged from $-.306$ to $-.319$), indicating that the HC items actually had less variation in responses. However, frequency of item endorsement also had moderately large negative correlations with centrality indices, whether assessed by the frequency with which participants reported the lowest or the two lowest response options (r s from $-.38$ to $-.48$). That is, HC items were less likely to be

responded to with one of the two lowest response options, suggesting that the HC items might simply assess those symptoms that were more commonly responded to as clearly present. We tested whether this issue might explain results in follow-up analyses. Conversely, we found moderately large positive correlations between the means of the nodes and their centrality indices (r s from .40 to .43 across imputations for closeness; r s from .46 to .47 across imputations for strength).

Treatment Response Prediction

Within-LSAS-network prediction. The latent change score model fit excellently (CFI = 1.00, TLI = 1.07, RMSEA = .00). The model, along with unstandardized estimates, is presented in Figure 3 (full output for the model is presented in the supplementary material). LSAS-MC change was predicted by both HC ($b^* = .46, p < .001$) and LC items ($b^* = .50, p < .001$). Contrary to our hypothesis, the HC items did not predict more strongly than the LC according to a chi-square test: $\chi^2(1) = 0.30, p = .587$. In other words, centrality was unrelated to how strongly change in a set of items predicted change in the other items.

Outside LSAS-network prediction of social anxiety severity. The model included the SIAS-S, BFNE-S, and SDS loading on a pre- and post-treatment latent severity factor. Consistent with hypothesis, post-treatment severity was predicted by pre-treatment severity ($b^* = 0.49, p < .001$) and change in HC items ($b^* = 0.46, p < .001$), but not change in LC items ($b^* = 0.05, p = .585$). However, this model fit poorly (CFI = .86, TLI = .76, RMSEA = .11), which was traced to the severity measures not showing a good fit in a two-factor (pre and post) model (CFI = .82, TLI = .66, RMSEA = .19). To ensure that the results for HC items were not due purely to this poor fit, an additional model tested the three severity measures as separate (but correlated) variables instead of a single factor; this model had close to excellent fit (CFI = .95, TLI = .93, RMSEA = .06). Eight participants were excluded from this analysis because they had a missing value for one or more of the pre-treatment severity measures. Again, supporting our hypotheses, change in the HC items predicted each of the three severity measures separately ($ps < .001$) whereas change in the LC items did not predict ($ps > .579$).

Follow-up Tests

Rationale. Hypotheses were not supported within the LSAS network, but were supported outside of the network. A stronger role for HC items outside the LSAS network than within it was not hypothesized. We conducted several follow-up analyses to attempt to determine why we might have obtained this puzzling result. During these tests we also addressed the possibility that these results might merely have been an artifact of item properties (primarily rates of endorsement). Although we intended to test item properties as a competing explanation, the specific form of these follow-up tests was not a priori. However, they do directly parallel the model provided by Robinaugh and colleagues (2016). We used Spearman correlations throughout because infrequency of endorsement had a non-normal distribution. In all of the following tests, we analyze node properties, such that the sample is the group of 48 LSAS items, with their properties drawn from the data sets as appropriate.

Correspondence of centrality indices between cross-sectional and treatment data.

One plausible explanation for our findings is that the cross-sectional centrality estimates do not relate to centrality in the treatment data. For example, the treatment data was a much smaller sample, and our results with the network sample suggest limited confidence in precision of centrality estimates with samples smaller than about 400 participants. We therefore estimated a network in the pre-treatment data in the same manner as our estimation in the cross-sectional data. Strength in the pre-treatment data was not significantly related to either strength ($r_s = .14$, $p = .341$) or closeness ($r_s = .15$, $p = .318$) from the cross-sectional data. However, closeness in the pre-treatment data was strongly related to both closeness ($r_s = .71$, $p < .001$) and strength ($r_s = .64$, $p < .001$) in the cross-sectional data. It should thus not be surprising that the HC items were among the items with the highest closeness in the pre-treatment data (with some additional items having equivalent levels of closeness). In contrast, only one of the HC items (fear of situation 6, giving a public speech) was among the items with the highest strength in the pre-treatment sample. The inconsistent correspondence between the centrality indices across datasets (cross-sectional versus treatment) was therefore examined as one possible explanation of the results, below. Notably, the count of how many participants responded to that item with a 0 or 1 rating (i.e., infrequency of endorsement) was related inversely to all of these indices ($|r_s| > .37$, $p_s < .02$; please note that infrequency predicted inversely such that more highly endorsed items had higher centrality) except for strength in the pre-treatment data ($r_s = -.12$, $p = .413$).

Analysis by node within LSAS network. Although our a priori analyses reported above were similar in intent to those of Robinaugh and colleagues (2016), we did not analyze across all nodes as they did (instead, above, we focused only on the HC and LC nodes). In the following analyses, as Robinaugh et al. (2016) did, we examined correlations between centrality indices (from the initial, cross-sectional network) and the strength of relationship between node change and change in the rest of the symptom network (here, the LSAS items across treatment). That is, first, we examined the correlation between the change of each symptom with the change of all other symptoms. For example, it could be the case that one symptom reduces across participants in a way that has nothing to do with change in the other symptoms: In such a case, the correlation would be low. Recalling the example of Figure 1, the low-centrality nodes should display a low correlation if their low centrality means that they have a low causal impact on the rest of the network. In contrast, a symptom with high centrality might change in a way across participants that closely tracks the change in other symptoms: In that case, the correlation would be high. Again, this should be true if high centrality means a node has a high causal impact on other nodes. We obtained the resulting correlation coefficient for each symptom. Second, we investigated the correlation between that coefficient and the centrality indices for each symptom, the idea being that items with higher centrality indices should also have higher correlation coefficients. This procedure therefore allowed a test of whether strength and closeness, as continuous variables, correlated with the strength of the connection between node change and change in the remainder of the network. These tests could then be extended to tests of whether strength and closeness were useful in predicting how node change would correlate with change in the other severity measures.

The raw data used in these analyses [can be obtained from the first author]. Results from the following analyses [can be obtained from the first author]. Consistent with our analyses above, neither strength nor closeness was associated with strength of connection between node and the rest of the network ($r_{s,s} < .20$, $ps > .10$). However, infrequency of endorsement was associated with strength of connection ($r_s = -.33$, $p = .024$), such that items that were more frequently endorsed showed a stronger connection with the rest of the network. Figure 3 shows the scatterplots and trend lines that correspond to these correlations.

As noted above, it is also possible to examine the same issue relative to strength and closeness in the pre-treatment data. Pre-treatment strength was not related to change in the rest of the network ($r_s = .10$, $p = .496$), but pre-treatment closeness was ($r_s = .37$, $p = .009$). However, when pre-treatment closeness and infrequency were both entered as predictors in a regression, closeness did not predict (part $r = .07$, $p = .627$) but infrequency did (part $r = -.30$, $p = .038$).

Analysis by node for other symptom measures. The tests across node for centrality indices could also be conducted in regard to how strongly change in node was correlated with change in the other severity measures. Again, we started with examination of the cross-sectional network indices. As would be expected given the results of our a priori tests, both strength and closeness had some significant associations with correlations with change in the three severity measures. More specifically, both strength and closeness trended toward predicting associations between change in node and change in the SIAS-S ($r_{s,s} > .23$, $ps < .10$), significantly predicted regarding the SDS ($r_{s,s} > .28$, $ps < .04$), but showed no tendency for the BFNE-S ($r_{s,s} < .23$, $ps > .10$). In contrast, infrequency strongly predicted these associations for all three measures ($|r_{s,s}| > .59$, $ps < .001$). As would be expected given these correlations, in follow-up regressions infrequency always predicted uniquely, whereas strength and closeness did not, whether each was compared to infrequency alone or both were compared to infrequency together.

These analyses were repeated using the pre-treatment centrality indices instead, with substantively similar results. Closeness predicted associations between change in node and change in the SIAS-S ($r_s = .29$, $p = .049$) and SDS ($r_s = .53$, $p < .001$), but not the BFNE ($r_s = .24$, $p = .106$), whereas strength did not predict any of these ($ps > .12$). In regressions, closeness did not predict above and beyond infrequency.

Additional tests. In [can be obtained from the first author], we present the by-node data for the above analyses such that other researchers can replicate our results and run additional tests. We also report, in [can be obtained from the first author], a large number of additional tests, all of which generally support the observation that infrequency of endorsement accounted for all effects that might be expected of centrality. The sole exception was strength derived from an additional network in which HC items were dropped completely, but even in this case, infrequency was a more robust predictor.

Figure Error5. Network models of five imputations of the Liebowitz Social Anxiety Scale items. The figure displays the network for each of the five imputed data sets. The green lines represent positive relations, whereas the red lines represent negative relations between items. Nodes of items numbered 1F through 24F refer to fear or anxiety of a given situation (blue) and nodes of items 1A through 24A refer to avoidance of the same situation (red). Placement of nodes was conducted by the estimation algorithm; please see Error6 for the same figures with the nodes constrained to their average position. Note that the most strongly related dyads are, in each case, the anxiety and avoidance for a specific situation.

Figure Error6. Network models of five imputations of the Liebowitz Social Anxiety Scale items. Unlike with Figure Error5, the node positions have been constrained to be an average of all five imputations. Note that, as a result, imputations are visibly very similar, although subtle differences in imputations do exist if the numerical values are examined.