How compliance with behavioural measures during the initial phase of a pandemic develops over time: a longitudinal COVID-19 study

Monique Chambon^{1,2*}, Jonas Dalege³, Denny Borsboom², Lourens J. Waldorp², Han L.J. Van der Maas² & Frenk van Harreveld^{1,2}

¹ National Institute for Public Health and the Environment (RIVM), the Netherlands

² Department of Psychology, University of Amsterdam, the Netherlands

³ Santa Fe Institute, United States of America

*Corresponding author information: Monique Chambon, National Institute for Public Health and the Environment (RIVM), P.O. Box 1, 3720 BA, Bilthoven, The Netherlands. E-mail: monique.chambon@rivm.nl

**This paper has been published in The British Journal of Social Psychology.
The Version of Record is available online at: <http://doi.org/10.1111/bjso.12572>**

Supplementary information can be found below the manuscript.

Data availability statement

Data and R code is made openly available in OSF at <https://osf.io/qu7p2/>.

Acknowledgements

We would like to thank Sacha Epskamp for contributing to the analyses. This research was funded by The Dutch Research Council (NWO grant 440.20.019). Jonas Dalege's work was supported by an EU Horizon 2020 Marie Curie Global Fellowship (no. 889682).

Conflict of interest

The author(s) declared no potential conflicts of interest.

Abstract

In this longitudinal research, we adopt a complexity approach to examine temporal dynamics of variables related to compliance with behavioural measures during the COVID-19 pandemic. Dutch participants ($N = 2399$) completed surveys with COVID-19-related variables five times over a period of 10 weeks (23 April – 30 June 2020). With this data, we estimated within-person COVID-19 attitude networks containing a broad set of psychological variables and their relations. These networks display variables' predictive effects over time between measurements and contemporaneous effects during measurements. Results show (1) bidirectional effects between multiple variables relevant for compliance, forming potential feedback loops, and (2) a positive reinforcing structure between compliance, support for behavioural measures, involvement in the pandemic, and vaccination intention. These results can explain why levels of these variables decreased throughout the course of the study. The reinforcing structure points towards potentially amplifying effects of interventions on these variables, and might inform processes of polarization. We conclude that adopting a complexity approach might contribute to understanding protective behaviour in the initial phase of pandemics by combining different theoretical models and modelling bidirectional effects between variables. Future research could build upon this research by studying causality with interventions and including additional variables in the networks.

Keywords: COVID-19 pandemic, compliance with behavioural measures, broad attitude networks, longitudinal, within-person temporal effects

During the initial phase of a pandemic, a vaccine is often absent and compliance with behavioural measures such as social distancing and isolation are often considered of pivotal importance to curtail the spread of the virus. Research during the COVID-19 pandemic shows that the majority of people reported compliance with behavioural measures but also indicates ample room for improvement (Hensel et al., 2022). Thus, greater insight into determinants of compliance is crucially important for improving our ability to understand, predict and improve compliance during epidemic outbreaks (Betsch et al., 2020).

The literature on compliance with behavioural guidelines during the COVID-19 pandemic is already immense, with numerous studies pointing towards factors contributing to compliance or the lack thereof, from a range of theoretical perspectives, such as the Theory of Planned Behaviour (Ajzen, 1991; Gibson et al., 2021), the Health Belief Model (Clark et al., 2020; Rosenstock, 1974) and Moral Foundations Theory (Chan, 2021; Graham et al., 2013). This body of work has generated important insight into the psychological underpinnings of compliance in the context of the COVID-19 pandemic, but theoretical frameworks like the ones just mentioned are inherently restrictive because of the limited number of variables they focus on. An influential perspective on compliance during pandemics that is less restrictive is that by Bish and Michie (2010), who provide an extensive review of the literature on this topic and discuss a wide range of determinants that play a role in predicting health behaviour during a pandemic. As such, most empirical work on this topic takes the perspective of a particular model and the specific elements in it, while the influential perspective by Bish and Michie takes a broader perspective but is theoretical in nature.

Our current aim is to deepen our understanding of compliance with behavioural measures during the initial phase of a pandemic by taking a complexity approach. We do so in accordance with the conceptual frameworks provided by Bish and Michie (2010). We add to this existing literature by conducting empirical research that includes not only the constructs of these conceptual frameworks, but also additional variables that more recent research showed to be relevant in the context of pandemics. Moreover, we employ a longitudinal design that provides insight into how relations between variables develop over time and thus indications of causal effects. Such longitudinal designs seem to be most lacking in the literature on pandemics thus far (Bish & Michie, 2010).

Psychological variables related to compliance during pandemics

Bish and Michie (2010) propose frameworks on health behaviour during a pandemic with different attitudinal and demographic determinants depending on the type of protective behaviour. In our current research we focussed on preventive behaviours (e.g., hygiene behaviours and vaccination) and avoidant behaviours (e.g., social distancing and working from home) since these resonate with the behavioural measures as advised by Dutch authorities during the initial phase of the pandemic. According to Bish and Michie (2010), attitudinal variables that are associated with preventive and avoidant behaviours include perceived severity, susceptibility, costs, efficacy and self-efficacy but

also social norms, cues to action, knowledge, trust in authorities and state anxiety. Moreover, they argue most of these variables are covered by three well-known models within psychology that have been used to explain relevant behaviour, namely the Theory of Planned Behaviour (TPB; Ajzen, 1991), the Health Belief Model (HBM; Rosenstock, 1974) and the Protection Motivation Theory (PMT; Rogers, 1975). The TPB predicts behaviour through intention, with the intention being influenced by attitudes, behavioural control, and behavioural norms. The HBM explains health-related behaviour through the perception of one's susceptibility and severity of the disease, benefits and barriers of the behaviour, self-efficacy, and cues to action. PMT seeks to understand behaviour through threat appraisal (i.e., severity of and vulnerability to the disease) and coping appraisal (i.e., response and self-efficacy). Research during the COVID-19 pandemic also associates components of these models with compliance with behavioural measures, namely attitudes (Bogg & Milad, 2020), perceived control and severity (Li et al., 2020), risk perception (Schneider et al., 2021), social norms (Tunçgenç et al., 2021) and self-efficacy (Shahnazi et al., 2020).

Bish and Michie (2010) argue that various psychological factors not included in these models are also relevant in the context of pandemics, such as knowledge about the disease, trust in authorities and state anxiety. The latter suggests that adverse effects of the pandemic on people's mental health should also be considered when aiming to understand behaviour in the context of pandemics. Recent research during the COVID-19 pandemic indeed associates compliance with behavioural measures with mental health, namely fear of the virus (Harper et al., 2021) and stress (Lieberoth et al., 2021). Additionally, potential consequences of behavioural measures such as changes in lifestyle and loneliness are associated with adverse mental and physical health effects (Blom et al., 2021; Leigh-Hunt et al., 2017). These findings suggest it is important to take changes in (mental) health into account when investigating compliance during a pandemic.

Thus, research points towards a wide range of interrelated variables relevant to health behaviour during a pandemic. Chambon, Dalege, Elberse, et al. (2022) corroborated this in the context of the COVID-19 pandemic by adopting a complex psychological systems approach to provide an overview of variables related to preventive behaviours in a study conducted in both the United Kingdom and the Netherlands. This resulted in a descriptive account of how these variables are interrelated and insight into differences between the countries. The study could not provide insight into the directions of relations between variables and their temporal effects due to its cross-sectional design (Chambon, Dalege, Elberse, et al., 2022).

Prior research suggests that temporal effects are an important subject of investigation because psychological factors associated with compliance with behavioural measures during pandemics can show substantive changes over time (Ibuka et al., 2010; van der Weerd et al., 2011). Likewise, research during the COVID-19 pandemic showed fluctuations in risk perception (Schneider et al., 2021), psychological distress (Daly & Robinson, 2021) and well-being (Wang et al., 2021). Such temporal changes in psychological factors associated with compliance highlight the importance of

longitudinal research into compliance during pandemics. Accordingly, the current research contributes to the existing literature by examining the temporal effects of variables and their interrelations. We focus on variables that can be expected to fluctuate during the initial phase of a pandemic and not relatively stable variables such as demographics, personality traits and political preferences. We recognize that such variables can be important for compliance as well and may moderate the relations between variables included in this study, but their effects fall outside of the scope of the present research.

To summarize, prior research showed that a broad set of variables can affect compliance with behavioural measures during pandemics. Also, there is a gap in the literature regarding longitudinal research that provides insight into the temporal effects of variables relevant to protective behaviours during a pandemic. Consequently, parsimonious theoretical models might not provide sufficient insight into the complexity of compliance during pandemics. Nevertheless, most research into compliance focussed on a rather limited set of variables examined in a cross-sectional research design. The current research therefore adopts a complexity approach in which a more diverse set of variables was included and employs a longitudinal research design. Importantly, the aim of this research is not to present a comprehensive overview or an exhaustive set of variables relevant for compliance, but to demonstrate that compliance during pandemics can be explained by a broad set of variables that extends beyond the traditional models (i.e., complexity approach) and to provide insight into directions of effects between these variables (i.e., temporal effects).

Complexity approach

We utilize a complexity approach to gain insight into how psychological variables organize and interact over time. A promising means to this goal is psychological network analysis, which (1) enables exploration of data with many variables of different types, (2) provides insight into patterns of unique relations that remain after controlling for effects of other variables (i.e., pairwise conditional dependencies) and (3) presents these statistical associations in powerful visualizations (Borsboom et al., 2021). A theoretically underpinned model of attitudes that adopts a complexity approach through network analysis is the Causal Attitude Network (CAN) model (Dalege et al., 2016). This model conceptualizes attitudes as complex systems of interacting cognitive, affective and behavioural factors. This systemic approach models properties of attitude dynamics, including their relation to behaviour. Because this approach naturally integrates many different factors relevant to behaviour, we adopt an extension of the CAN model to improve our understanding of compliance behaviour.

Attitude networks in the CAN model consist of nodes that represent cognitive, affective, and behavioural components of attitudes. Links between nodes represent interactions between these components. These links, also known as edges, represent linear relations between nodes and can be positive, indicating an excitatory relation, or negative, indicating an inhibitory relation. The strength of interactions between nodes can vary, resulting in a weighted network. In the CAN model, the

overall state of the system (i.e., the pattern of feelings, thoughts, and behaviours that characterize a person) arises out of these interactions between attitude components (Dalege et al., 2016). Figure 1 shows a simplified and hypothetical network of the attitude towards physical distancing. Through patterns of interaction, some node pairs will be more strongly aligned than others. These patterns can be identified using partial correlations, where the relation between two given nodes is conditional on all other nodes in the network (see Epskamp, 2020a; van Borkulo et al., 2014). Thus, via statistical analyses of data, empirical estimates of attitude network models can be obtained (Dalege et al., 2017). Representations of these models can yield important information to advance our understanding of behavioural variables. For example, the centrality of specific nodes may provide information on the relative importance of different nodes in the network organization, that is, nodes central to the network have more and stronger relations with other nodes. Ideally, network models are estimated on longitudinal data, as such data can harbour important clues about the causal organization of the network. This results in network models that include directions of effects between variables (see Figure 1).

Fig. 1. Simplified and hypothetical attitude network of the attitude towards physical distancing. Variables are displayed as nodes (circles) and linear relations between variables as edges (lines between circles). The arrows indicate predictive effects between nodes from one measurement to the next. The network consists of a behavioural element ('Keep distance from others'), a cognitive evaluation ('Physical distance is effective to prevent spread') and three affective evaluations ('Fear of disease', 'Feeling lonely' and 'Missing physical contact'). The edge width indicates the strength of the relations. In this example, feeling lonely is more strongly affected by keeping distance from others than by perceiving physical distance as effective to prevent the spread, as indicated by the different width of these edges in the network. Furthermore, relations can be unidirectional (e.g., between keeping distance from others and feeling lonely) or bidirectional (e.g., between feeling lonely and missing physical contact). Bidirectional effects can consist of edges with different strengths.

To summarize, the present research addresses gaps in the literature regarding complexity and temporal effects by shedding light on temporal dynamics of the system of psychological factors (i.e., broad attitudes networks) related to compliance with behavioural measures during the COVID-19 pandemic. Our means in doing so is conducting network analysis with longitudinal data gathered during the pandemic: participants completed a survey with COVID-19 related variables five times over a period of approximately 10 weeks (23 April – 30 June 2020).

Method

Participants and design

The Ethics Review Board of the University of Amsterdam approved this study (2020-SP-12194). Participants were recruited via a research panel (Ipsos). The sample of the first wave was representative of the Dutch population in terms of age, gender and residential area. Table 1 provides sample and descriptive information for the longitudinal sample in the column on Wave 5 (valid $N = 2399$), consisting of respondents that participated throughout the study and finished all surveys. This sample size far exceeds the advised sample size of approximately 500 participants for accurate network estimation of a moderately sized network, that is, an accurate representation of the true underlying network (Epskamp, 2017; van Borkulo et al., 2014).

Table 1. Methodological and sample information for each separate wave

	Measure	Wave 1	Wave 2	Wave 3	Wave 4	Wave 5
Sample formation						
Start data collection (2020)		23 rd April	13 th May	27 th May	10 th June	24 th June
End data collection (2020)		5 th May	18 th May	2 nd June	16 th June	30 th June
Failed attention check ¹	n	519	118	41	24	8
Sample (including missing values)	n	6219	4953	3754	2822	2449
Dropout ²	n (%)		1,266 (20.4%)	1,199 (24.2%)	932 (24.8%)	373 (11.4%)
Missing values ³	n (%)	126 (2.0%)	89 (1.0%)	70 (1.9%)	58 (2.1%)	50 (2.0%)
Valid N		6093	4864	3684	2764	2399
Demographics						
Valid N						
Gender	% female	51.4%	50.6%	50.1%	49.1%	49.4%
Age	Range (years)	16-89	16 – 89	18 – 89	18 – 89	18 – 89
	M (SD)	49.32 (16.72)	51.20 (16.32)	51.99 (16.30)	53.42 (16.06)	53.69 (15.84)
Education	% primary or secondary education	55.3%				54.6%
	% higher education	44.7%				45.4%
Illness	% confirmed	30.6%				34.9%
Smoking	% confirmed	17%				15.9%

¹ Two attention checks were included in each wave to ensure data quality. Failing both attention checks led to exclusion from further data collection and analysis in all subsequent waves.

² Comparing networks of the longitudinal sample and dropouts showed very few significant differences in relations between variables. More information is provided in the supplemental online material (S1.1).

³ The answer option 'I prefer not to answer' was included in items on education, illness and smoking, and was treated as a missing value. Participants with missing values for one or more nodes were deleted from network analysis, which is reasonable given the small number of missing values.

Measures

Data collection was conducted with a Dutch online survey. We first identified the most important constructs in the literature (see introduction) and then developed a survey with items based on these constructs (see online supplemental material S1.2 for a complete overview). Analysis for node construction was conducted with the largest and most diverse sample available, namely wave 1, as this wave included all participants that completed the first survey (i.e., featured no dropout). Psychological variables were constructed as nodes through either predetermined combinations of items or based on results of a reduction technique aimed at identifying components in data (i.e., Principal Axis Factoring [PAF]). Detailed descriptions of how items were combined to form nodes, including PAF results, are provided in Supplement S1.3. Resulting nodes are presented in Table 2.

Table 2. Nodes (psychological variables) based on items in the survey, including item examples and answer scales. Mean scores of items were calculated for nodes that were based on multiple items (except for Risk Perception as explained in text).

Node (items per node)	Examples of items per node (/ in the same text line means separate item in survey)	Scale
Compliance (5)	I comply with the corona measures. / Keep 1.5 metres away from others. / Wash your hands regularly with water and soap.	1 (<i>I do not display this behaviour more</i>) to 7 (<i>I display this behaviour much more now</i>)
Risk Perception (2)	How likely (/ severe) do you believe it is you will get infected with the coronavirus within the next year?	1 (<i>Extremely unlikely</i>) to 7 (<i>Extremely likely</i>)
Health Risk (2)	For me personally (/ my family and friends), I consider the health risk of an infection with the coronavirus...	1 (<i>Extremely small</i>) to 7 (<i>Extremely severe</i>)
Economic Consequences (2)	For me personally (/ my family and friends), I consider the economic consequences of the corona pandemic...	1 (<i>Extremely small</i>) to 7 (<i>Extremely severe</i>)
Self-exempting Beliefs (2)	I think I am already immune (protected) against the coronavirus. / I will not get infected with the coronavirus because I never get the seasonal flu either.	1 (<i>Strongly disagree</i>) to 7 (<i>Strongly agree</i>)
Negative Affect (8)	The corona pandemic is making me (feel)... (e.g., sad / frustrated / overwhelmed)	1 (<i>Strongly disagree</i>) to 7 (<i>Strongly agree</i>)
Compassion (1)	The corona pandemic is making me feel compassion.	1 (<i>Strongly disagree</i>) to 7 (<i>Strongly agree</i>)
Worries Virus^ (4)	I worry about.. (e.g., getting infected / infecting others / the health care system overloading)	1 (<i>Do not worry at all</i>) to 7 (<i>Worry a lot</i>)
Worries Measures^ (6)	I worry about.. (e.g., a recession / limited access to food / getting lonely)	1 (<i>Do not worry at all</i>) to 7 (<i>Worry a lot</i>)
Vaccination Intention^ (1)	If a vaccine becomes available, I would get it.	1 (<i>Strongly disagree</i>) to 7 (<i>Strongly agree</i>)
Measures Support (7)	I find the corona measures.. (Unnecessary-Necessary / Unfair-Fair / Unacceptable-Acceptable)	1 (Negative option) to 7 (Positive option)
Measures Ease (2)	I find the corona measures.. (Difficult-Easy / Unpleasant-Pleasant)	1 (Negative option) to 7 (Positive option)
Social Norm (2)	I think the majority of people (/ find it important that people) comply with the corona measures.	1 (<i>Strongly disagree</i>) to 7 (<i>Strongly agree</i>)
Control Infection (2)	For me personally (/my family and friends), avoiding an infection with the coronavirus in the current situation is..	1 (<i>Extremely difficult</i>) to 7 (<i>Extremely easy</i>)
Self-efficacy (1)	I know how to protect myself from the coronavirus.	1 (<i>Strongly disagree</i>) to 7 (<i>Strongly agree</i>)
Involvement (3)	How important is the topic of corona pandemic to you? / To what extent does the news about the corona pandemic have your attention?	1 (<i>Not at all</i>) to 7 (<i>Very much</i>)
Perceived Knowledge (1)	How much knowledge do you think you have about the corona pandemic?	1 (<i>Very little</i>) to 7 (<i>Very much</i>)

Trust (4)	I trust.. (e.g., the authorities / health care professionals) .. during the corona pandemic.	1 (<i>Strongly disagree</i>) to 7 (<i>Strongly agree</i>)
General Health (1)	In general, how would you rate your health?	1 (<i>Very poor</i>) to 7 (<i>Very good</i>)
Health change Physical (1)	How would you rate your physical health now as compared to before the corona pandemic?	-3 (<i>Much worse</i>) to 3 (<i>Much better</i>)
Health change Mental (1)	How would you rate your mental health now as compared to before the corona pandemic?	-3 (<i>Much worse</i>) to 3 (<i>Much better</i>)
Healthy Lifestyle (3)	I've been (exercising / eating / sleeping) in the past two weeks, compared to before the corona pandemic...	-3 (<i>Much [less healthy / less / worse]</i>) to 3 (<i>Much [healthier / more / better]</i>)
Mental Well-being ¹ (7)	I've been feeling useful. / I've been dealing with problems well. / I've been feeling close to other people.	1 (<i>Never</i>) to 5 (<i>Always</i>)
Loneliness ² (6)	There are plenty of people I can lean on when I have problems. / There are enough people I feel close to.	1 (<i>Not at all</i>) to 5 (<i>Very much</i>)
Complaints ³ (anxiety [6]; depressive [6]; somatic [6])	To what extent did you experience.. (e.g., feeling suddenly scared / fearful / tense; e.g., feeling lonely / hopeless / a loss of interest; e.g., difficulty breathing / numbness / feeling weak) during the past two weeks?	1 (<i>Not at all</i>) to 5 (<i>Very much</i>)

Note: The sections of the survey that referred to 'the corona measures' contained the following explanatory text 'By this, we mean the recommendations to prevent the spread of the coronavirus and thus prevent overloading the healthcare system, for example stay at home as much as possible, keep 1.5 metres of distance from others and wash your hands regularly with soap and water.'

¹ WHO Regional Office for Europe (2020), Multiple items (e.g., worries, vaccination intention) were adopted from the WHO protocol for COVID-19 monitoring

² Tennant et al. (2007), raw scores were converted to metric scores as required for the (S)WEMWBS

³ de Jong Gierveld and van Tilburg (2008), answer scale formally ranges from 1 (No!) to 5 (Yes!). Please note that answer scales were adjusted, and a 2-week time frame was specified for the current study.

⁴ Derogatis (2001), answer scale formally ranges from 0 (*Not at all*) to 4 (*Extremely*). Please note that we changed the BSI-18 answer scales and specified a 2-week time frame, possibly invalidating clinical interpretation of scores. Therefore, we use the term *complaints* instead of *symptoms*.

Compliance. Protective behaviours as recommended by the Dutch national government (Government of the Netherlands, 2020) provided the operationalization of compliance with behavioural measures in this study. We measured to what degree participants adopted protective behaviours as advised to the general public (i.e., physical distancing and hygiene behaviours), providing a self-reported measure for compliance. Items forming this node changed during the study because of changes in recommended protective behaviours.¹

Attitudes. Attitudes were measured in line with the multi-component model, namely consisting of cognitive, affective and behavioural attitude elements (Eagly & Chaiken, 1993). Items forming the cognitive attitudinal nodes Risk Perception, Health Risk ($r_{sb} = .59$)² and Economic Consequences ($r_{sb} = .75$) were based on prior research into psychological COVID-19 networks (Chambon, Dalege,

¹ The node Compliance consisted of five items in the first wave ($\alpha = .75$), and one of these items was adapted for the second wave. A sixth item was added in the third wave, leading to Compliance consisting of six items from the third wave onwards. Thus, this node contains behavioural items that applied during the administration of the survey.

² The scale reliability, although for some nodes lower than one would normally prefer, was interpreted as sufficient since our aim was to measure evaluations and not to design reliable measuring scales (Dalege et al., 2016).

Elberse, et al., 2022). Risk perception was measured with two items on perceptions of likelihood and severity of getting infected with the coronavirus. Risk Perception was the product of these two items (Wolff et al., 2019). Health Risk represents perceived health consequences due to an infection, and the node Economic Consequences contains items measuring expected consequences of the pandemic for the economy. Self-exempting Beliefs ($r_{sb} = .57$) tapped into one's conviction of not being susceptible to an infection with the coronavirus.

Items measuring affect surrounding the pandemic resulted in the nodes Negative Affect (e.g., anxiety, anger, and confusion; $a = .89$) and Compassion (single item). Items on worries also resulted in two nodes: Worries Virus ($a = .73$) and Worries Measures ($a = .67$), reflecting worries about events during the pandemic resulting directly from the coronavirus (e.g., losing someone they love), and events resulting from measures taken because of the virus (e.g., overloading the health care system), respectively.

Behavioural attitudinal nodes consisted of Vaccination Intention (single item), Measures Support ($a = .90$) and Measures Ease ($r_{sb} = .68$). Vaccination Intention represents one's intention to get a COVID-19 vaccine once available. The items covering attitudes towards the behavioural measures, consisting of both general items and semantic differential scale items, resulted in two nodes: Measures Support reflects support for the behavioural measures advised to prevent the spread of the coronavirus, and Measures Ease represents participants' perceived ease of complying with these measures.

Additional psychological factors. Social Norm ($r_{sb} = .76$) reflects prescriptive (what other people think one should do) and descriptive (what other people do) social norms regarding compliance with behavioural measures. Items on perceived control formed the nodes Control Infection ($r_{sb} = .61$) and Self-efficacy (single item). The former node represents to what degree participants felt able to avoid getting infected with the coronavirus. The latter node represents whether participants perceived to know how to protect themselves from the coronavirus. Involvement ($a = .84$) was comprised of items on how actively involved participants perceived themselves to be in the COVID-19 pandemic (e.g., watching the news). This node also represents a dimension of attitude strength. A single item measuring perceived knowledge about the pandemic formed the node Perceived Knowledge. The node Trust ($a = .86$) reflects general trust in the four actors relevant to the corona pandemic in the Netherlands: the authorities, the Dutch National Institute for Public Health and the Environment (RIVM), health care professionals and science.

Physical and mental health nodes. General Health reflects an overall self-reported evaluation of participants' health. Single items measuring participants' evaluation of their physical and mental health during the survey, compared to before the pandemic, formed the nodes Health change Physical and Health change Mental, respectively. Lifestyle items measured an improvement or deterioration in

exercise, diet, and sleep, resulting in Healthy Lifestyle ($\alpha = .49$).³ The sum of items from the Short Warwick Edinburgh Mental Well-being Scale ([S]WEMWBS; Tennant et al., 2007) formed Mental Well-being ($\alpha = .84$). The shortened version of the loneliness scale (de Jong Gierveld & van Tilburg, 2008) resulted in the node Loneliness ($\alpha = .76$). The Brief Symptom Inventory 18 (BSI-18; Derogatis, 2001) was adopted to measure psychological complaints. This scale contains 18 items measuring three areas of psychological distress (i.e., anxiety, depressive, and somatic symptoms). Three nodes were formed based on the BSI-18 items: Anxiety Complaints ($\alpha = .92$), Depressive Complaints ($\alpha = .90$) and Somatic Complaints ($\alpha = .84$).

Individual differences. The first survey also included questions on individual differences, namely demographics (i.e., age/gender/education) and health (i.e., illness/smoking). These variables provided descriptive information about the sample and were excluded from temporal analyses given that they were measured just once, in the first wave. The first survey also included items on relatively stable personality aspects (i.e., consideration of future consequences, resilience, and coping) that provided input for a different paper on interventions. Readers are referred to Chambon, Dalege, Waldorp, et al. (2022) for more information on (results of) these measures.

Procedure

Participants that subscribed to Ipsos' research panel received an invitation to participate via e-mail. Participants were informed about and asked to commit to the longitudinal research design beforehand. Only participants who finished the survey received invitations for subsequent waves. They received compensation in the form of points that can be spent at web shops.

Interventions were also included at the beginning of the third and last wave, to which participants were randomly assigned (see Supplement S1.4). These interventions are not the focus of the current paper and will be presented in a different paper (Chambon, Dalege, Waldorp, et al., 2022).

Network analysis

Networks were estimated with the *dlvm1* function (Lag-1 dynamic latent variable model for panel data) in *psychonetrics* (Epskamp, 2020a, 2020b). The R script is made available as a supplement (S1.5). We estimated two networks that provide information on average within-person effects on a population level: *temporal* and *contemporaneous* (Epskamp, 2020a) psychological networks.⁴ Temporal networks contain edges representing whether one node predicts other nodes in

³ See previous footnote.

⁴ The analysis normally also provides a between-person network, but estimating this network was not successful as the edges in this network had too high standard errors to be interpretable. Confidence intervals that are too wide can result in unstable (unreliable) edges; thus, this network was omitted in the current paper.

the next measurement, while controlling for all other nodes. These temporal effects are calculated by regressing each variable on all variables (including itself) on the previous measurement (i.e., lag-1) and therefore require repeated measures (i.e., waves). The resulting partial correlations are indicative of directed predictive effects, depicted by edges with arrows in the network (from wave t-1 to wave t). The weight of edges indicates their effect size and can be interpreted as one would interpret regular partial correlations. Edges between two nodes represent either a causal effect or are the result of a third (unknown) underlying cause. Contemporaneous networks are based on residuals of the temporal network (i.e., variance and covariance that cannot be explained by the modelled temporal effects). These edges can be interpreted as partial correlations between nodes in the same measurement after controlling for all other nodes in the same measurement and the previous measurement (i.e., temporal effects). These partial correlations thus represent undirected associations, depicted by edges without arrows in the network. Again, edge weights indicate effect size and can be interpreted as regular partial correlations. Readers are referred to Epskamp et al. (2018) for additional information on the type of networks presented here.

Centrality measures facilitate the interpretation of visualized networks. The centrality measure ‘strength’ is among the most commonly used centrality measure for psychological networks (Borsboom et al., 2021) and provides information on the conditional association between a node and other nodes in the network. It is calculated by the sum of the absolute edge weights of relations a specific node has with connected nodes. Two different types of strength can be distinguished in temporal networks, due to directed edges: InStrength for edges directed towards that specific node and OutStrength for edges directed from that specific node to other nodes. In other words, we distinguish between effects *to* a node and *from* a node.

Results

This section presents COVID-19 broad attitude networks containing temporal dynamics based on the longitudinal survey. In the current study, temporal effects indicate which nodes predict other nodes over a time frame of 2 - 3 weeks (i.e., from one survey of a wave to the next), whereas contemporaneous effects indicate which nodes predict other nodes within the same survey of a wave.

Preliminary analysis

Descriptive statistics of nodes are provided in Supplement S2.1, together with results of repeated measures ANOVA analyses to examine the effects of time on variables. Results indicated a significant change in all nodes over time. Furthermore, Figure 2 provides the context at the time of

data collection by showing a general timeline of the pandemic in the Netherlands (see Supplement S3 for underlying data).

Evaluation of the overall fit of the *dlvm1* model in which we included all edges in the COVID-19 broad attitude networks showed excellent fit (see Supplement S2.2). Generally speaking, confidence intervals of edge weights were not wide, indicating stable (reliable) edge estimates. The weights of edges discussed below are reported in parentheses (see Supplement S2.3 – S2.5 for a complete overview of edges in the COVID-19 networks).

Temporal COVID-19 network

Figure 3a shows temporal effects, standardized to partial directed correlations, in the COVID-19 broad attitude network. Edges indicate nodes' predictive value for the next measurement after controlling for all other nodes. Table 3 provides edge weights of all edges in the temporal network. The width of node borders represents the degree to which nodes were influenced by the same node in the previous wave (i.e., autoregression). Thicker node borders indicate more stable nodes. The most stable nodes in the temporal COVID-19 network were Vaccination Intention (.41), Compliance (.37), Measures Support (.25) and Healthy Lifestyle (.24).

Fig. 3. The temporal COVID-19 network. a) Visualized network – see Table 3 for edges weights. Nodes represent measured psychological factors. Edges represent relations between two nodes after controlling for other nodes in the network, with their weight indicating the strength of relations. Blue edges represent positive (excitatory) relations and red edges represent negative (inhibitory) relations. Edges with weights below .05 are omitted to facilitate readability. The network has a cut-off value of .10, meaning edges with weights below that value are depicted with similar width and colour density; b) Standardized strength measure. This measure represents direct effects of a specific node on the network and is calculated by the sum of the absolute edge weights, with InStrength for edges affecting that node and OutStrength for edges from that node affecting other nodes.

Table 3. Edge weights of partial directed correlations from the temporal COVID-19 network. Read rows (first column) as node from which the edge originates.

To \ From		Perceived knowledge	Compassion	Self-efficacy	Vaccine Intention	Health General	Health change Physical	Health change Mental	Risk Perception	Health Risk	Self-exempting Beliefs	Economic Consequences	Involvement	Negative Affect	Worries Virus	Worries Measures	Trust	Social norm	Measures Support	Measures Ease	Control Infection	Compliance	Somatic Complaints	Healthy Lifestyle	Depressive Complaints	Anxiety Complaints	Loneliness	Mental Well-being
Perceived knowledge		.09		.02	-.01	.01	.01	-.01	-.01	-.02	-.02	-.04	-.02	-.02	-.03	-.02	.01	-.04	.01	.03	.02	-.01		.02	-.02	-.03	-.01	.01
Compassion		-.01	.10			.01		-.02		.01	.01	.02	.05	.05	.03	.01	.01	.04	.01	-.02	.02	.04	-.02	-.01	.01	.01		-.02
Self-efficacy		.02	-.02	.08	-.02	-.02	-.01	.01	-.02		.04	-.01	-.03	.02	-.01	-.01	.01	-.01	-.01	.01	.01	-.03	-.01			.01	.01	-.01
Vaccine Intention			.05	.02	.41	-.01	.01		.04	.04	-.01	.02	.06	.03	.06	.01	.02	.04	.08	.01		.06	.01	-.01	.01			.01
Health General		-.01	.01	-.01	.01	.15	.03		.01	.01		.03	.02	.02	.01	.02	.02	.03		-.02	-.01			.01	.01		.02	
Health change Physical		.02	.02	.04	.02	.05	.14	.08	.01	.01	.02	.02	.02	.03	.02		-.01	.02	.04		.01	.02	-.02	.04	-.01			-.01
Health change Mental		-.04		-.01		.01	.06	.10		.02			-.01	-.02	.01	-.01			-.01	-.01		-.01	.03	.01	-.05	-.03	-.02	.02
Risk Perception		.01	.02	-.01		-.01	.02	-.01	.11	.04	-.02		.03	.01	.02	.02		.02		-.02	-.02	.01	.03	.03			.02	
Health Risk		-.03		.01	-.02	-.01	-.01	.02	.04	.09	-.02	.02	-.02	-.02	.02	-.02	.02	-.03	.02	.03	-.01		-.01	-.02		-.01	-.02	
Self-exempting Beliefs		-.02	.01	.02	.03	-.01	.01	.01	-.03	-.02	.06	.01	.04			.01	.01	.02	.02	.01	.02	.06	-.02	-.01	-.03	-.03		.01
Economic Consequences				.03	.01	.03	.02	-.01		.03	.01	.10		.01	.02	.03	-.01	-.01		-.01	.02	.02	-.02	-.02	-.02			-.03
Involvement		.03	.09		.07	.01	.01	-.04	.06	.03	.01	.02	.21	.06	.11	.07		.10	.09	-.01		.10	.01	-.02	.05	.05	.02	-.04
Negative Affect		-.03	.04		-.01	.02	.01	-.01			.02		.03	.19	.03	.05	-.04	-.02	-.06	-.05	-.02	-.03		.02	.03	.01	.02	-.01
Worries Virus			.07		.02	-.02	-.01		.06	.07		.04	.08	.08	.13	.03	.03	.03	.04		-.01	.06	.01	-.02	.04	.03	.04	-.01
Worries Measures		.01	.01	-.02		-.02	-.05		.03	-.02	.01	.06	.02	.02	.04	.12	.01	.01		-.02		.01	.01	-.02	.02		.05	-.02
Trust		.02	-.01		-.03	.02	-.02	.02	.01		-.01	-.03	-.02	-.03		-.01	.11	-.03	.04		.01	.01	-.02	.01				-.02
Social norm			.07	.02	.03	-.01	.01	-.01	.04	.02	.02	.04	.06	.06	.08	.05	.01	.16	.04	-.02	-.03	.07	.01	-.01	.02	.03	.02	-.02
Measures Support		-.01	.05	.02	.07	.04	.02		.04	.04	-.02	.03	.08	.02	.08	.03	.11	.08	.25	.06	-.02	.13	.06		.02	.03	.01	.01
Measures Ease		.03	-.01	.02	-.01	-.02	.02	.02		.02	-.01	-.02		-.04	-.02	-.03	.01	-.01	.02	.16	.01		-.01		-.02	-.03	-.01	.01
Control Infection				.02	.01	.01	-.02		-.03	-.02	.02	.01	-.01	-.01	-.01				-.03	-.01	.04	-.03	-.03		.01	.01		.03
Compliance			.07	.03	.06		.01	-.01	.02	.02		.04	.08	.03	.07	.06	.06	.12	.11		.04	.37	.01	.01	.03	.03	.04	-.01
Somatic Complaints		-.01	-.02	-.01	.01	-.03	-.03	.03	.02	.01	.02		-.01	.01	.01	.01	-.02		.02		-.02	-.01	.21	-.04	.03	.04	.02	-.01
Healthy Lifestyle		.02	.02	.02		.02	.07	.03	.01		.01	-.02		.02	-.01	-.01		-.01	-.01		.02	-.02	.24	.04	.01	.03	.02	
Depressive Complaints			.01	.03	.01	.02		-.07	.01	.01	-.02	.02	.03	.07	.06	.05		-.01	.01	-.01		-.01	.02		.18	.07	.09	-.05
Anxiety Complaints			.01	-.03	.03	-.03	.01	-.03	.02	.02	-.02	.02	.06	.04	.04	.04	.02	.03	.03		-.03	.04	.05	-.02	.10	.20	.02	-.05
Loneliness		-.02	-.01	-.01			.03	-.01	.01	.01	-.01	.02	.01	.06	.02	.07	-.02			-.03	-.01	.03	.03	.01	.06	.03	.14	-.04
Mental Well-being			-.03		.01	-.01		.01	.01	-.02	.01	-.01	-.03	-.03	-.02	-.03	.01	-.05		-.02	.01	-.03	.02		-.02		-.02	.09

Regarding nodes related to compliance, results showed bidirectional effects between Compliance and multiple other variables in the network after controlling for the effects of every other node in the network. Compliance was predicted by Measures Support (.13), Involvement (.10), Social Norm (.07), Vaccination Intention (.06) and Worries Virus (.06). Also, Compliance predicted Social Norm (.12), Measures Support (.11), Involvement (.08), Worries Virus (.07) and Vaccination Intention (.06). The former indicates that change in compliance with behavioural measures was predicted by the degree to which one supports the behavioural measures, was mentally involved in the COVID-19 pandemic, perceived social norms regarding compliance, intended to get vaccinated against COVID-19, and worried because of the virus. The latter indicates that change in compliance with behavioural measures was predictive of one's perceived social norms regarding compliance, support for the behavioural measures, mental involvement in the pandemic, worries because of the virus, and intention to get vaccinated against COVID-19. Such positive feedback loops indicate that compliance decreases over time if one of the aforementioned five variables decreases, and, vice versa, that these five variables decrease if compliance decreases. Figure 3a shows that the temporal network contains more of these patterns.

Moreover, results suggest bidirectional effects between three of the variables that showed bidirectional effects with compliance. That is, the nodes Measures Support, Involvement and Vaccination Intention had bidirectional relations among each other. Such a pattern of relations between Compliance, Measures Support, Involvement and Vaccination Intention suggests a reinforcing structure between these nodes.

Another interesting pattern in the temporal network was found between Depressive Complaints, Anxiety Complaints and Loneliness. Depressive Complaints predicted Anxiety Complaints (.07) and Loneliness (.09), and Anxiety Complaints and Loneliness predicted Depressive Complaints (.10 and .06, respectively). However, edges between Anxiety Complaints and Loneliness were relatively small (i.e., from Anxiety Complaints to Loneliness .02; from Loneliness to Anxiety Complaints .03). This pattern suggests a relatively central role for Depressive Complaints in this triangle with Anxiety Complaints and Loneliness.

Node centrality. Figure 3b shows the standardized centrality measure 'strength' for the temporal COVID-19 network (value is provided in text in parentheses). Results showed that Involvement, Compliance, Worries Virus and Social Norm had a central role in the COVID-19 network by both affecting and being affected by other nodes (OutStrength 2.48, 1.79, 1.05, 1.05; InStrength 1.87, 1.75, 1.72, 1.21, respectively). Interestingly, Measures Support predominantly affected other nodes (OutStrength 2.26) and was affected to a lesser extent (InStrength 0.97). Salient discrepancies between InStrength and OutStrength measures were found for Negative Affect, which was predominantly affected by other nodes in the COVID-19 network (InStrength 1.69; OutStrength 0.08) and Worries Measures, which was mostly affected by other nodes (InStrength 0.97; OutStrength - 0.22). Finally, nodes relevant to mental health were of relatively low strength, suggesting that these

nodes had a limited impact on the network. A table with node strength values for each node is provided in Supplement S2.4.

Contemporaneous COVID-19 network

Figure 4 shows (a) contemporaneous effects in the COVID-19 broad attitude network (left) and (b) the standardized centrality measure ‘strength’ for the contemporaneous COVID-19 network (right). Results concerning edges related to Compliance are comparable to the temporal level: the strongest edges with Compliance are Measures Support (.22), Social Norm (.13) and Involvement (.09). These edges indicate that when someone reports compliance, they are also likely to report support for the measures, social norms and involvement in the pandemic. This, together with the effects in the temporal network (see Figure 3a), implies that dynamics concerning compliance with behavioural measures are comparable between measurements over time (i.e., temporal) and within measurements (i.e., contemporaneous). Strength measures for the node Compliance differed between the COVID-19 broad attitude networks: this node played a more central role in the temporal network (InStrength 1.75; OutStrength 1.79) than in the contemporaneous network (0.32). This indicates that, relative to other nodes, compliance with behavioural measures is more predictive of and predicted by other nodes in the network in the next wave than that it was strongly connected with other nodes in the same survey.

Fig. 4. The contemporaneous COVID-19 network. See Fig. 3. for interpretation guidelines. a) Visualized network. Positive (negative) edges indicate that people who reported higher scores on one variable also reported higher (lower) scores on the other variable. See S2.4 for the corresponding edge weights table; b) Standardized centrality measures.

Furthermore, strength measures of the contemporaneous COVID-19 network showed that several variables with higher strength in the temporal network also played a central role here, namely Measures Support (2.16), Negative Affect (1.96), Worries Virus (1.56) and Involvement (1.36). This implies that these nodes are important not only for the COVID-19 broad attitude network over time between measurements (i.e., the nodes are predictive of and/or predicted by many other nodes in the network) but also within a measurement wave (i.e., these nodes are related to many other nodes in the same survey). Interestingly, the node Social Norm had relatively high strength on the temporal level (InStrength 1.21; OutStrength 1.05) but less on the contemporaneous level (-0.35). This suggests that social norms are more predictive of other nodes in the COVID-19 broad attitude network over time than that this node was strongly connected with other nodes in the same survey.

Finally, we conducted correlational analyses to examine coherence between the temporal and contemporaneous networks. Results showed a moderate to large correlation between the temporal and contemporaneous COVID-19 networks ($r = .47$, $z = 0.51$; see Supplement S2.6 for plot). This indicates a relation between edges representing effects over time between measurements (i.e., waves) and edges representing effects during measurement (i.e., within a survey).

Discussion

The current research explored the temporal dynamics of psychological factors related to compliance with behavioural measures during the initial phase of the COVID-19 pandemic. Results of this high-powered longitudinal study showed that nodes in the broad attitude networks were highly interconnected. This indicates that all included variables provided substantive information on the dynamics of adopting protective behaviours during a pandemic, justifying a complexity approach. Several insights can be drawn from this study.

Three interesting patterns were observed in the temporal network. First, results show a series of bidirectional effects between variables in the temporal network relevant for compliance with behavioural measures during the COVID-19 pandemic. We found that the degree to which people support behavioural measures and were mentally involved in the pandemic were most important for predicting compliance over time after controlling for every other variable in the network. This is in line with previous research showing relations between support for measures and compliance (Chambon, Dalege, Elberse, et al., 2022; van Rooij et al., 2020). Interestingly, the current study also found effects the other way around, namely that compliance predicted support for the behavioural measures and involvement in the pandemic over time. Similar results were obtained for vaccination intention and social norms: not only do these variables predict compliance with behavioural measures over time, but compliance also predicts these variables over time. These potential feedback loops between variables consist of positive relations between two nodes, indicating that effects strengthen as time progresses. More specifically, an increase in one variable leads to an increase in the other variable in the next measurement, which in turn leads to an increase in the first variable in the

measurement thereafter, etcetera. Insights regarding such potential feedback loops add to existing literature such as the conceptual frameworks of health behaviour during a pandemic proposed by Bish and Michie (2010) and underlying theoretical frameworks such as the TPB, HBM and PMT by moving from unidirectional effects to more complex interactions between variables. Moreover, such feedback loops provide possible explanations for observed effects. For instance, we observed a decrease in compliance with behavioural measures, support for these measures and involvement in the pandemic throughout the course of this study. According to the temporal network, a decrease in compliance could be explained by a decrease in support for the measures or involvement. The feedback loops suggest that this applies vice versa as well: a decrease in support for the measures or involvement could be explained by a decrease in compliance. These findings resonate with different theoretical models in which behaviour can both be determined by other variables (e.g., TPB; Ajzen, 1991) and determine other variables (e.g., dissonance theory; Festinger, 1957). Moreover, the findings suggest that (elements of) these models can apply simultaneously. Thus, combining different models with bidirectional effects between variables might improve our understanding of health behaviour in a pandemic beyond unidirectional effects between limited sets of variables based on singular theoretical models.

The second pattern observed in the temporal network is that several of these bidirectional effects combined formed a positive reinforcing structure. That is, three variables that showed positive feedback loops with compliance (i.e., support for behavioural measures, involvement in the pandemic, and vaccination intention) also showed positive feedback loops among each other. Such structures can potentially amplify change in these variables. That is, an increase (or decrease) in one variable from the reinforcing structure is likely to be accompanied by an increase (or decrease) in the remaining variables over time, further increasing (decreasing) the initially increased (decreased) variable and other variables over time, and so on.

Such positive reinforcing structures are important to acknowledge for different scenarios. First, considering interventions: if one aims to increase compliance, doing so via a variable that is not only directly related to compliance but also part of a positive reinforcing structure could amplify the effects of interventions over time. For instance, results suggest that an intervention aimed at positively influencing support for behavioural measures could increase not only compliance, but also involvement in the next time frame. This could increase compliance via multiple variables: in addition to the intervention initiating a feedback loop between compliance and support for the measures, the intervention's effect on involvement could also initiate the feedback loop between compliance and involvement. Thus, positive reinforcing structures can amplify the effects of interventions beyond bidirectional effects.

The second reason why reinforcement structures are important is that they can provide insight into processes of polarization during a pandemic. Reinforcement structures can enhance initial leanings towards one end of the evaluative spectrum and lead to movement towards the extremes of

that spectrum. This effect resembles the phenomena of polarization, in which people strengthen their attitude in the initial direction, a process that has been reported in relation to the initial phase of the COVID-19 pandemic (Graso et al., 2021; Kerr et al., 2021). The observation that involvement is part of a reinforcing structure with compliance, support for behavioural measures and vaccination intention corresponds with the Attitudinal Entropy (AE) framework grounded in network theory (Dalege et al., 2018). This framework explains polarization through attitudinal entropy, a state reflecting inconsistent and instable attitudes. Dalege et al. (2018) propose that attitudinal entropy is reduced by thinking about and turning attention towards attitudes objects, and involvement as included in the current study can be interpreted as such. This reduction in entropy results in strengthening attitudes in the initial direction, i.e., polarization. This is also in line with the mere thought effect on attitude polarization (Tesser & Conlee, 1975), proposing that thought can result in attitude polarization.

The third pattern observed in the temporal network is also relevant in the context of interventions. That is, the network contained a triangle of three nodes in which one node shows a bidirectional relation with both other nodes, but these other nodes show a weak or no relation with each other. Such a pattern was for instance observed between depressive complaints, anxiety complaints and loneliness, in which depressive complaints showed a bidirectional relation with both other nodes, but anxiety and loneliness showed a relatively weak relation. This suggests that intervening on depressive complaints can be effective to improve mental health.

Another insight from this study is that our results suggest a weak predictive effect between (nodes related to) mental health and compliance. One could have expected a predictive relation, for example by compliance in the form of adhering to social distancing and isolation guidelines to have an impact on mental health, but there were only weak predictive effects between these constructs when controlling for other variables. This could imply that relations between variables related to mental health (e.g., stress) and compliance observed in prior research (Harper et al., 2021; Lieberoth et al., 2021) might be explained by constructs other than included in those studies, for instance support for the behavioural measures or involvement in the pandemic. Another possible explanation could be that prior research focussed mainly on between-person effects and that relations between mental health and compliance are more pronounced on that level. Comparing psychological within-person and between-person effects in the context of compliance and mental health during pandemics would be the next step in this line of research to examine whether meaningful differences can be observed.

A final insight of this study is that the most important temporal effects concerning compliance with behavioural measures between measurements were comparable with effects found within the same measurement (i.e., contemporaneous effects). For instance, the positive reinforcing structure found between compliance, support for the behavioural measures, involvement and vaccination intention was found not only over time between measurements but also within the same measurement:

higher scores on compliance, support for the measures, involvement in the pandemic and vaccination intention tends to co-occur during a measurement. However, there were also important differences between these time frames, as corroborated by a moderate to large relation between the (edges in the) networks. These differences are also important to consider when designing interventions as these can be tailored based on their anticipated effects. For instance, social norms were found to be relatively important for the COVID-19 network over time, but not within the same measurement. This implies that the effects of interventions aimed at influencing social norms should ideally be studied over time, that is, a next measurement, and not within the same survey.

The current research enables us to formulate several recommendations. The first recommendation concerns future research into causality. Although interesting indications of causality can be drawn based on the approach adopted in this study, no firm statements on causality can be made. In order to do so, interventions should be studied (Kossakowski et al., 2021). Second, the ecological validity of the presented COVID-19 broad attitude networks is unknown. Although the COVID-19 broad attitude networks contain a diverse set of variables, more psychological variables can be relevant for compliance with behavioural measures during pandemics. Changes over time observed in this study could also be caused by predictors not included in the model. This also applies to the contemporaneous effects, which can also indicate that other variables have not (yet) been taken into account. Future research could include additional variables that are deemed relevant based on the scientific literature to further meet real-life complex psychological systems concerning pandemics. Third, although comparing networks of the longitudinal sample and other respondents showed very few significant differences in relations between variables, we cannot predict how respondent dropout affects the results presented here. Given the aim to demonstrate how this approach provides an overview of predictive relations between variables, and not to present a generalizable theoretical model or representative overview of node scores, we assume that dropout has no significant effect on the outcome given the aim of the current study. Fourth, future research could examine temporal dynamics of compliance during pandemics in different phases of a pandemic. The current study was conducted during the initial phase of the COVID-19 pandemic and we cannot predict to what degree results are generalizable beyond the first wave. Results therefore represent a particular period in time, and future research could focus on other phases such as the perseverance period in which adopting protective behaviours needs to be maintained. Fifth and finally, it should be noted that the time spacing of measurements is likely to affect results. For example, it is more likely that change in stable nodes such as compliance and intention to get vaccinated is detected when administering surveys weeks apart than hours apart. We think that the time frame adopted in this study (i.e., 2 to 3 weeks between surveys) is adequate to detect changes in this context; nevertheless, future research could adopt different time frames to examine whether this generates different patterns of effects.

In conclusion, the COVID-19 broad attitude networks obtained in this study show the added value of adopting a complexity approach to compliance in the context of pandemics. Moreover, the

adopted method provides insight into unique relations between a broad set of variables, and how relations between these variables develop over time. Finally, the results suggest that the network structure can provide important insights for explaining observed effects and designing effective interventions, providing an informed strategy grounded in network theory to influence compliance during pandemics.

References

- Ajzen, I. (1991). The Theory of Planned Behavior. *Organizational Behavior and the Human Decision Process*, 50, 179-211.
- Betsch, C., Wieler, L. H., & Habersaat, K. (2020). Monitoring behavioural insights related to COVID-19. *The Lancet*, 395(10232), 1255-1256. [https://doi.org/10.1016/S0140-6736\(20\)30729-7](https://doi.org/10.1016/S0140-6736(20)30729-7)
- Bish, A., & Michie, S. (2010). Demographic and attitudinal determinants of protective behaviours during a pandemic: A review. *British Journal of Health Psychology*, 15(4), 797-824. <https://doi.org/10.1348/135910710x485826>
- Blom, V., Lönn, A., Ekblom, B., Kallings, L. V., Väisänen, D., Hemmingsson, E., Andersson, G., Wallin, P., Stenling, A., Ekblom, Ö., Lindwall, M., Salier Eriksson, J., Holmlund, T., & Ekblom-Bak, E. (2021). Lifestyle Habits and Mental Health in Light of the Two COVID-19 Pandemic Waves in Sweden, 2020. *International Journal of Environmental Research and Public Health*, 18(6), 3313. <https://doi.org/10.3390/ijerph18063313>
- Bogg, T., & Milad, E. (2020). Demographic, personality, and social cognition correlates of coronavirus guideline adherence in a U.S. sample. *Health Psychology*, 39(12), 1026–1036. <https://doi.org/10.1037/hea0000891>
- Borsboom, D., Deserno, M. K., Rhemtulla, M., Epskamp, S., Fried, E. I., McNally, R. J., Robinaugh, D. J., Perugini, M., Dalege, J., Costantini, G., Isvoranu, A.-M., Wysocki, A. C., van Borkulo, C. D., van Bork, R., & Waldorp, L. J. (2021). Network analysis of multivariate data in psychological science. *Nature Reviews Methods Primers*, 1(1), 58. <https://doi.org/10.1038/s43586-021-00055-w>
- Chambon, M., Dalege, J., Elberse, J. E., & van Harreveld, F. (2022). A Psychological Network Approach to Attitudes and Preventive Behaviors During Pandemics: A COVID-19 Study in the United Kingdom and the Netherlands. *Social Psychological and Personality Science*, 13(1), 233-245. <https://doi.org/10.1177/19485506211002420>
- Chambon, M., Dalege, J., Waldorp, L. J., van der Maas, H. L. J., Borsboom, D., & van Harreveld, F. (2022). Tailored interventions into broad attitude networks towards the COVID-19 pandemic. <https://psyarxiv.com/fmu9w/>

- Chan, E. Y. (2021). Moral foundations underlying behavioral compliance during the COVID-19 pandemic. *Personality and Individual Differences*, 171, 110463.
<https://doi.org/10.1016/j.paid.2020.110463>
- Clark, C., Davila, A., Regis, M., & Kraus, S. (2020). Predictors of COVID-19 voluntary compliance behaviors: An international investigation. *Global Transitions*, 2, 76-82.
<https://doi.org/10.1016/j.glt.2020.06.003>
- Dalege, J., Borsboom, D., van Harreveld, F., van den Berg, H., Conner, M., & van der Maas, H. L. J. (2016). Toward a formalized account of attitudes: The Causal Attitude Network (CAN) model. *Psychological Review*, 123(1), 2–22. <https://doi.org/10.1037/a0039802>
- Dalege, J., Borsboom, D., van Harreveld, F., & van der Maas, H. L. J. (2017). Network Analysis on Attitudes: A Brief Tutorial. *Social Psychological and Personality Science*, 8(5), 528-537.
<https://doi.org/10.1177/1948550617709827>
- Dalege, J., Borsboom, D., van Harreveld, F., & van der Maas, H. L. J. (2018). The Attitudinal Entropy (AE) Framework as a General Theory of Individual Attitudes. *Psychological Inquiry*, 29(4), 175-193. <https://doi.org/10.1080/1047840X.2018.1537246>
- Daly, M., & Robinson, E. (2021). Longitudinal changes in psychological distress in the UK from 2019 to September 2020 during the COVID-19 pandemic: Evidence from a large nationally representative study. *Psychiatry Research*, 300, 113920.
<https://doi.org/10.1016/j.psychres.2021.113920>
- de Jong Gierveld, J., & van Tilburg, T. (2008). De ingekorte schaal voor algemene, emotionele en sociale eenzaamheid. *Tijdschrift voor Gerontologie en Geriatrie*, 39(1), 4-15.
<https://doi.org/10.1007/BF03078118>
- Derogatis, L. R. (2001). *BSI 18, Brief Symptom Inventory 18: Administration, scoring and procedures manual*. NCS Pearson, Incorporated.
- Eagly, A. H., & Chaiken, S. (1993). *The psychology of attitudes*. Harcourt Brace Jovanovich College Publishers.
- Epskamp, S. (2017). Brief Report on Estimating Regularized Gaussian Networks from Continuous and Ordinal Data. <https://arxiv.org/abs/1606.05771>
- Epskamp, S. (2020a). Psychometric network models from time-series and panel data. *Psychometrika*, 85(1), 206-231. <https://doi.org/10.1007/s11336-020-09697-3>
- Epskamp, S. (2020b). *Psychonetrics: Structural Equation Modeling and Confirmatory Network Analysis*. <https://cran.r-project.org/web/packages/psychonetrics/index.html>
- Epskamp, S., van Borkulo, C. D., van der Veen, D. C., Servaas, M. N., Isvoranu, A.-M., Riese, H., & Cramer, A. O. J. (2018). Personalized Network Modeling in Psychopathology: The Importance of Contemporaneous and Temporal Connections. *Clinical Psychological Science*, 6(3), 416-427. <https://doi.org/10.1177/2167702617744325>
- Festinger, L. (1957). *A theory of cognitive dissonance* (Vol. 2). Stanford University Press.

- Gibson, L. P., Magnan, R. E., Kramer, E. B., & Bryan, A. D. (2021). Theory of Planned Behavior Analysis of Social Distancing During the COVID-19 Pandemic: Focusing on the Intention–Behavior Gap. *Annals of Behavioral Medicine*, 55(8), 805-812.
<https://doi.org/10.1093/abm/kaab041>
- Government of the Netherlands. (2020). *Dutch measures against coronavirus: basic rules for everyone*. <https://www.government.nl/topics/coronavirus-covid-19/tackling-new-coronavirus-in-the-netherlands/basic-rules-for-everyone>
- Graham, J., Haidt, J., Koleva, S., Motyl, M., Iyer, R., Wojcik, S. P., & Ditto, P. H. (2013). Chapter Two - Moral Foundations Theory: The Pragmatic Validity of Moral Pluralism. In P. Devine & A. Plant (Eds.), *Advances in Experimental Social Psychology* (Vol. 47, pp. 55-130). Academic Press. <https://doi.org/10.1016/B978-0-12-407236-7.00002-4>
- Graso, M., Chen, F. X., & Reynolds, T. (2021). Moralization of Covid-19 health response: Asymmetry in tolerance for human costs. *Journal of Experimental Social Psychology*, 93, 104084. <https://doi.org/10.1016/j.jesp.2020.104084>
- Harper, C. A., Satchell, L. P., Fido, D., & Latzman, R. D. (2021). Functional Fear Predicts Public Health Compliance in the COVID-19 Pandemic. *International Journal of Mental Health and Addiction*, 19, 1875-1888. <https://doi.org/10.1007/s11469-020-00281-5>
- Hensel, L., Witte, M., Caria, A. S., Fetzer, T., Fiorin, S., Götz, F. M., Gomez, M., Haushofer, J., Ivchenko, A., Kraft-Todd, G., Reutskaja, E., Roth, C., Yoeli, E., & Jachimowicz, J. M. (2022). Global Behaviors, Perceptions, and the Emergence of Social Norms at the Onset of the COVID-19 Pandemic. *Journal of Economic Behavior & Organization*, 193, 473-496. <https://doi.org/10.1016/j.jebo.2021.11.015>
- Ibuka, Y., Chapman, G. B., Meyers, L. A., Li, M., & Galvani, A. P. (2010). The dynamics of risk perceptions and precautionary behavior in response to 2009 (H1N1) pandemic influenza. *BMC Infectious Diseases*, 10(1), 296. <https://doi.org/10.1186/1471-2334-10-296>
- Kerr, J., Panagopoulos, C., & van der Linden, S. (2021). Political polarization on COVID-19 pandemic response in the United States. *Personality and Individual Differences*, 179, 110892. <https://doi.org/10.1016/j.paid.2021.110892>
- Kossakowski, J. J., Waldorp, L. J., & van der Maas, H. L. J. (2021). The Search for Causality: A Comparison of Different Techniques for Causal Inference Graphs. *Psychological Methods*, 26(6), 719–742. <https://doi.org/10.1037/met0000390>
- Leigh-Hunt, N., Baggeley, D., Bash, K., Turner, V., Turnbull, S., Valtorta, N., & Caan, W. (2017). An overview of systematic reviews on the public health consequences of social isolation and loneliness. *Public Health*, 152, 157-171. <https://doi.org/10.1016/j.puhe.2017.07.035>
- Li, J., Yang, A., Dou, K., Wang, L., Zhang, M., & Lin, X. (2020). Chinese public's knowledge, perceived severity, and perceived controllability of COVID-19 and their associations with

- emotional and behavioural reactions, social participation, and precautionary behaviour: a national survey. *BMC Public Health*, 20, 1589. <https://doi.org/10.1186/s12889-020-09695-1>
- Lieberoth, A., Lin, S.-Y., Stöckli, S., Han, H., Kowal, M., Gelpi, R., Chrona, S., Tran, T. P., Jeftić, A., Rasmussen, J., Cakal, H., & Milfont, T. L. (2021). Stress and worry in the 2020 coronavirus pandemic: relationships to trust and compliance with preventive measures across 48 countries in the COVIDiSTRESS global survey. *Royal Society Open Science*, 8(2), 200589. <https://doi.org/10.1098/rsos.200589>
- Rogers, R. W. (1975). A Protection Motivation Theory of Fear Appeals and Attitude Change. *The Journal of Psychology*, 91(1), 93-114. <https://doi.org/10.1080/00223980.1975.9915803>
- Rosenstock, I. M. (1974). Historical Origins of the Health Belief Model. *Health Education Monographs*, 2(4), 328-335. <https://doi.org/10.1177/109019817400200403>
- Schneider, C. R., Dryhurst, S., Kerr, J., Freeman, A. L. J., Recchia, G., Spiegelhalter, D., & van der Linden, S. (2021). COVID-19 risk perception: a longitudinal analysis of its predictors and associations with health protective behaviours in the United Kingdom. *Journal of Risk Research*, 1-20. <https://doi.org/10.1080/13669877.2021.1890637>
- Shahnazi, H., Ahmadi-Livani, M., Pahlavanzadeh, B., Rajabi, A., Hamrah, M. S., & Charkazi, A. (2020). Assessing preventive health behaviors from COVID-19: a cross sectional study with health belief model in Golestan Province, Northern of Iran. *Infectious Diseases of Poverty*, 9(6), 91-99. <https://doi.org/10.1186/s40249-020-00776-2>
- Tennant, R., Hiller, L., Fishwick, R., Platt, S., Joseph, S., Weich, S., Parkinson, J., Secker, J., & Stewart-Brown, S. (2007). The Warwick-Edinburgh Mental Well-being Scale (WEMWBS): development and UK validation. *Health and Quality of Life Outcomes*, 5(1), 63. <https://doi.org/10.1186/1477-7525-5-63>
- Tesser, A., & Conlee, M. C. (1975). Some effects of time and thought on attitude polarization. *Journal of Personality and Social Psychology*, 31(2), 262-270. <https://doi.org/10.1037/h0076292>
- Tunçgenç, B., El Zein, M., Sulik, J., Newson, M., Zhao, Y., Dezechache, G., & Deroy, O. (2021). Social influence matters: We follow pandemic guidelines most when our close circle does. *British Journal of Psychology*, 112(3), 763-780. <https://doi.org/10.1111/bjop.12491>
- van Borkulo, C. D., Borsboom, D., Epskamp, S., Blanken, T. F., Boschloo, L., Schoevers, R. A., & Waldorp, L. J. (2014). A new method for constructing networks from binary data. *Scientific Reports*, 4(1), 5918. <https://doi.org/10.1038/srep05918>
- van der Weerd, W., Timmermans, D. R., Beaujean, D. J., Oudhoff, J., & van Steenberghe, J. E. (2011). Monitoring the level of government trust, risk perception and intention of the general public to adopt protective measures during the influenza A (H1N1) pandemic in The Netherlands. *BMC Public Health*, 11, 575. <https://doi.org/10.1186/1471-2458-11-575>

- van Rooij, B., de Bruijn, A. L., Reinders Folmer, C., Kooistra, E. B., Kuiper, M. E., Brownlee, M., Olthuis, E., & Fine, A. (2020). Compliance with COVID-19 Mitigation Measures in the United States. <https://doi.org/10.31234/osf.io/qymu3>
- Wang, Z., Luo, S., Xu, J., Wang, Y., Yun, H., Zhao, Z., Zhan, H., & Wang, Y. (2021). Well-Being Reduces COVID-19 Anxiety: A Three-Wave Longitudinal Study in China. *Journal of Happiness Studies*, 22, 3593–3610. <https://doi.org/10.1007/s10902-021-00385-2>
- WHO Regional Office for Europe. (2020). COVID-19 Snapshot Monitoring (COSMO Standard): Monitoring knowledge, risk perceptions, preventive behaviours, and public trust in the current coronavirus outbreak - WHO standard protocol. <https://www.psycharchives.org/handle/20.500.12034/2392>
- Wolff, K., Larsen, S., & Øgaard, T. (2019). How to define and measure risk perceptions. *Annals of Tourism Research*, 79, 102759. <https://doi.org/10.1016/j.annals.2019.102759>

Supplemental Material

1. Method

1.1 Dropped out participants longitudinal study

Demographic information was compared between the longitudinal sample and the participants that dropped out of the study before the final wave (total drop-out $n = 3694$). A chi square test of independence indicated that the proportion of participants that dropped out of the study differed by gender, $\chi^2(1, N = 6093) = 6.55, p = .011$, and illness, $\chi^2(1, N = 6093) = 34.53, p < .001$. Specifically, men and participants suffering from a condition that could worsen the consequences of COVID-19 were more likely to participate in the current study up and until the final wave. No significant differences between the longitudinal sample and drop-outs were found for education and smoking. A Mann-Whitney U test indicated that the age was significantly higher in the longitudinal sample ($Mdn = 56$) than in the group of participants that dropped out of the study ($Mdn = 47$), $U = 5,529,364.50, p < .001$, which implies that the average age of the longitudinal sample is higher than the average Dutch population.

The Network Comparison Test (van Borkulo et al., 2022) was conducted to formally test for differences in network structure at the first measurement between respondents from the longitudinal sample and respondents that dropped out of the study after the first measurement. Results revealed network invariance ($p < .001$), indicating that not all edges are equal between the networks of the first measurement of the longitudinal sample and respondents that dropped out. Further investigation into specific edge differences reveals that, after applying a Holm correction for multiple testing, only two edges differ significantly at the first measurement between the longitudinal and dropout sample. More specifically, the edges that differ significantly are somatic symptoms and self-exempting beliefs (i.e., weak positive edge is stronger in dropout sample), and between somatic symptoms and social norm (i.e., weak negative edge is absent in dropout sample). Finally, results indicated a significant difference in global strength (i.e., overall connectivity of the networks) between the longitudinal sample and respondents that dropped out of the study (global strength is 2.61, $p = .036$).

1.2 Survey items per node in the COVID-19 broad attitude networks

Factor	Survey items - (R) = reverse score	Answer scale
Compliance T1	I comply with the corona measures.	1 (<i>I never comply with the measures</i>) to 7 (<i>I always comply with the measures</i>)
	Stay at home as much as possible.	1 (<i>I do not display this behaviour more</i>) to 7 (<i>I display this behaviour much more now</i>)
	Keep 1.5 metres away from others.	
	Wash your hands regularly with soap and water.	
	Cough and sneeze into the inside of your elbow.	
Compliance T2	I comply with the corona measures.	1 (<i>I never comply with the measures</i>) to 7 (<i>I always comply with the measures</i>)
	Keep 1.5 metres away from others.	1 (<i>I do not display this behaviour more</i>) to 7 (<i>I display this behaviour much more now</i>)
	Wash your hands regularly with soap and water.	
	Cough and sneeze into the inside of your elbow.	
	Avoid crowds.	
Compliance T3 -T5	I comply with the corona measures.	1 (<i>I never comply with the measures</i>) to 7 (<i>I always comply with the measures</i>)
	Keep 1.5 metres away from others.	1 (<i>I do not display this behaviour more</i>) to 7 (<i>I display this behaviour much more now</i>)
	Wash your hands regularly with soap and water.	
	Cough and sneeze into the inside of your elbow.	
	Avoid crowds.	
	Stay close to home.	
Risk Perception	How likely do you think it is that u will get infected with the coronavirus in the upcoming year?	1 (<i>Extremely unlikely</i>) to 7 (<i>Extremely likely</i>)
	How severe do you think an infection with the coronavirus would be for you?	1 (<i>Not severe</i>) to 7 (<i>Very severe</i>)
Health Risk	For me personally, I consider the health risk of an infection with the coronavirus..	1 (<i>Extremely small</i>) to 7 (<i>Extremely severe</i>)
	For my family and friends, I consider the health risk of an infection with the coronavirus..	
Economic Consequences	For me personally, I consider the economic consequences of the corona pandemic ..	1 (<i>Extremely small</i>) to 7 (<i>Extremely severe</i>)
	For my family and friends, I consider the economic consequences of the corona pandemic are..	
Self-exempting Beliefs	I will not get infected with the coronavirus because I never get the seasonal flu (influenza) either.	1 (<i>Strongly disagree</i>) to 7 (<i>Strongly agree</i>)
	I think I am already immune (protected) against the coronavirus.	
Negative Affect	The corona pandemic is making me angry.	1 (<i>Strongly disagree</i>) to 7 (<i>Strongly agree</i>)
	The corona pandemic is making me sad.	
	The corona pandemic is making me feel confused.	

Compassion Worries Virus	The corona pandemic is making me feel uncertain.	
	The corona pandemic is making me feel overwhelmed.	
	The corona pandemic is making me feel frustrated.	
	The corona pandemic is making me feel fearful.	
	The corona pandemic is making me feel out of control.	
Worries Measures	The corona pandemic is making me feel compassion.	1 (<i>Strongly disagree</i>) to 7 (<i>Strongly agree</i>)
	I worry about getting infected with the coronavirus.	1 (<i>Don't worry at all</i>) to 7 (<i>Worry a lot</i>)
	I worry about infecting others with the coronavirus.	
	I worry about losing someone I love.	
	I worry about the health care system overloading.	
Vaccination Intention Measures Support	I worry about what staying at home a lot will do to my health.	1 (<i>Don't worry at all</i>) to 7 (<i>Worry a lot</i>)
	I worry about the schools closing.	
	I worry about a recession.	
	I worry about limited access to food.	
	I worry about losing my job.	
Measures Ease	I worry about getting lonely.	
	If a vaccine becomes available, I would get it.	1 (<i>Strongly disagree</i>) to 7 (<i>Strongly agree</i>)
	I find the corona measures...	1 (<i>Senseless</i>) to 7 (<i>Sensible</i>)
	I find the corona measures...	1 (<i>Useless</i>) to 7 (<i>Useful</i>)
	I find the corona measures...	1 (<i>Unnecessary</i>) to 7 (<i>Necessary</i>)
Social norm	I find the corona measures...	1 (<i>Unfair</i>) to 7 (<i>Fair</i>)
	I find the corona measures...	1 (<i>Unacceptable</i>) to 7 (<i>Acceptable</i>)
	I think the corona measures will prevent the spread of the coronavirus.	1 (<i>Strongly disagree</i>) to 7 (<i>Strongly agree</i>)
	As a society, we must do whatever is necessary to prevent the spread of the coronavirus.	
	I find the corona measures...	1 (<i>Unpleasant</i>) to 7 (<i>Pleasant</i>)
Control Infection	I find the corona measures...	1 (<i>Difficult</i>) to 7 (<i>Easy</i>)
	I think the majority of people comply with the corona measures.	1 (<i>Strongly disagree</i>) to 7 (<i>Strongly agree</i>)
	I think majority of people find it important that people comply with the corona measures.	
	For me personally, avoiding an infection with the coronavirus in the current situation is...	1 (<i>Extremely difficult</i>) to 7 (<i>Extremely easy</i>)
	For my family and friends, avoiding an infection with the coronavirus in the current situation is...	
Self-efficacy Involvement	I know how to protect myself from the coronavirus.	1 (<i>Strongly disagree</i>) to 7 (<i>Strongly agree</i>)
	How important is the topic of corona pandemic to you?	1 (<i>Very unimportant</i>) to 7 (<i>Very important</i>)
	To what extent does the news about the corona pandemic have your attention?	1 (<i>Not at all</i>) to 7 (<i>Very much</i>)
	How much do you think about the corona pandemic?	1 (<i>Very little</i>) to 7 (<i>Very much</i>)

Perceived Knowledge	How much knowledge do you think you have about the corona pandemic?	1 (<i>Very little</i>) to 7 (<i>Very much</i>)
Trust	I trust the authorities to adequately manage the corona pandemic. I trust RIVM during the corona pandemic. I trust health care professionals during the corona pandemic. I trust science during the corona pandemic.	1 (<i>Strongly disagree</i>) to 7 (<i>Strongly agree</i>)
Health General	In general, how would you rate your health?	1 (<i>Very poor</i>) to 7 (<i>Very good</i>)
Health Change Physical	How would you rate your physical health now as compared to before the corona pandemic?	-3 (<i>Much worse</i>) to 3 (<i>Much better</i>)
Health Change Mental	How would you rate your mental health now as compared to before the corona pandemic?	-3 (<i>Much worse</i>) to 3 (<i>Much better</i>)
Healthy Lifestyle	I've been exercising in the past two weeks, compared to before the corona pandemic...	-3 (<i>Much less</i>) to 3 (<i>Much more</i>)
	I've been eating in the past two weeks, compared to before the corona pandemic...	-3 (<i>Much less healthy</i>) to 3 (<i>Much healthier</i>)
	I've been sleeping for the past two weeks, compared to before the corona pandemic...	-3 (<i>Much worse</i>) to 3 (<i>Much better</i>)
Mental Wellbeing	I've been feeling optimistic about the future. I've been feeling useful. I've been feeling relaxed. I've been dealing with problems well. I've been thinking clearly. I've been feeling close to other people. I've been able to make up my own mind about things.	1 (<i>Never</i>) to 5 (<i>Always</i>)
Loneliness	I experience a general sense of emptiness. There are plenty of people I can lean on when I have problems. There are many people I can trust completely. I miss having people around me. There are enough people I feel close to. I often feel rejected.	1 (<i>Not at all</i>) to 5 (<i>Very much</i>)
Anxiety Complaints	To what extent did you experience nervousness during the past two weeks? To what extent did you experience feeling suddenly scared during the past two weeks? To what extent did you experience feeling fearful during the past two weeks? To what extent did you experience feeling tense during the past two weeks? To what extent did you experience anxiety or panic attacks during the past two weeks? To what extent did you experience restlessness during the past two weeks?	1 (<i>Not at all</i>) to 5 (<i>Very much</i>)
Depressive Complaints	To what extent did you experience a loss of the will to live during the past two weeks? To what extent did you experience feeling lonely during the past two weeks? To what extent did you experience feeling blue during the past two weeks? To what extent did you experience a loss of interest in things during the past two weeks?	1 (<i>Not at all</i>) to 5 (<i>Very much</i>)

	To what extent did you experience feeling hopeless about the future during the past two weeks?	
	To what extent did you experience a feeling of worthlessness during the past two weeks?	
Somatic Complaints	To what extent did you experience dizziness or faintness during the past two weeks?	1 (<i>Not at all</i>) to 5 (<i>Very much</i>)
	To what extent did you experience chest pain during the past two weeks?	
	To what extent did you experience nausea during the past two weeks?	
	To what extent did you experience difficulty breathing during the past two weeks?	
	To what extent did you experience numbness during the past two weeks?	
	To what extent did you experience feeling weak during the past two weeks?	

1.3 Node construction and Principal Axis Factoring

Analysis for node construction was conducted with the largest and most diverse sample available, namely wave 1, as this wave included all participants that completed the first survey (i.e., featured no drop-out). Survey items were combined to form nodes: the combination of items was either predetermined by validated scales or a fixed operationalization, or identified through Principal Axis Factoring (PAF; see Table 1). PAF was conducted with Oblimin rotation given the expected intercorrelation between items. Extraction of components was based on eigenvalues greater than one. The results are discussed below. Items presented in *italic* were excluded from the node.

Table 1 - Overview of nodes and the approach to the selection of items per node

Nodes	Approach to combining items in node
Compliance (T1)	Predetermined - Operationalized with the recommended behavioural measures, PAF confirmed single component
Risk Perception	Predetermined by product of likelihood and severity
Health Risk	Predetermined – Two item construct
Economic Consequences	Predetermined – Two item construct
Self-exempting Beliefs	PAF – Single component
Negative Affect	PAF – Component identified in items on affect
Compassion	PAF – Component identified in items on affect
Worries Virus	PAF – Component identified in items on worries
Worries Measures	PAF – Component identified in items on worries
Vaccination Intention	Predetermined single item node
Measures Support	PAF – Component identified in items on attitudes toward corona measures
Measures Ease	PAF – Component identified in items on attitudes toward corona measures
Social Norm	Predetermined – Two item construct
Control Infection	Predetermined – Two item construct
Self-efficacy	Predetermined single item node
Involvement	PAF – Single component
Perceived Knowledge	Predetermined single item node
Trust	PAF – Single component
Health General	Predetermined single item node
Health change Physical	Predetermined single item node
Health change Mental	Predetermined single item node
Healthy Lifestyle	Predetermined - Operationalized with basic items on diet, exercise and sleep, PAF confirmed single component
Mental Wellbeing	Predetermined - Validated scale
Loneliness	Predetermined - Validated scale
Somatic Complaints	Predetermined - Validated scale
Depressive Complaints	Predetermined - Validated scale
Anxiety Complaints	Predetermined - Validated scale

Compliance

Compliance was predetermined through operationalization with the recommended behavioural measures. Kaiser-Meyer-Olkin Measure of Sampling Adequacy was .78 and Bartlett's

test of sphericity was significant, $\chi^2(df = 10) = 8,429.90, p < .001$, supporting a rationale for performing PAF with the five items for *Compliance* in the first wave (i.e. a general item and four specific behaviours). The results suggested a single component for the items on compliance, resulting in the node *Compliance* with the four items shown below.

No.	Survey items	Compliance Component load
1	I comply with the corona measures.	.837
2	Stay at home as much as possible.	.613
3	Keep 1.5 metres away from others.	.735
4	Wash your hands regularly with soap and water.	.545
5	Cough and sneeze into the inside of your elbow.	.444

Self-exempting Beliefs

Kaiser-Meyer-Olkin Measure of Sampling Adequacy was .54 and Bartlett's test of sphericity was significant, $\chi^2(df = 3) = 1,320.13, p < .001$, not convincingly supporting a rationale for performing PAF with the three items for *Self-exempting Beliefs*. Nevertheless, the results suggested excluding the item 'I think infections are good for building up group immunity (protection)', because of its low component load. This is supported by an increase in scale reliability after deletion (three items $\alpha = .47$, if item deleted $\alpha = .57$; resulting in $r_{sb} = .57$). As a result, the node *Self-exempting Beliefs* consisted of item 1 and 2 (see below).

No.	Survey items	Self-exempting Beliefs Component load
1	I will not get infected with the coronavirus because I never get the seasonal flu (influenza) either.	.505
2	I think I am already immune (protected) against the coronavirus.	.780
3	<i>I think infections are good for building up herd immunity (protection).</i>	.242

Negative Affect and Compassion

Kaiser-Meyer-Olkin Measure of Sampling Adequacy was .90 and Bartlett's test of sphericity was significant, $\chi^2(df = 36) = 24,219.84, p < .001$, supporting a rationale for performing PAF with the nine items covering affect. Compassion was identified as a separate component. This resulted in two nodes: *Negative Affect*, consisting of the items 1 to 8 shown below, and the single-item (no. 9) node *Compassion*.

No.	Survey items	Negative Affect Component load	Compassion Component load
1	The corona pandemic is making me angry.	.729	
2	The corona pandemic is making me sad.	.475	
3	The corona pandemic is making me feel confused.	.678	
4	The corona pandemic is making me feel uncertain.	.649	
5	The corona pandemic is making me feel overwhelmed.	.472	
6	The corona pandemic is making me feel frustrated.	.890	

7	The corona pandemic is making me fearful.	.572	
8	The corona pandemic is making me feel out of control.	.628	
9	The corona pandemic is making me feel compassion.		.511

Worries Virus and Worries Measures

Kaiser-Meyer-Olkin Measure of Sampling Adequacy was .84 and Bartlett's test of sphericity was significant, $\chi^2(df = 55) = 13,097.02, p < .001$, supporting a rationale for performing PAF with 11 items covering worries. The results indicated two components for items on worries: items on worries about the virus and its effects (*Worries Virus*) and items on worries about the effects of the measures taken to manage the pandemic (*Worries Measures*). The item 'I worry about society becoming increasingly egoistic' was excluded because this item had a low Component load on both components. This resulted in two nodes: the node *Worries Virus* that consisted of item 1 to 4 shown below, and the node *Worries Measures* that consisted of items 5 to 10.

No.	Survey items	Worries Virus Component load	Worries Measures Component load
1	I worry about getting infected with the coronavirus.	.718	
2	I worry about infecting others with the coronavirus.	.511	
3	I worry about losing someone I love.	.695	
4	I worry about the health care system overloading.	.627	
5	I worry about what staying at home a lot will do to my health.		.716
6	I worry about the schools closing.		.326
7	I worry about a recession.		.301
8	I worry about limited access to food.		.384
9	I worry about losing my job.		.427
10	I worry about getting lonely.		.633
11	<i>I worry about society becoming increasingly egoistic.</i>	.269	.218

Measures Support and Measures Ease

Kaiser-Meyer-Olkin value was .90 and Bartlett's test of sphericity was significant, $\chi^2(df = 45) = 31,141.32, p < .001$, supporting a rationale for performing PAF with the 10 items on attitudes toward the corona measures. The item 'I think the corona measures will prevent the spread of the coronavirus' initially loaded on a third component, but due to the low eigenvalue for this component (1.003) we reran the PAF with two fixed factors, resulting in this item loading on the component *Measures Support*. Furthermore, Cronbach's alpha of the node *Measures Support* did not require the exclusion this item ($\alpha = .90$, if item deleted $\alpha = .91$). The item 'I find the corona measures... Insufficient - Overdone' was excluded, because this item has a low Component load on the *Measures Support* component. This resulted in two nodes: the node *Measures Support*, consisting of the items 1 to 7 shown below, and the node *Measures Ease*, consisting of the items 8 and 9.

No.	Survey items	Measures Support	Measures Ease
-----	--------------	---------------------	------------------

		Component load	Component load
1	I find the corona measures... Senseless – Sensible.	.869	
2	I find the corona measures... Useless – Useful.	.877	
3	I find the corona measures... Unnecessary – Necessary.	.901	
4	I find the corona measures... Unfair – Fair.	.647	
5	I find the corona measures... Unacceptable – Acceptable.	.776	
6	I think the corona measures will prevent the spread of the coronavirus.	.497	
7	As a society, we must do whatever is necessary to prevent the spread of the coronavirus.	.654	
8	I find the corona measures... Unpleasant – Pleasant.		.661
9	I find the corona measures... Difficult – Easy.		.774
10	<i>I find the corona measures... Insufficient – Overdone.</i>	-.242	

Involvement

Kaiser-Meyer-Olkin value was .73 and Bartlett's test of sphericity was significant, $\chi^2(df = 6) = 7,774.14, p < .001$, supporting a rationale for performing PAF with the four items on the degree to which participants are involved in the corona pandemic. The items loaded on one component, resulting in the node *Involvement* with the items 1 to 3 shown below. The item 'To what extent do you feel you can avoid the news about the corona pandemic?' was excluded, because this item has a low component load on the *Involvement* component.

No.	Survey items	Involvement Component load
1	How important is the topic of corona pandemic to you?	.832
2	To what extent does the news about the corona pandemic have your attention?	.807
3	How much do you think about the corona pandemic?	.754
4	<i>To what extent do you feel you can avoid the news about the corona pandemic?</i>	-.176

Trust

Kaiser-Meyer-Olkin value was .73 and Bartlett's test of sphericity was significant, $\chi^2(df = 6) = 14,759.76, p < .001$, supporting a rationale for performing PAF with the four items on trust in actor relevant to manage the corona pandemic. The results suggested a single component for the items on trust, resulting in the node *Trust* with the four items shown below.

No.	Survey items	Trust Component load
1	I trust the authorities to adequately manage the corona pandemic.	.883
2	I trust RIVM during the corona pandemic.	.901
3	I trust health care professionals during the corona pandemic.	.543
4	I trust science during the corona pandemic.	.767

Healthy Lifestyle

Healthy Lifestyle was predetermined through operationalization with basic items on diet, exercise and sleep.⁵ Kaiser-Meyer-Olkin value was .60 and Bartlett's test of sphericity was significant, $\chi^2(df = 3) = 1,171.04, p < .001$, not convincingly supporting a rationale for performing PAF with the three items on healthy lifestyle change compared to before the corona pandemic. Nevertheless, in accordance with the predetermined operationalization the results suggest a single component for the items. This is confirmed by the result that scale reliability statistics ($\alpha = .49$) decrease when excluding items. This resulted in the node *Healthy Lifestyle* consisting of the three items shown below.

No.	Survey items	Healthy Lifestyle Component load
1	I've been eating in the past two weeks, compared to before the corona pandemic... Less healthy – healthier.	.607
2	I've been exercising in the past two weeks, compared to before the corona pandemic... Less – More.	.521
3	I've been sleeping for the past two weeks, compared to before the corona pandemic... Worse – Better.	.404

⁵ The survey also contained items on smoking and alcohol consumption, but these items were excluded from further analysis due to the large amount of missing values. These items loaded on another component than *Healthy Lifestyle*.

1.4 Comparing results from intervention conditions

The interventions featured conditions aimed at both increasing and decreasing scores on the same nodes. Consequently, there were no significant differences between scores on nodes that were targeted by the interventions depending on whether participants were assigned to an intervention or control condition (see below).

The table on the next page provides the necessary Mann-Whitney test parameters to compare the scores on the nodes between the conditions participants were randomly assigned to for the interventions. Wave 3 and Wave 5 both contained five conditions: two conditions aimed at increasing and decreasing a central node, two conditions aimed at increasing and decreasing a peripheral node, and a control condition. The following table shows the results of comparing the scores on the nodes of respondents in the control condition with respondents in either of the four intervention condition. The scores on the nodes did not differ significantly depending on whether participants were assigned to one of the intervention conditions or the control condition. The only exception was *Vaccination Intention* in Wave 5: there was a significant difference between the control condition and the combined intervention conditions for this node. The interventions in this wave were aimed at the nodes *Measures Support* and *Economic Consequences*, therefore the interventions do not provide a plausible explanation for this difference.

Nodes ¹	Wave 3				Wave 5			
	<i>Mdn</i> Control condition (<i>n</i> = 750)	<i>Mdn</i> Intervention condition (<i>n</i> = 3004)	<i>U</i>	<i>p</i> -value	<i>Mdn</i> Control condition (<i>n</i> = 488)	<i>Mdn</i> Intervention condition (<i>n</i> = 1961)	<i>U</i>	<i>p</i> -value
Compliance	6	6	1,165,498.50	.14	5.83	5.83	458,036.00	.14
Risk Perception	16	16	1,150,719.50	.36	15	15	485,290.00	.63
Health Risk	4	4	1,145,925.00	.46	4	4	464,738.00	.32
Economic Consequences	4	4	1,133,674.00	.79	3.50	3.50	479,542.50	.94
Self-exempting Beliefs	1.50	1.50	1,108,881.00	.49	1.50	1.50	481,802.50	.81
Negative Affect	3.50	3.38	1,100,315.00	.32	3.13	3.13	478,979.00	.97
Compassion	5	5	1,115,263.50	.67	5	5	465,307.00	.34
Worries Virus	4	4	1,104,672.00	.41	3.75	3.75	466,229.50	.38
Worries Measures	3	3	1,142,642.50	.54	2.83	2.83	471,742.50	.63
Vaccination Intention	6	6	1,145,813.50	.44	7	7	447,726.50	.021*
Measures Support	5.71	5.71	1,129,523.00	.91	5.71	5.71	462,268.50	.25
Measures Ease	4	4	1,151,451.50	.34	4	4	461,460.50	.22
Social Norm	5	5	1,139,130.50	.63	4.50	4.50	457,933.00	.14
Control Infection	4.50	4.50	1,161,302.00	.18	4.50	4.50	463,272.00	.27
Self-efficacy	6	6	1,158,188.50	.21	6	6	468,057.50	.44
Involvement	5	4.67	1,091,875.50	.19	4.67	4.67	468,680.50	.48
Perceived Knowledge	5	5	1,111,568.00	.56	5	5	454,054.50	.07
Trust	5.75	5.75	1,158,710.00	.22	6	6	453,831.50	.08
Health General	6	6	1,125,179.50	.96	5	5	489,679.50	.40
Health change Physical	4	4	1,134,294.50	.71	4	4	490,732.50	.27
Health change Mental	4	4	1,149,268.50	.31	4	4	475,838.50	.82
Healthy Lifestyle	4	4	1,134,066.50	.77	4	4	497,116.50	.16
Mental Wellbeing~	24.11	25.03	1,163,825.50	.16	25.03	25.03	464,135.50	.30
Loneliness~	2.17	2.17	1,110,859.50	.56	2	2	503,637.00	.07
Somatic Complaints~	1	1	1,129,146.50	.91	1	1	482,149.00	.77
Depressive Complaints~	1.50	1.50	1,095,845.00	.24	1.33	1.33	482,215.50	.78
Anxiety Complaints~	1.33	1.33	1,088,371.00	.14	1.17	1.17	484,616.00	.65

¹ 7-point Likert-scale, unless marked ~, indicating a 5-point Likert-scale. Please note that *Risk Perception* is the product of two items and *Mental Wellbeing* is the metric sum score of all items.

* Significant difference ($p < .05$)

1.5 R script

See OSF (https://osf.io/qu7p2/?view_only=01a4c512371c460cab4919f2dcbb7bdc) for the .R file containing the R script (R Core Team, 2013). Besides the values specified in the R-script, the default values were used.

The script and the interpretation of the output was based on the tutorial paper of (Epskamp, 2020a). We used the package *qgraph* (Epskamp et al., 2012) to visualize the graphs, *bootnet* (Epskamp et al., 2018) for the stability and accuracy measures and *psychometrics* (Epskamp, 2020b) for the temporal analyses.

Please note that the layout of the network (i.e., position of the nodes) was determined by the average layout of both the temporal and contemporaneous networks. Subsequently, we manually adjusted the position of several nodes to improve readability, as specified in the script.

2. Results

2.1 Descriptive statistics and Repeated Measures ANOVA

The statistics relevant for the temporal dynamics of the COVID-19 broad attitude networks are presented in the table below (i.e., longitudinal sample of participants that completed all waves). The table also includes the results of a repeated measures ANOVA to compare the mean scores on the continuous variables for the different waves (i.e., T1 – T5; see column T5), which is discussed below.

The Kolmogorov-Smirnov value was significant for all variables ($p < .001$). Given the large sample size, we based the decisions concerning data transformation on the QQ-plots. The variables *Health Risk*, *Economic Consequences*, *Negative Affect*, *Worries Virus* and *Measures Ease* were treated as normally distributed and therefore not transformed. The rest of the variables were transformed with $\log_{10}(x)$, unless negatively skewed, in which case we used the $\log_{10}(k-x)$ transformation. The latter applied to the following variables: *Compliance*, *Compassion*, *Vaccination Intention*, *Measures Support*, *Social Norm*, *Control Infection*, *Self-efficacy*, *Involvement*, *Perceived Knowledge*, *Trust* and *Health General*. When the assumption of sphericity was violated, we report the lower-bound corrected results by default, unless otherwise specified. Post hoc analysis for all nodes was based on estimated marginal means with a Bonferroni adjustment for multiple comparisons.

The results indicate a significant effect of time on all nodes. A detailed description of the repeated measures ANOVA results and post hoc analysis with a specification of which waves significantly differ is provided below. *Compliance* with behavioural measures significantly decreased over time. All nodes on elements of attitudes relevant for the pandemic and the behavioural measures (e.g., perceived [health] risk, negative affect, worries and support for the measures) decreased in average score over time, with the exception of *Measures Ease*. With regard to the additional psychological nodes, the scores on *Social Norm* and *Involvement* decreased, whereas scores for *Control Infection* increased. *Trust* initially decreased, but showed an increase between the last two measurements. Participants reported an improvement in the health-related nodes: the majority of the health-related nodes decreased over time, with the exception of *Health change Mental*, *Healthy Lifestyle* and *Mental Wellbeing* that increased over time. The largest effect sizes were found for the nodes *Compliance* ($\eta^2 = .184$), *Worries Virus* ($\eta^2 = .120$), *Social Norm* ($\eta^2 = .109$) and *Involvement* ($\eta^2 = .103$), which all decreased over time.

Completed waves	Wave 1 - 5				
	T1	T2	T3	T4	T5
Nodes ¹	M (SD)	M (SD)	M (SD)	M (SD)	M (SD)
Compliance	6.11 (0.77)	6.10 (0.77)	5.92 (0.87)	5.79 (0.93)	5.68 (1)**
Risk Perception	18.35 (8.81)	17.88 (8.76)	17.39 (8.8)	17.09 (8.76)	16.41 (9.03)**
Health Risk	4.25 (1.12)	4.24 (1.15)	4.18 (1.2)	4.17 (1.22)	4.11 (1.27)**
Economic Consequences	3.99 (1.37)	3.79 (1.33)	3.73 (1.35)	3.65 (1.36)	3.48 (1.33)**
Self-exempting Beliefs	2.04 (1.10)	1.99 (1.10)	1.95 (1.09)	1.99 (1.12)	1.92 (1.09)**
Negative Affect	3.63 (1.26)	3.58 (1.31)	3.40 (1.35)	3.28 (1.41)	3.12 (1.43)**
Compassion	5.14 (1.35)	5 (1.41)	4.80 (1.46)	4.64 (1.49)	4.52 (1.53)**
Worries Virus	4.31 (1.17)	4.17 (1.18)	3.96 (1.20)	3.91 (1.22)	3.75 (1.25)**
Worries Measures	3.21 (0.99)	3.15 (1.01)	3.06 (1.02)	2.99 (1.03)	2.84 (1.05)**
Vaccination Intention	5.84 (1.64)	5.75 (1.73)	5.58 (1.83)	5.50 (1.86)	5.38 (1.95)**
Measures Support	5.74 (1.03)	5.59 (1.06)	5.51 (1.16)	5.36 (1.21)	5.40 (1.27)**
Measures Ease	3.72 (1.29)	3.81 (1.28)	3.85 (1.28)	3.84 (1.28)	3.92 (1.32)**
Social Norm	5.14 (1.06)	4.85 (1.10)	4.77 (1.15)	4.55 (1.23)	4.38 (1.27)**
Control Infection	4.52 (1.07)	4.50 (1.04)	4.51 (1.02)	4.52 (1.01)	4.57 (1.05)*
Self-efficacy	5.61 (1.18)	5.59 (1.13)	5.60 (1.12)	5.55 (1.12)	5.59 (1.11)*
Involvement	5.07 (1.15)	4.89 (1.19)	4.76 (1.23)	4.56 (1.27)	4.55 (1.29)**
Perceived Knowledge	4.77 (1.17)	4.75 (1.15)	4.76 (1.12)	4.78 (1.11)	4.80 (1.15)*
Trust	5.63 (1.07)	5.50 (1.08)	5.52 (1.12)	5.42 (1.15)	5.50 (1.16)**
Health General	5.30 (1.28)	5.28 (1.27)	5.25 (1.24)	5.24 (1.25)	5.23 (1.25)**
Health change Physical	4.14 (0.89)	4.04 (0.82)	4.01 (0.82)	4.01 (0.81)	4 (0.84)**
Health change Mental	3.88 (0.89)	3.87 (0.87)	3.87 (0.86)	3.89 (0.84)	3.94 (0.86)*
Healthy Lifestyle	3.86 (0.74)	3.85 (0.73)	3.89 (0.71)	3.89 (0.66)	3.91 (0.68)*
Mental Wellbeing~	24.45 (4.13)	24.64 (4.29)	24.78 (4.36)	24.92 (4.56)	25.26 (4.58)**
Loneliness~	2.32 (0.75)	2.30 (0.77)	2.26 (0.78)	2.22 (0.81)	2.16 (0.81)**
Somatic Complaints~	1.36 (0.54)	1.33 (0.52)	1.31 (0.53)	1.30 (0.54)	1.30 (0.54)**
Depressive Complaints~	1.90 (0.90)	1.86 (0.91)	1.80 (0.91)	1.77 (0.92)	1.69 (0.9)**
Anxiety Complaints~	1.77 (0.87)	1.73 (0.86)	1.67 (0.85)	1.65 (0.86)	1.58 (0.83)**

¹ 7-point Likert-scale, unless marked ~, indicating a 5-point Likert-scale. Please note that *Risk Perception* is the product of two items and *Mental Wellbeing* is the sum score of all items, after which raw scores were converted to metric scores.

* $p < .05$; ** $p < .001$; indicating a significant effect of time on the node. A detailed description of the repeated measures ANOVA results and post hoc analysis with a specification of which waves significantly differ is provided below.

Compliance

Mauchly's test indicated that the assumption of sphericity had been violated, $\chi^2(9) = 776.46$, $p < .001$. The lower-bound corrected results showed that there was a significant effect of time on *Compliance*, $F(1, 2448) = 553.41$, $p < .001$, $\eta^2 = .184$. Post hoc analysis showed that with the exception of T1 with T2 ($p = 1.00$), all pairwise comparisons differed significantly, $p < .001$. These results suggest that the degree to which participants indicated to comply with the behavioural measures decreased over time.

Risk Perception

Mauchly's test indicated a violation of the sphericity assumption, $\chi^2(9) = 330.08, p < .001$. There was a statistically significant effect of time on *Risk Perception*, $F(1, 2448) = 75.75, p < .001, \eta^2 = .030$. Post hoc analysis showed that with the exception of T3 with T4 ($p = .084$), all pairwise comparisons differed significantly (p -values $< .001$, with one exception of $p = .037$). These results suggest that participants' perception of the risk of getting infected with the coronavirus decreased over time.

Health Risk

Mauchly's test indicated that the assumption of sphericity had been violated, $\chi^2(9) = 173.94, p < .001$. The results showed that there was a significant effect of time on *Health Risk*, $F(1, 2398) = 17.37, p < .001, \eta^2 = .007$. Post hoc analysis showed that with the exception of T1 with T2 ($p = 1.00$) and T3 with T4 ($p = 1.00$), all pairwise comparisons differed significantly (p -value ranges from $< .001$ to $.039$). These results suggest that participants' perception of the health risk in case of an infection with the coronavirus decreased over time.

Economic Consequences

Mauchly's test indicated a violation of the sphericity assumption, $\chi^2(9) = 150.09, p < .001$. There was a statistically significant effect of time on *Economic Consequences*, $F(1, 2398) = 122.58, p < .001, \eta^2 = .049$. Post hoc analysis showed that with the exception of T2 with T3 ($p = .52$), all pairwise comparisons differed significantly (p -value ranges from $< .001$ to $.019$). These results suggest that participants' view on the economic consequences of the pandemic decreased over time.

Self-exempting Beliefs

Mauchly's test indicated that the assumption of sphericity had been violated, $\chi^2(9) = 101.68, p < .001$. The lower-bound corrected results showed that there was a significant effect of time on *Self-exempting Beliefs*, $F(1, 2448) = 14.63, p < .001, \eta^2 = .006$. Post hoc analysis showed that the following measurements differed significantly: T1 with T2 ($p = .031$), T1 with T3 ($p < .001$), T1 with T4 ($p = .007$), T1 with T5 ($p < .001$), T2 with T5 ($p = .001$) and T4 with T5 ($p = .001$). These results suggest that participants' self-exempting beliefs decreased over time.

Negative Affect

Mauchly's test indicated a violation of the sphericity assumption, $\chi^2(9) = 523.03, p < .001$. There was a statistically significant effect of time on *Negative Affect*, $F(1, 2398) = 259.65, p < .001, \eta^2 = .098$. Post hoc analysis showed that with the exception of T1 with T2 ($p = .072$), all pairwise comparisons differed significantly, $p < .001$. These results suggest that the degree to which participants felt that the pandemic caused negative emotions decreased over time.

Compassion

Mauchly's test indicated that the assumption of sphericity had been violated, $\chi^2(9) = 108.13, p < .001$. The lower-bound corrected results showed that there was a significant effect of time on *Compassion*, $F(1, 2448) = 190.71, p < .001, \eta^2 = .072$. Post hoc analysis showed that all pairwise comparisons differed significantly, $p < .001$. These results suggest that the degree to which the pandemic makes participants feel compassionate decreased over time.

Worries Virus

Mauchly's test indicated a violation of the sphericity assumption, $\chi^2(9) = 293.06, p < .001$. There was a statistically significant effect of time on *Worries Virus*, $F(1, 2398) = 328.16, p < .001, \eta^2 = .120$. Post hoc analysis showed that all pairwise comparisons differed significantly (p -value ranges from $< .001$ to $.002$). These results suggest that the degree to which participants worry about events related to the coronavirus (i.e. getting infected or infecting others, losing a loved one or the health care system getting overloaded) decreased over time.

Worries Measures

Mauchly's test indicated that the assumption of sphericity had been violated, $\chi^2(9) = 304.92, p < .001$. The lower-bound corrected results showed that there was a significant effect of time on *Worries Measures*, $F(1, 2448) = 231.89, p < .001, \eta^2 = .087$. Post hoc analysis showed that all pairwise comparisons differed significantly, $p < .001$. These results suggest that the degree to which participants worry about events related to the corona measures (e.g., health effects of staying home, schools closing, a recession) decreased over time.

Vaccination Intention

Mauchly's test indicated a violation of the sphericity assumption, $\chi^2(9) = 883.50, p < .001$. There was a statistically significant effect of time on *Vaccination Intention*, $F(1, 2448) = 118.27, p < .001, \eta^2 = .046$. Post hoc analysis showed that all pairwise comparisons differed significantly (p -values $< .001$, with one exception of $p = .009$). These results suggest that participants intention to get vaccinated (if available) decreased over time.

Measures Support

Mauchly's test indicated that the assumption of sphericity had been violated, $\chi^2(9) = 507.60, p < .001$. The lower-bound corrected results showed that there was a significant effect of time on *Measures Support*, $F(1, 2448) = 172.94, p < .001, \eta^2 = .066$. Post hoc analysis showed that all

pairwise comparisons differed significantly, $p < .001$. These results suggest that participants' support for the corona measures decreased over time.

Measures Ease

Mauchly's test indicated a violation of the sphericity assumption, $\chi^2(9) = 294.16, p < .001$. There was a statistically significant effect of time on *Measures Ease*, $F(1, 2398) = 24.25, p < .001, \eta^2 = .010$. Post hoc analysis showed that with the exception of T2 with T3 ($p = .429$), T2 with T4 ($p = 1.00$) and T3 with T4 ($p = 1.00$), all pairwise comparisons differed significantly (p -value ranges from $< .001$ to $.004$). These results suggest that participants' perceived ease of the corona measures increased over time (i.e. participants experienced the corona measures as easier at the end of the data collection compared to the first survey).

Social Norm

Mauchly's test indicated that the assumption of sphericity had been violated, $\chi^2(9) = 102.79, p < .001$. The lower-bound corrected results showed that there was a significant effect of time on *Social Norm*, $F(1, 2448) = 298.08, p < .001, \eta^2 = .109$. Post hoc analysis showed that all pairwise comparisons differed significantly (p -values $< .001$, with one exception of $p = .022$). These results suggest that the degree to which participants think other people (find it important to) comply with the corona measures decreased over time.

Control Infection

Mauchly's test indicated a violation of the sphericity assumption, $\chi^2(9) = 73.46, p < .001$. There was a statistically significant effect of time on *Control Infection*, $F(1, 2448) = 3.88, p = .049, \eta^2 = .002$. Post hoc analysis showed that T2 differed significantly from T5 ($p = .010$), T3 differed significantly from T5 ($p = .009$) and T4 differed significantly from T5 ($p = .022$). The remaining pairwise comparisons did not differ significantly (p -value ranges from $.536$ to 1.00). These result suggest an increase in participants' perceived control to avoid an infection with the coronavirus between the middle and last surveys.

Self-efficacy

Mauchly's test indicated that the assumption of sphericity had been violated, $\chi^2(9) = 147.67, p < .001$. The lower-bound corrected results showed that there was a significant effect of time on *Self-efficacy*, $F(1, 2448) = 5.43, p = .020, \eta^2 = .002$. Post hoc analysis showed that the following measurements differed significantly: T1 with T4 ($p < .001$), T2 with T4 ($p = .035$) and T3 with T4 ($p = .047$). The remaining pairwise comparisons did not differ significantly (p -value ranges from $.202$ to 1.00). These result suggest a decrease over time in participants' perceived ability to protect themselves, more specifically between the first three waves and the forth wave.

Involvement

Mauchly's test indicated a violation of the sphericity assumption, $\chi^2(9) = 368.75, p < .001$. There was a statistically significant effect of time on *Involvement*, $F(1, 2448) = 280.31, p < .001, \eta^2 = .103$. Post hoc analysis showed that with the exception of T4 with T5 ($p = 1.00$), all pairwise comparisons differed significantly ($p < .001$). These results suggest that the degree to which participants are involved in the corona pandemic decreased over time.

Perceived Knowledge

Mauchly's test indicated that the assumption of sphericity had been violated, $\chi^2(9) = 157.43, p < .001$. Since sphericity is violated ($\epsilon = .97$) and the lower-bound corrected were not significant, Huynh-Feldt corrected results are reported⁶. The results showed that there was a significant effect of time on *Perceived Knowledge*, $F(3.88, 9487.88) = 3.23, p = .013, \eta^2 = .001$. Post hoc analysis showed that T2 and T3 differed significantly from T5 ($p = .016$ and $p = .023$, respectively). The remaining pairwise comparisons did not differ significantly (p -value ranges from .457 to 1.00). These results suggest an increase over time in participants' perceived knowledge about the corona pandemic.

Trust

Mauchly's test indicated a violation of the sphericity assumption, $\chi^2(9) = 198.43, p < .001$. There was a statistically significant effect of time on *Trust*, $F(1, 2448) = 49.49, p < .001, \eta^2 = .020$. Post hoc analysis showed that with the exception of T2 with T3 ($p = .458$), T2 with T5 ($p = .589$) and T3 with T5 ($p = 1.00$), all pairwise comparisons differed significantly ($p < .001$). These results suggest an initial decrease over time in participants' trust in the actors relevant for the pandemic, followed by an increase between the last two surveys.

Health General

Mauchly's test indicated that the assumption of sphericity had been violated, $\chi^2(9) = 193.47, p < .001$. The lower-bound corrected results showed that there was a significant effect of time on *Health General*, $F(1, 2448) = 12.54, p < .001, \eta^2 = .005$. Post hoc analysis showed that the following pairwise comparisons differed significantly: T1 with T3 ($p < .001$), T1 with T4 ($p < .001$), T1 with T5 ($p < .001$), T2 with T3 ($p = .040$), T2 with T4 ($p = .003$) and T2 with T5 ($p = .001$). The remaining pairwise comparisons did not differ significantly (p -value ranges from .925 to 1.00). These results suggest that participants' rating of their health in general decreased over time.

⁶ We report the Huynh-Feldt corrected results given that epsilon was greater than .75.

Health change Physical

Mauchly's test indicated a violation of the sphericity assumption, $\chi^2(9) = 441.08, p < .001$. There was a statistically significant effect of time on *Health change Physical*, $F(1, 2448) = 20.65, p < .001, \eta^2 = .008$. Post hoc analysis showed that T1 differed significantly from all subsequent waves ($p < .001$), whereas the remaining pairwise comparisons did not differ significantly (p -value ranges from .126 to 1.00). These results suggest that participants initially experienced an improvement in their physical health compared to before the pandemic, but this improvement disappeared after the first survey. From there on participants viewed their physical health as comparable to their physical health before the pandemic.

Health change Mental

Mauchly's test indicated that the assumption of sphericity had been violated, $\chi^2(9) = 378.64, p < .001$. The lower-bound corrected results showed that there was a significant effect of time on *Health change Mental*, $F(1, 2448) = 5.99, p = .014, \eta^2 = .002$. Post hoc analysis showed that all waves differed significantly from T5 (p -value ranges from $< .001$ to .018), whereas the remaining pairwise comparisons did not differ significantly ($p = 1.00$). These results suggest that participants initially experienced a deterioration in their mental health compared to before the pandemic. Although they reported a deterioration in all surveys, the experienced deterioration in mental health compared to before the pandemic decreased in the last survey.

Healthy Lifestyle

Mauchly's test indicated a violation of the sphericity assumption, $\chi^2(9) = 517.18, p < .001$. There was a statistically significant effect of time on *Healthy Lifestyle*, $F(1, 2448) = 8.86, p = .003, \eta^2 = .004$. Post hoc analysis showed that the following pairwise comparisons differed significantly: T1 with T4 ($p = .032$), T1 with T5 ($p = .013$), T2 with T3 ($p = .008$), T2 with T4 ($p < .001$) and T2 with T5 ($p < .001$). The remaining pairwise comparisons did not differ significantly (p -value ranges from .493 to 1.00). These results suggest that participants initially experienced a decrease in their healthy lifestyle compared to before the pandemic. Although they reported a decrease in all surveys, the experienced decrease in healthy lifestyle compared to before the pandemic lessened over time.

Mental Wellbeing

Mauchly's test indicated that the assumption of sphericity had been violated, $\chi^2(9) = 137.26, p < .001$. The lower-bound corrected results showed that there was a significant effect of time on *Mental Wellbeing*, $F(1, 2448) = 37.10, p < .001, \eta^2 = .015$. Post hoc analysis showed that with the exception of T1 and T2 ($p = .265$), T2 and T3 ($p = .299$) and T3 with T4 ($p = .666$), all pairwise comparisons differed significantly (p -value ranges from $< .001$ to .001). These results suggest that participants' mental wellbeing improved between the start and the end of the study.

Loneliness

Mauchly's test indicated a violation of the sphericity assumption, $\chi^2(9) = 246.19, p < .001$. There was a statistically significant effect of time on *Loneliness*, $F(1, 2448) = 100.32, p < .001, \eta^2 = .039$. Post hoc analysis showed that with the exception of T1 with T2 ($p = .054$), all pairwise comparisons differed significantly ($p < .001$). These results suggest that the degree to which participants felt lonely decreased over time.

Somatic Complaints

Mauchly's test indicated that the assumption of sphericity had been violated, $\chi^2(9) = 388.48, p < .001$. The lower-bound corrected results showed that there was a significant effect of time on *Somatic Complaints*, $F(1, 2448) = 26.80, p < .001, \eta^2 = .011$. Post hoc analysis showed that T1 differed significantly from all subsequent waves ($p < .001$). Furthermore, the pairwise comparisons T2 with T4 and T2 with T5 differed significantly ($p = .036$ and $p = .008$, respectively), and T3 differed from T5 ($p = .005$). The remaining pairwise comparisons did not differ significantly (p -value ranges from .069 to 1.00). These results suggest that the degree to which participants experienced somatic complaints decreased over time.

Depressive Complaints

Mauchly's test indicated a violation of the sphericity assumption, $\chi^2(9) = 511.15, p < .001$. There was a statistically significant effect of time on *Depressive Complaints*, $F(1, 2448) = 142.73, p < .001, \eta^2 = .055$. Post hoc analysis showed that all pairwise comparisons differed significantly (p -value ranges from $< .001$ to .002). These results suggest a decrease over time in the degree to which participants experienced depressive complaints.

Anxiety Complaints

Mauchly's test indicated that the assumption of sphericity had been violated, $\chi^2(9) = 419.64, p < .001$. The lower-bound corrected results showed that there was a significant effect of time on *Anxiety Complaints*, $F(1, 2448) = 121.32, p < .001, \eta^2 = .047$. Post hoc analysis showed that all pairwise comparisons differed significantly (p -value ranges from $< .001$ to .011). These results suggest that the degree to which participants experienced anxiety complaints decreased over time.

2.2 Network analysis model fit evaluation

The values provided below present the model fit of the *dlvm1* model of the *psychonetrics* package. We used the *dlvm1* model of the *psychonetrics* package to estimate the model. We specified within latent (ggm [Gaussian graphical model]), between latent (chol [Cholesky]), within and between residuals (cov [covariance]), and estimator (FIML). Overall fit, including the temporal and contemporaneous networks, showed excellent fit according to RMSEA (root mean square error of approximation = .023 [95% CI 0.023–0.024]) and CFI (comparative fit index = .96), although the standard chi-square test, not accounting for the freedom, rejected the model, $\chi^2(7803) = 17930.86$, $p < .001$. See table on the next page for additional measures.

Measure	Value
logl	-373328.903
unrestricted.logl	-364363.473
baseline.logl	-499487.446
nvar	135
nobs	9315
npar	1512
df	7803
objective	63.124
chisq	17930.859
pvalue	0.000
baseline.chisq	270247.946
baseline.df	9045
baseline.pvalue	0.000
nfi	0.934
pnfi	0.805
tli	0.955
nnfi	0.955
rfi	0.923
ifi	0.961
rni	0.961
cfi	0.961
rmsea	0.023
rmsea.ci.lower	0.023
rmsea.ci.upper	0.024
rmsea.pvalue	1.000
aic.ll	749681.805
aic.ll2	754845.814
aic.x	2324.859
aic.x2	20954.859
bic	758425.410
bic2	753621.452
ebic.25	765842.185
ebic.5	773258.961
ebic.75	779192.381
ebic1	788092.512

2.3 COVID-19 broad attitude networks including all edges

Temporal

Contemporaneous

PKnow: Perceived Knowledge
 Compas: Compassion
 S_Eff: Self-efficacy
 Vacc: Vaccine Intention
 H_Gen: Health General
 Hc_Phy: Health change Physical
 Hc_Ment: Health change Mental
 RiskP: Risk Perception
 H_Risk: Health Risk
 S_Exem: Self-exempting Beliefs
 EconCon: Economic Consequences
 Invol: Involvement
 NegAff: Negative Affect
 Vwor_Vir: Worries Virus
 Wor_Meas: Worries Measures
 Trust: Trust
 Norm: Social norm
 MeasSup: Measures Support
 MeasEas: Measures Ease
 ContrInf: Control Infection
 Comply: Compliance
 Somat: Somatic Complaints
 LStyl: Healthy Lifestyle
 Depres: Depressive Complaints
 Anxiet: Anxiety Complaints
 Lone: Loneliness
 MWellb: Mental Wellbeing

2.4 Edge weights and node strength of COVID-19 broad attitude networks

Temporal COVID-19 network

Read rows (first column) as node from which the edge originates. Please note that the value .00 in the table does not indicate an edge was absent, because these values were rounded to two decimals.

From \ To	Perceived knowledge	Compassion	Self-efficacy	Vaccine Intention	Health General	Health change Physical	Health change Mental	Risk Perception	Health Risk	Self-exempting Beliefs	Economic Consequences	Involvement	Negative Affect	Worries Virus	Worries Measures	Trust	Social norm	Measures Support	Measures Ease	Control Infection	Compliance	Somatic Complaints	Healthy Lifestyle	Depressive Complaints	Anxiety Complaints	Loneliness	Mental Well-being
Perceived knowledge	.09	.00	.02	-.01	.01	.01	-.01	-.01	-.02	-.02	-.04	-.02	-.02	-.03	-.02	.01	-.04	.01	.03	.02	-.01	.00	.02	-.02	-.03	-.01	.01
Compassion	-.01	.10	.00	.00	.01	.00	-.02	.00	.01	.01	.02	.05	.05	.03	.01	.01	.04	.01	-.02	.02	.04	-.02	-.01	.01	.01	.00	-.02
Self-efficacy	.02	-.02	.08	-.02	-.02	-.01	.01	-.02	.00	.04	-.01	-.03	.02	-.01	-.01	.01	-.01	-.01	.01	.01	-.03	-.01	.00	.00	.01	.01	-.01
Vaccine Intention	.00	.05	.02	.41	-.01	.01	.00	.04	.04	-.01	.02	.06	.03	.06	.01	.02	.04	.08	.01	.00	.06	.01	-.01	.01	.03	.00	.01
Health General	-.01	.01	-.01	.01	.15	.03	.00	.01	.01	.00	.03	.02	.02	.01	.02	.02	.03	.00	-.02	-.01	.00	.00	.01	.01	.00	.02	.00
Health change Physical	.02	.02	.04	.02	.05	.14	.08	.01	.01	.02	.02	.02	.03	.02	.00	-.01	.02	.04	.00	.01	.02	-.02	.04	-.01	.00	.00	-.01
Health change Mental	-.04	.00	-.01	.00	.01	.06	.10	.00	.02	.00	.00	-.01	-.02	.01	-.01	.00	.00	-.01	-.01	.00	-.01	.03	.01	-.05	-.03	-.02	.02
Risk Perception	.01	.02	-.01	.00	-.01	.02	-.01	.11	.04	-.02	.00	.03	.01	.02	.02	.00	.02	.00	-.02	-.02	.01	.03	.03	.00	.00	.02	.00
Health Risk	-.03	.00	.01	-.02	-.01	-.01	.02	.04	.09	-.02	.02	-.02	-.02	.02	-.02	.02	-.03	.02	.03	-.01	.00	-.01	-.02	.00	-.01	-.02	.00
Self-exempting Beliefs	-.02	.01	.02	.03	-.01	.01	.01	-.03	-.02	.06	.01	.04	.00	.00	.01	.01	.02	.02	.01	.02	.06	-.02	-.01	-.03	-.03	.00	.01
Economic Consequences	.00	.00	.03	.01	.03	.02	-.01	.00	.03	.01	.10	.00	.01	.02	.03	-.01	-.01	.00	-.01	.02	.02	-.02	-.02	-.02	.00	.00	-.03
Involvement	.03	.09	.00	.07	.01	.01	-.04	.06	.03	.01	.02	.21	.06	.11	.07	.00	.10	.09	-.01	.00	.10	.01	-.02	.05	.05	.02	-.04
Negative Affect	-.03	.04	.00	-.01	.02	.01	-.01	.00	.00	.02	.00	.03	.19	.03	.05	-.04	-.02	-.06	-.05	-.02	-.03	.00	.02	.03	.01	.02	-.01
Worries Virus	.00	.07	.00	.02	-.02	-.01	.00	.06	.07	.00	.04	.08	.08	.13	.03	.03	.03	.04	.00	-.01	.06	.01	-.02	.04	.03	.04	-.01
Worries Measures	.01	.01	-.02	.00	-.02	-.05	.00	.03	-.02	.01	.06	.02	.02	.04	.12	.01	.01	.00	-.02	.00	.01	.01	-.02	.02	.00	.05	-.02
Trust	.02	-.01	.00	-.03	.02	-.02	.02	.01	.00	-.01	-.03	-.02	-.03	.00	-.01	.11	-.03	.04	.00	.01	.01	-.02	.01	.00	.00	.00	-.02
Social norm	.00	.07	.02	.03	-.01	.01	-.01	.04	.02	.02	.04	.06	.06	.08	.05	.01	.16	.04	-.02	-.03	.07	.01	-.01	.02	.03	.02	-.02
Measures Support	-.01	.05	.02	.07	.04	.02	.00	.04	.04	-.02	.03	.08	.02	.08	.03	.11	.08	.25	.06	-.02	.13	.06	.00	.02	.03	.01	.01
Measures Ease	.03	-.01	.02	-.01	-.02	.02	.02	.00	.02	-.01	-.02	.00	-.04	-.02	-.03	.01	-.01	.02	.16	.01	.00	-.01	.00	-.02	-.03	-.01	.01
Control Infection	.00	.00	.02	.01	.01	-.02	.00	-.03	-.02	.02	.01	-.01	-.01	-.01	.00	.00	.00	-.03	-.01	.04	-.03	-.03	.00	.01	.01	.00	.03
Compliance	.00	.07	.03	.06	.00	.01	-.01	.02	.02	.00	.04	.08	.03	.07	.06	.06	.12	.11	.00	.04	.37	.01	.01	.03	.03	.04	-.01
Somatic Complaints	-.01	-.02	-.01	.01	-.03	-.03	.03	.02	.01	.02	.00	-.01	.01	.01	.01	-.02	.00	.02	.00	-.02	-.01	.21	-.04	.03	.04	.02	-.01
Healthy Lifestyle	.02	.02	.02	.00	.02	.07	.03	.01	.00	.01	-.02	.00	.02	-.01	-.01	.00	-.01	-.01	.00	.00	.02	-.02	.24	.04	.01	.03	.02
Depressive Complaints	.00	.01	.03	.01	.02	.00	-.07	.01	.01	-.02	.02	.03	.07	.06	.05	.00	-.01	.01	-.01	.00	-.01	.02	.00	.18	.07	.09	-.05
Anxiety Complaints	.00	.01	-.03	.03	-.03	.01	-.03	.02	.02	-.02	.02	.06	.04	.04	.04	.02	.03	.03	.00	-.03	.04	.05	-.02	.10	.20	.02	-.05
Loneliness	-.02	-.01	-.01	.00	.00	.03	-.01	.01	.01	-.01	.02	.01	.06	.02	.07	-.02	.00	.00	-.03	-.01	.03	.03	.01	.06	.03	.14	-.04
Mental Well-being	.00	-.03	.00	.01	-.01	.00	.01	.01	-.02	.01	-.01	-.03	-.03	-.02	-.03	.01	-.05	.00	-.02	.01	-.03	.02	.00	-.02	.00	-.02	.09

Contemporaneous COVID-19 network

Contemporaneous COVID-19 network	Perceived knowledge	Compassion	Self-efficacy	Vaccine Intention	Health General	Health change Physical	Health change Mental	Risk Perception	Health Risk	Self-exempting Beliefs	Economic Consequences	Involvement	Negative Affect	Worries Virus	Worries Measures	Trust	Social norm	Measures Support	Measures Ease	Control Infection	Compliance	Somatic Complaints	Healthy Lifestyle	Depressive Complaints	Anxiety Complaints	Loneliness	Mental Wellbeing
Perceived knowledge	.01	.05	-.02	.02	.01	-.02	.01	.02	.04	-.01	.17	-.02	-.01	.00	.04	-.02	.00	.01	.02	-.01	.00	.00	-.02	-.01	-.03	.03	
Compassion		.01	.00	.03	.02	.02	-.01	.00	.00	.01	.10	.18	.08	.02	.04	.06	.05	-.01	.00	.06	-.01	-.02	.00	.00	-.02	.04	
Self-efficacy			.04	.03	.00	-.01	-.04	.00	.01	.00	.02	.01	-.01	-.03	.04	.01	.07	.02	.09	.03	-.02	.00	.00	-.01	.00	.02	
Vaccine Intention				.02	.02	-.01	.02	.02	.00	.00	.06	.02	.06	-.01	.05	.02	.10	.00	-.01	.06	.03	.03	.00	.00	.00	-.01	
Health General					.13	.02	.00	-.03	.01	.01	-.01	.01	.00	-.01	.03	.01	.01	.00	.05	.04	-.07	.03	.00	-.02	.01	.05	
Health change Physical						.19	-.01	-.02	.03	.02	.02	.01	.01	-.02	-.01	.04	.02	.04	.00	-.02	-.06	.16	.00	.02	.04	.00	
Health change Mental							.00	.01	.00	.01	-.03	-.03	.00	.02	.02	.00	.01	.03	-.01	-.01	-.02	.11	-.13	-.07	-.04	.06	
Risk Perception								.28	-.03	.05	.07	.05	.11	.02	-.01	.01	.04	.01	-.03	.01	.04	.03	.01	-.01	.01	.00	
Health Risk									-.05	.06	.00	.03	.13	-.02	.03	-.02	.00	.04	-.02	.01	.02	-.03	-.02	.02	.01	.00	
Self-exempting Beliefs										.02	-.02	.03	-.02	.04	-.03	.04	-.03	.01	.05	-.01	.02	.01	-.01	.01	-.01	-.01	
Economic Consequences											.06	.06	.04	.13	-.04	.00	.02	-.03	.02	.02	-.01	.01	-.01	.01	.02	-.01	
Involvement												.16	.12	.02	.04	.06	.10	-.03	.00	.09	-.01	-.01	.03	.02	-.01	-.02	
Negative Affect													.15	.19	-.04	-.01	-.07	-.05	-.01	.01	.02	.02	.09	.09	.04	-.04	
Worries Virus														.24	.02	.04	.08	-.03	-.04	.05	.02	-.01	.01	.02	-.01	-.01	
Worries Measures															.00	.02	-.01	-.04	.00	.00	.04	-.04	.09	-.01	.09	-.03	
Trust																	.09	.26	-.02	.04	.03	-.01	.01	-.02	-.01	-.01	.04
Social norm																		.12	-.02	.01	.13	.02	.00	.01	.01	.02	-.02
Measures Support																			.20	.00	.22	.02	.00	.00	.01	-.01	.01
Measures Ease																				.11	.01	.03	.01	-.02	-.03	-.02	.01
Control Infection																					.05	.01	.01	.02	-.01	-.02	.03
Compliance																						.00	.02	.03	.01	.02	.03
Somatic Complaints																							-.03	.07	.16	-.02	-.02
Healthy Lifestyle																								-.01	-.03	.01	.02
Depressive Complaints																									.34	.19	-.10
Anxiety Complaints																										.08	-.09
Loneliness																											-.15
Mental Wellbeing																											

Node strength

Node	Temporal network		Contemporaneous network
	InStrength	OutStrength	Strength
Perceived Knowledge	-1.23	-0.5	-1.19
Compassion	0.56	-0.55	-0.33
Self-efficacy	-0.96	-0.79	-1.25
Vaccine Intention	-0.32	0.42	-1.07
Health General	-0.95	-1.04	-1
Health change Physical	-0.32	-0.07	-0.04
Health change Mental	-0.66	-0.88	-0.07
Risk Perception	-0.14	-0.75	0.04
Health Risk	-0.29	-0.52	-0.09
Self-exempting Beliefs	-1.05	-0.47	-1.38
Economic Consequences	-0.1	-0.78	-0.85
Involvement	1.87	2.48	1.36
Negative Affect	1.69	0.08	1.96
Worries Virus	1.72	1.05	1.56
Worries Measures	0.97	-0.22	1.01
Trust	-0.49	-0.78	0.33
Social norm	1.21	1.05	-0.35
Measures Support	0.97	2.26	2.16
Measures Ease	-0.89	-0.61	-0.27
Control Infection	-1.23	-0.97	-0.97
Compliance	1.75	1.79	0.32
Somatic Complaints	-0.51	-0.46	-0.6
Healthy Lifestyle	-1.19	-0.6	-0.9
Depressive Complaints	0.47	0.66	1.14
Anxiety Complaints	-0.15	0.96	0.7
Loneliness	-0.21	-0.02	-0.1
Mental Wellbeing	-0.5	-0.76	-0.15

2.5 Edge accuracy of COVID-19 broad attitude networks

The figures below present the edge bootstrap confidence intervals, which are an indication of edge accuracy. Generally speaking, the confidence intervals of the edge weights were not wide, indicating stable (reliable) edges.

Edge accuracy temporal COVID-19 network

Please note that the edges in the temporal COVID-19 network presented in the manuscript represent partial directed correlations (PDC), which are *standardized* beta values. The figure presented here contains the beta values and their confidence intervals of the edges of the temporal network.

Edge accuracy contemporaneous COVID-19 network

2.6 Coherence within-person COVID-19 broad attitude networks

Plot of relation between edges of the temporal and contemporaneous COVID-19 networks ($r = .47, z = 0.51$). A Fisher's r -to- z transformation was included to approximate normal distribution.

3. Timeline COVID-19 in the Netherlands

Date	NL/international	Theme (statistics/safety measures/news/research)	Description of the event	Source
31-12-2019	International	Statistics	WHO is informed of cases of pneumonia of unknown cause detected in the city of Wuhan, in the Hubei Province, China.	WHO: http://www.euro.who.int/en/health-topics/health-emergencies/international-health-regulations/news/news/2020/2/novel-coronavirus-emerges-in-china
10-1-2020	International	Statistics	The WHO reports that a novel coronavirus has emerged in China. After the first reports of pneumonia came in China came in, it has been confirmed by Chinese authorities that this is caused by a novel coronavirus.	WHO: http://www.euro.who.int/en/health-topics/health-emergencies/international-health-regulations/news/news/2020/2/novel-coronavirus-emerges-in-china
30-1-2020	International	Safety measures	The WHO declares the coronavirus outbreak as a global emergency, demanding temporary safety measures to be made in all countries. Countries have to prove their healthcare safety measures are based on sound scientific reasoning. According to the WHO, declaring an international emergency helps strengthen global coordination, transparency and increases support for countries with less developed healthcare systems.	WHO: http://www.euro.who.int/en/health-topics/health-emergencies/international-health-regulations/news/news/2020/2/2019-ncov-outbreak-is-an-emergency-of-international-concern
31-1-2020	International	Statistics	First case of coronavirus confirmed in Spain.	CNN: https://edition.cnn.com/asia/live-news/coronavirus-outbreak-02-01-20-intl-hnk/h_afcf3a4665521aab11c721c8cc80dd03 ; El Pais: https://elpais.com/sociedad/2020/01/31/actualidad/1580509404_469734.html
2-2-2020	NL	News	Flight with over 15 Dutch people returns from Wuhan after waiting several days to be able to return.	Volkskrant: https://www.volkskrant.nl/nieuws-achtergrond/verslag-van-de-evacuatie-uit-wuhan-bij-de-douane-loop-je-een-rampenfilm-binnen~bb1dba1b/ ; Telegraaf: https://www.telegraaf.nl/nieuws/455667699/nederlanders-uit-coronastad-wuhan-moeten-eerst-tot-rust-komen

7-2-2020	International	News	Dr Wenliang, a Chinese doctor who tried to warn people about the possible spread of the virus, died from the coronavirus.	BBC: https://www.bbc.com/news/world-asia-china-51403795 ; The Guardian: https://www.theguardian.com/global-development/2020/feb/07/coronavirus-chinese-rage-death-whistleblower-doctor-li-wenliang
11-2-2020	International	News	The WHO announces official names for the novel coronavirus disease (COVID-19) and the virus that causes it (SARS-CoV-2).	WHO: https://www.who.int/emergencies/diseases/novel-coronavirus-2019/technical-guidance/naming-the-coronavirus-disease-(covid-2019)-and-the-virus-that-causes-it
23-2-2020	NL	Safety measures	After the virus starts to spread in other European countries like Italy, the Dutch government is not yet implementing extra safety measures in the Netherlands. Meanwhile, the Dutch Carnival has started in most southern parts of the Netherlands, with large crowds of people coming together to celebrate.	Volkskrant: https://www.volkskrant.nl/nieuws-achtergrond/uitbraak-coronavirus-in-noord-italie-geen-extra-maatregelen-in-nederland~b6518425/
27-2-2020	NL	Statistics	A first case of the new coronavirus has been confirmed by the National Institute for Public Health and Environment (RIVM). The patient is a 56 year old man living in Brabant, a southern province of the Netherlands. He most likely contracted the virus on a trip to the Lombardy region in Italy.	RIVM: https://www.rivm.nl/nieuws/patient-met-nieuw-coronavirus-in-nederland ; Volkskrant: https://www.volkskrant.nl/nieuws-achtergrond/burgemeester-tilburg-coronapatient-tilburg-was-niet-besmettelijk-tijdens-carnaval~b5510762/ ; Telegraaf: https://www.telegraaf.nl/nieuws/214089776/rivm-eerste-coronageval-in-nederland
28-2-2020	NL	Statistics	A second case of the coronavirus has been confirmed by the RIVM. The patient is a woman living in Amsterdam, who returned from a trip to the Lombardy region in Italy	RIVM: https://www.rivm.nl/nieuws/tweede-patient-in-nederland-met-covid-19 ; WHO: https://covid19.who.int/region/euro/country/nl
4-3-2020	NL	Safety measures	The RIVM advises for "only necessary travels" to all of Northern Italy. Many big companies in the Netherlands start to take measures against the virus, even though this is not yet required by authorities.	RIVM: https://www.rivm.nl/coronavirus-covid-19/actueel ; Volkskrant: https://www.volkskrant.nl/nieuws-achtergrond/veel-thuiswerken-en-niet-meer-reizen-internationale-bedrijven-nemen-maatregelen-tegen-corona~b773241c/ ;
4-3-2020	International	Safety measures	Italy announces it is closing down schools and universities nationwide until mid-march.	The Guardian https://www.theguardian.com/world/2020/mar/04/italy-orders-closure-of-schools-and-universities-due-to

				coronavirus; Independent: https://www.independent.co.uk/news/world/europe/coronavirus-italy-school-university-closed-outbreak-death-toll-health-latest-a9374801.html
5-3-2020	NL	Statistics	The number of positive tested patients in the Netherlands reportedly grew with 41, more than doubling the total amount to 82 patients. According to Bruno Bruins, minister of medical care, this number is due to a catch-up in conducted tests.	WHO: https://covid19.who.int/region/euro/country/nl ; RIVM: https://www.rivm.nl/coronavirus-covid-19/actueel ; AD: https://www.ad.nl/binnenland/aantal-coronapatienten-in-nederland-ruim-verdubbeld-naar-82~a886fdb8/?referrer=https://en.wikipedia.org/
6-3-2020	NL	Statistics	A first patient with COVID-19 dies in the Netherlands. The patient was an 86 year old man who lived in Hoeksche Waard, Zuid-Holland. Total cases grows to: 128	WHO: https://covid19.who.int/region/euro/country/nl ; RIVM: https://www.rivm.nl/nieuws/patient-met-nieuw-coronavirus-overleden ; Parool: https://www.parool.nl/nederland/eerste-dode-door-coronavirus-in-nederland~b8d72ed6/?referer=https%3A%2F%2Fnl.wikipedia.org%2F
6-3-2020	NL	Safety measures	RIVM advises people in Noord-Brabant to limit their social contacts in case of colds, coughs and fevers.	RIVM: https://www.rivm.nl/nieuws/covid-19-nieuwe-aanwijzing-voor-inwoners-noord-brabant
9-3-2020	International	Safety measures	Italy's prime minister Guiseppe Conte announces a nationwide forced quarantine on national television. Only people with a valid work or family reason are allowed to travel. All social events are cancelled.	BBC: https://www.bbc.com/news/world-europe-51810673 ; The Wall Street Journal: https://www.wsj.com/articles/italy-bolsters-quarantine-checks-after-initial-lockdown-confusion-11583756737
9-3-2020	NL	Safety measures	Prime minister Mark Rutte holds a nationally televised press conference with Jaap van Dissel, RIVM director of disease control, to inform people on the coronavirus. In this press conference, he advises people to stop shaking hands and to work from home if you live in Noord-Brabant. Earlier that day, the government also advised to regularly wash your hands, cough or sneeze in your elbow and to only make use of paper towels. After the press conference ended, the PM mistakenly shook hands with Jaap van Dissel.	NOS: https://nos.nl/video/2326490-rutte-stop-met-handen-schudden-en-werk-thuis.html ; Parool: https://www.parool.nl/nederland/rutte-reikt-rivm-de-hand-na-oproep-om-geen-handen-meer-te-schudden~ba6d39c9/ ; RIVM: https://www.rivm.nl/coronavirus-covid-19/actueel
11-3-2020	International	News	The WHO declares that COVID-19 can be characterized as a pandemic. This does not change the WHO's assessment of the threat posed by the virus, neither does it change what the WHO's actions are towards solving the situation. According to the WHO, the	WHO: https://www.who.int/dg/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19---11-march-2020

characterization as a pandemic serves to warn countries and to call for countries to 'take urgent and aggressive action'.

12-3-2020	NL	Statistics	The total number of patients that have tested positive has grown with 111, bringing the total number of cases to 614. The total number of patients that have died is 5.	WHO: https://covid19.who.int/region/euro/country/nl (retrieved on 29-06-2020)
12-3-2020	NL	Safety measures	The RIVM announces that the safety measures implemented in the province of Noord-Brabant now apply to the whole of the Netherlands. Events with over 100 people are banned at least until 31st of March, schools remain open.	RIVM: https://www.rivm.nl/nieuws/uitbreiding-maatregelen-coronavirus ; https://www.rijksoverheid.nl/onderwerpen/coronavirus-covid-19/nieuws/2020/03/12/nieuwe-maatregelen-tegen-verspreiding-coronavirus-in-nederland ; The Guardian: https://www.theguardian.com/world/2020/mar/14/spain-government-set-to-order-nationwide-coronavirus-lockdown ; Businessinsider: https://www.businessinsider.com/coronavirus-spain-nationalises-private-hospitals-emergency-covid-19-lockdown-2020-3?international=true&r=US&IR=T
14-3-2020	International	Safety measures	The Spanish government has formally declared a state of emergency over the coronavirus, placing the country in lockdown for the next weeks.	Government: https://www.rijksoverheid.nl/actueel/nieuws/2020/03/15/aanvullende-maatregelen-onderwijs-horeca-sport
15-3-2020	NL	Safety measures	In a press conference, minister Bruins and minister of education Arie Slob announces that schools are closed and teachers have to prepare remote teaching for the children at home. Restaurants, bars, sports clubs and more establishments have to close until April 8. People are asked to keep a distance of 1,5 meters to each other where possible.	WHO: https://covid19.who.int/region/euro/country/nl (retrieved on 29-06-2020)
16-3-2020	NL	Statistics	The total number of patients that have tested positive has grown with 278 and 4 patients have died. This brings the total number of cases to 1,413 and the number of patients that died to 24.	NOS: https://www.youtube.com/watch?v=D3QPmphUK_M ; https://nos.nl/artikel/2327309-de-komende-tijd-zal-een-groot-deel-van-de-bevolking-met-het-virus-besmet-raken.html
16-3-2020	NL	Safety measures	PM Rutte addresses the nation about the coronavirus. In his speech, he announced that the government will not resort to a full lockdown, but instead to opt for a method of controlled spread of the virus. This method aims to build group immunity, but Rutte also stated that a large proportion of the Dutch population will be infected because of this.	RTL: https://www.rtlnieuws.nl/nieuws/nederland/artikel/5057841/gebarentolk-irma-sluis-hamsteren-persconferentie-
16-3-2020	NL	News	There were positive reactions to the sign language interpreter, because of the gestures she made for the word 'hamsteren' (hoarding).	

17-3-2020	NL	News	In the evening of the 17th, many Dutch people applauded the people with necessary jobs, like care workers, garbage collectors and others.	coronavirus ; Volkskrant: https://www.volkskrant.nl/nieuws-achtergrond/voor-het-eerst-een-doventolk-bij-een-belangrijke-persconferentie-het-gebaar-voor-hamsteren-moest-ik-wel-even-opzoeken~be0bdbcc/ Parool: https://www.parool.nl/nederland/oproep-klap-je-handen-stuk-voor-iedereen-die-nederland-door-coronacrisis-slaat~b874fe2a/?referer=https%3A%2F%2Fwww.google.nl%2F ; https://www.nrc.nl/nieuws/2020/03/17/applaus-voor-zorgmedewerkers-a3994068
17-3-2020	NL	News	People have started hoarding certain products, like toilet paper and pasta, from supermarkets around the country.	Trouw: https://www.trouw.nl/economie/nederland-blijft-wc-papier-blikgroente-en-zeep-hamsteren~bce1b5ff/ ; NRC: https://www.nrc.nl/nieuws/2020/03/30/napersconferenties-van-rutte-begint-het-hamsteren-zien-ze-bij-jumbo-a3995329 ; FD: https://fd.nl/ondernemen/1338423/nederlandse-supermarkten-nemen-maatregelen-tegen-hamsteren-en-besmetting
17-3-2020	NL	News	In a newsletter to the public, the cabinet announces that they will take exceptional economic measures to accommodate the jobs and incomes of self-employed workers, entrepreneurs and large companies.	Government: https://www.rijksoverheid.nl/actueel/nieuws/2020/03/17/coronavirus-kabinet-neemt-pakket-nieuwe-maatregelen-voor-banen-en-economie
19-3-2020	NL	Statistics	The total number of patients that have tested positive has grown with 409 and the deaths have grown with 18. This brings the total number of cases to 2,460 and the number of people that died to 76.	WHO: https://covid19.who.int/region/euro/country/nl (retrieved on 29-06-2020)
19-3-2020	NL	News	Bruno Bruins resigns as minister of medical care due to the workload being too taxing. The minister collapsed the previous night during a debate on the coronavirus. Minister Hugo de Jonge takes over his coronavirus-related tasks.	RTL: https://www.rtlnieuws.nl/nieuws/politiek/artikel/5062871/bruins-minister-ontslag ; AD: https://www.ad.nl/politiek/plichtsgetrouwe-minister-bruins-stapt-op-coronacrisis-wordt-hem-nu-toch-te-machtig-br~a324a364/

20-3-2020	NL	News	King of the Netherlands Willem Alexander addressed the nation on national television, telling people that they cannot stop the coronavirus, but they can stop the 'loneliness-virus'.	NOS: https://nos.nl/artikel/2327794-koning-coronavirus-kunnen-we-niet-stoppen-eenzaamheidsvirus-wel.html
22-3-2020	NL	Safety measures	The Dutch government has sent an 'NL-alert' text message to a large proportion of Dutch citizens, telling them to keep 1.5m distance and remain at home if you are sick.	Government: https://www.rijksoverheid.nl/actueel/nieuws/2020/03/22/nl-alert-met-instructies-tegen-coronavirus
23-3-2020	NL	Statistics	The total number of patients that have tested positive has grown with 573 and the deaths have grown with 43. This brings the total number of cases to 4,204 and the number of people that died to 179.	WHO: https://covid19.who.int/region/euro/country/nl (retrieved on 29-06-2020)
23-3-2020	NL	Safety measures	In a nationally televised press conference, PM Rutte introduces new safety measures, calling it an 'intelligent lockdown'. This means that people are still allowed to go outside as long as they keep 1.5 meters distance. Other new safety measures include that all events are cancelled until June 1st and people are not allowed to meet more than 3 other people outside or in their homes. The government also gave more power to local authorities to hand out fines to people that do not follow the instructions (up to €390 for individuals).	NOS: https://nos.nl/video/2328097-premier-rutte-dit-is-een-intelligente-lockdown.html ; Parool: https://www.parool.nl/nederland/dit-was-het-belangrijkste-coronanieuws-van-23-maart~bf16457b/
24-3-2020	NL	Safety measures	In a letter to the house of representatives, the minister of education announces all central final exams in May are cancelled. This is the first time since 1945 that this has happened in the Netherlands.	Government: https://www.tweedekamer.nl/kamerstukken/brieven_regering/detail?id=2020Z05514&did=2020D11480
25-3-2020	NL	Safety measures	The government made guidelines for responsible shopping to instruct stores on how to keep their businesses running in a way that is safe for both employees and customers.	Government: https://www.rijksoverheid.nl/actueel/nieuws/2020/03/25/nieuwe-regels-voor-verantwoord-winkelen
26-3-2020	NL	Statistics	The total number of patients that have tested positive has grown with 852 and the deaths have grown with 80. This brings the total number of cases to 6,412 and the number of people that died to 365.	WHO: https://covid19.who.int/region/euro/country/nl (retrieved on 29-06-2020)
26-3-2020	NL	News	The Government starts a new campaign that aims to offer people a clear and calming message. The campaign is called 'Alleen samen' (Only together) and is displayed on various media platforms like television, social media, websites.	RTL: https://www.rtlnieuws.nl/editienl/artikel/5070981/campagne-slogan-rijksoverheid-coronacrisis-alleen-samen ; Marketing tribune: https://www.marketingtribune.nl/bureaus/nieuws/2020/03/alleen-samen/index.xml
26-3-2020	NL	News	The government's central planning office (CPB) made four scenarios for the economic impact of the coronavirus pandemic and states that a recession is inevitable.	CPB: https://www.cpb.nl/scenarios-coronacrisis

31-3-2020	NL	Statistics	The total number of patients that have tested positive has grown with 884 and the deaths have grown with 93. This brings the total number of cases to 11,750 and the number of people that died to 864. The RIVM reported that the number of reported patients is still growing, but growing less hard.	WHO: https://covid19.who.int/region/euro/country/nl (retrieved on 29-06-2020)
31-3-2020	NL	Safety measures	In a press conference with Hugo de Jonge, minister of health, Mark Rutte announces that all safety measures initiated on 15 March will be extended until at least 28 April.	Government: https://www.youtube.com/watch?v=m5gkmETnV0 ; NU.nl: https://www.nu.nl/coronavirus/6041538/noodmaatregel-en-verlengd-scholen-en-horeca-blijven-dicht-tm-28-april.html
2-4-2020	NL	Statistics	The total number of patients that have tested positive has grown with 1,019 and the deaths have grown with 134. This brings the total number of cases to 13,614 and the number of people that died to 1,173.	WHO: https://covid19.who.int/region/euro/country/nl (retrieved on 29-06-2020)
7-4-2020	NL	Safety measures	In a nationally televised press conference, PM Rutte says that the measures could possibly be loosened a bit after 28 April, however getting back to 'normal' will take considerable time.	Volkskrant: https://www.volkskrant.nl/nieuws-achtergrond/rutte-terug-naar-normaal-gaat-lang-duren~bd06f4f5/ ; Parool: https://www.parool.nl/wereld/dit-was-het-belangrijkste-coronanieuws-van-dinsdag-7-april~bfe1a094/
9-4-2020	NL	Statistics	The total number of patients that have tested positive has grown with 969 and the deaths have grown with 147. This brings the total number of cases to 20,549 and the number of people that died to 2,248.	WHO: https://covid19.who.int/region/euro/country/nl (retrieved on 29-06-2020)
11-4-2020	NL	News	The Ministry of Health, Welfare and Sport invites companies and experts to contribute ideas on the development and deployment of apps in the fight against the coronavirus. The apps can be used for tracing infections and reporting health problems.	Government: https://www.rijksoverheid.nl/actueel/nieuws/2020/03/17/coronavirus-kabinet-neemt-pakket-nieuwe-maatregelen-voor-banen-en-economie
15-4-2020	NL	News	PM Rutte said in a press conference that the cabinet is considering to loosen safety measures in the near future. This can only happen if the following conditions are met: the pressure on healthcare must be reduced, the elderly must be adequately protected and there must be 'insight into the spread of the virus'.	AD: https://www.ad.nl/politiek/rutte-het-loket-gaat-zeker-niet-in-een-keer-open~ada8a36a/?referrer=https://www.google.com/ ; Parool: https://www.parool.nl/nederland/rutte-het-loket-gaat-zeker-niet-in-een-keer-open~bda8a36a/
16-4-2020	NL	Statistics	The total number of patients that have tested positive has grown with 734 and the deaths have grown with 189. This brings the total	WHO: https://covid19.who.int/region/euro/country/nl (retrieved on 29-06-2020)

			number of cases to 28,153 and the number of people that died to 3,134.	
21-4-2020	NL	Statistics	The total number of patients that have tested positive has grown with 750 and the deaths have grown with 67. This brings the total number of cases to 33,405 and the number of people that died to 3,751.	WHO: https://covid19.who.int/region/euro/country/nl (retrieved on 29-06-2020)
21-4-2020	NL	Safety measures	In a press conference, PM Rutte announces that schools for children up to 12 can be partially reopened. Children up to 12 are allowed to play organized sports under supervision and children up to 18 can play team sports when keeping a distance. The other initial measures remain until 20 May. All events are cancelled until 1st of September.	Government: https://www.rijksoverheid.nl/actueel/nieuws/2020/04/21/maatregelen-corona-verlengd
23-4-2020	NL	Research	Start data collection wave 1	
24-4-2020	NL	News	The ministry of Finance announces that the national budget deficit will be 92 billion euros or 11.8 percent. This is the highest deficit since the second world war.	AD: https://www.ad.nl/politiek/coronacrisis-is-dreun-voor-de-schatkist-grootste-tekort-sinds-tweede-wereldoorlog~a8102f15/?referrer=https://www.google.com/ ; Telegraaf: https://www.telegraaf.nl/nieuws/308026519/duizelingwekkend-begrotingstekort-92-miljard
27-4-2020	NL	News	The government is working on an emergency law to increase the legal grounds of the corona measures.	NOS: https://nos.nl/artikel/2331870-kabinet-werkt-aan-spoedwet-nu-noodmaatregelen-langer-gaan-duren.html ; NRC https://www.nrc.nl/nieuws/2020/04/27/spoedwet-voor-noodmaatregelen-a3997955
28-4-2020	NL	Statistics	The total number of patients that have tested positive has grown with 400 and the deaths have grown with 43. This brings the total number of cases to 38,245 and the number of people that died to 4,518.	WHO: https://covid19.who.int/region/euro/country/nl (retrieved on 29-06-2020)
28-4-2020	NL	News	More than 100 people demonstrate in The Hague against the corona measures, more specifically the lockdown.	NOS: https://nos.nl/video/2332030-demonstratie-tegen-lockdownmaatregelen-van-het-nederlandse-kabinet.html ; Nu.nl: https://www.nu.nl/coronavirus/6047702/ruim-honderd-mensen-bij-demonstratie-tegen-coronamaatregelen-in-den-haag.html?redirect=1
5-5-2020	NL	Statistics	The total number of patients that have tested positive has grown with 199 and the deaths have grown with 26. This brings the total number of cases to 40,770 and the number of people that died to 5,082.	WHO: https://covid19.who.int/region/euro/country/nl (retrieved on 29-06-2020)

5-5-2020	NL	News	Several demonstrations are organized against the corona measures. An unauthorized demonstration with (estimated) 300 people in The Hague led to eighty arrests. The demonstration was not reported to the municipality.	NOS: https://nos.nl/artikel/2332855-tachtig-arrestaties-in-den-haag-na-beeindigen-anti-lockdowndemonstratie.html ; AD: https://www.ad.nl/binnenland/tachtig-anti-lockdownactivisten-aangehouden-bij-demonstratie-in-den-haag~aa966a30/
5-5-2020	NL	Research	End data collection wave 1	
6-5-2020	NL	Safety measures	In a press conference, PM Rutte announces a plan to loosen the safety measures step by step. The new message for the people is: "Stay home if you experience symptoms" and "avoid crowded areas".	Government: https://www.rijksoverheid.nl/onderwerpen/coronavirus-covid-19/vraag-en-antwoord/de-persconferentie-van-6-mei-in-eenvoudige-taal ; https://www.rijksoverheid.nl/actueel/nieuws/2020/05/06/vermijd-drukke-houd-15-meter-afstand
11-5-2020	NL	Safety measures	The first cautious step are being taken towards loosening the visiting arrangements for nursing homes. In certain nursing homes, they will allow one visitor per resident as a trial.	Government: https://www.rijksoverheid.nl/actueel/nieuws/2020/05/11/deelnemende-instellingen-versoepeling-bezoekregeling-verpleeghuizen-bekend
11-5-2020	NL	Safety measures	Hairdressers and beauty salons are allowed to reopen, but only if they can guarantee the safety measures are complied with.	RTL: https://www.rtlnieuws.nl/economie/business/artikel/5118746/kappers-open-corona-hygiene-toeslag-maatregelen-klanten ; Nu.nl: https://www.nu.nl/economie/6050458/kappers-weer-open-met-ruimere-openingsstijden-en-hogere-prijzen.html
12-5-2020	NL	Statistics	The total number of patients that have tested positive has grown with 161 and the deaths have grown with 16. This brings the total number of cases to 42,788 and the number of people that died to 5,456.	WHO: https://covid19.who.int/region/euro/country/nl (retrieved on 29-06-2020)
13-5-2020	NL	Research	Start data collection wave 2	
14-5-2020	NL	News	> 10,000 corona fines have been handed out (see 23-03-2020), resistance increases	NOS: https://nos.nl/artikel/2333914-fouten-bij-uitschrijven-coronaboetes-deel-kan-niet-geind-worden.html ; v Volkskrant: https://www.volkskrant.nl/nieuws-

15-5-2020	NL	News	First cases of infected pets in NL (3 cats and one dog).	achtergrond/groeiend-verzet-tegen-coronaboetes-al-ruim-10-duizend-bekeuringen-uitgedeeld~bc97c8efa/ NOS: https://nos.nl/artikel/2334019-coronavirus-vastgesteld-bij-drie-katten-en-een-hond.html ; Volkskrant: https://www.volkskrant.nl/nieuws-achtergrond/voor-het-eerst-in-nederland-huisdieren-besmet-met-corona-virus-aangetroffen-bij-katten-en-een-hond~b4d81036/
18-5-2020	NL	Research	End data collection wave 2	
19-5-2020	NL	Safety measures	The government announces that safety measures still apply, but certain measures can be made more flexible. Groups of people are allowed to meet outside, when keeping 1,5 m distance. Cafés/restaurants and secondary schools are partially reopened on June 1st as planned. Primary schools will be completely reopened on June 8. Not wearing a face mask in public transport will result in a €5 fine.	Government: https://www.rijksoverheid.nl/actueel/nieuws/2020/05/19/corona-aanpak-de-volgende-stap
19-5-2020	NL	Statistics	The total number of patients that have tested positive has grown with 146 and the deaths have grown with 14. This brings the total number of cases to 44,141 and the number of people that died to 5,694.	WHO: https://covid19.who.int/region/euro/country/nl (retrieved on 11-08-2020)
21-5-2020	NL	News	It is getting busier at outdoor locations (e.g., parks and beaches). Park gets evacuated due to it being overcrowded.	NOS: https://nos.nl/artikel/2334600-drukke-op-hemelvaartsdag-parken-ontruimd.html ; NU.nl: https://www.nu.nl/den-haag/6052846/zomerse-drukke-op-strand-bij-scheveningen-op-hemelvaartsdag.html
26-5-2020	NL	Statistics	The total number of patients that have tested positive has grown with 209 and the deaths have grown with 8. This brings the total number of cases to 45,445 and the number of people that died to 5,830.	WHO: https://covid19.who.int/region/euro/country/nl (retrieved on 11-08-2020)
27-5-2020	NL	Research	Start data collection wave 3	
1-6-2020	NL	Safety measures	From the 1st of June, every Dutch citizen with symptoms related to the coronavirus can get tested (free of charge).	Government: https://www.rijksoverheid.nl/actueel/nieuws/2020/05/27/vanaf-1-juni-testen-mogelijk-voor-iedereen-met-milde-klachten ; RIVM: https://www.rivm.nl/coronavirus-covid-19/testen

1-6-2020	NL	Safety measures	Bars, restaurants, museums and movie theaters partly reopen at 12:00 pm June 1st.	RTL: https://www.rtlnieuws.nl/nieuws/nederland/artikel/5139421/livestream-opening-terrassen-1-juni-juno-horeca ; Nu.nl: https://www.nu.nl/coronavirus/6055117/horeca-eindelijk-weer-open-was-wel-toe-aan-wat-anders-dan-een-tosti.html
1-6-2020	NL	News	A Black Lives Matter demonstration on the Dam square in Amsterdam has an attendance of 5,000 people. People criticized the local governments for not intervening, because this was a violation of safety measures.	Volkskrant: https://www.volkskrant.nl/nieuws-achtergrond/kritiek-op-niet-ingrijpen-van-burgemeester-bij-grote-black-lives-matter-betoging-in-amsterdam~b34ee186/ ; Telegraaf: https://www.telegraaf.nl/nieuws/1397493631/bomvolle-dam-in-amsterdam-voor-protest-tegen-amerikaans-geweld
2-6-2020	NL	Research	End data collection wave 3	
3-6-2020	NL	Statistics	The total number of patients that have tested positive has grown with 102 and the deaths have grown with 5. This brings the total number of cases to 46,647 and the number of people that died to 5,967.	WHO: https://covid19.who.int/region/euro/country/nl (retrieved on 11-08-2020)
3-6-2020	NL	Safety measures	PM Mark Rutte announces that holidays inside the country are allowed again and holidays to most European countries will be allowed from June 15th	Government: https://www.rijksoverheid.nl/onderwerpen/coronavirus-covid-19/vraag-en-antwoord/persconferentie-coronavirus-zomervakantie-2020-in-eenvoudige-taal
6-6-2020	NL	News	There has been a data breach on the RIVM "Infectieradar" platform, where Dutch people can report whether they had corona-related symptoms.	NOS: https://nos.nl/artikel/2336416-lek-in-rivm-coronasite-gegevens-van-gebruikers-makkelijk-in-te-zien.html ; AD: https://www.ad.nl/tech/lek-in-rivm-coronasite-gegevens-gebruikers-makkelijk-in-te-zien~ac0a3e6b/?referrer=https://www.google.com/
9-6-2020	NL	News	The previously announced emergency law has received much criticism. Critics say the law could violate citizens' fundamental rights.	NOS: https://nos.nl/artikel/2336696-nieuwe-coronawet-vol-verboden-krijgt-van-alle-kanten-kritiek.html ; Volkskrant: https://www.volkskrant.nl/nieuws-achtergrond/grote-zorgen-over-spoedwet-corona-terug-naar-de-tijd-van-willem-i~b67b3650/
10-6-2020	NL	Statistics	The total number of patients that have tested positive has grown with 164 and the deaths have grown with 15. This brings the total number of cases to 47,903 and the number of people that died to 6,031.	WHO: https://covid19.who.int/region/euro/country/nl (retrieved on 11-08-2020)

10-6-2020	NL	Research	Start data collection wave 4	
12-6-2020	NL	News	Many hospitality entrepreneurs want to get rid of the 1.5 meter rule in cafés, bars and terraces. They say the rule does not work well and customers should be able to decide for themselves.	RTL: https://www.rtlnieuws.nl/nieuws/politiek/artikel/5150926/15-meter-horeca-onder-vuur-dit-bezopen ; Parool: https://www.parool.nl/nederland/horeca-wil-van-1-5-meterregel-af~b8700006/ ; Trouw: https://www.trouw.nl/economie/horeca-wil-van-anderhalvemeter-af-we-krijgen-het-niet-meer-uitgelegd~bc111546/ Volkskrant: https://www.volkskrant.nl/nieuws-achtergrond/nederland-heeft-miljoenen-doses-van-het-oxfordvaccin-besteld-hoe-kansrijk-is-het~bb87f3f8/ ; RTL nieuws: https://www.rtlnieuws.nl/nieuws/politiek/artikel/5150956/vaccin-corona-astrazeneca-farmaceut NOS: https://nos.nl/artikel/2337304-nederlanders-mogen-weer-naar-buitenland-op-vakantie.html ; Nu.nl: https://www.nu.nl/coronavirus/6055355/kabinet-versoepelt-op-15-juni-reisadvies-voor-europese-bestemmingen.html
14-6-2020	NL	News	The Netherlands pre-orders millions of vaccines that are under development in Oxford.	Volkskrant: https://www.volkskrant.nl/economie/economie-krimpt-dit-jaar-met-6-procent-en-werkloosheid-verdubbelt-verwacht-cpb~b15485b0/ ; NOS: https://nos.nl/artikel/2337409-cpb-ongekende-krimp-van-6-procent-werkloosheid-verdubbelt.html AD: https://www.ad.nl/politiek/toch-onafhankelijk-onderzoek-naar-coronabeleid-kabinet~abafba3d/ ; Volkskrant: https://www.volkskrant.nl/nieuws-achtergrond/kamer-wil-toch-onafhankelijk-onderzoek-naar-coronabeleid~b33c364c/
15-6-2020	International	Safety measures	European Union loosens travel restrictions (within the EU).	
16-6-2020	NL	News	The Dutch economy is expected to shrink by 6 percent this year, and unemployment is expected to double.	
16-6-2020	NL	News	An independent mid-term evaluation of the corona policy is announced.	
16-6-2020	NL	Research	End data collection wave 4	

17-6-2020	NL	Statistics	The total number of patients that have tested positive has grown with 139 and the deaths have grown with 5. This brings the total number of cases to 49,087 and the number of people that died to 6,070.	WHO: https://covid19.who.int/region/euro/country/nl (retrieved on 11-08-2020)
19-6-2020	NL	News	Demonstration of the action group "Virus madness" is banned.	NOS: https://nos.nl/artikel/2337788-coronaprotest-in-den-haag-verboden-organisatoren-stappen-naar-rechter.html ; Telegraaf: https://www.telegraaf.nl/nieuws/489283767/den-haag-verbiedt-anti-lockdownprotest-virus-waanzin-naar-de-rechter
23-6-2020	NL	News	First time since March 14 that no corona related deaths have been reported.	WHO: https://covid19.who.int/region/euro/country/nl (retrieved on 11-08-2020)
24-6-2020	NL	Safety measures	Government announces further loosening of corona measures.	NOS: https://nos.nl/artikel/2338386-kijk-hier-de-voorlopig-laatste-coronapersconferentie-terug.html ; https://www.rijksoverheid.nl/onderwerpen/coronavirus-covid-19/vraag-en-antwoord/persconferentie-coronavirus-nieuwe-regels-per-1-juli-2020-in-eenvoudige-taal
24-6-2020	NL	Statistics	The total number of patients that have tested positive has grown with 64 and the deaths have grown with 5. This brings the total number of cases to 49,722 and the number of people that died to 6,095.	WHO: https://covid19.who.int/region/euro/country/nl
24-6-2020	NL	Research	Start data collection wave 5	
26-6-2020	NL	News	Demonstration of the action group "Virus madness" is banned again.	NOS: https://nos.nl/artikel/2338608-den-haag-verbiedt-opnieuw-demonstratie-viruswaanzin.html ; AD: https://www.ad.nl/binnenland/viruswaanzin-na-verbod-demonstratie-kom-morgen-niet-naar-malieveld~a93bb982/
30-6-2020	NL	Statistics	The total number of patients that have tested positive has grown with 76 and the deaths have grown with 2. This brings the total number of cases to 50,223 and the number of people that died to 6,107.	WHO: https://covid19.who.int/region/euro/country/nl (retrieved on 11-08-2020)
30-6-2020	NL	Research	End data collection wave 5	

4. References

- Epskamp, S. (2020a). Psychometric network models from time-series and panel data. *Psychometrika*, 85(1), 206-231. <https://doi.org/10.1007/s11336-020-09697-3>
- Epskamp, S. (2020b). *Psychonetrics: Structural Equation Modeling and Confirmatory Network Analysis*. <https://cran.r-project.org/web/packages/psychonetrics/index.html>
- Epskamp, S., Borsboom, D., & Fried, E. I. (2018). Estimating psychological networks and their accuracy: A tutorial paper. *Behavior Research Methods*, 50(1), 195-212. <https://doi.org/10.3758/s13428-017-0862-1>
- Epskamp, S., Cramer, A. O. J., Waldorp, L. J., Schmittmann, V. D., & Borsboom, D. (2012). qgraph: Network Visualizations of Relationships in Psychometric Data. *Journal of Statistical Software*, 48(4), 1-18. <http://www.jstatsoft.org/v48/i04/>
- R Core Team. (2013). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. <http://www.R-project.org/>
- van Borkulo, C. D., van Bork, R., Boschloo, L., Kossakowski, J. J., Tio, P., Schoevers, R. A., Borsboom, D., & Waldorp, L. J. (2022). Comparing network structures on three aspects: A permutation test. *Psychological Methods*, Advance online publication. <https://doi.org/10.1037/met0000476>