A Mathematical Model of Person Judgment

Part 1: Cue Emergence. Technical supplement

Daniel Leising*

René L. Schilling**

Our aim is to develop a fully parametrized mathematical model of person judgment, combining both (elements of) Kenny’s (1994) Weighted Average Model and Brunswik’s (1956) lens model. The first part of the presentation entirely focusses on the emergence of cues, i.e. average levels of so-called substance variables that may be observed and then used to inform judgements. As it turns out, our approach may well be understood as a temporally dynamic version of classical test theory (Novick, 1966; Lord & Novick, 1968). Among the key observations are (i) that measurements have to be taken over non-degenerate¹ intervals of time, a strict point measurement does not allow one to distinguish between the non-systematic noise and the true value of the observation, (ii) that the non-systematic noise must be modelled by a stochastic process, (iii) that one has to make a careful distinction between two possible sources of randomness, namely the underlying non-systematic noise and the randomness which is induced by the selection of the target(s) from $\{1, \dots, N\}$, and (iv) that these random effects must be discussed simultaneously.

Two further sources of randomness, namely the choice of the perceivers, and the non-systematic component in how perceivers make sense of their observations, are also relevant for person judgment, but will be ignored at this stage (cue emergence).

1 Model setup

A number N of **targets**, generically denoted by lower-case Greek letters $\tau, \tau' \in \{1, 2, \dots, N\}$, are observed by M **perceivers**, which are denoted by $\pi, \pi' \in \{1, 2, \dots, M\}$. Time $t \geq 0$ is represented by the positive real line $[0, \infty)$, the **standard interval** is $I_{\max} = [0, t_{\max})$ for some $t_{\max} > 0$. All possible observations take place in this interval, but the actual duration of the observations may be shorter than $t_{\max}$. In the standard interval $I_{\max}$, the targets are exposed to a sequence of **situations**, which are, in general, determined by the experimental set-up. The set of situations is $\Sigma = \{A, B, C, \dots\}$, and situations are written as capital Roman letters (with or without further indices). The situations may change over time. This is expressed by a map $\sigma : I_{\max} \rightarrow \Sigma$, whose value is the situation $\sigma(t) \in \Sigma$ which is present at time $t \in I_{\max}$. An example of several situations within $I_{\max}$ is shown in (the upper panel of) Fig. 1.

It is sometimes useful to concatenate infinitely many (appropriately shifted) copies of $I_{\max}$ to get an infinite time-horizon: $[0, t_{\max}) \cup [t_{\max}, 2t_{\max}) \cup [2t_{\max}, 3t_{\max}) \cup \dots$. The sequence of situations is repeated over each (shifted) interval, i.e. the function σ is extended periodically with period $t_{\max}$. This construction allows for infinitely long observation horizons, which is important if one wants to de-noise the measurements or if one wants to use “ergodicity”, i.e. the possibility to switch between “observation of many targets in a short-time interval” and “observation of single targets in many intervals”.²

*Prof. Dr. Daniel Leising, Faculty of Psychology, Technische Universität Dresden, 01062 Dresden, Germany. daniel.leising@tu-dresden.de

**Prof. Dr. René L. Schilling, Faculty of Mathematics, Department of Mathematical Stochastics, Technische Universität Dresden, 01062 Dresden, Germany. rene.schilling@tu-dresden.de

¹A non-degenerate interval I is an interval that consists of more than a single point and, thus, has strictly positive length $|I| > 0$.

²A similar assumption is already present in Novick’s classic paper [4].

Figure 1: Top panel: the function $\sigma(t)$ giving the situation at time t . Bottom panel: two intervals of observation $I, I' \subseteq I_{\max}$ and their relative position to $[I_1 \cup I_2 \cup I_3 \cup \dots \cup I_5] \subseteq I_{\max}$. The intervals $I_1, I_2, \dots$ are chosen in such a way that they contain exactly one situation.

When making observations, randomness plays a crucial role. There are two sources of randomness: At any point in time, each target is affected by an unsystematic noise $dW^\tau(t, \omega)$ which is, e.g., due to measurement errors, hidden variables or target-specific conditions. It is natural to assume that $dW^\tau(t, \omega)$ and $dW^{\tau'}(t, \omega)$ are stochastically independent (hence, uncorrelated), but behave statistically in a similar way – with target-specific parameters. This can be used to model the fact that some targets may be more easily “recognized” from their behavior at any given time than other targets. Notice that there are no cross-target effects of this part of the randomness. The other source is a statistical randomness, which arises from the random selection of the targets $\tau \in \{1, \dots, N\}$.

The Unsystematic noise $dW^\tau(t, \omega)$ and $W^\tau(t, \omega)$. In the absence of further information on the concrete situations and the measurements, it is reasonable to assume maximal uncertainty. A standard choice here is **white noise** $dW^\tau(t, \omega)$, resp. its cumulative version **Brownian motion** $W^\tau(t, \omega)$, since their probability distributions maximize entropy (which is a measure of uncertainty) for a fixed variance, see Schilling [5, Section 21.1]. Brownian motion can be completely described by the underlying probability distribution: at each point of time t , the noise level $W^\tau(t, \omega)$ up to time t is a normally distributed random variable with zero mean $\mathbb{E}W^\tau(t, \cdot) = 0$ and variance $\text{Var}W^\tau(t, \cdot) = \gamma^2(\tau)t$. The variance parameter $\gamma^2(\tau)$ may depend on the specific target τ . Two possible trajectories $t \mapsto W^\tau(t, \omega)$ are schematically shown in Fig. 2. Brownian motion is a mathematically well understood process, see e.g. Schilling [5].

The choice of Brownian motion as underlying unsystematic noise is somewhat arbitrary, but helps to simplify the presentation. One could easily replace $W^\tau(t, \omega)$ by a general second-

Figure 2: Two trajectories of Brownian motion (tagged with ω /red and ω' /blue, respectively, moving from the point $W^\tau(t_0, \omega) = 0$ at time $t_0 = 0$ in t_1 “seconds” to a random point $W^\tau(t_1, \omega)$ and $W^\tau(t_1, \omega')$ – the frequencies of the positions make up a normal distribution $N(0, \gamma^2(\tau)t_1)$ – and it moves from that position in $t_2 - t_1$ “seconds” to a further random position $W^\tau(t_2, \omega)$ or $W^\tau(t_2, \omega')$ (again normally distributed $N(0, \gamma^2(\tau)t_2)$ /black, if we view it from $W^\tau(t_0, \omega) = 0$ at $t_0 = 0$, or $N(W^\tau(t_1, \omega), \gamma^2(\tau)(t_2 - t_1))$ /red, resp., $N(W^\tau(t_1, \omega'), \gamma^2(\tau)(t_2 - t_1))$ /blue), if we view it from the intermediary time t_1 .

order stationary random process, see e.g. [4, Section 3], making the notation more cumbersome without real gain in generality. Let us point out that our discussion uses only the following features of the noise: (i) $W^\tau(0, \omega) = 0$. (ii) For any two fixed times $0 \leq s < t$ the difference $W^\tau(t, \omega) - W^\tau(s, \omega)$ is random variable with mean 0 and variance $\gamma^2(\tau)(t - s)$ depending only on the time increment (and, possibly, the target). (iii) For any two (or more) non-overlapping time increments, say $0 \leq t_1 < t_2 \leq t_3 < t_4$, the random variables $W^\tau(t_2, \omega) - W^\tau(t_1, \omega)$ and $W^\tau(t_4, \omega) - W^\tau(t_3, \omega)$ are stochastically independent. So, there is no interaction between two or more non-overlapping intervals, (iv) One may think of $W^\tau(t, \omega) - W^\tau(s, \omega)$ as an infinitesimal sum of spot noises $dW^\tau(u, \omega)$, i.e. $W^\tau(t, \omega) - W^\tau(s, \omega) = \int_s^t dW^\tau(u, \omega)$.³

If $W^\tau(t, \omega)$ is Brownian motion (i)–(iv) hold. In addition, $W^\tau(t, \omega)$ is **normally distributed** with mean zero and variance $\gamma^2(\tau)t$; the “spot” noise $dW^\tau(t, \omega)$ is white noise.

The statistical randomness. Choosing targets (randomly) from $\{1, \dots, N\}$ adds an additional source of randomness. It is due to the variation within the targets (each target has its own characteristic behavior in any one situation) which makes the collection of targets (resp., the sample of targets which we are observing) random. In fact, the probability to choose a specific target τ from the whole population $\{1, \dots, N\}$ is $\frac{1}{N}$. This leads to an empirical mean value and empirical standard deviation which will be denote by a tilde $\widetilde{}$.

Since the unsystematic and statistical randomness are independent, we can use a product construction of the respective probability spaces, and discuss both effects simultaneously.

³This is a naive point of view, which can be made rigorous within a well-defined mathematical theory (so-called Itô or stochastic differentials, see [5]).

2 Means, variances and covariances

As mentioned above, we will treat sample points depending on two types of randomness. If $X = X(\tau, t, \omega)$ is a (random) sample point, then τ describes the randomness incurred by the choice of targets, while ω indicates the presence of an unsystematic noise $W^\tau(t, \omega)$; t is the dependence on time which is suppressed in this section.

The basic operation is **taking means**, e.g. the mean value over the population $\{1, \dots, N\}$ or the expected value over the unsystematic noise:

$$\begin{aligned}\tilde{X}(\omega) = \tilde{X}(\cdot, \omega) &= \frac{1}{N} \sum_{\tau=1}^N X(\tau, \omega) && \text{mean value in } \tau \\ \mathbb{E}X(\tau) = \mathbb{E}X(\tau, \cdot) &&& \text{mean value in } \omega.\end{aligned}$$

Notice the convention that the variable, in which the mean is taken, either disappears from the sample point X or is replaced by a “dot”. Means are linear, i.e. $\mathbb{E}(\alpha X(\tau) + \beta Y(\tau)) = \alpha \mathbb{E}X(\tau) + \beta \mathbb{E}Y(\tau)$ (α, β are constants) and the mean of a constant $C = C(\tau)$ (i.e. not depending on ω) is just the constant itself: $\mathbb{E}C(\tau) = C(\tau)$; similar relations apply to the mean in τ .

Once the mean is defined, the **variance** is defined as the mean of the squared deviation of the sample points from their respective mean:

$$\begin{aligned}\widetilde{\text{var}} X(\omega) = \widetilde{\text{var}} X(\cdot, \omega) &= \frac{1}{N} \sum_{\tau=1}^N [X(\tau, \omega) - \tilde{X}(\omega)]^2, && \text{variance in } \tau; \\ \text{Var}X(\tau) = \text{Var}X(\tau, \cdot) &= \mathbb{E}([X(\tau) - \mathbb{E}X(\tau)]^2), && \text{variance in } \omega.\end{aligned}$$

Thus, adding or subtracting a constant from $X(\tau, \omega)$ does not influence $\text{Var}X(\tau)$ nor $\widetilde{\text{var}} X(\omega)$. Finally, **covariances** are defined as the mean of the product of the mean-deviations of each sample point. The covariances in ω and τ , respectively, are:

$$\begin{aligned}\widetilde{\text{cov}}(X(\omega), Y(\omega)) &= \widetilde{\text{cov}}(X(\cdot, \omega), Y(\cdot, \omega)) = \frac{1}{N} \sum_{\tau=1}^N (X(\tau, \omega) - \tilde{X}(\omega))(Y(\tau, \omega) - \tilde{Y}(\omega)), \\ \text{Cov}(X(\tau), Y(\tau)) &= \text{Cov}(X(\tau, \cdot), Y(\tau, \cdot)) = \mathbb{E}([X(\tau) - \mathbb{E}X(\tau)][Y(\tau) - \mathbb{E}Y(\tau)]).\end{aligned}$$

Notice that covariances are linear in each of its two arguments,⁴ and that covariances do not change if one adds to X (or Y or both) a constant (relative to the variable in which the covariance is calculated). This will be used later on.

The order in which means are taken is immaterial, i.e.

$$\widetilde{\mathbb{E}X} = \mathbb{E}\tilde{X} = \frac{1}{N} \sum_{\tau=1}^N \mathbb{E}X(\tau) = \frac{1}{N} \mathbb{E} \left[\sum_{\tau=1}^N X(\tau) \right],$$

consequently,

$$\begin{aligned}\widetilde{\text{var}} X &= \frac{1}{N} \sum_{\tau=1}^N \mathbb{E} \left([X(\tau) - \widetilde{\mathbb{E}X}]^2 \right) \\ \widetilde{\text{cov}}(X, Y) &= \frac{1}{N} \sum_{\tau=1}^N \mathbb{E} \left([X(\tau) - \widetilde{\mathbb{E}X}] [Y(\tau) - \widetilde{\mathbb{E}Y}] \right).\end{aligned}$$

⁴that is,

$$\begin{aligned}\text{Cov}(\alpha X(\tau) + \beta Y(\tau), \alpha' X'(\tau) + \beta' Y'(\tau)) \\ = \alpha \alpha' \text{Cov}(X(\tau), X'(\tau)) + \alpha \beta' \text{Cov}(X(\tau), Y'(\tau)) + \beta \alpha' \text{Cov}(Y(\tau), X'(\tau)) + \beta \beta' \text{Cov}(Y(\tau), Y'(\tau)) \\ \text{Cov}(X(\tau), Y(\tau)) = \text{Cov}(Y(\tau), X(\tau)) \quad \text{and} \quad \text{Cov}(X(\tau), X(\tau)) = \text{Var}X(\tau),\end{aligned}$$

where $\alpha, \beta, \alpha', \beta'$ are constants.

3 The concrete model

All relevant parameters of the following model are summarized in Table 1. A perceiver observes one (or several) target(s) τ in some intervals I and I' . The position of the intervals I and I' is free, i.e. they may or may not overlap, they may or may not be adjacent. If I, I' depend on the perceiver, this is indicated by $I(\pi)$. This feature will not be used in the present paper. In general, the model is set up in a way that the observation of the aggregate substance variable S does not depend on the perceiver, nor on the fact that it is being perceived, i.e. measurement/perception etc. does not influence the substance level.

We are interested in modelling the response of a target τ when exposed to different situations over time. This is expressed by a **substance variable** (i.e. a certain substance level, the notion of **variable** is in a traditional, non-mathematical sense denoting any quantity that may change its value), which will be given in differential form (so-called Itô differentials) as:

$$(1) \quad dS(\tau, t, \omega) = \theta(\tau, \sigma(t)) dt + dW^\tau(t, \omega).^5$$

Let us briefly discuss the terms appearing in (1): $dW^\tau(t, \omega)$ is the non-systematic spot noise (see below) and $\theta(\tau, \sigma(t))$ is the noise-free true value of the substance variable. It is useful to represent it as a sum of various differences. That is, for every interval I – comprising one or several situations – and every $t \in I$

$$(2) \quad \theta(\tau, \sigma(t)) - \tilde{\bar{\theta}}(I) = \underbrace{\left(\bar{\theta}(\tau, I) - \tilde{\bar{\theta}}(I)\right)}_{=\Delta\bar{\theta}(\tau, I)} + \underbrace{\left(\tilde{\theta}(\sigma(t)) - \tilde{\bar{\theta}}(I)\right)}_{=\Delta\tilde{\theta}(\sigma(t), I)} + \Delta\theta(\tau \times \sigma(t), I)$$

where we follow standard notation, using the superscript “bar” to indicate **temporal means**, i.e.

$$\bar{\theta}(\tau, I) = \frac{1}{|I|} \int_I \theta(\tau, \sigma(t)) dt, \quad \tilde{\bar{\theta}}(I) = \frac{1}{N} \sum_{\tau=1}^N \bar{\theta}(\tau, I), \quad \tilde{\theta}(\sigma(t)) = \frac{1}{N} \sum_{\tau=1}^N \theta(\tau, \sigma(t)),$$

and $\Delta\theta(\tau \times \sigma(t), I)$ is given by

$$(3) \quad \Delta\theta(\tau \times \sigma(t), I) = \theta(\tau, \sigma(t)) - \Delta\bar{\theta}(\tau, I) - \Delta\tilde{\theta}(\sigma(t), I) - \tilde{\bar{\theta}}(I).$$

In (2), (3) the interval I is completely arbitrary. The most interesting special case where $I = I_{\max}$ is discussed below in (33) and (34). Observe that $\Delta\theta(\tau \times \sigma(t), I_{\max})$ is the **interaction effect**, $\Delta\bar{\theta}(\tau, I_{\max})$ is the **target effect** and $\Delta\tilde{\theta}(\sigma(t), I_{\max})$ is the **situation effect**, while $\tilde{\bar{\theta}}(I_{\max})$ is the **grand mean**. The essential point is that we may later make assertions about these quantities in the standard interval $I_{\max}$ based on observations in shorter intervals, say $I, I' \subset I_{\max}$.

⁵This is a model with **additive noise**. It is possible to introduce a further time-, target-, perceiver- and even substance-specific weight w , replacing $dW^\tau(t, \omega)$ by $w(t, \tau, \pi, S(\tau, \sigma(t), \omega))dW^\tau(t, \omega)$; while the τ and π -dependence is rather simple to handle, the time- and substance-dependence will make essential changes necessary, leading to so-called **multiplicative noise models**. These are not discussed here.

Table 1: Overview of the parameters

Parameter	Meaning	Ref.
t	time	
τ	target	
π	perceiver	
$\sigma(t)$	situation at time t	Fig. 1
ω	unsystematic randomness	
$I_{\max}$	standard interval	Fig. 1
I	interval of observation	Fig. 1
$ I $	length of the interval I	
$\theta(\tau, \sigma(t))$	noiseless score of target τ at time t	(1), (2)
$\tilde{\theta}(\sigma(t))$	the average target's noiseless score for the situation $\sigma(t)$ at time t	
$\bar{\theta}(\tau, I)$	noiseless cue level of target τ for the interval I	(17), (36)
$\tilde{\bar{\theta}}(I) = \tilde{\bar{\theta}}(I)$	grand mean of the substance variable across I	(29)
$\Delta\tilde{\theta}(\sigma(t), I)$	$\tilde{\theta}(\sigma(t)) - \tilde{\bar{\theta}}(I)$; situation effect if $I = I_{\max}$	(33), (2)
$\Delta\bar{\theta}(\tau, I)$	$\bar{\theta}(\tau, I) - \tilde{\bar{\theta}}(I)$; target effect if $I = I_{\max}$	(33), (2)
$\Delta\theta(\tau \times \sigma(t), I)$	$\theta(\tau, \sigma(t)) - \Delta\bar{\theta}(\tau, I) - \Delta\tilde{\theta}(\sigma(t), I) - \tilde{\bar{\theta}}(I)$ target by situation interaction effect if $I = I_{\max}$	(34), (3)
$W^\tau(t, \omega)$	cumulative noise level up to time t for target τ , modelled by a Brownian motion with mean zero and variance $\gamma^2(\tau)t$	
$\gamma(\tau)\sqrt{t}$	standard deviation of the noise at time t for target τ	Fig. 2
$dS(\tau, t, \omega)$	rate of the (noisy) substance level of target τ at time t	(1)
$dW^\tau(t, \omega)$	rate of the random error for target τ at time t ; "white noise"	(1)
$\bar{S}(\tau, I, \omega)$	observable cue level of target τ for the interval I	(10), (35)
$\widetilde{}, \widetilde{\text{var}}, \widetilde{\text{cov}}$	mean, variance and covariance taken only in targets (τ)	Sec. 2
$\mathbb{E}, \text{Var}, \text{Cov}$	mean value, variance and covariance taken only in the noise (ω)	Sec. 2
$\widetilde{\text{Var}}, \widetilde{\text{Cov}}$	variance and covariance taken in targets (τ) and noise (ω) simultaneously	Sec. 2

There are several equivalent forms of the quantity $\Delta\theta(\tau \times \sigma(t))$

$$\begin{aligned}
(4) \quad & \Delta\theta(\tau \times \sigma(t), I) \\
& = \theta(\tau, \sigma(t)) - \left(\bar{\theta}(\tau, I) - \tilde{\theta}(I) \right) - \left(\tilde{\theta}(\sigma(t)) - \tilde{\theta}(I) \right) - \tilde{\theta}(I) \\
(5) \quad & = \frac{1}{N} \sum_{\tau'=1}^N \frac{1}{|I|} \int_I \left\{ \left[\theta(\tau, \sigma(t)) - \theta(\tau, \sigma(s)) \right] + \left[\theta(\tau', \sigma(s)) - \theta(\tau', \sigma(t)) \right] \right\} ds \\
(6) \quad & = \frac{1}{|I|} \int_I \left\{ \left[\theta(\tau, \sigma(t)) - \theta(\tau, \sigma(s)) \right] + \left[\tilde{\theta}(\sigma(s)) - \tilde{\theta}(\sigma(t)) \right] \right\} ds \\
(7) \quad & = \frac{1}{N} \sum_{\tau'=1}^N \left\{ \left[\theta(\tau, \sigma(t)) - \bar{\theta}(\tau, I) \right] + \left[\bar{\theta}(\tau', I) - \theta(\tau', \sigma(t)) \right] \right\} \\
(8) \quad & = \left[\theta(\tau, \sigma(t)) - \bar{\theta}(\tau, I) \right] + \left[\tilde{\theta}(I) - \tilde{\theta}(\sigma(t)) \right].
\end{aligned}$$

In the above formulae τ and t denote a fixed target and a fixed point of time t whereas τ' is any member of the target population $\{1, 2, \dots, N\}$ and s is any time in the observation interval I .

The noise component, $dW^\tau(t, \cdot)$ is the differential of Brownian motion – this is frequently called **white noise** in the physics and engineering literature – i.e. an object that, if it is integrated, gives

$$(9) \quad W^\tau(t, \omega) = \int_0^t dW^\tau(s, \omega)$$

and $W^\tau(t, \cdot)$ is a Brownian motion with standard deviation $\gamma(\tau)\sqrt{t}$, see above.

Formula (1) is given in a **differential form** which can be interpreted as the infinitesimal change $S(\tau, t, \omega)$ at time t . This quantity cannot be observed as such, but any observation has to be done over some interval of time $I = [a, b]$ leading to observable quantities like

$$S(\tau, b, \omega) - S(\tau, a, \omega) = \int_a^b dS(\tau, t, \omega) dt = \int_a^b \theta(\tau, \sigma(t)) dt + W^\tau(b, \omega) - W^\tau(a, \omega).$$

In order to eliminate the effect of observation time, one should normalize by the length of the observation and divide everything by $|I| = b - a$; otherwise we would get a cumulative measurement.

This leads naturally to the **cue level** for the interval I

$$(10) \quad \bar{S}(\tau, I, \omega) = \frac{1}{|I|} \int_I \theta(\tau, \sigma(t)) dt + \frac{1}{|I|} \int_I dW^\tau(t, \omega).$$

For clarity, if $I = [a, b]$, then

$$(11) \quad \bar{S}(\tau, [a, b], \omega) = \frac{1}{b-a} \int_a^b \theta(\tau, \sigma(t)) dt + \frac{1}{b-a} \int_a^b dW^\tau(t, \omega).$$

The bar $\bar{}$ indicates again that one takes the mean over time in the given interval I .

In the case of the unit interval $I = [0, 1]$, one recovers the standard discrete model

$$(12) \quad \bar{S}(\tau, \omega) = \bar{\theta}(\tau) + \epsilon(\tau, \omega),$$

with a target's true score $\bar{\theta}(\tau) = \int_0^1 \theta(\tau, \sigma(t)) dt$ and a normally distributed unsystematic error $\epsilon(\tau, \omega) = W^\tau(1, \omega)$; the normal distribution appears since the underlying process is Brownian motion, but this can be easily changed.

It is straightforward to calculate the ω -mean. Since θ does not depend on ω , it is not affected by $\mathbb{E}$, so

$$(13) \quad \mathbb{E}(\bar{S}(\tau, [a, b])) = \frac{1}{b-a} \int_a^b \theta(\tau, \sigma(t)) dt + \mathbb{E} \left(\frac{1}{b-a} \int_a^b dW^\tau(t) \right)$$

$$(14) \quad = \frac{1}{b-a} \int_a^b \theta(\tau, \sigma(t)) dt + \frac{1}{b-a} \mathbb{E}(W^\tau(b) - W^\tau(a))$$

$$(15) \quad = \frac{1}{b-a} \int_a^b \theta(\tau, \sigma(t)) dt$$

$$(16) \quad = \bar{\theta}(\tau, [a, b]).$$

If in the interval $I = [a, b]$ there is only one situation, i.e. $\sigma(t) = A$ (say) is constant for all $t \in I = [a, b]$, and one has

$$(17) \quad \bar{\theta}(\tau, [a, b]) = \frac{1}{b-a} \int_a^b \theta(\tau, \sigma(t)) dt$$

$$(18) \quad = \frac{1}{b-a} \int_a^b \theta(\tau, A) dt$$

$$(19) \quad = \frac{\theta(\tau, A)}{b-a} \int_a^b dt = \theta(\tau, A).$$

The interpretation is two-fold: first, $\theta(\tau, \sigma(t))$ will be the noiseless, i.e. unperturbed, response of target τ in situation $\sigma(t)$ (here $\sigma(t) = A$, yielding $\theta(\tau, A)$). In this sense, $\theta(\tau, \sigma(t))$ denotes the true substance variable, but this cannot, in general, be observed. Moreover, the calculation shows that if one can make many observations of the target τ inside a one-situation interval, then one obtains a very good statistical estimate for $\theta(\tau, \sigma(t))$, since the noise will be reduced by averaging across the observations.

Let us discuss the role of the length of the observation interval $|I| = b - a$. Two types of observations are particularly interesting: long-term observations when $b - a \rightarrow \infty$ and short-term observations when $b - a \rightarrow 0$. Consider first $b - a \rightarrow \infty$. The variance in ω is given by

$$(20) \quad \mathbb{V}\text{ar}(\bar{S}(\tau, [a, b])) = \mathbb{V}\text{ar} \left(\frac{1}{b-a} \int_a^b dW^\tau(t, \cdot) \right)$$

$$(21) \quad = \frac{1}{(b-a)^2} \mathbb{V}\text{ar} \left(\int_a^b dW^\tau(t, \cdot) \right)$$

$$(22) \quad = \frac{1}{(b-a)^2} \mathbb{V}\text{ar}(W^\tau(b, \cdot) - W^\tau(a, \cdot))$$

$$(23) \quad = \frac{1}{(b-a)^2} (b-a) \gamma^2(\tau)$$

$$(24) \quad = \frac{\gamma^2(\tau)}{b-a} \xrightarrow{b-a \rightarrow \infty} 0$$

This means that **the longer the observation, the less influence of random noise we have** in the substance level tied to this interval (= cue). This is due to the effect of the factor $1/(b-a) = 1/|I|$, which pulls out from under the variance in a quadratic manner while the noise contributes only linearly with $|I| = b - a$.

Now consider $|I| = b - a \rightarrow 0$. This gives an even better idea of the normalization by $1/|I|$. Recall first the fundamental theorem of integral calculus. One has

$$(25) \quad \frac{1}{b-a} \int_a^b f(x) dx \xrightarrow{b \rightarrow a} f(a)$$

This means that $b - a$ normalizes the observation $\int_a^b dS(\tau, t, \omega)$, the latter is an integral over $I = [a, b]$, and one obtains, in the mean as $b \rightarrow a$ or $b - a \rightarrow 0$:

$$(26) \quad \mathbb{E}\bar{S}(\tau, [a, b]) = \frac{1}{b-a} \int_a^b \theta(\tau, \sigma(t)) dt \xrightarrow{b \rightarrow a} \theta(\tau, \sigma(a))$$

i.e. very short observations yield a point (or spot) observation of the true substance level $\theta(\tau, \sigma(a))$ at time $t = a$. However, this is statistically not feasible, since at the same time the ω -variance becomes infinite (same calculation as before):

$$(27) \quad \text{Var}\bar{S}(\tau, [a, b]) = \frac{1}{(b-a)^2} \text{Var}(W^\tau(b) - W^\tau(a)) = \frac{\gamma^2(\tau)}{(b-a)} \xrightarrow{b \rightarrow a} \infty.$$

This has the following neat interpretation: in a point measurement it is impossible to distinguish between true substance and non-systematic noise.

Infinite time horizons In order to cancel out the noise completely, it is necessary to have an infinite time-horizon. This can be (formally) achieved by copying the standard observation interval $I_{\max} = [0, t_{\max})$ and concatenating the (appropriately shifted) copies. The function $\sigma(t)$ is assumed to be the same on each copy (i.e. it is periodically extended from $I_{\max}$ to $[0, \infty)$), cf. Section 1. On each copy the noise will be structurally the same, nevertheless we have different, statistically independent noise levels.

This procedure allows one to speak of a target grand mean $\bar{\theta}(\tau, [0, \infty))$ and an overall grand mean $\tilde{\theta}([0, \infty))$ which are given by

$$(28) \quad \bar{\theta}(\tau, [0, \infty)) = \lim_{|I| \rightarrow \infty} \frac{1}{|I|} \int_I \theta(\tau, \sigma(t)) dt = \lim_{T \rightarrow \infty} \frac{1}{T} \int_{[0, T)} \theta(\tau, \sigma(t)) dt$$

and

$$(29) \quad \tilde{\theta}([0, \infty)) = \lim_{|I| \rightarrow \infty} \frac{1}{|I|} \int_I \tilde{\theta}(\sigma(t)) dt = \lim_{T \rightarrow \infty} \frac{1}{T} \int_{[0, T)} \tilde{\theta}(\sigma(t)) dt.$$

Since $\sigma(t)$ is periodic, one also has

$$(30) \quad \begin{aligned} \bar{\theta}(\tau, [0, \infty)) &= \frac{1}{|I_{\max}|} \int_{I_{\max}} \theta(\tau, \sigma(t)) dt = \bar{\theta}(\tau, I_{\max}), \quad \text{and} \\ \tilde{\theta}([0, \infty)) &= \frac{1}{|I_{\max}|} \int_{I_{\max}} \tilde{\theta}(\sigma(t)) dt = \tilde{\theta}(I_{\max}). \end{aligned}$$

Therefore, the grand means coincide with the temporal resp. target-temporal means taken over the standard observation interval $I_{\max}$.

The noise $dW^\tau(t, \omega)$ is only statistically periodic, and gives for each repetition of the interval $I_{\max}$ independent contributions. Therefore, the strong law of large numbers applies and yields

$$(31) \quad \lim_{|I| \rightarrow \infty} \frac{1}{|I|} \int_I S(\tau, t, \omega) dt = \lim_{|I| \rightarrow \infty} \frac{1}{|I|} \int_I \theta(\tau, \sigma(t)) dt + \lim_{|I| \rightarrow \infty} \frac{1}{|I|} \int_I dW^\tau(t, \omega) = \bar{\theta}(\tau, I_{\max}).$$

Similarly,

$$(32) \quad \lim_{|I| \rightarrow \infty} \frac{1}{|I|} \int_I \tilde{S}(t, \omega) dt = \tilde{\theta}(I_{\max}).$$

This conforms to the intuition that sufficiently long observations minimize (and, in the limit, even eradicate) the influence of noise, cf. also (20)–(24).

Moreover, the above construction allows one to use in the decomposition (2), (3) the interval $I = [0, \infty)$ to get for every $t \in [0, \infty)$

$$(33) \quad \theta(\tau, \sigma(t)) - \tilde{\bar{\theta}}(I_{\max}) = \underbrace{\left(\bar{\theta}(\tau, I_{\max}) - \tilde{\bar{\theta}}(I_{\max}) \right)}_{=\Delta\bar{\theta}(\tau, I_{\max})} + \underbrace{\left(\tilde{\bar{\theta}}(\sigma(t)) - \tilde{\bar{\theta}}(I_{\max}) \right)}_{=\Delta\tilde{\bar{\theta}}(\sigma(t), I_{\max})} + \Delta\theta(\tau \times \sigma(t), I_{\max})$$

where $\Delta\theta(\tau \times \sigma(t), I_{\max})$ is given by

$$(34) \quad \Delta\theta(\tau \times \sigma(t), I_{\max}) = \theta(\tau, \sigma(t)) - \Delta\bar{\theta}(\tau, I_{\max}) - \Delta\tilde{\bar{\theta}}(\sigma(t), I_{\max}) - \tilde{\bar{\theta}}(I_{\max}).$$

4 Calculation of the variances and co-variances

Recall that $\text{Var}, \widetilde{\text{var}}, \mathbb{V}\widetilde{\text{ar}}$ and $\text{Cov}, \widetilde{\text{cov}}, \mathbb{C}\widetilde{\text{ov}}$ denote the variance and co-variance w.r.t. ω, τ and (ω, τ) . Throughout this section I, I' are any two intervals of observation, no assumption on their length, position and mutual position is made at the moment. Recall from (2) that

$$(35) \quad \bar{S}(\tau, I, \omega) \stackrel{\text{def}}{=} \frac{1}{|I|} \int_I \theta(\tau, \sigma(t)) dt + \frac{1}{|I|} \int_I dW^\tau(t, \omega) =: \bar{\theta}(\tau, I) + \bar{W}^\tau(I, \omega),$$

$$(36) \quad \bar{\theta}(\tau, I) \stackrel{\text{def}}{=} \frac{1}{|I|} \int_I \theta(\tau, \sigma(t)) dt,$$

$$(37) \quad \bar{W}^\tau(I, \omega) \stackrel{\text{def}}{=} \frac{1}{|I|} \int_I dW^\tau(t, \omega),$$

$$(38) \quad \tilde{\bar{\theta}}(I) \stackrel{\text{def}}{=} \frac{1}{N} \sum_{\tau=1}^N \bar{\theta}(\tau, I),$$

and so one arrives at the following decomposition

$$(39) \quad \bar{S}(\tau, I, \omega) = \tilde{\bar{\theta}}(I) + \underbrace{(\bar{\theta}(\tau, I) - \tilde{\bar{\theta}}(I))}_{=\Delta\bar{\theta}(\tau, I)} + \bar{W}^\tau(I, \omega).$$

Mean values. One has

$$(40) \quad \mathbb{E}\bar{S}(\tau, I) = \bar{\theta}(\tau, I) + \mathbb{E}\bar{W}^\tau(I, \cdot) = \bar{\theta}(\tau, I)$$

$$(41) \quad \tilde{\bar{S}}(I, \omega) = \tilde{\bar{\theta}}(I) + \underbrace{\widetilde{\bar{W}}^\bullet(I, \omega)}_{\text{cannot be simplified}}$$

and, for the mean in both τ and ω ,

$$(42) \quad \mathbb{E}\tilde{\bar{S}}(I) = \frac{1}{N} \sum_{\tau=1}^N \mathbb{E}\bar{S}(\tau, I) = \frac{1}{N} \sum_{\tau=1}^N \bar{\theta}(\tau, I) = \tilde{\bar{\theta}}(I).$$

Variances. One has

$$(43) \quad \text{Var}\bar{S}(\tau, I) = \text{Var}\bar{W}^\tau(I) = \frac{1}{|I|^2} \text{Var} \int_I dW^\tau(t) = \frac{\gamma^2(\tau)|I|}{|I|^2} = \frac{\gamma^2(\tau)}{|I|};$$

since the θ -part has no randomness, it behaves like a constant for Var . Moreover,

$$(44) \quad \widetilde{\text{var}} \bar{S}(I, \omega) = \frac{1}{N} \sum_{\tau=1}^N \left(\bar{S}(\tau, I, \omega) - \widetilde{\bar{S}}(I, \omega) \right)^2$$

$$(45) \quad = \frac{1}{N} \sum_{\tau=1}^N \left(\bar{S}(\tau, I, \omega) - \widetilde{\theta}(I) - \widetilde{\bar{W}}^\bullet(I) \right)^2$$

$$(46) \quad = \frac{1}{N} \sum_{\tau=1}^N \left(\left[\bar{\theta}(\tau, I) - \widetilde{\theta}(I) \right] + \left[\bar{W}^\tau(I, \omega) - \widetilde{\bar{W}}^\bullet(I, \omega) \right] \right)^2$$

and, finally, the variance in both arguments

$$(47) \quad \mathbb{V}\widetilde{\text{ar}} \bar{S}(I) = \frac{1}{N} \sum_{\tau=1}^N \mathbb{E} \left(\bar{S}(\tau, I, \omega) - \mathbb{E} \widetilde{\bar{S}}(I, \omega) \right)^2$$

$$(48) \quad = \frac{1}{N} \sum_{\tau=1}^N \mathbb{E} \left(\bar{\theta}(\tau, I) + \bar{W}^\tau(I, \cdot) - \widetilde{\theta}(I) \right)^2$$

$$(49) \quad = \frac{1}{N} \sum_{\tau=1}^N \mathbb{E} \left(\bar{\theta}(\tau, I) - \widetilde{\theta}(I) + \bar{W}^\tau(I, \cdot) \right)^2$$

$$(50) \quad = \frac{1}{N} \sum_{\tau=1}^N \left(\bar{\theta}(\tau, I) - \widetilde{\theta}(I) \right)^2 + \frac{1}{N} \sum_{\tau=1}^N \mathbb{E} (\bar{W}^\tau(I, \cdot)^2)$$

$$(51) \quad = \widetilde{\text{var}} \bar{\theta}(I) + \left(\frac{1}{N} \sum_{\tau=1}^N \gamma^2(\tau) \right) \frac{1}{|I|}$$

Since the variance of any quantity X is the same as the co-variance of X with itself, it holds that

$$\begin{aligned} \text{Var} \bar{S}(\tau, I) &= \text{Cov} (\bar{S}(\tau, I), \bar{S}(\tau, I)) , \quad \widetilde{\text{var}} \bar{S}(I, \omega) = \widetilde{\text{cov}} (\bar{S}(I, \omega), \bar{S}(I, \omega)) , \\ \text{and} \quad \mathbb{V}\widetilde{\text{ar}} \bar{S}(I) &= \text{Cov} (\bar{S}(I), \bar{S}(I)) . \end{aligned}$$

One should compare the analogy of the variance formulae with the following formulae for the covariance.

Covariances if I and I' are non-overlapping. Consider two targets $\tau, \tau^* \in \{1, \dots, N\}$ which may be different $\tau \neq \tau^*$ but need not be different. Since the observation intervals I, I' are non-overlapping, $\bar{W}^\tau(I, \omega)$ and $\bar{W}^{\tau^*}(I', \omega)$ are independent, hence uncorrelated. This leads to a considerable simplification.

$$(52) \quad \text{Cov} (\bar{S}(\tau, I), \bar{S}(\tau, I')) = \text{Cov} (\bar{W}^\tau(I), \bar{W}^\tau(I')) = 0 \quad \left(= |I \cap I'| \right)$$

Now use that $\bar{S}(\tau, I, \omega) = \bar{\theta}(\tau, I) + \bar{W}^\tau(I, \omega)$; by the linearity of the covariance in any argument, one has

$$(53) \quad \widetilde{\text{cov}} (\bar{S}(I, \omega), \bar{S}(I', \omega)) = \widetilde{\text{cov}} (\bar{\theta}(I), \bar{\theta}(I'))$$

$$(54) \quad + \widetilde{\text{cov}} (\bar{\theta}(I), \bar{W}^\bullet(I', \omega))$$

$$(55) \quad + \widetilde{\text{cov}} (\bar{\theta}(I'), \bar{W}^\bullet(I, \omega))$$

$$(56) \quad + \widetilde{\text{cov}} (\bar{W}^\bullet(I, \omega), \bar{W}^\bullet(I', \omega))$$

The last equality is quite interesting: if the noise W^τ **does not depend on the target** τ – i.e. if $\gamma(\tau) = \gamma$ for every τ –, then the last three terms vanish. The same effect happens **always** for the joint (τ, ω) -covariance. This has an interesting reason: although $W^\tau(t, \omega)$ depends on the individual target, the non-correlation assumption kills off these terms. To wit – note that $\mathbb{E}\tilde{S}(I') = \tilde{\theta}(I)$ is the joint (τ, ω) -mean –

$$(57) \quad \text{Cov}(\bar{S}(I), \bar{S}(I'))$$

$$(58) \quad = \frac{1}{N} \sum_{\tau=1}^N \mathbb{E} \left[\left(\bar{\theta}(\tau, I) + \bar{W}^\tau(I, \cdot) - \tilde{\theta}(I) \right) \left(\bar{\theta}(\tau, I') + \bar{W}^\tau(I', \cdot) - \tilde{\theta}(I') \right) \right]$$

$$(59) \quad = \frac{1}{N} \sum_{\tau=1}^N \mathbb{E} \left[\left(\bar{\theta}(\tau, I) - \tilde{\theta}(I) + \bar{W}^\tau(I, \cdot) \right) \left(\bar{\theta}(\tau, I') - \tilde{\theta}(I') + \bar{W}^\tau(I', \cdot) \right) \right]$$

$$(60) \quad = \frac{1}{N} \sum_{\tau=1}^N \left(\bar{\theta}(\tau, I) - \tilde{\theta}(I) \right) \left(\bar{\theta}(\tau, I') - \tilde{\theta}(I') \right)$$

$$(60) \quad = \text{cov}(\bar{\theta}(I), \bar{\theta}(I'));$$

this is intuitively clear as the noise is uncorrelated, so there must be no effect on the covariance from the noise.

Covariances if I and I' are overlapping. Assume that $I \cap I' \neq \emptyset$ and that they have more in common than just the end-point. It is not assumed that I and I' have the same length. We introduce a factor which expresses the extent of the overlap of $|I|$ and $|I'|$.

$$(61) \quad w_{I,I'} = \frac{|I \cap I'|}{|I|}, \quad \text{and so} \quad \frac{|I \setminus I'|}{|I|} = \frac{|I| - |I \cap I'|}{|I|} = 1 - w_{I,I'}.$$

Note that $w_{I,I'} \neq w_{I',I}$, unless $|I| = |I'|$. In fact,

$$(62) \quad w_{I',I} = \frac{|I' \cap I|}{|I'|}, \quad \text{and so} \quad \frac{|I' \setminus I|}{|I'|} = \frac{|I'| - |I \cap I'|}{|I'|} = 1 - w_{I',I}.$$

The number $w_{I,I'} \in [0, 1]$ gives the relative size of the overlap $I \cap I'$ with respect to I ; $w_{I',I}$ is the analogous quantity, but now relative to I' . If $w = 0$, then there is **no overlap** (or an overlap in at most one point), if $w = 1$ there is **total overlap**. The quantity $w_{I,I'}$ can be used to re-scale the observed substance level **if one splits the observation interval**. This is shown in the bottom panel of Fig. 1. One has

$$\begin{aligned} \bar{S}(\tau, I, \omega) &= \frac{1}{|I|} \int_I dS(\tau, t, \omega) \\ &= \frac{1}{|I|} \left(\int_{I \cap I'} dS(\tau, t, \omega) + \int_{I \setminus I'} dS(\tau, t, \omega) \right) \\ &= \frac{1}{|I|} (|I \cap I'| \cdot \bar{S}(\tau, I \cap I', \omega) + |I \setminus I'| \cdot \bar{S}(\tau, I \setminus I', \omega)) \\ &= w_{I,I'} \bar{S}(\tau, I \cap I', \omega) + (1 - w_{I,I'}) \bar{S}(\tau, I \setminus I', \omega). \end{aligned}$$

Combining this formula with the linearity of the covariance in each argument, it is easy to calculate various covariances. To keep notation simple, we write in the following calculations

$w = w_{I,I'}$ and $w' = w_{I',I}$.

$$(63) \quad \text{Cov}(\bar{S}(\tau, I), \bar{S}(\tau, I'))$$

$$(64) \quad = \text{Cov}(w\bar{S}(\tau, I \cap I') + (1-w)\bar{S}(\tau, I \setminus I'), w'\bar{S}(\tau, I' \cap I) + (1-w')\bar{S}(\tau, I' \setminus I))$$

$$(65) \quad = ww' \text{Cov}(\bar{S}(\tau, I \cap I'), \bar{S}(\tau, I' \cap I))$$

$$(66) \quad = ww' \text{Var}\bar{S}(\tau, I \cap I') = ww' \frac{\gamma^2(\tau)}{|I \cap I'|} = \frac{\gamma^2(\tau)|I \cap I'|}{|I| \cdot |I'|}$$

Note that the three other terms which involve the covariances over $(I \cap I')$ & $(I \setminus I')$ resp. $(I \cap I')$ & $(I' \setminus I)$ resp. $(I \setminus I')$ & $(I' \setminus I)$ are all zero because of the disjointness of the intervals.

The formula for $\text{cov}(\bar{S}(I, \omega), \bar{S}(I', \omega))$ can also be worked out explicitly, but it is rather messy, containing 16 different terms, and does not really seem to be helpful.

Let us finally work out the joint (τ, ω) -covariance

$$(67) \quad \text{Cov}(\bar{S}(I), \bar{S}(I'))$$

$$= \text{Cov}(w\bar{S}(I \cap I') + (1-w)\bar{S}(I \setminus I'), w'\bar{S}(I' \cap I) + (1-w')\bar{S}(I' \setminus I))$$

$$(68) \quad = ww' \text{Cov}(\bar{S}(I \cap I'), \bar{S}(I' \cap I))$$

$$+ w(1-w') \text{Cov}(\bar{S}(I \cap I'), \bar{S}(I' \setminus I))$$

$$+ (1-w)w' \text{Cov}(\bar{S}(I \setminus I'), \bar{S}(I' \cap I))$$

$$+ (1-w)(1-w') \text{Cov}(\bar{S}(I \setminus I'), \bar{S}(I' \setminus I))$$

$$(69) \quad = ww' \text{Var}(\bar{S}(I \cap I'))$$

$$+ w(1-w') \text{cov}(\bar{\theta}(I \cap I'), \bar{\theta}(I' \setminus I))$$

$$+ (1-w)w' \text{cov}(\bar{\theta}(I \setminus I'), \bar{\theta}(I' \cap I))$$

$$+ (1-w)(1-w') \text{cov}(\bar{\theta}(I \setminus I'), \bar{\theta}(I' \setminus I))$$

$$(70) \quad = ww' \text{var}(\bar{\theta}(I \cap I')) + ww' \left[\frac{1}{N} \sum_{\tau=1}^N \gamma^2(\tau) \right] |I \cap I'|$$

$$+ w(1-w') \text{cov}(\bar{\theta}(I \cap I'), \bar{\theta}(I' \setminus I))$$

$$+ (1-w)w' \text{cov}(\bar{\theta}(I \setminus I'), \bar{\theta}(I' \cap I))$$

$$+ (1-w)(1-w') \text{cov}(\bar{\theta}(I \setminus I'), \bar{\theta}(I' \setminus I))$$

$$(71) \quad = \text{cov}(\bar{\theta}(I), \bar{\theta}(I')) + ww' \left[\frac{1}{N} \sum_{\tau=1}^N \gamma^2(\tau) \right] \frac{1}{|I \cap I'|}$$

$$(72) \quad = \text{cov}(\bar{\theta}(I), \bar{\theta}(I')) + \left[\frac{1}{N} \sum_{\tau=1}^N \gamma^2(\tau) \right] \frac{|I \cap I'|}{|I| \cdot |I'|}.$$

The covariance (in τ) of two intervals does not depend on the fact whether these intervals overlap or do not overlap. This is intuitively clear – $\text{cov}(\bar{\theta}(I), \bar{\theta}(I'))$ does not depend on time and is, thus, time homogeneous. Yet, this requires formal proof. To do so, consider four non-overlapping intervals K_1, K_2, K'_2, K_3 where K_2, K'_2 have the same length $|K_2| = |K'_2|$ and are such that $\sigma(t)$, hence $\theta(\tau, \sigma(t))$, shows identical behavior on K_2 and K'_2 resulting in $\bar{\theta}(\tau, K_2) = \bar{\theta}(\tau, K'_2)$. Therefore,

$$\begin{aligned} \bar{\theta}(\tau, K_2 \cup K_3) &= \frac{|K_2|}{|K_2 \cup K_3|} \bar{\theta}(\tau, K_2) + \frac{|K_3|}{|K_2 \cup K_3|} \bar{\theta}(\tau, K_3) \\ &= \frac{|K'_2|}{|K'_2 \cup K_3|} \bar{\theta}(\tau, K'_2) + \frac{|K_3|}{|K'_2 \cup K_3|} \bar{\theta}(\tau, K_3) = \bar{\theta}(\tau, K'_2 \cup K_3) \end{aligned}$$

and one sees from this, that

$$\widetilde{\text{cov}}(\bar{\theta}(K_1 \cup K_2), \bar{\theta}(K_2 \cup K_3)) = \widetilde{\text{cov}}(\bar{\theta}(K_1 \cup K_2), \bar{\theta}(K'_2 \cup K_3)).$$

Notice that the first expression means that $I = K_1 \cup K_2$ and $I' = K_2 \cup K_3$ overlap at K_2 , the second expression covers the case of non-overlapping $I = K_1 \cup K_2$ and $I' = K'_2 \cup K_3$, but under the same experimental conditions.

The relation obtained in line (72),

$$(73) \quad \text{Cov}(\bar{S}(I), \bar{S}(I')) = \widetilde{\text{cov}}(\bar{\theta}(I), \bar{\theta}(I')) + \left[\frac{1}{N} \sum_{\tau=1}^N \gamma^2(\tau) \right] \frac{|I \cap I'|}{|I| \cdot |I'|},$$

or the normalized version which takes values in the interval $[-1, 1]$,

$$(74) \quad \text{Corr}(\bar{S}(I), \bar{S}(I')) = \frac{\widetilde{\text{cov}}(\bar{\theta}(I), \bar{\theta}(I')) + \left[\frac{1}{N} \sum_{\tau=1}^N \gamma^2(\tau) \right] \frac{|I \cap I'|}{|I| \cdot |I'|}}{\sqrt{\text{Var}(\bar{S}(I))} \sqrt{\text{Var}(\bar{S}(I'))}},$$

are the **most important formulae in the present analysis**. They describe the extent to which the observable differences in the targets' behavior (the “cues” in the lens model) will remain stable across two observation intervals, allowing for intervals of any length and any degree of overlap. For a more detailed discussion as to why this matters so much for personality judgment, check the main paper.

Let us finally discuss the covariance $\widetilde{\text{cov}}(\bar{\theta}(I), \bar{\theta}(I'))$. Observe that one can split $\bar{\theta}(\tau, I)$ as follows:

$$(75) \quad \bar{\theta}(\tau, I) = \underbrace{\bar{\theta}(\tau, I) - \bar{\theta}(\tau, I_{\max})}_{=\delta\bar{\theta}(\tau, I, I_{\max})} + \underbrace{\bar{\theta}(\tau, I_{\max}) - \tilde{\bar{\theta}}(I_{\max})}_{=\Delta\bar{\theta}(\tau, I_{\max})} + \tilde{\bar{\theta}}(I_{\max})$$

The notation $\delta\bar{\theta}(\tau, I, I_{\max})$ is used here, since $\Delta\bar{\theta}(\tau, I)$ has a different meaning, see (2). The difference $\delta\bar{\theta}(\tau, I, I_{\max})$ comprises the situation effect $\Delta\bar{\theta}(\sigma(t), I)$; since the latter quantity is the same for all targets, this component of $\delta\bar{\theta}(\tau, I, I_{\max})$ does not contribute towards the covariance across all targets τ . An analogous decomposition holds for $\bar{\theta}(\tau, I')$. Since $\tilde{\bar{\theta}}(I_{\max})$ does not contribute to $\widetilde{\text{cov}}$, one has

$$(76) \quad \widetilde{\text{cov}}(\bar{\theta}(I), \bar{\theta}(I'))$$

$$(77) \quad = \widetilde{\text{cov}}(\Delta\bar{\theta} + \delta\bar{\theta}(I), \Delta\bar{\theta} + \delta\bar{\theta}(I'))$$

$$(78) \quad = \widetilde{\text{cov}}(\Delta\bar{\theta}, \Delta\bar{\theta}) + \widetilde{\text{cov}}(\delta\bar{\theta}(I), \delta\bar{\theta}(I')) + \widetilde{\text{cov}}(\Delta\bar{\theta}, \delta\bar{\theta}(I')) + \widetilde{\text{cov}}(\delta\bar{\theta}(I), \Delta\bar{\theta})$$

$$(79) \quad = \widetilde{\text{var}}(\Delta\bar{\theta}) + \widetilde{\text{cov}}(\delta\bar{\theta}(I), \delta\bar{\theta}(I')) + \widetilde{\text{cov}}(\Delta\bar{\theta}, \delta\bar{\theta}(I')) + \widetilde{\text{cov}}(\delta\bar{\theta}(I), \Delta\bar{\theta}).$$

In order to keep the above formulae readable, we have used the following shorthand notation $\Delta\bar{\theta} = \Delta\bar{\theta}(I_{\max})$ and $\delta\bar{\theta}(I) = \delta\bar{\theta}(I, I_{\max})$, $\delta\bar{\theta}(I') = \delta\bar{\theta}(I', I_{\max})$.

5 Test reliability

This part of the model follows the set-up of classical test theory, see Novick (1966) and Lord & Novick (1968), but we have to distinguish between various means and quantities (which is not always properly done in the literature). In particular, we see that the present model generalizes classical test theory to time intervals and non-spot observations.

We begin with a simple observation. Assume that I and I' are two intervals of the same length $|I| = |I'|$ and such that σ has the same form in I and in I' – that is, I and I' have

the same line-up of situations. We will call such intervals **situation-similar** or, for short, **σ -similar**. In particular

$$(80) \quad \theta(\tau, I) = \theta(\tau, I') \quad \text{if } I, I' \text{ are } \sigma\text{-similar}$$

for any target τ , no matter how I and I' are located relative to each other (i.e. overlapping, non-overlapping). In general, the true value $\theta(\tau, I)$ is not accessible, and observable with noise only. This makes the next definition problematic – at first sight. The **test reliability** is defined as

$$(81) \quad \text{rel}(I) := \frac{\widetilde{\text{var}}(\bar{\theta}(I))}{\widetilde{\text{Var}}(\bar{S}(I))}.$$

Of course, $\text{rel}(I)$ depends on the interval I . One of the main results of classical test theory is to express rel in terms of **observable** quantities, i.e. one has to get rid of $\bar{\theta}(\tau, I)$ in the above formula.

First of all, (57)–(60) show that for non-overlapping intervals I, I' one has

$$(82) \quad \mathbb{C}\widetilde{\text{ov}}(\bar{S}(I), \bar{S}(I')) = \mathbb{c}\widetilde{\text{ov}}(\bar{\theta}(I), \bar{\theta}(I')) \quad \text{whenever } I \cap I' = \emptyset$$

Moreover, from $\bar{\theta}(\tau, I) = \bar{\theta}(\tau, I')$ (this requires the one-situation assumption) one gathers that

$$(83) \quad \mathbb{c}\widetilde{\text{ov}}(\bar{\theta}(I), \bar{\theta}(I')) = \widetilde{\text{var}}(\bar{\theta}(I)) = \widetilde{\text{var}}\bar{\theta}(I') \quad \text{whenever } I, I' \text{ are } \sigma\text{-similar}.$$

Finally, if I, I' are σ -similar, (47)–(51) yield that

$$(84) \quad \widetilde{\text{Var}}(\bar{S}(I)) = \widetilde{\text{Var}}(\bar{S}(I')) \quad \text{if } I, I' \text{ are } \sigma\text{-similar}.$$

This uses that Brownian motion is temporally homogeneous, i.e. depends only on the length, but not the position of I and I' . Combining the last three equalities and inserting everything into the definition of $\text{rel}(I)$, one obtains for σ -similar and non-overlapping intervals I, I' that

$$(85) \quad \text{rel}(I) \stackrel{\text{def}}{=} \frac{\widetilde{\text{var}}\bar{\theta}(I)}{\widetilde{\text{Var}}(\bar{S}(I))}$$

$$(86) \quad = \frac{\mathbb{C}\widetilde{\text{ov}}(\bar{S}(I), \bar{S}(I'))}{\widetilde{\text{Var}}(\bar{S}(I))}$$

$$(87) \quad = \frac{\mathbb{C}\widetilde{\text{ov}}(\bar{S}(I), \bar{S}(I'))}{\sqrt{\widetilde{\text{Var}}(\bar{S}(I))}\sqrt{\widetilde{\text{Var}}(\bar{S}(I'))}}$$

$$(88) \quad = \frac{\mathbb{C}\widetilde{\text{ov}}(\bar{S}(I), \bar{S}(I'))}{\sqrt{\widetilde{\text{Var}}(\bar{S}(I))}\sqrt{\widetilde{\text{Var}}(\bar{S}(I'))}}$$

$$(89) \quad = \mathbb{C}\widetilde{\text{orr}}(\bar{S}(I), \bar{S}(I')).$$

This is a familiar formula. Note that this formula even holds, if each target has his own “private” noise. This is due to the fact that the derivation relied only on (57)–(60), and not the formulae (53)–(56).

We close our discussion with the role of the length of the observations. The starting point is, once again, the definition of $\text{rel}(I)$, making no further assumptions on I . One has

$$(90) \quad \text{rel}(I) \stackrel{\text{def}}{=} \frac{\widetilde{\text{var}}\bar{\theta}(I)}{\widetilde{\text{Var}}(\bar{S}(I))}$$

$$(91) \quad \stackrel{(51)}{=} \frac{\widetilde{\text{var}}\bar{\theta}(I)}{\widetilde{\text{var}}\bar{\theta}(I) + \left(N^{-1} \sum_{\tau=1}^N \gamma^2(\tau)\right) / |I|}.$$

This shows that test reliability increases to 1 if the length of observation $|I| \rightarrow \infty$. On the downside, note that too short observation intervals $|I|$ affect the reliability, which tends to 0 if $|I| \rightarrow 0$. Moreover, the above formula shows how the variance of the unsystematic noise, $\gamma^2(\tau)$, enters into test reliability. The smaller the noise variance, the better the reliability. If $\gamma^2(\tau) = \gamma^2$, i.e. if the unsystematic noise is (statistically) the same for each target, then

$$(92) \quad \text{rel}(I) \stackrel{\text{def}}{=} \frac{\widetilde{\text{var}} \bar{\theta}(I)}{\widetilde{\text{var}}(\bar{S}(I))} = \frac{\widetilde{\text{var}} \bar{\theta}(I)}{\widetilde{\text{var}} \bar{\theta}(I) + \frac{\gamma^2}{|I|}}.$$

References

- [1] Brunswik, E.: *Perception and the representative design of psychological experiments*. University of California Press, Berkeley (CA) 1956.
- [2] Kenny, D.A.: *Interpersonal perception. A social relations analysis*. The Guilford Press, New York (NY) 1994.
- [3] Lord, F.M., Novick, M.R.: *Statistical theories of mental test scores*. Addison-Wesley Publishing Company, Reading (MA) 1968.
- [4] Novick, M.R.: The axioms and principal results of classical test theory. *Journal of Mathematical Psychology* **3**(1) (1966) 1–18.
- [5] Schilling, R.L.: *Brownian Motion* (3rd ed). De Gruyter, Berlin 2021.