**The role of orthographic neighbourhood effects in lateralized lexical decision:
A replication study and meta-analysis.**

Adam J. Parker^{1*}

Ciara Egan^{1,2}

Jack H. Grant^{1,3}

Sophie Harte^{1,4}

Brad T. Hudson^{1,5}

Zoe V. J. Woodhead¹

Word count: 7,913

Affiliations:

¹Department of Experimental Psychology, University of Oxford

²School of Psychology, Bangor University

³School of Psychology, University of Lincoln

⁴Deafness, Cognition and Language Research Centre, University College London

⁵Department of Psychology, Lancaster University

*Correspondence regarding this article should be addressed to Adam J. Parker, Department of Experimental Psychology, University of Oxford, Anna Watts Building, Woodstock Road, Oxford, OX2 6GG. Email: adam.parker@psy.ox.ac.uk.

Abstract

The effect of orthographic neighbourhood size (N) on lexical decision reaction time differs when words are presented in the left or right visual fields. Evidence suggests a facilitatory N effect (i.e., faster reaction times for words with larger neighbourhoods) in the left visual field. However, the N effect in the right visual field remains controversial: it may have a weaker facilitative role or it may even be inhibitory. In a pre-registered online experiment, we replicated the interaction between N and visual field and provided support for an inhibitory N effect in the right visual field. We subsequently conducted a pre-registered systematic review and meta-analysis to synthesise the available evidence and determine the direction of N effects across visual fields. Based on the evidence, it would seem the effect is inhibitory in the right visual field. Furthermore, the size of the N effect is considerably smaller in the right visual field. Both studies revealed considerable heterogeneity between participants and studies, and we consider the implications of this for future work.

Keywords: Orthographic neighbourhood effects, lateralized presentation, laterality, cerebral hemisphere, replication, systematic review and meta-analysis

1 The orthographic neighbourhood of a word is the set of words that can be formed by
2 substituting a single letter in a target word. For example, the neighbourhood of *sleet* is *fleet*,
3 *sheet*, *skeet*, *sweet*, *slept*, *sleek*, and *sleep*. The size of this neighbourhood (i.e. the number of
4 words in the neighbourhood; referred to as '*N*' hereafter; Coltheart, Davelaar, Jonasson, &
5 Besner, 1977) has been reported to influence word identification. Several studies using the
6 lexical decision paradigm have indicated that, for words presented in central/foveal vision, a
7 larger *N* is associated with more accurate and faster responses (for reviews see Andrews,
8 1997; Grainger, 1992; Mathey, 2001).

9 This facilitative *N* effect may seem counterintuitive when considered within a Parallel
10 Distributed Processing framework, such as the Interactive Activation model (McClelland &
11 Rumelhart, 1981), which posits that there is lateral inhibition between competing lexical
12 representations. This might predict that words with more neighbours would experience more
13 lateral inhibition, resulting in slower reaction times. It has been theorised instead that the
14 facilitation may arise due to excitatory feedback connections from the word to the letter level.
15 When *N* is larger, partial activation of orthographic neighbours at the word level may thereby
16 increase letter level activation and speed up word recognition (Andrews, 1989, 1992;
17 Coltheart & Rastle, 1994; Whitney & Lavidor, 2005).

18 The role of *N* during lexical access has also been examined using lateralized
19 presentation, yielding similarly counterintuitive results. During these visual half-field
20 experiments, words are presented in the left or right visual hemifield. Words presented to the
21 left visual field (LVF) will initially be projected to the visual cortex in the right cerebral
22 hemisphere. Words presented to the right visual field (RVF) will initially be projected to the
23 visual cortex of the left cerebral hemisphere. It has been shown that in the LVF, there is a
24 facilitative *N* effect so that words with more neighbours are responded to more quickly and
25 accurately. By contrast, in the RVF the *N* effect is weaker or even inhibitory (Ellis, 2004;

Lavidor & Ellis, 2001, 2002; Perea, Acha, & Fraga, 2008; Whitney, 2004). This pattern of results is surprising, as the LVF shows the same direction of effect as the central visual field (i.e., facilitation); whereas the RVF, which is characterised as being more efficient for word processing due to its direct access to the language dominant hemisphere, shows an atypically weak or reversed N effect.

One specific computational model of letter-encoding that makes predictions about differential orthographic processing in each hemisphere is the SERIOL (sequential encoding regulated by inputs to oscillations within letter units) model (Whitney, 2001). The SERIOL model makes specific predictions about how word identification occurs from early retinotopic visual processing via abstract feature, letter and bigram levels, to a lexical word form level (see Figure 1). At the earliest levels of the model (the retina and edge level processing in V1/V2), there is an activation gradient related to acuity, where activity is highest at central fixation and falls off into the periphery. In the left hemisphere (receiving inputs from the RVF) activity is graded from left-to-right, whereas in the right hemisphere (receiving inputs from the LVF) it is graded right-to-left. In order to preserve left-to-right serial order in the abstract letter representations, the acuity gradient in the right hemisphere has to be inverted. This gradient inversion is achieved in two stages: (1) bottom-up excitation from the edge to the feature level, which boosts activation in the right hemisphere relative to the left; and (2) left-to-right lateral inhibition at the feature level in the right hemisphere, but not the left.

[Insert Figure 1 here]

Whitney (2004; Whitney & Lavidor, 2005) proposed that differences between the left hemisphere and right hemisphere processing up to the letter level may explain the differential effects of N across hemispheres. Under visual half field viewing conditions, words presented to the LVF extend further than normal into parafoveal vision, and the usual processes of excitation and lateral inhibition may be insufficient to fully invert the acuity gradient. As a

result, the spatial gradient of letters for LVF words may be imprecise, and require greater than usual top-down support from the word level to resolve word recognition. As a result, there will be a greater than usual facilitatory N effect, as words with more neighbours will produce stronger top-down support from the word to letter level. Put more simply, the SERIOL model suggests that there is relatively poor orthographic encoding in the LVF, which is offset by increased top-down orthographic facilitation from the word level back to the letter level. More top-down orthographic facilitation means a stronger N effect, and hence we observe a facilitative N effect in the right hemisphere (Whitney, 2004).

In contrast, the SERIOL model posits that for words presented in the RVF, letter processing is more efficient as the acuity gradient matches the serial order of letters in the word. This results in more efficient orthographic encoding in the left hemisphere. This means that top-down excitatory feedback has a weaker effect in the left hemisphere (Whitney & Lavidor, 2005), making N effects smaller or negligible.

Given SERIOL's predictions for the role of N in each hemisphere, we would expect empirical studies which examine effects of N and visual field on accuracy and reaction time to yield a significant interaction. This interaction would point towards a facilitative N effect in the LVF but not in the RVF. Indeed, several studies have reported facilitative effects of N in the LVF (Ellis, 2004; Lavidor & Ellis, 2001, 2002; Lavidor, Johnston, & Snowling, 2006; Mano, Patrick, Andresen, Capizzi, Biagioli, & Osmon, 2010; Perea, Acha, & Fraga, 2008; Whitney & Lavidor, 2005)¹. Out of these studies, some report weaker facilitatory effects in the RVF (Lavidor & Ellis, 2001; Lavidor et al., 2006; Whitney & Lavidor, 2005), while others report inhibitory effects (Lavidor & Ellis, 2002; Mano et al., 2010; Perea et al., 2008).

¹Accuracy in these studies is typically high and variability is low. While some studies report effects of visual field on accuracy (with greater accuracy in the RVF relative to the LVF; Perea et al., 2008), others do not (Lavidor & Ellis, 2001). There is also little evidence to conclude that N, or the interaction between N and visual field, yield strong effects on accuracy.

The largest inhibitory N effect in the RVF to date was reported by Perea et al. (2008), who reported a very small (-2 ms) facilitative effect in the central visual field (CVF), a much larger facilitative effect in the LVF (-23 ms), and a strong inhibitory effect in the RVF (28 ms). This contrasts the equally sizeable -24 ms facilitatory effect² reported by Lavidor et al. (2006) for typical readers.

The extent to which the N effect is weakly facilitatory or inhibitory in the RVF (i.e. left hemisphere) is of interest then to models that make predictions based on each hemisphere's involvement in word recognition. To further examine the role of N under lateralized presentation, and to add weight to the claim that N effects are inhibitory in the RVF, the current work describes two studies of N effects during lateralized lexical decision. The first was an online replication of Perea et al. (2008), who reported facilitative effects of N in the LVF and inhibitory effects in the RVF. In anticipation of the results, we replicate Perea et al.'s findings and report that N effects differ across visual fields in a sample of English speaking participants. We subsequently conducted a systematic review and meta-analysis of the published literature to determine both the direction and size of N effects in the RVF.

Study 1: An online replication study

We first set out to examine whether we could replicate the pattern of results reported by Perea, Acha, and Fraga (2008) in an online replication study with English speaking participants. The experiment was pre-registered on the Open Science Framework prior to the commencement of data collection. We pre-registered three criteria for a successful replication: (1) reduced errors and faster reaction times in the CVF relative to the LVF and RVF for word targets; (2) reduced errors and faster reaction times in the RVF relative to the

²This effect size was estimated from Figure 1 in Lavidor et al. (2006) using the metagear package (version 0.6; Lajeunesse, 2020).

LVF for word targets; (3) a facilitative N effect in the LVF and an inhibitory effect in the RVF in reaction times for word targets. The registration form, materials, data, and R scripts can be found at osf.io/vknbp.

Method

Participants

A priori power analysis using pilot data and the SimR package (version 1.0.5; Green & McLeod, 2016) in R (R Development Core Team, 2016) suggested that 30 participants would provide sufficient power (power > 80%) to detect the critical N by visual field interaction in reaction time. The critical effect size used in these simulations was 15 ms, which is more conservative than the effect size reported by Perea et al. (2008). To reach this sample size, 45 native English speakers were recruited through Prolific Academic (<https://www.prolific.co/>); a web-based platform for recruiting and screening paid research participants according to experimenter-specified criteria. Participants were right-handed, had normal or corrected-to-normal vision, no significant hearing loss, history of neurological disease, head injury, epilepsy, or diagnosis of language and/or reading impairment. Thirteen participants were removed from the study as their score on the Edinburgh Handedness Inventory was below 80 (Oldfield, 1971). This cut-off was chosen to match Perea et al. (2008). Two further participants were removed from the study as they scored below chance when completing the lexical decision task in at least one of the three visual field conditions. The remaining 30 participants (18 female) had a mean age of 29.0 years ($SD_{age} = 8.12$ years).

This study was approved by the University of Oxford's Medical Sciences Inter-divisional Research Ethics Committee (reference number: R67385/RRE001) and conducted in accordance with the principles of the Declaration of Helsinki. Participants gave informed written consent and received compensation at a rate of £10/hour for their participation.

Materials

Stimuli were selected using the same restrictions as Perea et al. (2008). We selected 120 word and 120 non-word targets. All targets had two syllables and were 5 letters in length. For word targets, 60 had few substitution neighbours ($M_N = 0.7$, range_N: 0-1) and 60 had many substitution neighbours ($M_N = 9.5$, range_N: 7-12). N was acquired using the English Lexicon Project (Balota et al., 2007). Word targets did not differ in zipf frequency scores (low N: $M_{\text{zipf}} = 3.1$, $SD_{\text{zipf}} = 0.82$; high N: $M_{\text{zipf}} = 3.1$, $SD_{\text{zipf}} = 0.77$; Mann Whitney $U = 1768.00$, $p = .869$), obtained from the SUBTLEX-UK database (van Heuven, Mandera, Keuleers, & Brysbaert, 2014). One hundred and twenty non-word targets were selected from the English Lexicon Project (Balota et al., 2007) and had either no substitution neighbours or one substitution neighbour.

Design and Procedure

The experiment consisted of two tasks: (1) the Edinburgh Handedness Inventory, (2) the visual half-field task. The visual half-field task was within-subjects and utilised a 2 (lexicality: word, non-word) x 2 (N: low, high) x 3 (visual field: RVF, LVF, CVF) design.

The study was administered via Gorilla (<https://gorilla.sc/>; Anwyl-Irvine, Massonnié, Flitton, Kirkham, & Evershed, 2020). Participants were instructed to sit approximately 50 cm from the screen. Each trial began with a fixation cross presented centrally for 400 ms. Participants were instructed to keep their eyes on the central fixation point. Then, a target was presented for 150 ms to the left, right, or at the position of the fixation point. Previous attempts at using Gorilla to run similar visual half-field tasks involving rapid presentation of stimuli yielded good test-retest reliability (Parker, Woodhead, Thompson, & Bishop, 2020).

Stimuli were presented in black lowercase Tahoma font on a white background. Gorilla's scaling tool was used to ensure that stimulus size and position were kept consistent despite differences in participants' screen sizes. Font was scaled to 250% equating to 28pt font, and stimuli were presented at a displacement of 2.5° of visual angle when presented to

the left or right of fixation. Participants were instructed to press one of two buttons on the keyboard to indicate whether the string was a legitimate English word or not (“S” for word or “K” for non-word). They were instructed to do so as quickly and accurately as possible. The next trial started contingent on the participant’s response. Stimuli were divided into three counterbalanced lists with each participant responding to 120 words and 120 non-words in a random order. The whole session took approximately 12 minutes.

Data analysis

Error rates were analysed using a generalized linear mixed-effects model (GLMM) fitted to binomial data using the *glmer()* function and logit link from the lme4 package (version 1.1-21; Bates et al., 2019). Reaction times were analysed with linear mixed-effects models (LMM) using the *lmer()* function from the lme4 package. Both models adopted an identical structure: $dv \sim VF + N + VF:N + (1 | participant) + (1 | word)$. The *contr.sum* function was used to implement summed-to-zero contrasts for N, constituting a main N effect. The *contr.helmert()* function was used to implement Helmert contrasts for visual-field. This enabled us to contrast means for the LVF and RVF directly before contrasting the combined mean of LVF and RVF with the mean for the CVF. This series of contrasts resulted in the intercept corresponding to the grand mean. For each model, we report regression coefficients (*b*), 95% confidence intervals (*CI*), standard errors (*SE*), and *t/z* values. We consider as statistically significant those cases where $|t/z| > 1.96$.

While we considered all trials for error rate analysis, only correct responses were considered for reaction time analysis. Trials were excluded if reaction times were less than 250 ms or greater than 1200 ms³. Following the removal of these cases, we identified and removed outliers using Hoaglin and Iglewicz’s (1987) procedure. For each participant (across

³ Removal procedures were optimised using pilot data prior to pre-registration. After applying these removal procedures and log transformation, RTs approximated a gaussian distribution.

all conditions) we calculated the difference between first and third quartiles of the response time data. Outliers were then defined as those where reaction times were 1.65 times this difference above the third and below the first quartiles. This enabled us to exclude trials with particularly long responses which may have resulted from delayed loading times. These procedures led to the removal of 12.3% of trials.

Results

Error rates

Error rates are shown in Table 2. The GLMM fitted to error rates indicated that there were fewer errors in the CVF relative to the mean of the RVF and LVF (see Table 2). There were fewer errors in the RVF than the LVF. The remaining fixed effects did not significantly impact error rates.

[Insert Table 1 here]

[Insert Table 2 here]

Reaction times

The LMM fitted to log-transformed reaction times indicated that reaction times were faster in the CVF relative to the mean of the RVF and LVF (meeting our first criterion for successful replication). Furthermore, reaction times were faster in the RVF than the LVF (our second criterion for replication). While the main N effect was not significant, the interaction between N and visual field (LVF – RVF) indicated that the N effect differed in the RVF and LVF. In the LVF, high N targets were responded to 24.7 ms faster than low N targets. In contrast, in the RVF high N targets were responded to 10.5 ms slower than low N targets (see Figure 2A). This pattern of effects was near identical to that reported by Perea et al. (2008; see Figure 2B). The second N by visual field interaction (CVF – LVF and RVF) did not significantly impact reaction time. This indicates that when N effects were summated across the LVF and RVF, the N effect did not statistically differ from the N effect in the CVF.

[Insert Figure 2 here]

Exploratory Analyses

In addition to our pre-registered analyses, we conducted exploratory analyses looking at the N effect at the participant level and how lateralisation on the visual half-field task correlated with the magnitude of the N effect.

To date, no study investigating the N effect during lateralized lexical decision has examined the extent to which individual participants show facilitatory or inhibitory effects of N in each visual field. Out of the 30 participants in the current study, 20 showed a facilitatory effect in the LVF, 14 showed a facilitatory effect in the CVF, and 11 showed a facilitatory effect in the RVF. The reaction time differences for individual participants are shown in Figure 3A, with the numerical difference for each participant reported in the Supplemental Materials (Table S1). Furthermore, only 12 out of the 30 participants showed both a facilitatory N effect in the LVF and an inhibitory effect in the RVF, indicating that while this pattern of results is consistent at the population level, there are likely to be subtle individual differences in N effects across hemispheres.

[Insert Figure 3 here]

One potential explanation for the observed differences in the magnitude of the N effect across each visual field is that the N effect is influenced by the extent to which participants are lateralized for language. It seems plausible that those who are left hemisphere language dominant may show one pattern of N effects while those who show bilateral, or even right hemisphere dominance, may show weaker effects of N or even a reversed pattern. To explore this potential explanation, we calculated laterality indices (LI) using correct responses to both high and low N words in either the LVF or RVF: we assume that participants who are left-hemisphere dominant for language should show a RVF advantage

(and therefore would respond faster to RVF words), whereas participants who are right hemisphere dominant would show a LVF advantage.

LIs were calculated using the formula: $100 \times (\text{Right} - \text{Left}) / (\text{Right} + \text{Left})$, where Right refers to the mean reaction time in the RVF and Left refers to the mean reaction time in the LVF. A negative LI on this scale indicates lateralisation to the left hemisphere and a positive LI indicates lateralisation to the right hemisphere. The mean LI was -2.8 ($SD = 4.27$) and a one-sample t -test indicated that there was significant lateralisation to the left hemisphere at the population level, $t(29) = -3.26$, 95% $CI = [-4.42, -1.23]$, $p = 0.001$. A series of Pearson's product-moment correlations indicated that the N effect was not correlated with LIs in the LVF, $r = .04$, 95% $CI = [-.32, .40]$, $p = .818$, CVF, $r = -.02$, 95% $CI = [-.38, .34]$, $p = .908$, or RVF, $r = .07$, 95% $CI = [-.30, .42]$, $p = .727^4$. Figure 3B shows scatterplots of these results.

Study 1: Discussion

In Study 1 we conducted a replication of Perea, Acha, and Fraga (2008) with a sample of English speaking participants. Analysis of error rates and reaction times met our three criteria for successful replication. We reported: (1) reduced errors and faster reaction times in the central visual field relative to the left and right visual fields for word targets; (2) reduced errors and faster reaction times in the right visual field relative to the left visual field for word targets; (3) a facilitative N effect in the left visual field and an inhibitory effect in the right visual field in reaction times for word targets. This pattern of results replicates several prior studies (Lavidor & Ellis, 2002; Mano et al., 2010; Perea et al., 2008) and confirms predictions from the SERIOL model regarding the interaction between N and visual fields.

⁴ We repeated these analyses when including participants with scores below 80 on the Edinburgh Handedness Inventory to increase variability in LIs. Again, a series of Pearson's product-moment correlations indicated that the N effect was not significantly correlated with LIs in the LVF, $r = -.14$, 95% $CI = [-.42, .17]$, $p = .382$, CVF, $r = -.04$, 95% $CI = [-.27, .33]$, $p = .819$, or RVF, $r = .07$, 95% $CI = [-.37, .24]$, $p = .657$.

The observation of a visual field by N interaction at a population level can be interpreted within the context of the SERIOL model. Facilitatory effects of N in the LVF (i.e. right hemisphere) occur as top-down orthographic activation offsets poorer encoding. This implies weak bottom-up input from the letter to word level and increased top-down input to letter nodes due to global activation in the lexicon. By contrast, more efficient letter encoding in the RVF (i.e. left hemisphere) results in a reduced need for top-down input from the word to letter levels.

However, this explanation from the SERIOL model fails to capture the individual differences that we observed in the current study when we considered N effects in each visual field. Our data suggest that only 12/30 participants show strictly facilitatory effects of N in the LVF and inhibitory effects in the RVF, indicating that participant-level characteristics must influence the pattern of this effect. To start to examine which participant-level characteristics influence the strength of the N effect, we explored the extent to which lateralisation on our visual half-field task influenced the size of the N effect across visual fields. The data suggest that there is no statistically reliable relationship. However, it is important to note here that the study was not designed for the assessment of individual differences as participants completed different versions (counterbalanced lists) of the task meaning that the obtained correlations likely reflect some between item variation. Thus, this is one potential avenue for future research.

It is important to note that the -8 ms facilitatory N effect in the CVF we observed is considerably smaller than N effects typically reported under central presentation. For instance, Pollatsek, Perea, and Binder (1999) reported a -26 ms facilitative N effect under central presentation. Several explanations for these observed differences are possible. First, the current study presented stimuli for 150 ms. By contrast, in lexical decision experiments involving only central presentation, stimuli are typically visible until a response is made. As

such, the N effect may be reduced when stimulus presentation is shorter. Such an explanation would explain why Perea et al. (2008) reported a small -2 ms N effect in the CVF. However, such a possibility would be difficult to reconcile with the large -33 ms facilitative CVF N effect reported by Lavidor and Ellis (2002) which adopted the same 150 ms stimulus presentation duration. A second explanation for this reduced N effect in the CVF relates to stimuli selection. In a review of the published literature, Andrews (1997) concluded that facilitatory effects of N are restricted to low-frequency items. Thus selecting a relatively high proportion of high-frequency items has the potential to diminish N effects under central presentation. Indeed, when applying a median split to zipf frequency, the N effect was more pronounced for low-frequency words (-13.7 ms) than for high-frequency words (-1.0 ms).

In comparison to the published literature on lateralized N effects, the reaction times in the current study were considerably longer; as illustrated in Figure 2B. The same is true when comparing our results to other lateralized lexical decision experiments (e.g. Lavidor & Ellis, 2001). Such differences in reaction times between lab- and web-based studies have been observed across a range of experimental paradigms (e.g. Hilbig, 2016; Semmelmann & Weigelt, 2017). Several explanations for this observation exist and relate to environmental factors such as hardware and/or software. First, participants engaging in online studies may be using slower computers than those taking part in lab-based studies and this might influence the absolute millisecond-based measurements negatively. Second, such reaction time differences may be caused by inaccuracies in the browser engine, as they are not optimised for stimulus presentation and the recording of response times. Regarding population level effects, it is reassuring to know that Hilbig (2016) reported near identical frequency effects in a lexical decision task that was carried out under lab- and web-based conditions. This suggests that although reaction times collected online may be longer, they can still accurately capture well-established effects at the millisecond level.

In sum, our replication study adds weight to the claim that effects of N differ across visual fields. However, one question that remains is whether the N effect is reliably different from zero in the RVF. As the current replication study was powered to detect the interaction between N and visual fields, we decided to conduct a systematic review and meta-analysis to address this quandary.

Study 2: A systematic review and meta-analysis

Despite the consensus that N effects differ across visual fields, the direction and magnitude of the N effect appear far more consistent in the LVF than the RVF. While most studies report a facilitatory effect in the LVF, the nature of the effect seems somewhat controversial in the RVF. Therefore, we conducted a systematic review and meta-analysis to synthesise the available data and provide the first statistical analysis examining whether the N effect was reliably different from zero in the LVF and RVF.

Method

The protocol for the systematic review and meta-analysis was pre-registered using the Non-Interventional, Reproducible, and Open (NIRO) Systematic Reviews guidelines (Pickering et al., 2020). No deviations were made from the registered protocol. The registration form, data, and R scripts can be found at osf.io/7ekyw.

Literature Search

The search of the literature followed the PRISMA guidelines (Moher et al., 2009) and covered all articles published between 1st January 1997 and 1st June 2020. Searches were conducted on the 18th June 2020 using several databases containing published and unpublished literature. The following bibliographic databases of published literature were searched: ProQuest, ScienceDirect, and Web of Science. The grey/unpublished literature were searched using: Ethos, OpenGrey.eu, Open Science Framework Pre-Prints, PsychArXiv, and DissOnline.de. Specifically, we searched the following terms in the title,

topic, and abstract fields of each database: (“orthographic neighbo*” AND “hemisphere”), (“orthographic neighbo*” AND “hemifield”), (“orthographic neighbo*” AND “laterali*”), (“orthographic neighbo*” AND “visual field”), (“orthographic neighbo*” AND “visual-field”), (“orthographic neighb*” AND “Visual half field”), and (“orthographic neighb*” AND “visual half-field”). The reference list of key papers were also checked (Andrews, 1997; Ellis, 2004; Lavidor & Ellis, 2001, 2002; Perea, Acha, & Fraga, 2008; Whitney, 2004).

The study selection process is illustrated in Figure 4. The abstracts of articles identified through database searches were screened for duplicates. All duplicates were then removed. Six reviewers then screened titles and abstracts of 232 unique articles using the metagear package (version 0.6; Lajeunesse, 2020), which was also used to store and manage bibliographic data throughout the review process. Reviewers assessed whether the article looked potentially eligible for inclusion and responded with yes, no, or maybe. Each article was screened twice by a different reviewer. The agreement rate was 94.4%. In the case of no agreement, a review was conducted by a third reviewer. From the 232 unique articles, seven were considered eligible for further screening. Four articles were rated as 'maybe' relevant by reviewers, which prompted screening of the methods sections. Of these four, one article was considered eligible for inclusion. Thus, there were eight eligible articles in total.

Each of the eight articles was then evaluated against the inclusion criteria presented in the Supplemental Materials. In short, the studies had to examine the N effect during a lateralized lexical decision and include reaction time as a dependent variable. At this stage, one article was removed as it was a doctoral thesis where the relevant chapters had been published and the corresponding article had already been deemed eligible for inclusion. In addition to reviewing the full text, we checked the Retraction Watch Database (<http://retractiondatabase.org/>) to ensure these articles had not been retracted. None of the articles that passed a full-text review were indicated to have been retracted.

For the 7 included papers we conducted a forward search of the cited literature. In total, 237 studies were identified. After removing duplicates and articles that had previously been screened, 41 additional unique articles were identified. Each article was screened twice by a set of four reviewers. The agreement rate was 93%. For 3 articles where there was disagreement, a third screen was conducted. None of the 41 articles were rated as eligible for inclusion in the meta-analysis.

To summarise, our initial search and screening strategies led to seven articles being identified for use in the current meta-analysis (Fiset & Arguin, 1999; Lavidor & Ellis, 2001, 2002; Lavidor et al., 2006; Mano et al., 2010; Perea et al., 2008; Whitney & Lavidor, 2005).

[Insert Figure 4 here]

Data extraction and effect size calculation

Of these seven articles, two reported multiple experiments (Lavidor & Ellis, 2002; Whitney & Lavidor, 2005). One of these (Lavidor & Ellis, 2002) included a between items manipulation of word length. Rather than aggregate across this manipulation, we treated each level of the manipulation as a separate experiment. Thus, data extraction was conducted for 11 experiments by two reviewers. In addition to extracting values such as sample size, language of stimuli, etc., reviewers extracted mean reaction times for low- and high-N words in both the LVF and RVF. This could be completed for all but one experiment (Fiset & Auguin, 1999). We contacted the authors of the experiment in an attempt to gain summary statistics and/or raw data. However, the authors could not locate the data, thus the experiment was removed from the meta-analysis. For the remaining experiments, both reviewers extracted identical mean reaction times across conditions and experiments.

As each experiment adopted a within-subjects design, it was necessary to obtain an estimate of the correlation between reaction times for high- and low-N words in each visual field. As none of the experiments included in the meta-analysis reported correlations between

reaction times for low- and high-N words across visual fields, we contacted the first, last, and corresponding authors for each of the six articles requesting both the full and aggregate data. The authors indicated that they no longer had access to the data or we received no reply within one month. Thus, we could not obtain an estimate of the correlation between mean reaction times in the high- and low-N condition in either visual field from the published literature. As such, we decided to use our replication study to derive estimates of these correlations. From this dataset, the correlation between reaction times in the low- and high-N condition was $r = .90$ in the LVF and $r = .79$ in the RVF. It is important to note that these correlations are stronger in magnitude than those typically observed in psychological science. This will of course have implications for the outcome of our meta-analysis; as the correlation coefficient increases, the estimate of the variance for each study decreases and precision increases. As such, we felt it necessary to repeat the meta-analysis assuming a number of values for r and report these in the Supplemental Materials.

To estimate the size of the N effect across visual fields, reaction times in the low N condition were subtracted from scores in the high N condition. The variance of the effect size was calculated using Formulas 12.8-12.9 from Borenstein, Hedges, Higgins, and Rothstein (2009). The variance of each study (V) was calculated as:

$$V = \frac{S_{dif}^2}{n}$$

where S_{dif} is the standard deviation of the mean difference for that study and n is its sample size. S_{dif} was calculated as:

$$S_{dif} = \sqrt{S_1^2 + S_2^2 - 2 \times r \times S_1 \times S_2}$$

where S_1 is the standard deviation of the high N condition, S_2 is the standard deviation of the low N condition, and r is the correlation between the means in the high and low N conditions. A summary of the effect sizes for the included experiments are shown in Table 3.

[Insert Table 3 here]

Publication bias

We assessed publication bias by presenting effect sizes against the inverse of their standard error in a funnel plot (Sterne et al., 2011). Symmetrical funnel plots indicate an absence of bias while asymmetrical funnels plots indicate the presence of bias. For the LVF, the funnel plot indicated no clear evidence of asymmetry/publication bias (see Supplemental Materials, Figure S1). To statistically assess asymmetry we conducted Egger's regression (Egger, Smith, Schneider, & Minder, 1997), which indicated no significant evidence for funnel plot asymmetry in the LVF ($t(9) = -.23, p = .821$). Inspection of the funnel plots for the RVF indicated potential asymmetry (see Supplemental Materials, Figure S2). However, the Egger's regression for the RVF indicated no significant evidence for funnel plot asymmetry ($t(9) = -2.06, p = .070$).

Quality assessment

The quality of each study was assessed by two reviewers via the Appraisal Tool for Cross-Sectional Studies (AXIS; Downes, Brennan, Williams, & Dean, 2016). The AXIS tool contains 20 questions grouped under introduction, method, results, discussion, and other. As some of the questions appeared more relevant to clinical work, we used only 15 items here. Ratings for each of the six published articles appear in the Supplemental Materials (S2). Scores ranged from 6-11 for Reviewer 1 and from 7-11 for Reviewer 2.

It is noteworthy that each study used an incredibly similar methodology. All studies relied on right-handers drawn from a university population. Furthermore, the experimental setup was highly consistent across studies. For instance, the presentation time and displacement of stimuli did not vary considerably between studies. Most studies presented stimuli for 150 ms at 2.5° eccentricity.

The studies differed in how they operationalise high- and low-N. Lavidor and Ellis (2002) reported that words had a mean N of 9.5 and 1.0 in their high- and low-N condition. Perea, Acha, and Fraga (2008) reported that the mean N of words in the high- and low-N conditions were 8.4 and 0.3 respectively. It may, therefore, be of interest to examine how the between-study differences in N thresholds influence the magnitude of the effect size across visual fields.

Data analysis

Meta-analysis. Data were synthesised using random-effects meta-analyses. The analyses were conducted in R using the *rma()* function from the metafor package (version 2.4-0; Viechtbauer, 2020) and compared the N effect in each visual field separately. This enabled us to test the size of the N effect directly in each visual field, rather than compare studies for the presence of an interaction as a meta-regression would permit.

For each model, we used the Sidik-Jonkman estimator (Sidik & Jonkman, 2002) to estimate the between-study variance and applied the Hartung-Knapp adjustment (Hartung & Knapp, 2001) when estimating effect sizes and confidence intervals. To statistically quantify the magnitude of heterogeneity, we examined the I^2 and Cochrane's Q. Because the correlation between reaction times for high- and low-N words in each visual field were estimated based on one study, we additionally conducted sensitivity analyses with alternative estimates of this correlation, where $r = .2$, $.5$, and $.8$ constituting to a small, medium, and large effect and report these in the Supplemental Materials. In sum, the outcome of the meta-analysis was consistent irrespective of the size of the correlation coefficient used to calculate the variance of each study.

Meta-regression. We conducted exploratory meta-regression using the *rma()* function from the metafor package to examine how the strength of the N manipulation influenced N effects across visual fields.

Results

Meta-analysis

The results from the meta-analysis are shown in Table 4 and Figure 5. The estimated N effect in the LVF was -31.8 ms, indicating that the N effect is facilitatory in the right hemisphere. In contrast, the estimated effect in the RVF was 10.8 ms, indicating that N is inhibitory in the left hemisphere. Not only was the direction of the effect different across visual fields, so was the size of the effect. Scrutiny of the I^2 statistic indicated moderate heterogeneity between studies comparing N effects in the LVF (Higgins, Thompson, Deeks, & Altman, 2003). The I^2 statistic was larger for studies comparing the N effect in the RVF, indicating increased variation in outcomes for the RVF relative to the LVF. This is an issue we return to in our meta-regression.

[Insert Table 4 here]

[Insert Figure 5 here]

Meta-regression

A series of exploratory meta-regressions were conducted to examine how the strength of the N manipulation influenced the N effect in each visual field. The strength of the N size manipulation was calculated by subtracting the mean low N rating for stimuli from the high N rating. For example, Lavidor and Ellis (2002) report that words had a mean N of 9.5 and 1.0 in their high- and low-N condition. Thus, the strength of the N manipulation was defined as $9.5 - 1.0 = 8.5$. We fit a meta-regression model to the LVF data which indicated that the strength of the N manipulation did not influence the N effect in the LVF, $b = .78$, 95% CI = [-3.28, 4.85], $SE = 2.07$, $p = .706$. An identical model was fit to the RVF data, which yielded a similar pattern of results where the strength of the N manipulation did not influence the N effect, $b = -3.34$, 95% CI = [-7.13, .46], $SE = 1.94$, $p = .085$.

Study 2: Discussion

This meta-analysis aimed to synthesise the available evidence on orthographic neighbourhood size (N) during lateralized lexical decision to estimate whether the N effect is facilitatory or inhibitory in the LVF and RVF. Based on the data, we can conclude that the N effect is facilitatory for words presented in the LVF (i.e. right hemisphere) and inhibitory for words presented in the RVF (i.e. left hemisphere). This indicates that orthographic information is processed differently across the two cerebral hemispheres. We will return to a discussion of the implications of these results in the General Discussion.

What is surprising from this meta-analysis is the level of heterogeneity, despite all studies adhering to a very similar methodology. Inspection of the Cochrane's Q and I^2 indicated substantial to large heterogeneity around the estimates of N effects in each visual field. When conducting the systematic review, it became clear that the strength of the N manipulation fluctuated between studies. However, the size of the N manipulation between studies did not significantly influence the N effect across visual fields.

The source of between study heterogeneity may instead result from differences in stimulus selection. For instance, the size of the N effect in each study will be influenced by factors such as the presence of stimuli with body neighbours and/or a word's frequency of occurrence in natural language. Body neighbours are orthographic neighbours that differ in relation to the first letter position (e.g. *house* / *mouse*) (Andrews, 1997). Body neighbours typically rhyme and have a facilitative effect on lexical decision times (Ziegler & Perry, 1998). Thus, a study that has many words with body neighbours may encounter different N effects than a study with fewer words with body neighbours. Similarly, as N effects appear stronger for low-frequency items (Andrews, 1997) it may be that studies which use a higher proportion of low-frequency items N effects of a different magnitude than a study using few low-frequency items. Other sources of variation may relate to environmental factors such as

contrast and luminosity that are not maintained across experiments. Below, in the General Discussion, we speculate on several participant-level explanations for this.

General Discussion

Over the last two decades, there have been a number of studies on orthographic neighbourhood effects under lateralized lexical decision, but with inconclusive results. Despite the apparent evidence of a facilitatory effect of having many neighbours in the LVF, the direction and magnitude of N effects in the RVF has been somewhat controversial. While almost all studies report an interaction between N and visual field, some studies report the N effect to be facilitatory in the RVF (e.g. Lavidor et al., 2006) while others report it to be inhibitory (e.g. Mano et al., 2010; Perea et al., 2008). The novel contributions of our current work can be summarised in two general points. First, in a replication study, we reported evidence of an interaction between N and visual field confirming that the main N effect differs between the left and right visual fields. Second, in a systematic review and meta-analysis, we reported the N effect to be strongly facilitative in the LVF and inhibitory in the RVF. We discuss these findings within the context of models of word recognition and consider future directions for this line of research.

These findings are compatible with the SERIOL model (Whitney, 2004). The SERIOL model proposes that letter encoding for words presented in the LVF is inefficient or imprecise, because there is a mismatch between the acuity gradient (based on the distance from fixation) and the serial order of letters in the word (see Figure 1). As a result, top-down support from the word level is required to disambiguate the word. Spreading activation among potential word level targets results in more top-down support for words with larger neighbours: hence, N size has a facilitative effect for words presented in the LVF. By comparison, letter encoding is efficient in the RVF, as the letters' serial order matches their proximity to fixation. Because letter identification is rapid in the RVF, there is less chance for

top-down processing to influence letter encoding; explaining negligible effects of N in the RVF. Instead, lateral inhibition between neighbours at the word level may result in competition, and slower reaction times for high N words. The results reported here support the view that the N effect is inhibitory in the RVF at the population level, with Study 2 showing for the first time that this effect is reliably different from zero.

The Multiple Read-Out Model (MROM; Grainger & Jacobs, 1996) also offers an explanation for these findings. Under this account words in a lexical decision task are identified when the critical activation level is reached on one of the appropriate response criteria, namely M or Σ . The M criterion refers to activation of the specific whole-word orthographic representation, whereas Σ refers to overall activation in the orthographic lexicon. According to the MROM the N effect is dependent on which criteria is used to generate most responses, with M leading to an inhibitory effect, and Σ leading to a facilitative effect. Due to the differential processing efficiencies for orthographic input in the two hemispheres, it may be that different criteria are preferentially used across the visual fields (cf. Perea et al., 2008). More specifically, the M criteria may be used more frequently for words presented in the RVF due to more efficient processing in the left hemisphere. In contrast, the less efficient processing in the right hemisphere (LVF) may lead to greater use of the Σ criterion. This would then lead to N having an inhibitory effect for words presented in the RVF and a facilitative effect for words presented in the LVF, which is consistent with the results of the meta-analysis.

It is worth highlighting that although our findings indicate a facilitative N effect in the LVF and an inhibitory effect in the RVF at the *population* level, this pattern was rarely observed within individual participants. In fact, there was considerable heterogeneity in the pattern of N effects seen at the *participant* level. This suggests that there are additional factors that influence the N by visual field interaction, which vary between individuals. This

may reflect the fact that the two effects (facilitation in the LVF, and inhibition in the RVF) result from two different mechanisms (inefficiency of letter encoding, and lateral inhibition at the word level). It is plausible that individual differences may affect these two mechanisms independently.

One important avenue for future research is the extent to which the precision and redundancy of participants' orthographic representations influence word processing under lateralized presentation. Research into lexical quality (as indexed by reading and spelling ability) in other areas of psycholinguistics has broadened our understanding of how participant level characteristics (typically reading and spelling ability) influence response times in lexical decision (Andrews, 2012; Andrews & Lo, 2013; Andrews & Hersch, 2010) and eye movements during reading (Drieghe, Veldre, Fitzsimmons, Ashby, & Andrews, 2019; Parker & Slattery, 2020; Veldre & Andrews, 2014). While no study has explicitly examined the influence of lexical quality on N effects, Andrews and Hersch (2010) investigated how masked neighbour priming effects are influenced by written language proficiency. They reported that poor spellers showed facilitation from a higher frequency orthographic neighbour prime (e.g. *jury*, *FURY*), whilst good spellers showed inhibition. These results indicate that high-quality lexical representations (as indexed by spelling ability) afford rapid activation of the briefly presented primes, which inhibit orthographically similar representations. However, it remains unclear if the precise orthographic knowledge indexed by spelling ability will influence the N effect in a divided visual half-field paradigm. It seems plausible that those with the most robust lexical representations may show inhibitory effects regardless of visual field while those with weaker lexical representations may show inhibition and facilitation, thus explaining some of the between-participant differences in the pattern of N effect in the LVF and RVF. An investigation of the N by visual field interaction in skilled and less-skilled readers would not only inform our interpretation of the between-participant

differences in N effects, but would inform computational models of word recognition that have classically ignored individual differences in the processing of lexical information across cerebral hemispheres.

Alternatively, it may be the case that individual differences in the N size effect in left and right visual fields relates to strength or direction of language lateralization. While approximately 90% of right-handers are left hemisphere dominant for language (Knecht et al., 2000a), a substantial number of individuals demonstrate bilateral or right language dominance (Knecht et al., 2000b). Investigating the relationship between language lateralization and N effects may be informative in understanding the locus of the N effect: if the N effect relates solely to the match between visual acuity and serial letter order (as predicted by the SERIOL model), it should not vary according to language lateralization. However, if it relates instead to the delay caused by callosal transfer from one hemisphere to another (the mechanism used to explain the RVF advantage), lateralization should have an influence on the strength or direction of the N effect.

Finally, we would like to note that as Study 1 was conducted online, we could not control for various environmental factors such as luminance, contrast, the participants' attention or distance from the screen, and these may explain the variability in our data. It is reassuring nonetheless that we managed to replicate the previous literature despite potential room for increased noise. One way to add credibility to our novel findings would be to examine the variability between participants with data collected under lab condition.

Conclusions

Two decades ago it was first reported that words with many orthographic neighbours were processed differently in each visual field (Lavidor & Ellis, 2001). Since then, numerous studies have investigated this interaction with many claiming that the N effect is facilitatory in the LVF. However, the N effect in the RVF remains somewhat controversial. The size of the

587 N effect in the RVF has differed considerably, with estimates ranging from -23.9 to 28.0 ms.
588 However, until now no study has sought to provide a synthesis of the data to determine
589 whether the effect is reliably different from zero. Evidence from our meta-analysis clearly
590 indicates that this effect does exist in the RVF; however, it is less sizable than the N effect in
591 the LVF. These results will have important implications for models of word recognition that
592 specify different contributions of each cerebral hemisphere in word processing. Additionally,
593 we provide the first look at individual differences in the N effect and report that only a subset
594 of participants show both a facilitative N effect in the LVF and an inhibitory N effect in the
595 RVF. This opens an avenue for future work to explore the role of individual differences in the
596 processing of N under lateralized presentation.

597

598

Declarations of Interest

599

The authors declare that they have no known competing financial interests or personal

600

relationships that could have appeared to influence the work reported in this paper.

601

Acknowledgments

602

Portions of this work were presented at the 2020 online meeting of the Experimental

603

Psychology Society and the 61st Annual Meeting of the Psychonomic Society. We would

604

like to thank Dorothy Bishop for continued input and guidance with this body of work.

605

This work was supported by an advanced grant awarded by the European Research Council (project 694189—Cerebral Asymmetry: New directions in Correlates and Etiology [CANDICE]).

References

- Andrews, S. (1989). Frequency and neighborhood effects on lexical access: Activation or search? *Journal of Experimental Psychology: Learning, Memory, and Cognition*, 15(5), 802–814. <https://doi.org/10.1037/0278-7393.15.5.802>
- Andrews, S. (1992). Frequency and neighborhood effects on lexical access: Lexical similarity or orthographic redundancy? *Journal of Experimental Psychology: Learning, Memory, and Cognition*, 18(2), 234–254. <https://doi.org/10.1037/0278-7393.18.2.234>
- Andrews, S. (1997). The effect of orthographic similarity on lexical retrieval: Resolving neighborhood conflicts. *Psychonomic Bulletin & Review*, 4(4), 439–461. <https://doi.org/10.3758/bf03214334>
- Andrews, S. (2012). Individual differences in skilled visual word recognition and reading: The role of lexical quality. In J. S. Adelman (Ed.), *Visual word recognition* (Vol. 2, pp. 151–172). Hove, United Kingdom: Psychology Press.
- Andrews, S., & Hersch, J. (2010). Lexical precision in skilled readers: Individual differences in masked neighbor priming. *Journal of Experimental Psychology: General*, 139(2), 299–318. <https://doi.org/10.1037/a0018366>
- Andrews, S., & Lo, S. (2012). Not all skilled readers have cracked the code: Individual differences in masked form priming. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, 38(1), 152–163. <https://doi.org/10.1037/a0024953>
- Anwyl-Irvine, A. L., Massonnié, J., Flitton, A., Kirkham, N., & Evershed, J. K. (2020). Gorilla in our midst: An online behavioral experiment builder. *Behavior Research Methods*, 52(1), 388–407. <https://doi.org/10.3758/s13428-019-01237-x>
- Balota, D. A., Yap, M. J., Hutchison, K. A., Cortese, M. J., Kessler, B., Loftis, B., ... & Treiman, R. (2007). The English lexicon project. *Behavior Research Methods*, 39(3), 445–459. <https://doi.org/10.3758/bf03193014>

631 Bates, D., Maechler, M., Bolker, B., Walker, S., Bojesen Christensen, R. H., Singmann, H.,
 632 ... & Fox, J. (2019). Package 'lme4'. Reference manual. Available at [https://cran.r-](https://cran.r-project.org/web/packages/lme4/lme4.pdf)
 633 [project.org/web/packages/lme4/lme4.pdf](https://cran.r-project.org/web/packages/lme4/lme4.pdf) [Verified 1 September 2020].
 634 Borenstein, M., Hedges, L. V., Higgins, J. P., & Rothstein, H. R. (2009). *Introduction to*
 635 *meta-analysis*. John Wiley & Sons.
 636 Coltheart, M., Davelaar, E., Jonasson, J. F., & Besner, D. (1977). Access to the
 637 internal lexicon. In S. Dornic (Ed.), *Attention and Performance VI* (pp. 535–555).
 638 Hillsdale, NJ: Erlbaum.
 639 Coltheart, M. & Rastle, K. (1994). Serial processing in reading aloud: Evidence for dual-
 640 route models of reading. *Journal of Experimental Psychology: Human Perception and*
 641 *Performance*, 20, 1197–1211. <https://doi.org/10.1037/0096-1523.20.6.1197>
 642 Crossman, D. L., & Polich, J. (1988). Hemispheric differences for orthographic and
 643 phonological processing. *Brain and Language*, 35(2), 301–312.
 644 [https://doi.org/10.1016/0093-934X\(88\)90114-9](https://doi.org/10.1016/0093-934X(88)90114-9)
 645 Downes, M. J., Brennan, M. L., Williams, H. C., & Dean, R. S. (2016). Development of a
 646 critical appraisal tool to assess the quality of cross-sectional studies (AXIS). *BMJ*
 647 *Open*, 6(12), e011458. <https://doi.org/10.1136/bmjopen-2016-011458>
 648 Drieghe, D., Veldre, A., Fitzsimmons, G., Ashby, J., & Andrews, S. (2019). The influence of
 649 number of syllables on word skipping during reading revisited. *Psychonomic Bulletin*
 650 *& Review*, 26(2), 616–621. <https://doi.org/10.3758/s13423-019-01590-0>
 651 Egger, M., Smith, G. D., Schneider, M., & Minder, C. (1997). Bias in meta-analysis detected
 652 by a simple, graphical test. *BMJ*, 315(7109), 629–634.
 653 <https://doi.org/10.1136/bmj.315.7109.629>

654 Ellis, A. W. (2004). Length, formats, neighbours, hemispheres, and the processing of words
655 presented laterally or at fixation. *Brain and Language*, 88(3), 355–366.
656 [https://doi.org/10.1016/S0093-934X\(03\)00166-4](https://doi.org/10.1016/S0093-934X(03)00166-4)

657 Fiset, S., & Arguin, M. (1999). Case alternation and orthographic neighborhood size effects
658 in the left and right cerebral hemispheres. *Brain and Cognition*, 40(1), 116–118.

659 Grainger, J. (1992). Orthographic neighborhoods and visual word recognition. In R. Frost &
660 L. Katz (Eds.), *Orthography, morphology, and meaning* (pp. 131-146). Amsterdam:
661 Elsevier.

662 Grainger, J., & Jacobs, A. M. (1996). Orthographic processing in visual word recognition: A
663 multiple read-out model. *Psychological Review*, 103(3), 518–565.
664 <https://doi.org/10.1037/0033-295X.103.3.518>

665 Green, P., & MacLeod, C. J. (2016). SIMR: An R package for power analysis of generalized
666 linear mixed models by simulation. *Methods in Ecology and Evolution*, 7(4), 493–
667 498. <https://doi.org/10.1111/2041-210X.12504>

668 Hartung, J., & Knapp, G. (2001). A refined method for the meta-analysis of controlled
669 clinical trials with binary outcome. *Statistics in Medicine*, 20(24), 3875–3889.
670 <https://doi.org/10.1002/sim.1009>

671 Higgins, J. P. T., Thompson, S. G., Deeks, J. J., & Altman, D. G. (2003). Measuring
672 inconsistency in meta-analyses. *British Medical Journal*, 327(7414), 557–560.
673 <https://doi.org/10.1136/bmj.327.7414.557>

674 Hoaglin, D. C., & Iglewicz, B. (1987). Fine-tuning some resistant rules for outlier labeling.
675 *Journal of the American Statistical Association*, 82(400), 1147–1149.
676 <https://doi.org/10.1080/01621459.1987.10478551>

677 Knecht, S., Deppe, M., Dräger, B., Bobe, L., Lohmann, H., Ringelstein, E. B., & Henningsen,
678 H. (2000a). Language lateralization in healthy right-handers. *Brain*, 123(1), 74–81.
679 <https://doi.org/10.1093/brain/123.1.74>

680 Knecht, S., Dräger, B., Deppe, M., Bobe, L., Lohmann, H., Flöel, A., ... & Henningsen, H.
681 (2000b). Handedness and hemispheric language dominance in healthy humans. *Brain*,
682 123(12), 2512–2518. <https://doi.org/10.1093/brain/123.12.2512>

683 Lajeunesse, M. J. (2016). Facilitating systematic reviews, data extraction and meta-analysis
684 with the metagear package for R. *Methods in Ecology and Evolution*, 7(3), 323–330.
685 <https://doi.org/10.1111/2041-210X.12472>

686 Lavidor, M., & Ellis, A. W. (2001). Mixed-case effects in lateralized word recognition. *Brain*
687 *and Cognition*, 46(1-2), 192–195. [https://doi.org/10.1016/S0278-2626\(01\)80063-4](https://doi.org/10.1016/S0278-2626(01)80063-4)

688 Lavidor, M., & Ellis, A. W. (2002). Orthographic neighborhood effects in the right but not in
689 the left cerebral hemisphere. *Brain and Language*, 80(1), 63–76.
690 <https://doi.org/10.1006/brln.2001.2570>

691 Lavidor, M., Hayes, A., Shillcock, R., & Ellis, A. W. (2004). Evaluating a split processing
692 model of visual word recognition: Effects of orthographic neighborhood size. *Brain*
693 *and Language*, 88(3), 312–320. [https://doi.org/10.1016/S0093-934X\(03\)00164-0](https://doi.org/10.1016/S0093-934X(03)00164-0)

694 Lavidor, M., Johnston, R., & Snowling, M. J. (2006). When phonology fails: Orthographic
695 neighbourhood effects in dyslexia. *Brain and Language*, 96(3), 318–329.
696 <https://doi.org/10.1016/j.bandl.2005.06.009>

697 Mano, Q. R., Patrick, C. J., Andresen, E. N., Capizzi, K., Biagioli, R., & Osmon, D. C.
698 (2010). Re-examining format distortion and orthographic neighbourhood size effects
699 in the left, central and right visual fields. *Journal of Research in Reading*, 33(4), 356-
700 373.

701 Mathey, S. (2001). L'influence du voisinage orthographique lors de la reconnaissance des
 702 mots écrits. *Revue Canadienne de Psychologie Expérimentale*, 55(1), 1–23.
 703 <https://doi.org/10.1037/h0087349>

704 Matsuki, K. (2016). RTSurvival: Survival analysis and divergence point estimation methods
 705 for reaction time data. *R package version 0.1*. Retrieved from
 706 <https://github.com/matsukik/RTsurvival>

707 McClelland, J. L., & Rumelhart, D. E. (1981). An interactive activation model of
 708 context effects in letter perception: I. An account of basic findings. *Psychological*
 709 *Review*, 88(5), 375. <https://doi.org/10.1037/0033-295X.88.5.375>

710 Moher, D., Liberati, A., Tetzlaff, J., Altman, D. G., & The PRISMA Group. (2009). Preferred
 711 reporting items for systematic reviews and meta-analyses: The PRISMA statement.
 712 *PLoS Med*, 6(7), e1000097. <https://doi.org/10.1371/journal.pmed.1000097>

713 Oldfield, R. C. (1971). The assessment and analysis of handedness: The Edinburgh
 714 Inventory. *Neuropsychologia*, 9(1), 97–113. [https://doi.org/10.1016/0028-](https://doi.org/10.1016/0028-3932(71)90067-4)
 715 [3932\(71\)90067-4](https://doi.org/10.1016/0028-3932(71)90067-4)

716 Parker, A. J., & Slattery, T. (2020). Spelling ability influences early letter encoding during
 717 reading: Evidence from return-sweep eye movements. *Quarterly Journal of*
 718 *Experimental Psychology*. <https://doi.org/10.1177/1747021820949150>

719 Parker, A. J., Woodhead, Z. V. J., Thompson, P. A., & Bishop, D. V. M. (2020). Assessing
 720 the reliability of an online behavioural laterality battery: A pre-registered study.
 721 *Laterality*. DOI: 10.1080/1357650X.2020.1859526

722 Perea, M. (1998). Orthographic neighbours are not all equal: Evidence using an identification
 723 technique. *Language and Cognitive Processes*, 13(1), 77–90.
 724 <https://doi.org/10.1080/016909698386609>

725 Perea, M., Acha, J., & Fraga, I. (2008). Lexical competition is enhanced in the left
726 hemisphere: Evidence from different types of orthographic neighbors. *Brain and*
727 *Language*, 105(3), 199–210. <https://doi.org/10.1016/j.bandl.2007.08.005>

728 Pickering, J. S., Topor, M., Barbosa Mendes, A., Bishop, D. V. M., Büttner, F. C.,
729 Henderson, E. L., ... Kothe, E. J. (2020). Non-Interventional, Reproducible, and Open
730 (NIRO) Systematic Review guidelines v0.1.
731 <https://doi.org/10.17605/OSF.IO/F3BRW>

732 R Development Core Team. (2016). *R: A language and environment for statistical*
733 *computing*. Vienna, Austria.

734 Reingold, E. M., Reichle, E. D., Glaholt, M. G., & Sheridan, H. (2012). Direct lexical control
735 of eye movements in reading: Evidence from a survival analysis of fixation durations.
736 *Cognitive Psychology*, 65(2), 177-206.
737 <https://doi.org/10.1016/j.cogpsych.2012.03.001>

738 Reingold, E. M., & Sheridan, H. (2014). Estimating the divergence point: A novel
739 distributional analysis procedure for determining the onset of the influence of
740 experimental variables. *Frontiers in Psychology*, 5, 1432.
741 <https://doi.org/10.3389/fpsyg.2014.01432>

742 Reingold, E. M., & Sheridan, H. (2018). On using distributional analysis techniques for
743 determining the onset of the influence of experimental variables. *Quarterly Journal of*
744 *Experimental Psychology*, 71(1), 260–271.
745 <https://doi.org/10.1080/17470218.2017.1310262>

746 Rutherford, B. J., & Mathesius, J. R. (2012). The brain's hemispheres and controlled search
747 of the lexicon: Evidence from fixated words and pseudowords. *Brain and Cognition*,
748 79(3), 188–199. <https://doi.org/10.1016/j.bandc.2012.03.004>

749 Shillcock, R., Ellison, T. M., & Monaghan, P. (2000). Eye-fixation behavior, lexical storage,
 750 and visual word recognition in a split processing model. *Psychological Review*,
 751 107(4), 824. <https://doi.org/10.1037/0033-295x.107.4.824>.
 752 Sidik, K., & Jonkman, J. N. (2002). A simple confidence interval for meta-analysis. *Statistics*
 753 *in Medicine*, 21(21), 3153–3159. <https://doi.org/10.1002/sim.1549>
 754 Sterne, J. A., Sutton, A. J., Ioannidis, J. P., Terrin, N., Jones, D. R., Lau, J., ... & Tetzlaff, J.
 755 (2011). Recommendations for examining and interpreting funnel plot asymmetry in
 756 meta-analyses of randomised controlled trials. *BMJ*, 343, d4002.
 757 <https://doi.org/10.1136/bmj.d4002>
 758 van Heuven, W. J., Mandera, P., Keuleers, E., & Brysbaert, M. (2014). SUBTLEX-UK: A
 759 new and improved word frequency database for British English. *Quarterly Journal of*
 760 *Experimental Psychology*, 67(6), 1176–1190.
 761 <https://doi.org/10.1080/17470218.2013.850521>
 762 Veldre, A., & Andrews, S. (2014). Lexical quality and eye movements: Individual
 763 differences in the perceptual span of skilled adult readers. *Quarterly Journal of*
 764 *Experimental Psychology*, 67(4), 703–727.
 765 <https://doi.org/10.1080/17470218.2013.826258>
 766 Viechtbauer W (2010). Conducting meta-analyses in R with the metafor package. *Journal of*
 767 *Statistical Software*, 36(3), 1–48. <https://www.jstatsoft.org/v36/i03/>.
 768 Whitney, C. (2001). How the brain encodes the order of letters in a printed word: The
 769 SERIOL model and selective literature review. *Psychonomic Bulletin & Review*, 8(2),
 770 221–243. <https://doi.org/10.3758/bf03196158>
 771 Whitney, C. (2004). Hemisphere-specific effects in word recognition do not require
 772 hemisphere-specific modes of access. *Brain and Language*, 88(3), 279–293.
 773 [https://doi.org/10.1016/S0093-934X\(03\)00160-3](https://doi.org/10.1016/S0093-934X(03)00160-3)

774 Whitney, C., & Lavidor, M. (2005). Facilitative orthographic neighborhood effects: The
775 SERIOL model account. *Cognitive Psychology*, 51(3), 179–213.
776 <https://doi.org/10.1016/j.cogpsych.2005.07.001>
777 Ziegler, J. C., & Perry, C. (1998). No more problems in Coltheart's neighborhood: Resolving
778 neighborhood conflicts in the lexical decision task. *Cognition*, 68(2), B53-B62.
779 [https://doi.org/10.1016/S0010-0277\(98\)00047-X](https://doi.org/10.1016/S0010-0277(98)00047-X)
780

Figures

Figure 1. A schematic illustration of the SERIOL model, based on descriptions in Whitney (2004) and Whitney & Lavidor (2005). Here, fixation falls between ‘O’ and ‘W’ with letters in each visual field projecting to the contralateral hemisphere. An acuity gradient is observed at the retina and the edge level (in V1 / V2). In the right hemisphere (but not the left), there is increased bottom-up excitation and left-to-right lateral inhibition, which reverses the acuity gradient to a spatial gradient at the feature level (in V4). Feature information is transferred from the right to left hemispheres, so that at the letter level (in the ventral occipitotemporal cortex, vOT) the serial order of letters in the word is represented by the activation for each letter. Activity spreads across potential matches at the word level (in the left temporal lobe), and top-down excitation back to the letter level facilitates word recognition.

Figure 2. Plots for pre-registered analyses. (A) Violin plots for RT across the six experimental conditions in the current study. The horizontal band reflects the mean in each experimental condition. (B) Means across experimental conditions in the current study and Perea et al. (2008). Both datasets show a facilitatory N effect in the left visual field and an inhibitory N effect in the right visual field. Solid black lines correspond to low N and dashed grey lines correspond to high N. LVF: left visual field, CVF: central visual field; RVF: right visual field.

Figure 3. Plots for exploratory analyses. (A) Reaction time differences (ms) [high N - low N] in each visual field. Dots represent individual participants' data while the boxplot shows the median and interquartile ranges. The horizontal dashed black line represents zero, or no N effect. (B) Scatterplots for the participants' reaction time difference (ms) and laterality index in each visual field. The solid and dashed lines represent the relationship between the two variables. The shaded grey region represents 95% CIs.

Figure 4. A flowchart of the literature search process. The flowchart on the left shows our initial search of the literature. The flowchart on the right shows our forward search of the cited literature from eligible studies.

Figure 5. Forest plot of the N effect in the (A) left visual field and the (B) right visual field. Plotted are the observed study means and 95% CI. The size of squares is proportional to the weight of each study in the analysis (i.e. the inverse of the within-study variance of the sampling distribution).

Table 1. Mean error rates and reaction times for word targets as a function of neighbourhood size (N) and visual field.

	Low N			High N		
	LVF	CVF	RVF	LVF	CVF	RVF
Error Rates (%)	43.2 (49.57)	16.0 (36.69)	26.7 (44.26)	47.3 (49.97)	19.7 (39.78)	30.0 (45.86)
Reaction times (ms)	774.0 (158.10)	684.5 (165.98)	715.6 (178.09)	749.2 (162.94)	676.0 (157.63)	726.1 (185.81)

Note. Standard deviations are shown in parentheses.

Table 2. (G)LMM results for error rates and reaction time analysis.

Measure	Fixed effect	<i>b</i>	95% <i>CI</i>	<i>SE</i>	<i>z/t</i>
Error rates	(intercept)	-1.124	[-1.436, -0.812]	0.159	-7.06
	VF (LVF – RVF)	-0.471	[-0.568, -0.373]	0.050	-9.46
	VF (CVF – LVF and RVF)	-0.418	[-0.483, -0.352]	0.033	-12.53
	N	-0.140	[-0.344, 0.064]	0.104	-1.34
	VF (LVF – RVF) × N	-0.037	[-0.133, 0.060]	0.049	-0.74
	VF (CVF – LVF and RVF) × N	-0.033	[-0.098, 0.031]	0.033	-1.00
log-RT	(intercept)	2.859	[2.836, 2.882]	0.011	250.23
	VF (LVF – RVF)	-0.014	[-0.018, -0.009]	0.002	-5.93
	VF (CVF – LVF and RVF)	-0.015	[-0.017, -0.012]	0.001	-12.04
	N	0.002	[-0.004, 0.007]	0.003	0.59
	VF (LVF – RVF) × N	-0.005	[-0.009, -0.001]	0.002	-2.00
	VF (CVF – LVF and RVF) × N	-0.001	[-0.003, 0.007]	0.001	-0.55

Note. Significant model terms are shown in **bold**.

Table 3. Experiments included in the meta-analysis.

Study	Language	Sample size	Word/cond.	Visual field	
				LVF	RVF
Lavidor & Ellis (2001)	English	24	20	-49.0	-1.0
Lavidor & Ellis (2002), Exp 1	English	24	20	-35.0	1.0
Lavidor & Ellis (2002), Exp 2, 4-letters	English	24	16	-29.0	19.0
Lavidor & Ellis (2002), Exp 2, 5-letters	English	24	16	-13.0	14.0
Lavidor & Ellis (2002), Exp 2, 6-letters	English	24	16	-44.0	20.0
Whitney & Lavidor (2005), Exp 1	English	19	13	-25.0	-3.0
Whitney & Lavidor (2005), Exp 2	English	25	13	-22.0	3.0
Lavidor et al. (2006)	English	11	20	-36.1	-23.9
Perea et al. (2008)	Spanish	30	20	-23.0	28.0
Mano et al. (2010)	English	26	20	-46.0	15.0
Parker et al. (2020) ¹	English	30	20	-24.7	10.5

Note. Positive effects indicate inhibitory effects for N while negative effects indicate facilitatory effects. Lavidor & Ellis, Exp 2 contained targets of different word length. Rather than aggregate over each word length, we included each as a separate study.

¹Study 1 this article.

Table 4. Effect size estimate of the N effect in the LVF and RVF and the 95% confidence intervals from the meta-analysis model. The I^2 and Q values statistically quantify the heterogeneity of the studies included.

Visual field	k	Mean ES (ms)	95% CI	p -value	I^2 (%)	Q
LVF	11	-31.8	[-39.82, -23.69]	< .001	59.43	$Q(10)=38.11, p< .001$
RVF	11	10.8	[2.84, 18.84]	.013	81.75	$Q(10)=69.35, p< .001$

Note. LVF: left visual field; RVF: right visual field; k: number of studies included in the analysis; Q : Cochrane's Q ; I^2 : percentage of variance that can be attributed to between-study heterogeneity.

Supplemental Materials

The role of orthographic neighbourhood effects in lateralized lexical decision:

A replication study and meta-analysis.

Parker, A. J., Egan, C., Grant, J. H., Harte, S., Hudson, B. T., and Woodhead, Z. V. J.

N effects across visual fields for each participant

Table S1. Individual reaction time differences (ms) for the high N - low N stimuli across each visual field. A negative value indicates a facilitatory effect and a positive value indicates an inhibitory effect.

Participant	Visual field		
	Left	Central	Right
1	27.2	19.0	8.9
2	-11.8	-63.2	78.9
3	-19.7	-33.9	76.2
4	-24.7	30.9	77.1
5	-162.8	-66.0	-102.8
6	-7.8	-41.1	25.3
7	30.1	12.2	36.4
8	86.5	-7.2	-62.0
9	-93.5	80.3	-41.7
10	44.8	88.0	119.6
11	3.2	4.4	21.1

12	-55.3	-8.9	13.2
13	3.8	20.3	-60.4
14	-6.4	-66.8	35.3
15	-105.7	-30.4	56.1
16	-23.2	10.0	-36.2
17	139.3	-98.8	124.8
18	-72.5	-33.1	10.7
19	-13.8	10.1	7.6
20	9.3	17.0	10.8
21	-31.1	54.6	7.3
22	98.7	56.6	2.1
23	-6.6	15.2	103.9
24	-132.5	-13.6	106.3
25	-114.0	18.9	-19.8
26	-36.2	-29.9	-55.6
27	-135.6	-4.0	-27.9
28	4.3	10.1	-64.6
29	-70.6	-37.0	-17.6
30	-1.2	32.2	-35.1

Inclusion criteria for the meta-analysis:

- The study investigated the effect of experimentally manipulated orthographic neighbourhood size during a lateralized lexical decision task. Manipulations of N and visual field must be within subjects and the visual field manipulation must include at least left and right visual field presentation.*
- The study reported lexical decision reaction times.
- The study did not involve priming.
- The study used an alphabetic language that is read from left to right.
- Participants were native readers of the language used for the lexical decision task.
- People with history language impairment or dyslexia, or any neurological condition such as epilepsy were excluded. If there is no explicit mention of these factors, then it will be assumed they did not apply.*
- Participants in the studies were aged between 18 and 50 years old.

* Note. Studies which include additional manipulations will be included. We will aggregate data across these conditions so that we have cell means for the 2 (Nsize) by 2 (visual field) manipulation, so long as the manipulation was within item.

** If studies include control groups (i.e. those without language impairment or dyslexia), we will include data for the control group in our analysis.

Visualisation of publication bias

Figure S1. Funnel plots of the studies included in the meta-analysis for the LVF. In the absence of bias, more precise studies are expected to appear narrowly at the top of the plot, whereas less precise studies are expected to scatter more widely at the bottom of the plot.

Figure S2. Funnel plots of the studies included in the meta-analysis for the RVF. In the absence of bias, more precise studies are expected to appear narrowly at the top of the plot, whereas less precise studies are expected to scatter more widely at the bottom of the plot.

AXIS scoring

Table S2. AXIS score for each article included in the meta-analysis.

Article	AXIS score	
	Reviewer 1	Reviewer 2
Lavidor & Ellis (2001)	10	10
Lavidor & Ellis (2002)	9	9
Lavidor et al. (2006)	6	7
Perea et al. (2008)	9	8
Mano et al. (2010)	11	11
Whitney & Lavidor (2005)	8	8

Note. Scores could range between 0 and 15.

Sensitivity Analysis

None of the reviewed studies reported the correlation coefficient for the relationship between reaction times in the high- and low-N conditions in any visual field. Thus, we calculated the variance associated with the observed effect in each study by assuming that the correlation coefficient would be identical to that observed in our replication study. To examine the influence of this estimate on our conclusions, we conducted sensitivity analysis where we repeated the meta-analysis under the assumption that $r = .2$, $.5$, or $.8$. The outcome of these meta-analyses were consistent; the N effect was always facilitatory in the LVF and always inhibitory in the RVF, irrespective of the size of the correlation coefficient used to calculate the variance of each study. The estimates produced when using variable r values are shown in Table S3 along with their corresponding p -values.

Table S3. Estimates (ms) of the N effect in the left visual field (LVF) and right visual field (RVF) when varying the correlation coefficient (r) used to obtain variances and effect sizes in the meta-analysis.

	$r = .2$		$r = .5$		$r = .8$	
	estimate	p -value	estimate	p -value	estimate	p -value
LVF	-32.7	< .001	-32.4	< .001	-31.7	< .001
RVF	16.2	.001	14.6	.002	11.9	.008