1 Awareness is bliss: How acquiescence affects exploratory factor analysis

2 E. Damiano D'Urso, Jesper Tijmstra, Jeroen K. Vermunt, Kim De Roover

3 Tilburg University, The Netherlands

Abstract

Assessing the measurement model (MM) of self-report scales is crucial to obtain valid measurement of individuals' latent psychological constructs. This entails evaluating the number of measured constructs and determining which construct is measured by which item. Exploratory factor analysis (EFA) is the most-used method to evaluate these psychometric properties, where the number of measured constructs (i.e., factors) is assessed, and, afterwards, rotational freedom is resolved to interpret these factors. This study assessed the effects of an acquiescence response style (ARS) on EFA for unidimensional and multidimensional (un)balanced scales. Specifically, we evaluated (i) whether ARS is captured as an additional factor, (ii) the effect of different rotation approaches on the recovery of the content and ARS factors, and (iii) the effect of extracting the additional ARS factor on the recovery of factor loadings. ARS was often captured as an additional factor in balanced scales when it was strong. For these scales, ignoring (i.e., not extracting) this additional ARS factor, or rotating to simple structure when extracting it, harmed the recovery of the original MM by introducing bias in loadings and cross-loadings. These issues were avoided by using informed rotation approaches (i.e., target rotation), where (part of) the MM is specified *a priori*. Not extracting the additional ARS factor did not affect the loading recovery in unbalanced scales. Researchers should consider the potential presence of an additional ARS factor when assessing the psychometric properties of balanced scales, and use informed rotation approaches when suspecting that an additional factor is an ARS factor.

Awareness is bliss: How acquiescence affects exploratory factor analysis

1 Introduction

Self-report scales are ubiquitous in behavioral sciences for assessing individuals with regard to latent constructs (e.g., self-esteem), and evaluating their psychometric properties is crucial for the validity of these assessments. These scales are generally composed of various questionnaire items and, for each item, the respondents rate how much they agree to this item by selecting one response option on a Likert scale. Assessing the psychometric properties of these scales entails, among other things, evaluating the measurement model (MM). The latter indicates the number of latent constructs or factors measured by the items, and which factor is measured by which items. Also, it needs to be determined whether items are good measurements of latent constructs (i.e., how strongly they load on factors), and whether they measure more than one latent construct at the same time (i.e., load on multiple factors).

The most frequently used method to unravel the psychometric properties of newly developed scales is exploratory factor analysis (EFA). Without imposing an assumed structure on the factor loadings, except (possibly) the number of factors, EFA identifies the relations between factors and items by analyzing the item correlations. Because of its advantageous exploratory nature as well as its popularity, EFA is often considered a mandatory step in the context of scale construction (Howard, 2016; Goretzko, Pham, & Bühner, 2019).

An important limitation of self-report scales is that, despite their widespread use, they might not always sufficiently capture the psychological trait being measured (Van Vaerenbergh & Thomas, 2013). In fact, subject responses might not always be consistent with the measured psychological construct (Bolt & Johnson, 2009). These inconsistencies, generally defined as response styles (RSs) or response bias, can be viewed as systematic or stylistic tendencies in the manner respondents use a rating scale when responding to self-report items (Paulhus, 1991). One well-known response style is the acquiescent one, which is a tendency to agree with items regardless of their content (Van Vaerenbergh & Thomas, 2013).

Failing to take into account acquiescence response style (ARS) can harm psychometric analyses in many ways. For instance, ARS can inflate observed means and correlations (Van Vaerenbergh & Thomas, 2013), increase or decrease the strength of relations between factors and items (Ferrando & Lorenzo-Seva, 2010) and result in an additional factor (Billiet & McClendon, 2000). These potential artifacts not only interfere with the psychometric assessment of the properties of a scale but can also invalidate the interpretation of subjects' scale scores (Bolt & Johnson, 2009).

When the scale has been previously validated, the number of factors to be measured, and their zero-loading structure are known *a priori*. In such cases, ARS can be explicitly included in the MM as an additional factor. Previous research has demonstrated how ARS can be easily incorporated in the context of confirmatory factor analysis (Billiet & McClendon, 2000), item response theory (Falk & Cai, 2016) and latent class analysis (Morren, Gelissen, & Vermunt, 2011). One crucial limitation of these confirmatory approaches, however, is the need for *a-priori* knowledge regarding the MM, which is, of course, lacking when the goal is to determine this MM in the first place.

The assessment of a scale's MM can, therefore, be difficult when ARS causes distortions. In EFA, the number of factors is usually evaluated and, upon resolving rotational freedom, an additional factor could be erroneously interpreted as a dimension of the psychological construct of interest, while it is merely a consequence of ARS. In addition, when not taking ARS into account in the rotation, items may seem to measure more than one factor at the same time, or seem to be a bad measurement of a factor (i.e., low loading), which might lead researchers to drop these seemingly malfunctioning items from the scale. Furthermore, in the most extreme case in which most, or all, items are heavily affected by ARS, the whole scale may seem to be dysfunctional.

While some methods have been proposed to reduce the effects of ARS on EFA (Ferrando, Lorenzo-Seva, & Chico, 2003; Lorenzo-Seva & Rodríguez-Fornells, 2006; Ferrando, Morales-Vives, & Lorenzo-Seva, 2016) only a few papers examined the impact of ignoring or being unaware of ARS on the recovery of factor loadings (Ferrando & Lorenzo-Seva, 2010, Savalei & Falk, 2014). The latter studies, however, have mostly dealt with scales

measuring only a single content factor (i.e., unidimensional scales) measured by continuous items or items that may be treated as such (i.e., items with more than 5 categories; Rhemtulla, Brosseau-Liard, & Savalei, 2012), which only partially mirror the features of commonly used self-report scales and preclude investigating the influence of rotation. In addition, none of these studies investigated to what extent ARS is retrieved as an additional factor by commonly used model selection criteria (e.g., Bayesian Information Criterion; Schwarz et al., 1978), which, in empirical practice, would generally precede any further investigation of the loadings. Drawing upon these existing gaps in current research, this paper aims to extensively study the impact of ARS on the assessment of the psychometric properties of self-report scales, as well as strategies to account for ARS when using EFA. This investigation comprises a simulation study on unidimensional and multidimensional scales for two types of data (i.e., ordinal and approximately continuous data). In addition, we simulated a null scenario (i.e., without an ARS factor) that served as a point of comparison. By means of this simulation study, we will assess: (i) how often and in which conditions different model selection criteria retain the additional ARS factor, (ii) the effect of different rotation approaches on the recovery of the content and ARS factors when the additional ARS factor is retained, and (iii) the effect of (not) retaining the ARS factor on the recovery of the (properly rotated) factor loadings and correlations.

The remainder of the paper proceeds as follows: in Section 2 we provide a general introduction to EFA and how ARS can affect some of its main steps, namely: dimensionality assessment and factor rotation. For factor rotation, we discuss two types of rotation, namely rotation to simple structure (i.e., as one usually does when unaware of a potential ARS) and informed rotation approaches (e.g., rotation to a partially specified target that takes the potential ARS factor into account). Section 3 focuses on a simulation study that evaluates the performance of EFA in assessing the psychometric properties of unidimensional and multidimensional scales (with and without the presence of ARS). Finally, in Section 4, recommendations are formulated based on the results of the simulation study along with limitations of the current study and future research directions.

2 Theoretical Framework

2.1 Factor analysis model with ARS

Consider that continuous responses by N subjects on J items are collected in a data matrix $\mathbf{X}$. Let us assume that each item response is a measure of three common factors: (i) two intended-to-be-measured (i.e., content) factors $\boldsymbol{\eta}_1$ and $\boldsymbol{\eta}_2$, and (ii) an ARS factor $\boldsymbol{\eta}_{ARS}$. A factor analysis model describes the response x_{ij} of subject i on item j as:

$$x_{ij} = \nu_j + \lambda_{j1}\eta_{i1} + \lambda_{j2}\eta_{i2} + \lambda_{jARS}\eta_{iARS} + \epsilon_{ij} \quad (1)$$

where ν_j is an item-specific intercept, λ_{j1} , λ_{j2} and λ_{jARS} are the loadings on item j on the three factors, η_{i1} , η_{i2} and η_{iARS} are the factors scores of subject i , respectively, and ϵ_{ij} is the residual. Factors are assumed to be multivariate normally distributed $\sim \text{MVN}(\boldsymbol{\alpha}, \boldsymbol{\Phi})$ ¹, independently of $\boldsymbol{\epsilon}$, which are $\sim \text{MVN}(0, \boldsymbol{\Psi})$, with $\boldsymbol{\Psi}$ containing the unique variances ψ_j on the diagonal and zeros on the off-diagonal.

When using exploratory factor analysis (EFA; Lawley & Maxwell, 1962) as a first step in assessing the psychometric properties of a scale, the factors in (1) are not (yet) labeled (i.e., researchers do not have or impose *a priori* assumptions on whether a factor corresponds to a certain content factor or an ARS). Also, the assumption of continuous item responses often cannot be safely made, especially in the case of ordered-categorical variables (e.g., a Likert-scale item with “disagree”, “neither agree nor disagree”, and “agree” as response options). In that case, it is better to assume that the data matrix $\mathbf{X}$ is composed of polytomously scored responses that can take on C possible values with $c = \{0, 1, 2, \dots, C - 1\}$. In a categorical EFA model, it is assumed that each of the observed responses is obtained from a discretization of a continuous unobserved response variable x_{ij}^* via some thresholds parameters $\tau_{j,c}$. The threshold parameters indicate the separation between the response categories, where the first and last thresholds are defined as $\tau_{j,0} = -\infty$ and $\tau_{j,C} = \infty$, respectively. In formal terms:

¹This distribution might not be realistic for the ARS factor η_{iARS} if one keeps in mind that a score < 0 would indicate a tendency to disagree. A more suitable distribution for ARS will be considered when generating the data in the simulation study section.

$$x_{ij} = c, \quad \text{if } \tau_{j,c} < x_{ij}^* < \tau_{j,c+1} \quad c = 0, 1, 2, \dots, C - 1. \quad (2)$$

136 A categorical EFA model for the vector of scores $\mathbf{x}_i^*$ of subject i can be specified as:

$$\mathbf{x}_i^* = \boldsymbol{\nu}^* + \mathbf{\Lambda}\boldsymbol{\eta}_i + \boldsymbol{\epsilon}_i \quad (3)$$

137 where $\boldsymbol{\nu}^*$ is a J -dimensional vector of latent intercepts (i.e., intercepts of the unobserved
138 response variables in $\mathbf{x}_i^*$), $\mathbf{\Lambda}$ is a $J \times Q$ matrix of factor loadings, $\boldsymbol{\eta}_i$ is a Q -dimensional
139 vector of scores on the Q factors, $\boldsymbol{\epsilon}_i$ is a J -dimensional vector of residuals. Gathering the
140 loadings of the unlabeled factors in a matrix $\mathbf{\Lambda}$, the model implied covariance matrix $\boldsymbol{\Sigma}$
141 is obtained as:

$$\boldsymbol{\Sigma} = \mathbf{\Lambda}\boldsymbol{\Phi}\mathbf{\Lambda} + \boldsymbol{\Psi}. \quad (4)$$

142 Polychoric correlations are generally used as the input for categorical EFA, where the
143 correlation between ordinal items is computed as the correlation of the standard bi-
144 variate normal distribution of their latent response variables x_{ij}^* (Ekström, 2011). Fur-
145 thermore, they are known to produce unbiased parameters estimates in factor analysis
146 models (Babakus, Ferguson Jr, & Jöreskog, 1987; Rigdon & Ferguson Jr, 1991), whereas
147 with Pearson correlations, which are commonly used for estimating EFA with continuous
148 item responses, the correlations among ordered-categorical items are commonly underes-
149 timated (Bollen & Barb, 1981).

150 2.2 Potential effects of ARS on factor rotation

151 Factors obtained from EFA have rotational freedom (i.e., rotating them does not affect
152 model fit; Browne, 2001), which should be resolved to obtain an interpretable solution.
153 Commonly, the goal is to strive for a factor structure that adheres to the rules of simple
154 structure, for which different criteria can be applied to minimize the variable complexity
155 (i.e., number of non-zero loadings per variable), the factor complexity (i.e., number of non-
156 zero loadings per factor) or a combination of both (Schmitt & Sass, 2011). In this paper,
157 we focus on minimizing the variable complexity by means of oblique simple structure
158 rotation (i.e., allowing the factors to become correlated) because there are little to no

theoretical reasons to assume that: (i) the content factors are uncorrelated in case of multidimensional constructs, and that (ii) the ARS factor is not correlated with the content factor(s) (Weijters, Geuens, & Schillewaert, 2010). Note that minimizing the variable complexity matches the idea of non-ambiguous items that are clear measurements of only one factor (Rao, 1997).

Simple structure can be pursued with uninformed or informed rotation approaches, where the former applies no a priori assumptions on the MM structure and the latter involves rotating to a (partially) specified target based on such *a priori* assumptions. To exemplify how (un)informed simple structure rotation can be affected by the presence of an ARS, we make use of an illustrative example, whose loadings are displayed in Table 1. Specifically, the top part of the table displays the (partially) specified targets for the informed rotation approaches, while the bottom part displays the different sets of rotated loadings. Moreover, the values in the target original matrix were used as the population values of the loadings to generate the data with $N = 10,000$ - implying that the estimated loadings are likely very close to the population values. A visual representation of this model is depicted in Figure 1, where $X_1 - X_{12}$ represent item responses².

2.2.1 Uninformed rotation. Uninformed simple structure rotation tries to achieve simple structure by minimizing a rotation criterion, without applying any user-specified expectations regarding the MM. Several oblique rotation criteria are available. One is (Direct) oblimin (Clarkson & Jennrich, 1988), which is widely-used and offered by popular statistical packages (e.g., SPSS, STATA); others are promax (Hendrickson & White, 1964), promin (Lorenzo-Seva, 1999) and geomin (the default in Mplus; Asparouhov & Muthén, 2009; Yates, 1988).

In the example, we rotated the estimated unrotated loadings using oblimin, and the

²Note that the multidimensional factor model depicted in Figure 1 is substantively different from a bi-factor model (i.e., with a general factor; Holzinger & Swineford, 1937) due to the differences in sign for the loadings that are negative for the content factors and positive for the ARS factor, while for unbalanced scales (i.e., only positive loadings) the model in Figure 1 will be mathematically equivalent to a bi-factor model. Since both types of scales (i.e., balanced and unbalanced) will be addressed in this paper, bi-factor rotation approaches will not be discussed.

results are displayed in the bottom part of Table 1. The oblimin rotated loadings illustrate how, by using uninformed simple structure rotation, the original factor structure is not recovered. For example, item 4 and item 8 load moderately on all factors, and, without further investigations, one might decide to erroneously discard these two items from the scale. This result is not surprising, since previous research already established that, in the case of items loading on multiple factors (here due to the ARS factor), uninformed simple structure rotation criteria perform sub-optimally (Lorenzo-Seva, 1999; Ferrando & Seva, 2000; Schmitt & Sass, 2011). It is interesting to observe how, in order to pursue simple structure, the rotation tries to separate the positive and negative poles of the two content factors. However, with only three factors this cannot be achieved, and, as a result, it produces many small and moderate crossloadings that seem to correspond with such a tendency to separate the different poles of each content factor. For example, the loadings of η_1 that are negative in the population (i.e., items 3 and 7) become primary loadings on the third factor, whereas the negative loadings on η_2 (i.e., items 4 and 8) become moderate loadings on all factors.

2.2.2 Informed rotation. In informed rotation approaches (e.g., target rotation, Browne, 2001) assumptions regarding the MM are made explicit in a user-specified target loading matrix. The loadings are, then, rotated to approximate this target loading matrix, which does not need to be fully specified (i.e., some elements may be unspecified). The specified elements can be zero or take on any value for the non-zero loadings, but, in many practical applications, it is recommended to specify only the zero loadings since precise values for the non-zero loadings are rarely, if ever, known prior to estimating the model (Browne, 2001). Furthermore, some studies have highlighted the robustness of partially (or semi-) specified target rotation when the zero target values are left unspecified and the non-zero target values are misspecified (Myers, Ahn, & Jin, 2013; Myers, Jin, Ahn, Celimli, & Zopluoglu, 2015), but the generalizability of these results to fully specified target rotation as well as to misspecification of the zero loadings (e.g., erroneously specifying a non-zero loading as zero) remains unclear (Garcia-Garzon, Abad, & Garrido, 2019).

In the top part of Table 1, two different fully-specified target matrices are displayed, that is, one with the data-generating values, and one in which the structure was specified using zeros and ones (as is often done in practice), and the corresponding rotated loadings are shown below these target matrices. In both cases, the rotated factor loadings as well as the factor correlations are well recovered, which highlights the suitability of informed rotation approaches in the presence of violations of simple structure, for instance, due to an ARS factor.

In practice researchers rarely know the full structure of the MM *a priori*, and, in order to avoid misspecification of the unknown elements in the target, semi-specified target rotation can be used, where the unknown target elements are left unspecified. Table 1 displays a semi-specified target matrix, specifying only the zero loadings, and the corresponding rotated loadings at the top and bottom part, respectively. The semi-specified target rotated loadings clearly show how zero and non-zero loadings as well as the factor correlations can be accurately recovered by specifying only part of the assumed factor structure in the target. Note that the loadings are recovered as well as with the rotation towards the fully-specified target matrices.

2.3 Potential effects of ARS on dimensionality assessment

Until now it was assumed that the additional ARS factor is retained, which might not always be the case in empirical applications. In fact, in EFA, the number of factors needs to be determined, and this decision generally relies on both “objective” criteria and subjective judgment (i.e., interpretability). A popular objective criterion for maximum likelihood (ML) factor analysis is the Bayesian Information Criterion (BIC; Schwarz et al., 1978), which is a function of how well a model fits the data (i.e., log-likelihood) and the model’s complexity (i.e., number of freely estimated parameters). For a model M , the BIC is calculated as

$$BIC = -2\text{LogLikelihood}(M) + fp \ln(N). \quad (5)$$

where fp indicates the number of free (or estimated) parameters. Even though this criterion is commonly used in empirical practice to determine the number of factors, it

may malfunction if multivariate normality cannot be safely assumed like in the case of ordered-categorical data, and in such cases other approaches might be preferred. One of these alternative approaches is parallel analysis (PA; Horn, 1965), which takes sampling variability into account when selecting the number of factors. In PA, the eigenvalues of the factors estimated from an empirical (polychoric) correlation matrix are compared to the distribution of the eigenvalues estimated from a number of randomly generated (polychoric) correlation matrices (e.g., 20) of the same size as the empirical ones. Afterwards, a factor is retained if its eigenvalue is larger than a given cut-off in the distribution of the eigenvalues obtained from the randomly generated data. Another flexible procedure to determine the numbers of factors is the CHull procedure (Ceulemans & Kiers, 2006; Lorenzo-Seva, Timmerman, & Kiers, 2011), which can be considered as a generalization of the scree test (Cattell, 1966) that aims to balance model fit and complexity. This goal is achieved by first creating a plot of a goodness-of-fit measure against the degree of freedom and, then, selecting the solution which is on or close to the elbow of the higher boundary (convex hull) of the plot by means of a scree test. Lorenzo-Seva et al. (2011) suggested to use the *common part accounted for* index (CAF; Lorenzo-Seva et al., 2011) as a goodness-of-fit measure. The CAF index expresses the degree to which the extracted factor(s) capture the common variance in the data. To calculate the CAF, first the Kaiser-Meyer-Olkin (KMO; Kaiser, 1970; Kaiser & Rice, 1974) index is calculated on the estimated residual correlation matrix Ψ_q of a factor model with q factors. Then, the CAF for a model with q factors is obtained as $CAF_q = 1 - \text{KMO}(\Psi_q)$. The values of the CAF index range from 0 to 1, where values close to 1 indicate that no substantial amount of common variance is left in the residual matrix after extracting q factors. A crucial advantage of the CAF compared to other goodness-of-fit measures is that it can be calculated for a model with no factors, in which case the residual correlation matrix is equal to the empirical correlation matrix. For a detailed overview of “objective” model selection criteria we refer the reader to Lorenzo-Seva et al. (2011). Note that the results obtained from these criteria should be supplemented with substantive knowledge of the measured psychological construct (Henson & Roberts, 2006; Lorenzo-Seva et al., 2011).

Different aspects might play a role in retaining (i.e., selecting) an ARS as an additional factor. For example, various studies suggest that an ARS factor can be conceptualized as a weak factor (i.e., with items showing weak to moderate loadings; Ferrando, Condon, & Chico, 2004; Danner, Aichholzer, & Rammstedt, 2015), potentially making it harder to capture by “objective” model selection criteria. Furthermore, scales that are unbalanced (i.e., with only positively worded items) or partially balanced (i.e., with few negatively worded items) might hamper the detection of an additional ARS factor since it would either be more difficult to differentiate it from the content factor(s), or even impossible in the case of unbalanced unidimensional scales (Ferrando & Lorenzo-Seva, 2010; Savalei & Falk, 2014).

Equally important, an ARS might seriously affect the assessment of the MM regardless of it being retained (i.e., an additional factor selected) in the model selection step or not. In fact, as shown in the illustrative example in Section 2.2, conclusions with regard to the MM are misleading if the ARS factor is retained and the loadings are rotated using uninformed simple structure rotation approaches. Alternatively, failure to select the ARS factor could result in biased loadings on the content factor(s) and bias in the factor correlations. An example of the latter is presented in Figure 2, where, after generating data using the model in Figure 1, a two-factor model was estimated (i.e., ignoring the ARS factor) and the estimated loadings were rotated using oblimin. The results displayed in Figure 2 indicate that not taking the ARS factor into account caused most loadings to be under/overestimated.

3 Simulation study

To evaluate the impact of an ARS on the assessment of the psychometric properties of unidimensional and multidimensional scales using EFA, a simulation study was conducted. The psychometric properties of interest in this simulation study were: (i) the selected number of dimensions (i.e., number of factors), (ii) the recovery of factor loadings and correlations when ARS was taken into count (i.e., extracted), and (iii) the recovery of factor loadings and correlations when ARS was ignored (i.e., not extracted). As a point

of comparison a null scenario (i.e., without an ARS factor) was simulated, the results of which are reported in the Appendix.

In particular, the following 6 factors were manipulated:

- The number of subjects N at 2 levels: 250, 500;
- The number of categories C for each item at 3 levels: 3, 5, 7;
- The type of scale at 2 levels: balanced, unbalanced;
- The number of content factors Q at 2 levels: 1, 2;
- The number of items J per factor at 2 levels: 12, 24;
- The strength of the ARS factor at 3 levels: small, medium and large.

The sample size of 250 is in line with the recommended minimal sample for obtaining precise factor loading estimates in the presence of moderate item communalities (Fabrigar, Wegener, MacCallum, & Strahan, 1999; MacCallum, Widaman, Zhang, & Hong, 1999). Furthermore, the manipulated levels for the number of categories were chosen to represent: (i) items that should be treated as ordinal (i.e., 3 categories), (ii) items that can be treated as continuous (i.e., 7 categories), and (iii) items that can be treated both as ordinal and continuous (i.e., 5 categories) (Rhemtulla et al., 2012). In addition, both balanced and unbalanced scales were included, since the former are generally suggested and preferred to detect ARS (Ferrando & Lorenzo-Seva, 2010; Van Vaerenbergh & Thomas, 2013), whereas the latter is representative of most empirical applications (Ferrando & Lorenzo-Seva, 2010). Finally, both unidimensional and multidimensional scales were simulated. A full-factorial design was used with 2 (number of subjects) $\times$ 3 (number of categories) $\times$ 2 (type of scale) $\times$ 2 (number of factors) $\times$ 2 (number of items) $\times$ 3 (strength of ARS) = 144 conditions. For each condition 100 replications were generated resulting in 14400 data sets.

3.1 Methods

3.1.1 Data Generation. To be able to use the *mirt* package (Chalmers, 2012) to generate the data we used a Q -dimensional normal ogive graded response model (no-

GRM) as the data generating model. This model allowed us to more flexibly generate data with varying numbers of categories while not substantially deviating from a factor model. In fact, parameters in the noGRM are directly related to those of a categorical factor model like the one described in Equation (2) and (3) (Takane & De Leeuw, 1987; Kamata & Bauer, 2008). Specifically, for a unidimensional noGRM, a discrimination parameter α_j can be converted to a factor loading λ_j as $\lambda_j = \frac{\alpha_j}{\sqrt{1+\alpha_j^2}}$ (Kamata & Bauer, 2008). The population values of the model parameters reparametrized in a categorical confirmatory factor analysis fashion are displayed in Table 2 both for unidimensional and multidimensional scales.

To simulate balanced scales, for the content factor(s) half of the loadings were positive (i.e., indicative items), and the other half were negative (i.e., contra-indicative items), whereas all loadings were positive to simulate unbalanced scales. Furthermore, as can be observed from the population values displayed in Table 2, the distance between the first threshold of the easiest and the most difficult item was 2 standard deviations (e.g., for items with three categories, first threshold of item 1 = 0, and first threshold of item 12 = 2). To avoid estimation issues (e.g., non-convergence), we only accepted data sets where each category for each item contains at least a single observation. In the rare cases where a category was not present among the generated scores for a specific item, the entire data generation process was repeated until all response categories were observed. To match the idea that an ARS is conceptualized as a tendency to agree with items regardless of their content, the ARS factor scores were sampled from a right-censored normal distribution. With this distribution we could simulate subjects who either did or did not show an ARS (i.e., have a positive or zero factor score on the ARS dimension), without allowing for scores representing a negative ARS (i.e., disagreeing tendency). Furthermore, with regard to the three levels of the ARS factor, the values of the loadings for the small, medium and large ARS scenarios were .218, .343 and .506, respectively³. The effects on

³The loading values are converted from discrimination parameters of .38, .62 and 1, which were chosen such that the ARS factor affected the item responses drastically less than, less than or as much as (one of) the content factors, respectively.

the items' univariate distribution of a small, medium and large ARS for the generated data are illustrated by the example shown in Figure 3, where data were generated for an item with 5 categories, 10,000 observations, and, using the same thresholds of the seventh item in Table 2, where $\tau_j = \{-3.091, -1.091, -0.909, -2.909\}$. This example clearly shows how the higher categories (i.e., 4 and 5) are more often selected as the strength of the ARS increases.

3.1.2 Data Analysis. The analyses proceeded as follows: first, for each generated dataset, we estimated EFA models with up to three factors, in the case of unidimensional scales, and up to four factors, in the case of multidimensional scales. Furthermore, to study the effects of ARS when treating the data as ordinal or continuous (e.g., ordinal for 3 categories or approximately continuous for 7 categories), the EFA models were estimated both for Pearson correlations and polychoric correlations.

Afterwards, three model selection criteria were considered to evaluate the number of dimensions (i.e., select among the three/four factor models), namely: BIC, Parallel Analysis (PA), and the CHull using the CAF index as a goodness-of-fit measure (See Section 2.3)⁴. For PA, we retained a factor if its eigenvalue was larger than a given 95th percentile in the distribution of eigenvalues obtained from the randomly generated data. Specifically, we used the 95th percentile as the selected cut-off, since it is commonly used in practice (Lorenzo-Seva et al., 2011).

Next, irrespective of the results of the model selection procedures, the loadings for the models with and without the ARS factor were rotated using uninformed and informed rotation approaches. Oblimin was used for uninformed rotation, while, for informed rotations, we used fully specified target (FST) and semi-specified target (SST) rotations, and the target matrices are displayed in Table 3. FST and SST were used both when ARS factor was retained or not in the conditions with multidimensional scales, whereas

⁴Note that the *multichull* package imposes a minimal proportional increase in fit for a more complex model to be included in the hull (see Vervloet, Wilderjans, Durieux, & Ceulemans, 2017 for more details). By default, this minimal increase is set to 0.01. For the simulation study, we lowered it to 0.001, because this minimal value was also not used in Lorenzo-Seva et al. (2011) and a value of 0.01 left the CHULL insensitive to small ARS factors.

only FST was used when the ARS factor was retained for unidimensional scales. For FST rotation, the elements of both the content and the ARS factor were fully specified in the target matrices using ones and zeros for the non-zero and zero loadings, respectively, whereas the zero loadings on the content factor were specified for SST⁵. Oblique Procrustes rotation was used for each target rotation. For the oblimin rotated loadings, the sign of the estimated oblimin factor loadings was reflected to match the ones used to generate the data for the purpose of evaluating the loadings recovery. That is, if the first half of the factor loadings was negative and the second half was positive, the sign of these two halves was reversed (i.e. the factor as a whole was reflected).

3.1.3 Outcome measures. The performance of the different model selection criteria in selecting the number of factors was assessed by calculating the true positive rate (TPR) for the BIC, PA and CHull, both for the models estimated using polychoric correlations and Pearson correlations. Here, the TPR represents the proportion of selecting a two- or three-factor model for unidimensional and multidimensional scales, respectively - that is, the proportion of selecting the additional ARS factor.

Furthermore, the root mean square error (RMSE) between the estimated and true values of the factor loadings was calculated as $RMSE_{loadings} = \sqrt{\frac{1}{JQ} \sum_{q=1}^Q \sum_{j=1}^J (\hat{\lambda}_{jq} - \lambda_{jq})^2}$. Note that this was computed twice for each generated data set: that is, for the model excluding the ARS factor and the model including it (i.e., regardless of the number of factors suggested by the different model selection criteria), and averaged across all replications in a cell of the factorial design. Specifically, the $RMSE_{loadings}$ was calculated for one- and two-factor models for scales with only one content factor, and for two- and three-factors models for scales with two content factors. Then, an $RMSE$ was obtained for the content factor(s) ($RMSE_{loadingsC}$) and the ARS factor⁶ ($RMSE_{loadingsARS}$) when

⁵SST rotations towards the ARS factor, for one- and two-dimensional scales, as well as SST towards the content factor for unidimensional scales were also considered. However, they were discarded due to distorted results in most conditions.

⁶Note that because a right-censored distribution was used to generate the ARS factor scores, the magnitude of the loadings on the ARS factor in the generating model are not directly comparable to the estimated loadings. In fact, for model identification purposes, in EFA the variance of each factor

ARS was extracted, and only for the content factor(s) when ARS was not extracted. In addition to the recovery of the loadings assessed by the $RMSE_{loadings}$, we evaluated whether ARS could cause items to load on more than one factor simultaneously (i.e., cross-loadings), which would cause researchers to conclude that these items are not pure measurements of only one factor. Therefore, for multidimensional scales the recovery of the loadings that are zero in the data-generating model (i.e., on the content factors) was also assessed by calculating the mean maximum absolute bias (MMAB). Specifically, we first selected, for each rotation approach, the item with the maximum absolute difference between the estimated and the “true” (zero) loading, and then we averaged across data sets⁷. In addition, the recovery of the factor correlations between content factors was calculated as $RMSE_{FactorCorr} = \sqrt{(\hat{\phi}_{\eta_1\eta_2} - \phi_{\eta_1\eta_2})^2}$. Similarly to the factor loadings, this measure was computed twice for each generated data set in the conditions with multidimensional scales (i.e., for the model excluding the ARS factor and for the model including it), and averaged across all data sets in a cell of the factorial design.

The null scenario results, i.e., results for the model selection and recovery of factor loadings and factor correlations when ARS was not simulated, are reported in the Appendix (Tables A1 - A4) since assessing the performance of the model selection and rotation approaches in non-ARS conditions is not the goal of this study but only serves as a comparison. In short, their performance was generally satisfactory in all conditions, with a TPR - in this case as the proportion of selecting the correct number of content factors -

is restricted to 1. However, the variance of a right-censored normal distribution is smaller than the variance of a normal distribution, which implies that the loadings are underestimated when imposing a variance of 1. Therefore, when calculating the $RMSE_{loadingsARS}$, the results of which are reported in the Appendix, the values of the estimated loadings on the ARS factor were not subtracted from the values of the original loadings, but from the values of the original loadings rescaled by the variance of a right-censored normal distribution. That is, we multiplied the value of the original loadings on the ARS factor by the standard deviation of a right-censored normal distribution, which is $\approx .583$. This resulted in loadings on the ARS factor of .128, .200 and .295 for the small, medium and large ARS conditions, respectively.

⁷The MMAB was also calculated for the non-zero loadings, and the results are reported in the Appendix.

that was always at or above .90 for all model selection criteria, and the $RMSE_{loadingsC}$ and $RMSE_{FactorCorr}$ were $< .1$ for all rotation approaches.

3.1.4 Data simulation, softwares and packages. The data were simulated and analyzed using R (R Core Team, 2013). Specifically, for generating the data, the R package *mirt* was used (Chalmers, 2012), while EFA and PA were conducted using the *psych* package (Revelle & Revelle, 2015). The CHull procedure was performed using the *multichull* package (Vervloet et al., 2017). For target rotation, we used a function based on Jennrich (2002), which, unlike the one in the popular R package *psych*, does not rescale the factors to improve agreement to the target. In fact, rescaling the factors would undesirably distort the FST rotated loadings, that is, both zero and non-zero loadings are rescaled, and thus increased to achieve agreement with the potentially misspecified values for the non-zero loadings.

3.2 Results

3.2.1 Unidimensional scales.

3.2.1.1 Dimensionality assessment. The TPR results for the different model selection criteria in the small, medium and large ARS conditions are displayed in Table⁸ 4. Overall, the performance of the model selection criteria was mostly affected by type of scale (i.e., balanced and unbalanced) and the strength of the ARS. In fact, none of the model selection criteria suggested to retain the additional ARS factor when unbalanced scales were simulated, which could be due to difficulties in distinguishing between a one-factor model with only positive loadings and a two-factor model where all items load positively on both factors (since the difference between the two lies only in the factor correlation of the two-factor model). For balanced scales, the strength of the ARS was the most important design factor in deciding to extract the additional ARS factor or not. As the strength of the ARS increased, the model selection criteria captured the additional ARS factor more frequently, especially when using Pearson-based PA. However, polychoric-based PA rarely suggested to retain the additional ARS factor in the low

⁸The complete results can be found in the appendix in Table A5 to A7

and medium ARS conditions, which is in line with previous research that showed that polychoric-based PA generally underestimates the number of dimensions (Cho, Li, & Bandalos, 2009).

3.2.1.2 Bias with the additional ARS dimension. The $RMSE_{loadings_C}$ results using balanced and unbalanced scales are displayed in Table 5 and Table 6, respectively, whereas we reported the $RMSE_{loadings_{ARS}}$ results for both balanced and unbalanced scales (Tables A8 - A9) in the Appendix. For balanced scales, FST rotation outperformed oblimin regardless of the strength of the ARS, with an $RMSE_{loadings_C}$ that was always $< .1$, and especially lower when EFA was estimated using polychoric correlations. Importantly, oblimin generally resulted in a $RMSE_{loadings_C} \approx .2$, which is not particularly surprising since uninformed rotation approaches are known to perform sub-optimally when simple structure is violated (Lorenzo-Seva, 1999, Ferrando & Seva, 2000; Schmitt & Sass, 2011). For unbalanced scales, $RMSE_{loadings_C}$ was especially large for FST (e.g., $\approx .5$ in the conditions with large ARS when polychoric correlations were used to estimate EFA), which may be due to the difficulties in distinguishing between the content and the ARS factors since they are both specified with high loadings (i.e., 1) in the target matrix. Oblimin rotation often resulted in $RMSE_{loadings_C} > .1$, but its performance was overall better than in the conditions with balanced scales. This is due to the fact that, similarly to what happened in the example discussed in Section 2.2.1, oblimin tries to separate the positive and negative pole of the content factor, whereas in the unbalanced case this cannot happen and it pursues simple structure by reducing all (or most) loadings on the ARS factor to 0.

3.2.1.3 Bias without the additional ARS dimension. The $RMSE_{loadings_C}$ results without the additional ARS dimension using balanced and unbalanced scales are displayed in Table 7. The loadings were generally accurately recovered when ignoring the additional ARS factor, as indicated by an $RMSE_{loadings_C}$ that was always $< .1$. Additionally, taking into account the ordinal nature of the items by estimating the EFA models using polychoric correlations resulted in a lower $RMSE_{loadings_C}$ compared to Pearson-based EFA, except in the conditions with large ARS and unbalanced scales.

3.2.2 Multidimensional scales.

3.2.2.1 Dimensionality assessment. Table 8 displays the TPR results for the dimensionality assessment in multidimensional scales with small, medium and large ARS⁹. The results mostly overlapped with those observed in the conditions with unidimensional scales, where the type of scale and strength of the ARS were the most impactful factors in choosing whether or not the additional ARS factor is retained. The ARS factor was almost never retained in the conditions with unbalanced scales as indicated by the close-to-zero TPRs. One possible explanation is that, by allowing cross-loadings among the factors, the additional ARS factor is easily absorbed by the content factors, and thus difficult to distinguish in the model selection step. For balanced scales, the additional ARS factor was mostly selected in the conditions with medium and large ARS, where both Pearson-based PA and CHull were equally sensitive or more sensitive than the BIC to this additional factor¹⁰. Similarly to the conditions with unidimensional scales, polychoric-based PA was less sensitive to the ARS factor compared to pearson-based PA in the conditions with a medium ARS.

3.2.2.2 Bias with the additional ARS dimension.

3.2.2.2.1 Factor loadings.

The results of the $RMSE_{loadingsC}$ for balanced scales and unbalanced scales are displayed in Table 9 and Table 10, respectively, whereas the $RMSE_{loadingsARS}$ results can be found in the Appendix (Table A14). For balanced scales, informed rotation approaches (i.e., FST and SST) outperformed oblimin, where the latter resulted in an $RMSE_{loadingsC}$ always higher than $>.1$ when large ARS was simulated. Additionally, in the conditions with

⁹The complete results, for all combinations of the manipulated factors, are displayed in Table A10 to Table A12 in the Appendix.

¹⁰Note that, for the CHull, we visually inspected the cases where a solution could not be selected because the hull contained only two points. This happened in around 25% of the cases for the conditions with large ARS and balanced scales, and it was due to a slight decrease in the CAF index in the models with four factors in comparison to the three-factor models, which, thus, were not included in the hull. Visual inspection of these cases showed that the elbow was quite visible for the model with three factors, and thus we regarded these cases as having selected the correct number of factors.

unbalanced scales, all rotation approaches performed sufficiently well (i.e., $RMSE_{loadingsC} < .1$), with the exception of FST in some conditions with small ARS (e.g., $N = 250$ and $C = 3$) - again indicating that the rotation has trouble distinguishing the ARS factor. As noted before, in balanced scales, oblimin rotation tends to separate the positive and negative poles of the content factor, which heavily affects its performance in terms of loadings recovery compared to the unbalanced scales conditions.

Table 11 and Table 12 display the MMAB results for the zero loadings when the ARS factor is extracted for balanced and unbalanced scales, respectively.¹¹ For balanced scales, the MMAB was below .2 for informed rotation approaches, but not for oblimin rotation, for which MMAB was often $> .2$ in conditions with medium ARS and always $> .3$ in conditions with large ARS, and thus is larger than the commonly used cut-off of .2 for “non-ignorable” cross-loadings (Stevens, 1992). Differently, in unbalanced scales, the MMAB was $< .2$ for oblimin rotation and SST, but not for FST, which often resulted in a MMAB $> .3$, especially for small ARS.

3.2.2.2.2 *Factor Correlations.*

The $RMSE_{FactorCorr}$ results for balanced and unbalanced scales are displayed in Table 13. The $RMSE_{FactorCorr}$ was $< .1$ for SST in all conditions, whereas FST had a $RMSE_{FactorCorr} > .1$ only in the conditions with small ARS and unbalanced scales when using Pearson correlations. Additionally, when large ARS was simulated, factor correlations using oblimin rotation resulted in an $RMSE_{FactorCorr}$ of .204 and .207 for Pearson and polychoric correlations, respectively.

3.2.2.3 *Bias without the additional dimension.*

3.2.2.3.1 *Factor loadings.*

The $RMSE_{loadingsC}$ results for multidimensional scales when the ARS factor was not retained are reported in Tables 14 and 15. The $RMSE_{loadingsC}$ was $< .1$ in all conditions and for both uninformed and informed rotation approaches, which suggests that ignoring (i.e., not extracting) the ARS factor did not strongly affect the recovery of factor loadings. Moreover, when comparing the rotation approaches, FST and SST generally performed

¹¹The results for the non-zero loadings are displayed in Table A15 and Table A16 in the Appendix.

as well as or better than oblimin, and, again, the loadings were more accurately recovered when the EFA models were estimated using polychoric correlations.

The MMAB results for the zero loadings in balanced and unbalanced scales are displayed in Table 16 and Table¹² 17. For all rotation approaches, the MMAB was $<.2$ when small or medium ARS was simulated in balanced scales. However, when large ARS was simulated in these scales, all rotation approaches resulted in a MMAB larger than this commonly used cut-off for “non-ignorable” cross-loadings (Stevens, 1992). In contrast, ignoring ARS did not increase the MMAB in the conditions with unbalanced scales as indicated by the MMAB always $<.2$. In fact, in comparison to Table 12 (i.e., when extracting the ARS factor), MMAB is now smaller (when using oblimin and FST) or equally small (when using SST).

3.2.2.3.2 *Factor correlations.*

The $RMSE_{FactorCorr}$ results for both balanced and unbalanced scales are displayed in Table 18. The recovery of the factor correlations was generally satisfactory. Specifically, $RMSE_{FactorCorr} <.1$ in most conditions, except for oblimin rotation in case of a large ARS, which for unbalanced scales resulted in a $RMSE_{FactorCorr}$ of .230 and .235 when using Pearson and polychoric correlations, respectively.

3.3 Conclusions

The simulation study assessed the performance of EFA with regard to the number of suggested factors as well as the recovery of factor loadings and correlations in the presence of ARS both when retaining the ARS as an additional factor or not. The results indicated that, in terms of model selection, the type of scale as well as the strength of the ARS were particularly impactful on the suggested number of factors to retain. In fact, for both unidimensional and multidimensional scales, the additional ARS factor was almost never captured when unbalanced scales were simulated. In the conditions with balanced scales, the additional ARS factor was mostly selected when its strength was medium or large, especially by Pearson-based PA and to a lesser extent by the BIC and the CHull. Thus,

¹²The results for the non-zero loadings are displayed in Table A17 and Table A18 in the Appendix.

in case of balanced scales, selecting an additional factor that may be an ARS factor is a realistic scenario one should be aware of.

In terms of factor rotation when the ARS factor was extracted, the type of scale (i.e., balanced or unbalanced) is an important consideration when choosing how to rotate. For balanced scales, rotating to simple structure (i.e., oblimin) resulted in biased loadings, and the maximal bias on the zero loadings was particularly large. The latter results are relevant for empirical practice, where trying to pursue simple structure in balanced scales with an additional (but unacknowledged) ARS factor might lead to (i) exclude items that seem to measure multiple factors (i.e., with cross-loadings), or (ii) under/overestimate how well the items measure a content factor (i.e., biased primary loading). In contrast, factor loadings were accurately recovered in balanced scales when using informed rotation approaches (i.e., fully- and semi-specified target rotation), which shows that it pays off to be aware of the fact that an additional factor may be an ARS factor. For unbalanced scales, when ARS was extracted as an additional factor in unidimensional scales especially fully-specified informed rotation approaches often failed to accurately recover the size of the loadings (i.e., $RMSE_{loadingC} > .1$) and, in multidimensional scales, only fully-specified target resulted in large cross-loadings (i.e., $MMAB > .2$). Taken together, these findings suggests that, when ARS is extracted as an additional factor, using semi-specified target rotation toward the assumed MM suffices to accurately assess the MM of multidimensional scales regardless of their type (i.e., balanced or unbalanced). Not extracting an additional ARS factor did not affect the factor loading recovery in unbalanced scales. However, ignoring the ARS factor in multidimensional balanced scales generally resulted in large cross-loadings (irrespective of the rotation). Hence, in empirical practice, researchers should be aware of the fact that not retaining an additional ARS factor might lead to erroneous conclusions on the psychometric properties of the questionnaire items in a balanced scale.

4 Discussion

Assessing the psychometric properties of self-report scales is essential to obtain valid measurements of individuals' latent psychological constructs (i.e., factors). This requires investigating the measurement model (MM) by determining the number of factors, their structure (i.e., which factor is measured by which item) and whether items are pure measurements of one factor. These psychometric properties are commonly assessed by exploratory factor analysis (EFA), where it is necessary to (i) evaluate the number of factors to retain, and (ii) solve rotational freedom to enhance the interpretability of these retained factors. By means of a simulation study, we showed that these two aspects are affected by an acquiescence response style (ARS) among the respondents, and that these effects are more severe for balanced rather than unbalanced scales. In what follows, we discuss the implications of these results for empirical practice for the two types of scales separately.

For balanced scales, especially large ARS often resulted in selecting an additional factor. For these scales, when retained, it is crucial to realize that this additional factor may be an ARS factor and to take this into account in the rotation step. In fact, we showed that naively rotating towards simple structure (i.e., assuming that each item measures only one factor) resulted in biased loadings as well as "non-ignorable" cross-loadings. The latter might drive researchers using balanced scales to draw erroneous conclusions when assessing whether items are non-ambiguous measures of a single factor, and whether they should be excluded from the scale (or replaced). This is avoided by using informed rotation approaches, where the additional ARS factor is taken into account by fully or partially specifying *a priori* assumptions or expectations regarding the MM in a target rotation matrix, and specifying the additional factor as a factor with high loadings for all items or leaving it unspecified. Furthermore, in multidimensional balanced scales, not extracting a large ARS factor often resulted in large cross-loadings, irrespective of the rotation. Thus, to properly assess the psychometric properties of a balanced scale, we not only recommend to use informed rotation if an additional factor is extracted but we even advise to extract the additional factor irrespective of whether the model selection criteria

suggests to do so and compare this solution (upon informed rotation) to the one without this additional factor. Note that this result is also relevant to researchers that aim to use exploratory structural equation modeling (ESEM; Asparouhov & Muthén, 2009), where the number of factors is commonly assumed to be known *a priori*, and one, thus, likely disregards the potential presence of an ARS factor.

For unbalanced scales, the additional ARS factor was almost never selected in the model selection step. This may be due to the use of EFA, where the cross-loadings between the factors allow for a lot of flexibility so that the additional ARS factor is easily “absorbed” by the content ones, and thus hardly ever (or never) retained as an additional factor. Furthermore, not extracting ARS as an additional factor did not impact the factor loadings and correlation much, and, thus, when evaluating these psychometric properties, researchers can simply ignore the potential factor. Nevertheless, one should not conclude that ignoring an additional ARS factor in unbalanced scales is completely harmless. It is important to bear in mind that ARS might influence individual estimates with regard to the measured factors (i.e., factor scores), which, however, were not part of our investigation.

In summary, these findings indicate that it is crucial for researchers to beware of ARS and, for balanced scales, it is best to extract this as an additional factor and take its nature into account when rotating the factors. For the latter, our advise is to use semi-specified target rotation since it proved to perform well, and it avoids the potential influence of miss-specifying the size of the primary loadings - even though such an influence was not found in the current paper (Myers et al., 2013; Myers et al., 2015).

While providing useful insights on the effects of ARS on EFA, the generalisability of these results is subject to certain limitations. For instance, in this study we only considered fully balanced or unbalanced scales but not semi-balanced scales. The latter are not uncommon in psychological research since, for some psychological constructs, contra-indicative items may be harder to formulate without facing the risk of measuring something else (Van Vaerenbergh & Thomas, 2013). Moreover, de la Fuente and Abad (2020) recently assessed the effects of ARS on both EFA and random intercept factor analysis (RIFA;

Maydeu-Olivares & Coffman, 2006) with partially unbalanced scales, and showed that factor loadings were severely affected when using EFA (but not RIFA), especially when the size of the loadings differed strongly between indicative and contra-indicative items. However, whether the additional ARS factor was suggested in the model selection step was not investigated by them, and, in future research, it would certainly be interesting to investigate whether the ARS factor would be suggested in the model selection step. An additional limitation of our study is that the data were simulated under conditions where the MMs did not include cross-loadings among the content factors. However, this does not entirely correspond to empirical practice, where cross-loadings are frequently encountered (Li, Wen, Hau, Yuan, & Peng, 2020). Cross-loadings can have an important impact, not only on the number of factors to retain in EFA (Li et al., 2020) but also on the performance of uninformed rotation approaches (Lorenzo-Seva, 1999; Ferrando & Seva, 2000; Schmitt & Sass, 2011).

Open practices: The data and the analysis scripts are freely available and have been posted at <https://osf.io/bn63u/>

5 References

- Asparouhov, T., & Muthén, B. (2009). Exploratory structural equation modeling. *Structural equation modeling: a multidisciplinary journal*, 16(3), 397–438.
- Babakus, E., Ferguson Jr, C. E., & Jöreskog, K. G. (1987). The sensitivity of confirmatory maximum likelihood factor analysis to violations of measurement scale and distributional assumptions. *Journal of marketing research*, 24(2), 222–228.
- Billiet, J. B., & McClendon, M. J. (2000). Modeling acquiescence in measurement models for two balanced sets of items. *Structural equation modeling*, 7(4), 608–628.
- Bollen, K. A., & Barb, K. H. (1981). Pearson's r and coarsely categorized measures. *American Sociological Review*, 232–239.
- Bolt, D. M., & Johnson, T. R. (2009). Addressing score bias and differential item functioning due to individual differences in response style. *Applied Psychological Measurement*, 33(5), 335–352.
- Browne, M. W. (2001). An overview of analytic rotation in exploratory factor analysis. *Multivariate behavioral research*, 36(1), 111–150.
- Cattell, R. B. (1966). The scree test for the number of factors. *Multivariate behavioral research*, 1(2), 245–276.
- Ceulemans, E., & Kiers, H. A. (2006). Selecting among three-mode principal component models of different types and complexities: A numerical convex hull based method. *British journal of mathematical and statistical psychology*, 59(1), 133–150.
- Chalmers, R. P. (2012). mirt: A multidimensional item response theory package for the R environment. *Journal of Statistical Software*, 48(6), 1–29.
- Cho, S.-J., Li, F., & Bandalos, D. (2009). Accuracy of the parallel analysis procedure with polychoric correlations. *Educational and Psychological Measurement*, 69(5), 748–759.
- Clarkson, D. B., & Jennrich, R. I. (1988). Quartic rotation criteria and algorithms. *Psychometrika*, 53(2), 251–259.
- Danner, D., Aichholzer, J., & Rammstedt, B. (2015). Acquiescence in personality questionnaires: Relevance, domain specificity, and stability. *Journal of Research in*

679 *Personality*, 57, 119–130.

680 de la Fuente, J., & Abad, F. J. (2020). Comparing methods for modeling acquiescence
681 in multidimensional partially balanced scales. *Psicothema*, 32(4), 590–597.

682 Ekström, J. (2011). A generalized definition of the polychoric correlation coefficient.

683 Fabrigar, L. R., Wegener, D. T., MacCallum, R. C., & Strahan, E. J. (1999). Evaluat-
684 ing the use of exploratory factor analysis in psychological research. *Psychological*
685 *methods*, 4(3), 272.

686 Falk, C. F., & Cai, L. (2016). A flexible full-information approach to the modeling of
687 response styles. *Psychological Methods*, 21(3), 328.

688 Ferrando, P. J., Condon, L., & Chico, E. (2004). The convergent validity of acquies-
689 cence: An empirical study relating balanced scales and separate acquiescence scales.
690 *Personality and individual differences*, 37(7), 1331–1340.

691 Ferrando, P. J., & Lorenzo-Seva, U. (2010). Acquiescence as a source of bias and model
692 and person misfit: A theoretical and empirical analysis. *British Journal of Mathe-*
693 *matical and Statistical Psychology*, 63(2), 427–448.

694 Ferrando, P. J., Lorenzo-Seva, U., & Chico, E. (2003). Unrestricted factor analytic proce-
695 dures for assessing acquiescent responding in balanced, theoretically unidimensional
696 personality scales. *Multivariate Behavioral Research*, 38(3), 353–374.

697 Ferrando, P. J., Morales-Vives, F., & Lorenzo-Seva, U. (2016). Assessing and controlling
698 acquiescent responding when acquiescence and content are related: A comprehen-
699 sive factor-analytic approach. *Structural Equation Modeling: A Multidisciplinary*
700 *Journal*, 23(5), 713–725.

701 Ferrando, P. J., & Seva, U. L. (2000). Unrestricted versus restricted factor analysis of
702 multidimensional test items: Some aspects of the problem and some suggestions.
703 *Psicológica*, 21(2), 301–323.

704 Garcia-Garzon, E., Abad, F. J., & Garrido, L. E. (2019). Improving bi-factor exploratory
705 modeling. *Methodology*.

706 Goretzko, D., Pham, T. T. H., & Bühner, M. (2019). Exploratory factor analysis: Current
707 use, methodological developments and recommendations for good practice. *Current*

708 *Psychology*, 1–12.

709 Hendrickson, A. E., & White, P. O. (1964). Promax: A quick method for rotation to
710 oblique simple structure. *British journal of statistical psychology*, 17(1), 65–70.

711 Henson, R. K., & Roberts, J. K. (2006). Use of exploratory factor analysis in published
712 research: Common errors and some comment on improved practice. *Educational
713 and Psychological measurement*, 66(3), 393–416.

714 Holzinger, K. J., & Swineford, F. (1937). The bi-factor method. *Psychometrika*, 2(1),
715 41–54.

716 Horn, J. L. (1965). A rationale and test for the number of factors in factor analysis.
717 *Psychometrika*, 30(2), 179–185.

718 Howard, M. C. (2016). A review of exploratory factor analysis decisions and overview
719 of current practices: What we are doing and how can we improve? *International
720 Journal of Human-Computer Interaction*, 32(1), 51–62.

721 Jennrich, R. I. (2002). A simple general method for oblique rotation. *Psychometrika*,
722 67(1), 7–19.

723 Kaiser, H. F. (1970). A second generation little jiffy. *Psychometrika*, 35(4), 401–415.

724 Kaiser, H. F., & Rice, J. (1974). Little jiffy, mark iv. *Educational and psychological
725 measurement*, 34(1), 111–117.

726 Kamata, A., & Bauer, D. J. (2008). A note on the relation between factor analytic and
727 item response theory models. *Structural Equation Modeling: A Multidisciplinary
728 Journal*, 15(1), 136–153.

729 Lawley, D. N., & Maxwell, A. E. (1962). Factor analysis as a statistical method. *Journal
730 of the Royal Statistical Society: Series D (The Statistician)*, 12(3), 209–229.

731 Li, Y., Wen, Z., Hau, K.-T., Yuan, K.-H., & Peng, Y. (2020). Effects of cross-loadings
732 on determining the number of factors to retain. *Structural Equation Modeling: A
733 Multidisciplinary Journal*, 27(6), 841–863.

734 Lorenzo-Seva, U. (1999). Promin: A method for oblique factor rotation. *Multivariate
735 Behavioral Research*, 34(3), 347–365.

736 Lorenzo-Seva, U., & Rodríguez-Fornells, A. (2006). Acquiescent responding in balanced

multidimensional scales and exploratory factor analysis. *Psychometrika*, 71(4), 769–777.

Lorenzo-Seva, U., Timmerman, M. E., & Kiers, H. A. (2011). The hull method for selecting the number of common factors. *Multivariate behavioral research*, 46(2), 340–364.

MacCallum, R. C., Widaman, K. F., Zhang, S., & Hong, S. (1999). Sample size in factor analysis. *Psychological methods*, 4(1), 84.

Maydeu-Olivares, A., & Coffman, D. L. (2006). Random intercept item factor analysis. *Psychological methods*, 11(4), 344.

Morren, M., Gelissen, J. P., & Vermunt, J. K. (2011). Dealing with extreme response style in cross-cultural research: A restricted latent class factor analysis approach. *Sociological Methodology*, 41(1), 13–47.

Myers, N. D., Ahn, S., & Jin, Y. (2013). Rotation to a partially specified target matrix in exploratory factor analysis: How many targets? *Structural Equation Modeling: A Multidisciplinary Journal*, 20(1), 131–147.

Myers, N. D., Jin, Y., Ahn, S., Celimli, S., & Zopluoglu, C. (2015). Rotation to a partially specified target matrix in exploratory factor analysis in practice. *Behavior research methods*, 47(2), 494–505.

Paulhus, D. L. (1991). Measurement and control of response bias.

R Core Team. (2013). R: A language and environment for statistical computing [Computer software manual]. Vienna, Austria. Retrieved from <http://www.R-project.org/>

Rao, C. R. (1997). *Statistics and truth: putting chance to work*. World Scientific.

Revelle, W., & Revelle, M. W. (2015). Package ‘psych’. *The comprehensive R archive network*.

Rhemtulla, M., Brosseau-Liard, P. E., & Savalei, V. (2012). When can categorical variables be treated as continuous? a comparison of robust continuous and categorical sem estimation methods under suboptimal conditions. *Psychological Methods*, 17(3), 354.

- Rigdon, E. E., & Ferguson Jr, C. E. (1991). The performance of the polychoric correlation coefficient and selected fitting functions in confirmatory factor analysis with ordinal data. *Journal of marketing research*, 28(4), 491–497.
- Savalei, V., & Falk, C. F. (2014). Recovering substantive factor loadings in the presence of acquiescence bias: A comparison of three approaches. *Multivariate behavioral research*, 49(5), 407–424.
- Schmitt, T. A., & Sass, D. A. (2011). Rotation criteria and hypothesis testing for exploratory factor analysis: Implications for factor pattern loadings and interfactor correlations. *Educational and Psychological Measurement*, 71(1), 95–113.
- Schwarz, G., et al. (1978). Estimating the dimension of a model. *The annals of statistics*, 6(2), 461–464.
- Stevens, J. (1992). *Applied multivariate statistics for social sciences*. hillsdale, nj: Erlbaum.
- Takane, Y., & De Leeuw, J. (1987). On the relationship between item response theory and factor analysis of discretized variables. *Psychometrika*, 52(3), 393–408.
- Van Vaerenbergh, Y., & Thomas, T. D. (2013). Response styles in survey research: A literature review of antecedents, consequences, and remedies. *International Journal of Public Opinion Research*, 25(2), 195–217.
- Vervloet, M., Wilderjans, T., Durieux, J., & Ceulemans, E. (2017). Multichull: A generic convex-hull-based model selection method. *Computer software manual*. <https://CRAN.R-project.org/package=multichull> (R package version 1.0.0).
- Weijters, B., Geuens, M., & Schillewaert, N. (2010). The individual consistency of acquiescence and extreme response style in self-report questionnaires. *Applied Psychological Measurement*, 34(2), 105–121.
- Yates, A. (1988). *Multivariate exploratory data analysis: A perspective on exploratory factor analysis*. Suny Press.

Table 1

(Semi-) specified targets (top), and rotated loadings using uninformed and informed rotation approaches (bottom) of an EFA model with 12 items and three factors for an illustrative example.

Target Matrices												
				Target Original			Target			Semi-specified target		
				η_1	η_2	ARS	η_1	η_2	ARS	η_1	η_2	ARS
X_1				0.506	0	0.295	1	0	1	NA	0	NA
X_2				0	0.506	0.295	0	1	1	0	NA	NA
X_3				-0.506	0	0.295	-1	0	1	NA	0	NA
X_4				0	-0.506	0.295	0	-1	1	0	NA	NA
X_5				0.506	0	0.295	1	0	1	NA	0	NA
X_6				0	0.506	0.295	0	1	1	0	NA	NA
X_7				-0.506	0	0.295	-1	0	1	NA	0	NA
X_8				0	-0.506	0.295	0	-1	1	0	NA	NA
X_9				0.506	0	0.295	1	0	1	NA	0	NA
X_{10}				0	0.506	0.295	0	1	1	0	NA	NA
X_{11}				-0.506	0	0.295	-1	0	1	NA	0	NA
X_{12}				0	-0.506	0.295	0	-1	1	0	NA	NA
Rotated loadings												
Oblimin				Target Original			Target			Semi-specified target		
				η_1	η_2	ARS	η_1	η_2	ARS	η_1	η_2	ARS
X_1	0.548	-0.071	-0.099	0.483	-0.011	0.294	0.481	-0.008	0.290	0.490	0.010	0.278
X_2	0.144	0.551	0.126	0.009	0.484	0.280	0.007	0.488	0.288	-0.015	0.485	0.288
X_3	-0.145	-0.101	0.520	-0.476	0.009	0.279	-0.477	0.014	0.283	-0.469	-0.012	0.294
X_4	0.254	-0.373	0.303	-0.011	-0.480	0.298	-0.013	-0.476	0.290	0.004	-0.478	0.290
X_5	0.537	-0.068	-0.130	0.497	-0.005	0.265	0.495	-0.002	0.262	0.502	0.003	0.250
X_6	0.108	0.566	0.123	-0.016	0.506	0.258	-0.018	0.510	0.266	0.011	0.508	0.266
X_7	-0.145	-0.079	0.520	-0.475	-0.012	0.276	-0.476	-0.007	0.279	-0.468	0.010	0.290
X_8	0.238	-0.397	0.271	0	-0.493	0.260	-0.001	-0.490	0.252	-0.006	-0.492	0.252
X_9	0.520	-0.085	-0.120	0.476	0.012	0.265	0.474	0.015	0.261	0.482	-0.014	0.250
X_{10}	0.109	0.548	0.115	-0.010	0.490	0.250	-0.012	0.493	0.258	0.005	0.492	0.258
X_{11}	-0.149	-0.085	0.511	-0.472	-0.004	0.267	-0.473	0	0.271	-0.465	0.002	0.282
X_{12}	0.217	-0.403	0.260	-0.008	-0.493	0.238	-0.009	-0.490	0.230	0.003	-0.492	0.231
Factor correlations												
Oblimin				Target Original			Target			Semi-specified target		
				η_1	η_2	ARS	η_1	η_2	ARS	η_1	η_2	ARS
η_1	1.000	0.072	-0.060	1.000	-0.003	-0.010	1.000	0	0.004	1.000	-0.001	0.001
η_2	0.072	1.000	0.063	-0.003	1.000	0.022	0	1.000	0.008	-0.001	1.000	0
ARS	-0.060	0.063	1.000	-0.010	0.022	1.000	0.004	0.008	1.000	0.001	0	1.000

Note. The "Target Original " loadings are the data-generating loadings, and, except for oblimin, the rotated loadings (below) are obtained by rotating towards the target specified in the corresponding columns of the top part of the table.

Table 2

Population values used in the simulation study

	Loadings			Thresholds											
	One factor	Two factors	3 categories	5 categories				7 categories							
item	λ	λ_{C1}	λ_{C2}	τ_1	τ_2	τ_1	τ_2	τ_3	τ_4	τ_1	τ_2	τ_3	τ_4	τ_5	τ_6
X_1	0.506	0.506	0	0	-2.000	0.875	-0.375	-1.625	-2.875	2.125	0.875	-0.375	-1.625	-2.875	-4.125
X_2	0.506	0	0.506	0.182	-1.818	1.057	-0.193	-1.443	-2.693	2.307	1.057	-0.193	-1.443	-2.693	-3.943
X_3	0.506	(-)0.506	0	0.364	-1.636	1.239	-0.011	-1.261	-2.511	2.489	1.239	-0.011	-1.261	-2.511	-3.761
X_4	0.506	0	(-)0.506	0.545	-1.455	1.420	0.170	-1.080	-2.330	2.670	1.420	0.170	-1.080	-2.330	-3.580
X_5	0.506	0.506	0	0.727	-1.273	1.602	0.352	-0.898	-2.148	2.852	1.602	0.352	-0.898	-2.148	-3.398
X_6	0.506	0	0.506	0.909	-1.091	1.784	0.534	-0.716	-1.966	3.034	1.784	0.534	-0.716	-1.966	-3.216
X_7	(-)0.506	(-)0.506	0	1.091	-0.909	1.966	0.716	-0.534	-1.784	3.216	1.966	0.716	-0.534	-1.784	-3.034
X_8	(-)0.506	0	(-)0.506	1.273	-0.727	2.148	0.898	-0.352	-1.602	3.398	2.148	0.898	-0.352	-1.602	-2.852
X_9	(-)0.506	0.506	0	1.455	-0.545	2.330	1.080	-0.170	-1.420	3.580	2.330	1.080	-0.170	-1.420	-2.670
X_{10}	(-)0.506	0	0.506	1.636	-0.364	2.511	1.261	0.011	-1.239	3.761	2.511	1.261	0.011	-1.239	-2.489
X_{11}	(-)0.506	(-)0.506	0	1.818	-0.182	2.693	1.443	0.193	-1.057	3.943	2.693	1.443	0.193	-1.057	-2.307
X_{12}	(-)0.506	0	(-)0.506	2.000	0	2.875	1.625	0.375	-0.875	4.125	2.875	1.625	0.375	-0.875	-2.125

Table 3

Target matrices

Target Matrices								
Unidimensional scales			Multidimensional scales					
FST			FST			SST		
	η_1	ARS	η_1	η_2	ARS	η_1	η_2	ARS
X_1	1	1	1	0	1	NA	0	NA
X_2	1	1	0	1	1	0	NA	NA
X_3	1	1	(-)1	0	1	NA	0	NA
X_4	1	1	0	(-)1	1	0	NA	NA
X_5	1	1	1	0	1	NA	0	NA
X_6	1	1	0	1	1	0	NA	NA
X_7	(-)1	1	(-)1	0	1	NA	0	NA
X_8	(-)1	1	0	(-)1	1	0	NA	NA
X_9	(-)1	1	1	0	1	NA	0	NA
X_{10}	(-)1	1	0	1	1	0	NA	NA
X_{11}	(-)1	-1	(-)1	0	-1	NA	0	NA
X_{12}	(-)1	1	0	(-)1	1	0	NA	NA

Note. FST = Fully-specified target; SST = Semi-specified target.

Table 4

Main effects on model selection TPR for unidimensional scales in function of strength of the ARS and the simulated conditions

Model Selection Unidimensional Scales																		
Small ARS							Medium ARS						Large ARS					
Pearson			Polychoric				Pearson			Polychoric			Pearson			Polychoric		
	CHull	BIC	PA	CHull	BIC	PA	CHull	BIC	PA	CHull	BIC	PA	CHull	BIC	PA	CHull	BIC	PA
N = 250	0.014	0	0.038	0.012	0.001	0	0.084	0.005	0.318	0.092	0.039	0.041	0.440	0.476	0.502	0.420	0.496	0.498
N = 500	0.012	0	0.083	0.013	0	0	0.232	0.129	0.481	0.229	0.277	0.007	0.458	0.499	0.506	0.448	0.501	0.498
C = 3	0.009	0	0.041	0.010	0.001	0	0.081	0.004	0.380	0.099	0.191	0.015	0.418	0.471	0.509	0.385	0.496	0.496
C = 5	0.016	0	0.071	0.019	0	0	0.164	0.099	0.379	0.156	0.134	0.006	0.443	0.492	0.501	0.437	0.499	0.499
C = 7	0.015	0	0.069	0.009	0	0	0.230	0.099	0.440	0.227	0.149	0.050	0.487	0.499	0.501	0.480	0.500	0.500
Balanced	0.020	0	0.105	0.015	0.001	0	0.312	0.134	0.782	0.315	0.315	0.048	0.893	0.975	0.995	0.856	0.996	0.997
Unbalanced	0.006	0	0.016	0.010	0	0	0.005	0	0.017	0.007	0.001	0	0.005	0	0.012	0.012	0.001	0
J = 12	0.021	0	0.065	0.019	0.001	0	0.103	0.006	0.352	0.096	0.073	0.012	0.417	0.475	0.507	0.405	0.497	0.497
J = 24	0.006	0	0.056	0.006	0	0	0.214	0.128	0.447	0.225	0.243	0.036	0.481	0.500	0.501	0.463	0.500	0.500

Note. CHull = convex hull based on the Common Part Accounted For (CAF) index; BIC = Bayesian Information Criterion; PA = parallel analysis.

Table 5

RMSE_{loadingsC} in unidimensional balanced scales when the ARS factor is extracted in function of the simulated conditions

Unidimensional balanced scales - $RMSE_{loadingsC}$ with ARS factor														
Pearson								Polychor						
Small ARS				Medium ARS				Large ARS				Small ARS		
N	J	C	Oblimin	FST	Oblimin	FST	Oblimin	FST	Oblimin	FST	Oblimin	FST	Oblimin	FST
250	12	3	0.178	0.035	0.176	0.036	0.219	0.032	0.148	0.024	0.147	0.026	0.170	0.032
		5	0.148	0.019	0.181	0.026	0.226	0.080	0.134	0.011	0.163	0.007	0.209	0.053
		7	0.167	0.023	0.183	0.020	0.235	0.051	0.155	0.011	0.179	0.011	0.232	0.040
	24	3	0.175	0.051	0.194	0.023	0.234	0.078	0.120	0.014	0.153	0.042	0.206	0.022
		5	0.161	0.028	0.200	0.016	0.227	0.063	0.135	0.007	0.181	0.018	0.208	0.037
		7	0.157	0.035	0.226	0.055	0.218	0.058	0.146	0.023	0.218	0.042	0.210	0.046
500	12	3	0.176	0.065	0.226	0.065	0.241	0.080	0.141	0.012	0.199	0.011	0.214	0.024
		5	0.174	0.024	0.219	0.056	0.268	0.036	0.151	0.007	0.202	0.030	0.232	0.010
		7	0.165	0.037	0.217	0.030	0.284	0.044	0.162	0.024	0.208	0.017	0.275	0.032
	24	3	0.177	0.050	0.265	0.067	0.234	0.051	0.124	0.011	0.242	0.015	0.200	0.011
		5	0.174	0.039	0.295	0.039	0.295	0.049	0.149	0.014	0.283	0.016	0.305	0.023
		7	0.184	0.042	0.242	0.041	0.270	0.026	0.178	0.030	0.237	0.029	0.270	0.014

Note. FST = fully-specified target.

Table 6

$RMSE_{loadingsC}$ in unidimensional unbalanced scales when the ARS factor is extracted in function of the simulated conditions

Unidimensional unbalanced scales - $RMSE_{loadingsC}$ with ARS factor														
Pearson									Polychoric					
			Small ARS		Medium ARS		Large ARS		Small ARS		Medium ARS		Large ARS	
N	J	C	Oblimin	FST	Oblimin	FST	Oblimin	FST	Oblimin	FST	Oblimin	FST	Oblimin	FST
250	12	3	0.172	0.265	0.187	0.206	0.159	0.306	0.140	0.464	0.181	0.394	0.153	0.526
		5	0.155	0.300	0.144	0.348	0.145	0.395	0.149	0.389	0.124	0.449	0.116	0.503
		7	0.153	0.356	0.139	0.327	0.126	0.464	0.156	0.401	0.132	0.372	0.115	0.518
	24	3	0.162	0.244	0.158	0.238	0.136	0.335	0.104	0.438	0.108	0.433	0.084	0.553
		5	0.165	0.236	0.149	0.278	0.082	0.440	0.137	0.320	0.125	0.370	0.042	0.560
		7	0.175	0.207	0.141	0.300	0.045	0.511	0.163	0.247	0.130	0.346	0.048	0.563
500	12	3	0.160	0.255	0.175	0.261	0.158	0.357	0.128	0.451	0.128	0.462	0.133	0.587
		5	0.134	0.311	0.109	0.408	0.097	0.440	0.132	0.401	0.094	0.516	0.108	0.553
		7	0.130	0.394	0.133	0.344	0.093	0.436	0.117	0.441	0.126	0.389	0.088	0.490
	24	3	0.124	0.259	0.099	0.297	0.081	0.328	0.051	0.459	0.043	0.506	0.024	0.547
		5	0.107	0.279	0.070	0.321	0.058	0.363	0.083	0.368	0.039	0.414	0.031	0.467
		7	0.103	0.265	0.056	0.362	0.025	0.407	0.094	0.308	0.046	0.407	0.015	0.456

Note. FST = fully-specified target.

Table 7

$RMSE_{loadingsC}$ in unidimensional scales when the ARS factor is not extracted in function of the simulated conditions

Unidimensional scales - $RMSE_{loadingsC}$ without ARS factor														
Balanced scales									Unbalanced scales					
			Small ARS		Medium ARS		Large ARS		Small ARS		Medium ARS		Large ARS	
N	J	C	Pearson	Polychoric	Pearson	Polychoric	Pearson	Polychoric	Pearson	Polychoric	Pearson	Polychoric	Pearson	Polychoric
250	12	3	0.043	0.015	0.043	0.018	0.042	0.025	0.045	0.019	0.063	0.008	0.034	0.037
		5	0.028	0.006	0.033	0.007	0.091	0.065	0.032	0.009	0.021	0.015	0.009	0.031
		7	0.030	0.018	0.026	0.016	0.069	0.066	0.016	0.006	0.023	0.010	0.021	0.035
	24	3	0.054	0.012	0.025	0.040	0.083	0.029	0.042	0.020	0.044	0.019	0.016	0.054
		5	0.031	0.007	0.018	0.016	0.068	0.042	0.043	0.017	0.030	0.004	0.025	0.057
		7	0.038	0.026	0.057	0.045	0.063	0.051	0.052	0.039	0.022	0.010	0.043	0.058
500	12	3	0.071	0.017	0.070	0.016	0.087	0.032	0.044	0.017	0.042	0.020	0.013	0.056
		5	0.029	0.006	0.060	0.035	0.048	0.021	0.026	0.005	0.010	0.036	0.016	0.046
		7	0.042	0.029	0.034	0.021	0.058	0.047	0.004	0.016	0.015	0.004	0.014	0.028
	24	3	0.051	0.009	0.070	0.019	0.054	0.010	0.035	0.026	0.024	0.041	0.017	0.053
		5	0.041	0.016	0.043	0.021	0.055	0.033	0.028	0.005	0.015	0.014	0.010	0.031
		7	0.044	0.031	0.043	0.031	0.032	0.022	0.032	0.019	0.003	0.013	0.014	0.028

Table 8

Main effects on model selection TPR for multidimensional scales in function of strength of the ARS and the simulated conditions

Model Selection Multidimensional Scales																		
	Small ARS						Medium ARS						Large ARS					
	Pearson			Polychoric			Pearson			Polychoric			Pearson			Polychoric		
	CHull	BIC	PA	CHull	BIC	PA	CHull	BIC	PA	CHull	BIC	PA	CHull	BIC	PA	CHull	BIC	PA
N = 250	0.044	0	0.020	0.033	0	0	0.266	0.001	0.363	0.260	0.087	0.088	0.507	0.453	0.501	0.504	0.497	0.498
N = 500	0.031	0	0.032	0.030	0.002	0	0.402	0.175	0.479	0.379	0.358	0.039	0.510	0.499	0.500	0.511	0.500	0.497
C = 3	0.035	0	0.020	0.031	0.002	0	0.328	0.029	0.411	0.284	0.298	0.104	0.510	0.469	0.501	0.511	0.502	0.500
C = 5	0.038	0	0.018	0.031	0	0	0.320	0.100	0.409	0.316	0.175	0.052	0.509	0.460	0.500	0.504	0.492	0.494
C = 7	0.040	0	0.041	0.032	0	0	0.355	0.135	0.444	0.359	0.195	0.035	0.507	0.500	0.500	0.507	0.500	0.499
Balanced	0.051	0	0.044	0.042	0	0	0.638	0.176	0.838	0.616	0.444	0.128	0.998	0.952	0.999	0.996	0.995	0.995
Unbalanced	0.024	0	0.008	0.021	0.002	0	0.030	0	0.005	0.023	0.001	0	0.020	0	0.002	0.019	0.002	0
J = 12	0.052	0	0.028	0.038	0.002	0	0.280	0.033	0.367	0.263	0.141	0.042	0.512	0.452	0.501	0.512	0.497	0.495
J = 24	0.023	0	0.025	0.025	0	0	0.388	0.142	0.476	0.376	0.304	0.086	0.505	0.500	0.500	0.502	0.500	0.500

Note. CHull = convex hull based on the Common Part Accounted For (CAF) index; BIC = Bayesian Information Criterion; PA = parallel analysis.

Table 9

RMSE_{loadingsC} in multidimensional balanced scales when the ARS factor is extracted in function of the simulated conditions

Multidimensional balanced scales - $RMSE_{loadingsC}$ with ARS factor																			
Pearson										Polychoric									
Small ARS					Medium ARS					Large ARS					Small ARS				
N	J	C	Oblimin	FST	SST	Oblimin	FST	SST		Oblimin	FST	SST		Oblimin	FST	SST	Oblimin	FST	SST
250	12	3	0.081	0.016	0.052	0.095	0.021	0.043		0.132	0.034	0.040		0.046	0.041	0.015	0.066	0.032	0.010
		5	0.065	0.023	0.029	0.086	0.039	0.040		0.142	0.043	0.045		0.049	0.027	0.013	0.075	0.035	0.025
		7	0.064	0.015	0.033	0.084	0.033	0.024		0.149	0.026	0.023		0.057	0.019	0.025	0.076	0.031	0.016
	24	3	0.070	0.036	0.050	0.088	0.037	0.043		0.153	0.059	0.058		0.037	0.013	0.013	0.053	0.026	0.006
		5	0.052	0.023	0.035	0.067	0.021	0.026		0.163	0.046	0.044		0.035	0.014	0.018	0.051	0.012	0.008
		7	0.047	0.027	0.032	0.076	0.027	0.031		0.151	0.038	0.038		0.040	0.024	0.024	0.069	0.021	0.023
	500	3	0.087	0.036	0.062	0.101	0.042	0.047		0.150	0.061	0.062		0.055	0.011	0.024	0.076	0.048	0.007
		5	0.053	0.026	0.026	0.083	0.033	0.041		0.122	0.035	0.034		0.038	0.023	0.009	0.065	0.019	0.022
		7	0.049	0.011	0.023	0.084	0.034	0.030		0.128	0.033	0.034		0.042	0.011	0.014	0.077	0.033	0.022
	24	3	0.056	0.039	0.044	0.093	0.050	0.053		0.137	0.045	0.044		0.020	0.013	0.006	0.064	0.021	0.014
		5	0.038	0.027	0.029	0.070	0.024	0.025		0.128	0.033	0.032		0.020	0.012	0.011	0.055	0.017	0.006
		7	0.037	0.026	0.029	0.072	0.029	0.032		0.146	0.027	0.027		0.030	0.018	0.021	0.065	0.021	0.024

Note. FST = fully-specified target; SST = semi-specified target.

Table 10

$RMSE_{loadingsC}$ in multidimensional unbalanced scales when the ARS factor is extracted in function of the simulated conditions

Multidimensional unbalanced scales - $RMSE_{loadingsC}$ with ARS factor																				
Pearson												Polychoric								
Small ARS					Medium ARS			Large ARS			Small ARS			Medium ARS			Large ARS			
N	J	C	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST
250	12	3	0.072	0.327	0.047	0.074	0.132	0.047	0.049	0.047	0.027	0.035	0.122	0.011	0.033	0.030	0.012	0.022	0.096	0.029
		5	0.046	0.017	0.022	0.047	0.025	0.030	0.038	0.054	0.016	0.028	0.026	0.012	0.031	0.044	0.009	0.020	0.078	0.013
		7	0.053	0.013	0.028	0.041	0.024	0.024	0.026	0.068	0.017	0.044	0.016	0.024	0.037	0.031	0.023	0.021	0.082	0.023
	24	3	0.049	0.389	0.036	0.047	0.204	0.033	0.033	0.053	0.022	0.012	0.033	0.010	0.010	0.028	0.013	0.031	0.094	0.032
		5	0.054	0.322	0.041	0.027	0.030	0.016	0.019	0.063	0.019	0.035	0.054	0.022	0.014	0.043	0.014	0.039	0.093	0.043
		7	0.057	0.454	0.039	0.031	0.011	0.017	0.019	0.070	0.020	0.048	0.396	0.032	0.021	0.009	0.012	0.026	0.083	0.029
500	12	3	0.071	0.292	0.050	0.060	0.026	0.044	0.055	0.034	0.031	0.038	0.019	0.012	0.024	0.039	0.006	0.019	0.071	0.015
		5	0.042	0.013	0.026	0.033	0.022	0.017	0.026	0.048	0.009	0.026	0.023	0.011	0.018	0.036	0.014	0.011	0.073	0.018
		7	0.033	0.013	0.017	0.041	0.010	0.019	0.016	0.050	0.007	0.025	0.022	0.008	0.037	0.014	0.011	0.010	0.062	0.012
	24	3	0.037	0.026	0.029	0.037	0.022	0.030	0.025	0.038	0.018	0.014	0.029	0.019	0.010	0.026	0.015	0.028	0.073	0.032
		5	0.026	0.022	0.019	0.025	0.021	0.019	0.011	0.009	0.007	0.007	0.012	0.004	0.008	0.019	0.005	0.016	0.030	0.020
		7	0.026	0.022	0.021	0.014	0.009	0.010	0.011	0.045	0.006	0.017	0.012	0.013	0.008	0.009	0.004	0.012	0.052	0.011

Note. FST = fully-specified target; SST = semi-specified target.

Table 11

Main effects on MMAB for zero loadings in multidimensional balanced scales when the ARS factor is extracted in function of the simulated conditions

Multidimensional balanced scales - MMAB with ARS factor																			
Small ARS						Medium ARS						Large ARS							
Pearson			Polychoric			Pearson			Polychoric			Pearson			Polychoric				
	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST	
N = 250	0.182	0.150	0.121	0.192	0.158	0.131	0.225	0.169	0.121	0.243	0.188	0.131	0.372	0.146	0.117	0.403	0.148	0.126	
N = 500	0.123	0.104	0.087	0.134	0.108	0.093	0.194	0.120	0.085	0.208	0.133	0.091	0.324	0.099	0.082	0.345	0.103	0.088	
C = 3	0.160	0.131	0.109	0.179	0.143	0.122	0.214	0.142	0.105	0.242	0.175	0.119	0.336	0.132	0.104	0.381	0.137	0.118	
C = 5	0.149	0.129	0.102	0.157	0.131	0.110	0.201	0.140	0.102	0.214	0.152	0.109	0.338	0.124	0.100	0.361	0.126	0.105	
C = 7	0.148	0.122	0.101	0.152	0.125	0.104	0.213	0.152	0.102	0.220	0.155	0.105	0.370	0.111	0.096	0.381	0.114	0.098	
J = 12	0.158	0.125	0.095	0.169	0.129	0.102	0.210	0.156	0.096	0.227	0.174	0.105	0.335	0.125	0.091	0.360	0.123	0.098	
J = 24	0.147	0.130	0.113	0.157	0.137	0.122	0.208	0.133	0.109	0.224	0.147	0.117	0.361	0.120	0.108	0.389	0.128	0.117	

Note. FST = fully-specified target; SST = semi-specified target.

Table 12

Main effects on MMAB for zero loadings in multidimensional unbalanced scales when the ARS factor is extracted in function of the simulated conditions

Multidimensional unbalanced scales - MMAB with ARS factor																		
	Small ARS						Medium ARS						Large ARS					
	Pearson			Polychoric			Pearson			Polychoric			Pearson			Polychoric		
	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST
N = 250	0.171	0.532	0.120	0.181	0.398	0.128	0.171	0.324	0.124	0.181	0.281	0.131	0.168	0.277	0.119	0.178	0.297	0.125
N = 500	0.116	0.277	0.085	0.124	0.233	0.093	0.115	0.234	0.084	0.126	0.242	0.089	0.117	0.227	0.084	0.126	0.244	0.093
C = 3	0.148	0.511	0.104	0.167	0.310	0.116	0.151	0.350	0.105	0.168	0.283	0.117	0.150	0.253	0.105	0.170	0.290	0.120
C = 5	0.143	0.337	0.100	0.151	0.272	0.108	0.139	0.244	0.107	0.147	0.259	0.107	0.140	0.243	0.098	0.143	0.260	0.103
C = 7	0.139	0.366	0.105	0.140	0.364	0.108	0.139	0.241	0.100	0.145	0.243	0.105	0.137	0.259	0.101	0.142	0.262	0.105
J = 12	0.146	0.365	0.094	0.155	0.289	0.103	0.146	0.282	0.099	0.158	0.266	0.103	0.150	0.257	0.095	0.159	0.276	0.103
J = 24	0.141	0.445	0.111	0.150	0.343	0.118	0.140	0.275	0.109	0.150	0.257	0.117	0.135	0.246	0.107	0.144	0.266	0.115

Note. FST = fully-specified target; SST = semi-specified target.

Table 13

Main effects on $RMSE_{FactorCorr}$ in function of the strength of the ARS and the simulated conditions when ARS is extracted

$RMSE_{FactorCorr}$ with ARS factor																		
	Small ARS						Medium ARS						Large ARS					
	Pearson			Polychoric			Pearson			Polychoric			Pearson			Polychoric		
	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST
N = 250	0.025	0.122	0.017	0.025	0.080	0.020	0.048	0.075	0.026	0.047	0.059	0.028	0.113	0.050	0.044	0.115	0.049	0.050
N = 500	0.034	0.069	0.012	0.035	0.053	0.013	0.049	0.053	0.016	0.050	0.051	0.020	0.100	0.046	0.030	0.100	0.046	0.035
C = 3	0.029	0.123	0.014	0.029	0.065	0.017	0.041	0.076	0.020	0.043	0.056	0.023	0.108	0.048	0.041	0.113	0.048	0.050
C = 5	0.028	0.082	0.018	0.032	0.059	0.019	0.062	0.059	0.022	0.062	0.055	0.027	0.096	0.050	0.039	0.095	0.048	0.040
C = 7	0.031	0.082	0.012	0.029	0.076	0.013	0.041	0.057	0.021	0.041	0.054	0.022	0.116	0.047	0.031	0.115	0.046	0.038
Balanced	0.008	0.018	0.011	0.011	0.008	0.013	0.008	0.015	0.013	0.009	0.009	0.018	0.009	0.006	0.034	0.009	0.003	0.038
Unbalanced	0.050	0.173	0.019	0.049	0.125	0.019	0.088	0.113	0.029	0.088	0.101	0.030	0.204	0.091	0.039	0.207	0.092	0.048
J = 12	0.034	0.092	0.017	0.036	0.066	0.020	0.048	0.070	0.026	0.049	0.062	0.028	0.106	0.052	0.041	0.103	0.052	0.043
J = 24	0.024	0.100	0.012	0.024	0.068	0.013	0.048	0.058	0.015	0.048	0.048	0.020	0.108	0.044	0.033	0.113	0.043	0.042

Note. FST = fully-specified target; SST = semi-specified target.

Table 14

$RMSE_{loadings_C}$ in multidimensional balanced scales when the ARS factor is not extracted in function of the simulated conditions

RMSE _{loadingsC} with ARS factor																				
Pearson												Polychoric								
Small ARS					Medium ARS			Large ARS			Small ARS			Medium ARS			Large ARS			
N	J	C	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST
250	12	3	0.039	0.036	0.037	0.046	0.041	0.045	0.078	0.054	0.073	0.020	0.015	0.015	0.022	0.011	0.020	0.065	0.037	0.056
		5	0.025	0.032	0.024	0.044	0.053	0.042	0.083	0.064	0.077	0.016	0.022	0.013	0.030	0.042	0.027	0.072	0.051	0.064
		7	0.028	0.025	0.025	0.030	0.039	0.029	0.097	0.083	0.083	0.023	0.020	0.020	0.024	0.034	0.024	0.098	0.084	0.079
	24	3	0.046	0.045	0.045	0.045	0.044	0.045	0.089	0.078	0.090	0.014	0.011	0.011	0.019	0.016	0.016	0.070	0.063	0.069
		5	0.031	0.031	0.030	0.028	0.026	0.027	0.085	0.080	0.084	0.016	0.017	0.014	0.015	0.009	0.010	0.084	0.081	0.082
		7	0.029	0.031	0.029	0.034	0.033	0.033	0.088	0.076	0.085	0.023	0.026	0.023	0.028	0.026	0.027	0.089	0.078	0.086
500	12	3	0.053	0.052	0.053	0.055	0.053	0.054	0.095	0.080	0.093	0.018	0.014	0.015	0.029	0.026	0.026	0.073	0.047	0.066
		5	0.022	0.032	0.021	0.045	0.034	0.044	0.064	0.058	0.063	0.009	0.021	0.008	0.028	0.026	0.027	0.058	0.050	0.056
		7	0.019	0.019	0.018	0.039	0.037	0.037	0.063	0.047	0.059	0.013	0.012	0.011	0.033	0.031	0.030	0.061	0.041	0.055
	24	3	0.042	0.044	0.042	0.055	0.055	0.054	0.062	0.062	0.062	0.007	0.012	0.006	0.020	0.021	0.018	0.049	0.048	0.047
		5	0.028	0.030	0.028	0.028	0.027	0.028	0.059	0.055	0.053	0.011	0.014	0.011	0.014	0.012	0.012	0.052	0.046	0.042
		7	0.029	0.029	0.028	0.035	0.034	0.034	0.070	0.060	0.070	0.021	0.021	0.021	0.027	0.026	0.026	0.076	0.065	0.076

Note. FST = fully-specified target; SST = semi-specified target.

Table 15

$RMSE_{loadings_C}$ in multidimensional unbalanced scales when the ARS factor is not extracted in function of the simulated conditions

RMSE _{loadings_C} with ARS factor																				
Pearson												Polychoric								
Small ARS					Medium ARS			Large ARS			Small ARS			Medium ARS			Large ARS			
N	J	C	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST
250	12	3	0.031	0.039	0.031	0.035	0.040	0.035	0.019	0.060	0.015	0.014	0.023	0.014	0.015	0.022	0.015	0.047	0.051	0.046
		5	0.011	0.020	0.011	0.013	0.044	0.011	0.015	0.051	0.014	0.014	0.019	0.014	0.018	0.037	0.017	0.034	0.045	0.034
		7	0.009	0.015	0.009	0.019	0.024	0.019	0.030	0.042	0.030	0.006	0.012	0.006	0.023	0.026	0.023	0.040	0.045	0.040
	24	3	0.028	0.029	0.028	0.024	0.025	0.024	0.019	0.063	0.018	0.016	0.017	0.016	0.020	0.021	0.020	0.048	0.051	0.048
		5	0.033	0.034	0.033	0.012	0.044	0.011	0.031	0.041	0.031	0.015	0.018	0.015	0.021	0.037	0.021	0.053	0.053	0.054
		7	0.035	0.035	0.035	0.012	0.020	0.012	0.028	0.048	0.028	0.027	0.027	0.027	0.006	0.016	0.006	0.038	0.047	0.039
500	12	3	0.037	0.038	0.037	0.028	0.049	0.027	0.016	0.057	0.015	0.007	0.010	0.007	0.018	0.031	0.018	0.033	0.040	0.033
		5	0.013	0.019	0.013	0.010	0.011	0.010	0.012	0.046	0.012	0.013	0.016	0.013	0.024	0.024	0.024	0.033	0.041	0.033
		7	0.004	0.014	0.004	0.010	0.022	0.010	0.019	0.037	0.019	0.011	0.016	0.011	0.007	0.019	0.007	0.029	0.038	0.029
	24	3	0.025	0.033	0.025	0.025	0.035	0.025	0.015	0.056	0.014	0.025	0.027	0.025	0.021	0.025	0.021	0.039	0.042	0.039
		5	0.016	0.020	0.016	0.014	0.037	0.014	0.007	0.021	0.007	0.007	0.012	0.007	0.009	0.027	0.009	0.024	0.027	0.024
		7	0.015	0.022	0.015	0.005	0.019	0.004	0.010	0.049	0.010	0.006	0.017	0.006	0.007	0.017	0.007	0.017	0.043	0.017

Note. FST = fully-specified target; SST = semi-specified target.

Table 16

Main effects on MMAB for zero loadings in multidimensional balanced scales when the ARS factor is not extracted in function of the simulated conditions

Multidimensional balanced scales - MMAB without ARS factor																		
Small ARS							Medium ARS						Large ARS					
Pearson			Polychoric				Pearson			Polychoric			Pearson			Polychoric		
	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST
N = 250	0.136	0.148	0.134	0.146	0.155	0.144	0.140	0.156	0.137	0.150	0.164	0.148	0.227	0.275	0.214	0.252	0.309	0.237
N = 500	0.093	0.104	0.092	0.100	0.107	0.099	0.100	0.106	0.099	0.108	0.113	0.107	0.150	0.175	0.144	0.171	0.203	0.163
C = 3	0.118	0.128	0.117	0.133	0.140	0.132	0.122	0.132	0.120	0.140	0.147	0.137	0.189	0.223	0.179	0.223	0.271	0.210
C = 5	0.114	0.129	0.113	0.121	0.132	0.119	0.117	0.131	0.116	0.125	0.136	0.123	0.177	0.208	0.170	0.196	0.234	0.188
C = 7	0.111	0.120	0.110	0.115	0.122	0.114	0.119	0.130	0.118	0.123	0.133	0.122	0.201	0.242	0.190	0.216	0.262	0.202
J = 12	0.109	0.122	0.107	0.117	0.127	0.114	0.119	0.136	0.117	0.129	0.143	0.126	0.189	0.230	0.178	0.211	0.260	0.196
J = 24	0.120	0.129	0.120	0.129	0.136	0.129	0.120	0.126	0.120	0.129	0.134	0.128	0.188	0.220	0.181	0.212	0.251	0.204

Note. FST = fully-specified target; SST = semi-specified target.

Table 17

Main effects on MMAB for zero loadings in multidimensional unbalanced scales when the ARS factor is not extracted in function of the simulated conditions

Multidimensional unbalanced scales - MMAB without ARS factor																		
Small ARS							Medium ARS						Large ARS					
Pearson			Polychoric				Pearson			Polychoric			Pearson			Polychoric		
	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST
N = 250	0.131	0.143	0.130	0.141	0.149	0.139	0.134	0.154	0.132	0.143	0.156	0.141	0.132	0.170	0.129	0.142	0.156	0.139
N = 500	0.091	0.103	0.091	0.098	0.105	0.098	0.092	0.115	0.091	0.099	0.112	0.098	0.094	0.138	0.093	0.101	0.123	0.100
C = 3	0.114	0.128	0.113	0.129	0.136	0.128	0.115	0.137	0.114	0.131	0.142	0.130	0.119	0.176	0.117	0.135	0.150	0.133
C = 5	0.111	0.122	0.110	0.118	0.125	0.117	0.113	0.142	0.111	0.118	0.137	0.117	0.109	0.140	0.108	0.115	0.128	0.114
C = 7	0.108	0.119	0.107	0.111	0.120	0.111	0.111	0.124	0.110	0.113	0.124	0.112	0.111	0.147	0.109	0.114	0.140	0.112
J = 12	0.106	0.119	0.104	0.114	0.123	0.113	0.109	0.131	0.108	0.117	0.131	0.115	0.111	0.153	0.109	0.119	0.137	0.117
J = 24	0.116	0.127	0.116	0.125	0.131	0.124	0.116	0.138	0.116	0.124	0.138	0.124	0.115	0.156	0.114	0.123	0.141	0.122

Note. FST = fully-specified target; SST = semi-specified target.

Table 18

Main effects on $RMSE_{FactorCorr}$ in function of the strength of the ARS and the simulated conditions when ARS is not extracted

$RMSE_{FactorCorr}$ without ARS factor																		
	Small ARS						Medium ARS						Large ARS					
	Pearson			Polychoric			Pearson			Polychoric			Pearson			Polychoric		
	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST	Oblimin	FST	SST
N = 250	0.026	0.020	0.007	0.026	0.009	0.008	0.053	0.025	0.029	0.053	0.022	0.029	0.137	0.095	0.096	0.138	0.145	0.096
N = 500	0.035	0.017	0.016	0.035	0.008	0.017	0.052	0.017	0.019	0.053	0.014	0.021	0.116	0.035	0.050	0.121	0.070	0.046
C = 3	0.030	0.024	0.015	0.030	0.008	0.016	0.045	0.027	0.019	0.045	0.017	0.020	0.132	0.052	0.089	0.134	0.116	0.089
C = 5	0.031	0.021	0.016	0.031	0.010	0.017	0.071	0.023	0.036	0.070	0.026	0.037	0.108	0.064	0.044	0.113	0.105	0.040
C = 7	0.031	0.010	0.004	0.031	0.008	0.005	0.042	0.012	0.018	0.043	0.012	0.019	0.138	0.079	0.086	0.141	0.101	0.083
Balanced	0.006	0.020	0.004	0.005	0.007	0.004	0.006	0.019	0.004	0.005	0.015	0.005	0.022	0.078	0.008	0.024	0.072	0.009
Unbalanced	0.056	0.017	0.020	0.056	0.010	0.021	0.100	0.022	0.044	0.101	0.022	0.045	0.230	0.051	0.138	0.235	0.142	0.133
J = 12	0.036	0.017	0.010	0.035	0.008	0.011	0.052	0.027	0.021	0.053	0.021	0.023	0.127	0.064	0.044	0.130	0.100	0.041
J = 24	0.026	0.020	0.014	0.026	0.009	0.014	0.053	0.014	0.027	0.053	0.016	0.027	0.126	0.065	0.102	0.128	0.114	0.101

Note. FST = fully-specified target; SST = semi-specified target.

Figure 1. A multidimensional factor model with an ARS factor, where the two content factors are defined as η_1 and η_2 , and ARS stands for the ARS factor. The zero and non-zero loadings are indicated by normal and dashed lines, respectively, and the residuals are omitted for visual clarity.

Figure 2. A multidimensional factor model in which the ARS factor is ignored. The dotted lines indicate the zero loadings, the elements in grey were not included in the estimation, and the residuals are omitted for visual clarity.

Figure 3. Effects of the ARS manipulations on a 5 categories item