Better Understanding the Psychologists Who Use Questionable Research Practices

Nicholas Fox, Nate Honeycutt, & Lee Jussim
Rutgers University

INTRODUCTION

- The use of questionable research practices (QRPs) has been considered a significant pathway for false-positive findings to enter the scientific literature.
- There is no consensus on the prevalence of QRP use in psychology – estimates of QRP use over entire careers have been measured as both high (>60%) and low (~10%), depending on estimating methods used. It is unknown what the current, proximal prevalence of QRP use is (Agnoli, Wicherts, Veldkamp, Albiero, & Cubelli, 2017; Fiedler & Schwarz, 2016; John, Loewenstein, & Prelec, 2012), and which estimating techniques are most relevant for measuring this hidden population.
- Furthermore, the relationship QRP users have with their peers has so far been unstudied.
- Better understanding the size and social situation of this subpopulation of psychologists can provide useful insights for the development of interventions aimed at reducing false-positive findings in the literature.

Questionable Research Practice	Example
1 Failing to report all of a study's dependent measures Collecting more data after looking to see if the results were significant	After measuring baseline anxiety with three different scales, only reporting one Collecting 25 subjects' worth of data, achieving p = 0.07, then recruiting 25 more subjects.
2	
3 Failing to report all of a study's conditions	Collecting data on three experimental conditions (low/high anxiety), but only reporting two (no/high anxiety)
4 Stopping data collection earlier than planned because one found the result one was looking for	After collecting half the final number of subjects, achieving p < 0.05, then stopping recruitment
5 Rounding off p values to achieve significance	Achieving p = 0.054, then reporting it as p = 0.05 or p < 0.05
6 Selectively reporting studies that "worked"	Running eight studies, but only reporting the 5 that had significant findings
7 Deciding whether to exclude observations after seeing the effect of doing so on the results	Removing potential outliers based on their effect on the data and not on an a priori guideline
8 Reporting unexpected findings as being predicted from the start	Predicting a relationship between A and B, but finding a relationship being A and C, reporting the A-C relationship was predicted. Also known as Hypothesizing After the Results are Known (HARKing)
9 Reporting results are unaffected by demographics when actually unsure or not tested	Reporting no gender differences when gender was not collected or relationship not tested

METHODS

- For QRP population sizing, three estimating techniques were used:
 - Direct Estimate:** Psychologists were directly asked if they had used a QRP in the past 12 months. N = 308.
- Unmatched Count:** Participants were randomized into either a Control group or a Sensitive Item group. Participants were asked to count the number of items that applied to them (i.e., I am a vegetarian, I own a dog, etc.). The Sensitive Item condition contained one extra item – “I have used at least one QRP in the past 12 months”. The mean difference between the two list conditions equals the proportion estimate of participants who identified with this sensitive item. N = 279.

$$\rho_2 = \frac{\sum x_y^s}{n^s} - \frac{\sum x_y^i}{n^i}$$

Item	List
1 I am a vegetarian.	1 & 2
2 I own a dog.	1 & 2
3 I work on a computer nearly every day.	1 & 2
4 I have a dishwasher in my kitchen.	1 & 2
5 I can drive a motorcycle.	1 & 2
6 My job allows me to work from home at least once a week.	1 & 2
7 I jog at least four times a week.	1 & 2
8 I enjoy modern art.	1 & 2
9 I have attended a professional soccer match.	1 & 2
10 I have used at least one QRP in the past 12 months.	2 only

- Generalized Network Scale Up:** Participants were asked questions to estimate the size of their professional social network (how many psychologists they know). They were also asked one question to estimate how many psychologists they know who have used at least one QRP in the past 12 months. This proportion summed across all participants produces the first part of the network scale-up estimate. Simultaneously, self-identified QRP users were asked about members of their professional social networks and which network members knew about their QRP-use identity. This information was used to determine the transmissibility of the QRP-use identity through the user's social network and was used to adjust the network scale-up to produce the generalized network scale-up. N = 587.

$$\rho_4 = \frac{\sum y_i}{\sum d_i} * \frac{1}{\tau} * \frac{1}{\delta}$$

Common	Uncommon	Rare
David	Alan	Dean
James	Donald	Seth
Jennifer	Sandra	Janet
Susan	Cynthia	Wendy
John	Keith	Guy
Mark	Gary	Jansen
Elizabeth	Emily	Susanne
Lisa	Andrea	Chantel

- Participants who did not self-identify as a QRP user were asked about their views on what most researchers think about QRP-using psychologists. Four domains of stigma were of interest: Attribution (internal vs. external), Fear, Power, and Defiance of Social Norms (Link & Phaelan, 2001). The outcome variable was QRP Stigma, a combined score from four items based on similar stigma outcomes focused on cigarette smokers (Stuber, Galea, & Link, 2008).
- Beyond assessing attitudes towards QRP users, stigma-associated behaviors were also measured in self-identified QRP users. Specifically, social withdrawal and selective disclosure, two stigma management behaviors, were directly observed (Illic et al., 2014).

Approximately 18% - 24% of American psychologists have used a questionable research practice in the past year.

These individuals are stigmatized by their academic peers.

Use the QR code below for additional information:
A summary video
A link to this poster
The preprint manuscript for this work
The full dissertation in which this work was included

Scan the QR code with your phone, or go to the following: https://www.youtube.com/r_Kb7e7-f_I

Scan me

To scan, just open your camera app and point the camera at the box above!

RESULTS

- The Generalized Network Scale Up estimated 24.4% [10.93% - 58.74%] of American psychologists have used at least one QRP in the past year. This corresponds to 1,733 psychologists. The direct estimate was 18.18% [13.96%, 22.40%], which corresponds to 1,291 psychologists. The unmatched count estimate was 10.46%, which corresponds to 743 psychologists, but had a confidence interval that crossed zero.
- The estimated transmissibility of QRP use from a QRP user to a social network partner (from a QRP-using “ego” to an “alter” in their social network) was six percent. This means that the vast majority of psychologists do not know the identity of their peers who use questionable research practices. It is a highly concealed identity. QRP users are also more likely to disclose their identity to other QRP users. However, QRP-using psychologists have social networks similar in size to the general population of psychologists (average network size = 180 psychologists).

- Participants who believed QRP use was internally attributed (“QRP user is due to weak character”) held more stigmatizing views of QRP users. This was also true for participants who believed QRP users were a threat to their students (the operationalization of the Fear domain). Participants who believed QRP users were breaking academic social norms held more stigmatizing views of QRP users. Interestingly, younger participants also held more stigmatizing views towards QRP users.

- Furthermore, if a QRP user knows a peer in their social network is also a QRP user, they are 95 times more likely to share their QRP use identity compared to an ambiguous social network member.

	In-group	Out-group
Alter knows participant's QRP-use	58	16
Alter doesn't know participant's QRP-use	42	1114

Odds ratio (OR): (58/16) / (42/1114) = 95.39

CONCLUSIONS

- Knowing the population size of psychologists who currently use questionable research practices is important for crafting future interventions that aim to reduce the publication of false-positive findings in the academic literature.
- Understanding that QRP using psychologists are stigmatized by the general population is critical: interventions can focus on either ramping up stigma (i.e., cigarette smoking) or reduce stigma (i.e., sexual orientation).
- While preregistration and Registered Reports are changing how scientists do science, this baseline estimate of QRP use will be important when measuring their impact on published science in the future.