Fear of COVID-19 predicts vaccination willingness 14 months later

Gaëtan Mertens¹, Paul Lodder^{1, 2}, Tom Smeets², & Stefanie Duijndam²

¹Department of Medical and Clinical Psychology, Tilburg University, Tilburg, the
Netherlands

²Department of Methods and Statistics, Tilburg University, Tilburg, the Netherlands

Word count (excluding the title page, abstract and references): 4946

Correspondence concerning this article should be addressed to Gaëtan Mertens,
Department of Medical and Clinical Psychology, Warandelaan 2, room T526, Tilburg
University, 5037AB Tilburg, the Netherlands.

E-mail: g.mertens@tilburguniversity.edu

Tel: +31134663757

Acknowledgements: The research reported in this paper was funded by a NWO fast-track coronavirus grant (grant number: 440-20-037) awarded to Gaëtan Mertens.

Abstract

Vaccines are an important tool for governments and health agencies to contain and curb the Coronavirus Disease-19 (COVID-19) pandemic. However, despite their effectiveness and safeness, a substantial portion of the population worldwide is hesitant to get vaccinated. In the current study, we examined whether fear of COVID-19 predicts vaccination willingness. In a longitudinal study ($N = 938$), fear for COVID-19 was assessed in April 2020 and vaccination willingness was measured in June 2021. Approximately 11% of our sample indicated that they were not willing to get vaccinated. Results of a logistic regression showed that increased fear of COVID-19 predicts vaccination willingness 14 months later, even when controlling for several anxious personality traits, infection control perceptions, risks for loved ones, self-rated health, previous infection, media use, and demographic variables. These results show that fear of COVID-19 is a relevant construct to consider for predicting and possibly influencing vaccination willingness. Nonetheless, sensitivity and specificity of fear of COVID-19 to predict vaccination willingness were quite low and only became slightly better when fear of COVID-19 was measured concurrently. This indicates that other potential factors, such as perceived risks of the vaccines, probably also play a role in explaining vaccination willingness.

Keywords: Fear; COVID-19; vaccination; hesitancy

1. Introduction

The Coronavirus Disease-19 (COVID-19) pandemic is one of the most devastating public health crises of modern times. Besides the more than five million official deaths and 200 million confirmed cases (Dong et al., 2020), governments across the world have also imposed strict and invasive policies to reduce the spread of the virus, including lockdowns and curfews (Hale et al., 2021). Unsurprisingly, this pandemic has brought about major issues such as economic decline, overburdened healthcare systems, and mental health problems (Galea et al., 2020; Miller et al., 2020; Park et al., 2020).

One major tool for governments to contain and curb the spread of COVID-19 is vaccination. The first vaccine against the SARS-CoV-2 virus (i.e., Pfizer-BioNTech) was approved for use in early December 2020 in the United Kingdom and since then different vaccines have become widely available in most countries. The four vaccines approved for use in most Western countries (i.e., Pfizer-BioNTech, Moderna, AstraZeneca, and Johnson & Johnson) have been shown to be both effective in preventing severe COVID-19 disease symptoms (Sadoff et al., 2021; Tenforde et al., 2021; Voysey et al., 2021) and show an acceptable safety profile (Barda et al., 2021). Nonetheless, not everyone is willing to get vaccinated. For example, a survey conducted in July and August 2020 by the market research company IPSOS among 19519 adults found that approximately 1 in 4 individuals worldwide were unwilling to be vaccinated (Boyon, 2020). Moreover, actual vaccination rates differ widely across countries and are typically lower than 90% (Mathieu et al., 2021), indicating that a non-trivial proportion of the population is unwilling to get vaccinated (Sallam, 2021). This is a problematic issue not only for the health of the unvaccinated portion of the population, but also for effectively containing the

pandemic and preventing the emergence of novel virus strains that may worsen or prolong the pandemic (Rella et al., 2021).

Hence, an important challenge for behavioral science is to uncover the psychological mechanisms of vaccination hesitancy and to develop interventions to increase vaccination willingness (Brewer, 2021). One potential reason for vaccination willingness is fear of COVID-19. That is, presumably people are more willing to get vaccinated if they perceive COVID-19 to be a threat to their health or the health of their loved ones. Indeed, risk perceptions and fear have been found to predict health-related behaviors both in the context of the COVID-19 pandemic and other diseases (Harper et al., 2020; Milne et al., 2000; Musche et al., 2020). Similarly, several recent cross-sectional studies found that fear of COVID-19 is positively related to vaccination willingness (Bono et al., 2021; Gagneux-Brunon et al., 2021; Willis et al., 2021). Nonetheless, to establish a causal connection, longitudinal and experimental studies are required. To our knowledge, no study so far has investigated whether fear of COVID-19 longitudinally predicts vaccination willingness.

To examine whether fear of COVID-19 is indeed a relevant predictor of vaccination willingness, we used the data of a longitudinal study investigating levels of fear of COVID-19 (Mertens, Duijndam, et al., 2020). This study was initiated in April 2020 and consisted of regular assessments of fear of COVID-19 and other relevant constructs over a period of 14 months in an international sample of 2000 participants. At the last measurement point (i.e., June 2021), participants were also asked about their vaccination willingness (i.e., whether they had been vaccinated or planned to get vaccinated). Using this data, the current study examined whether fear of COVID-19 at the beginning of the pandemic, while controlling for several potential

confounders (e.g., demographic variables, anxious personality, etc.), can predict vaccination willingness 14 months later.

Additionally, besides fear of COVID-19 predicting vaccination willingness, it is likely that vaccination status also had an impact on fear of COVID-19. Particularly, those participants who have been vaccinated may show lower levels of fear of COVID-19. This is of interest, given that fear of COVID-19 has been related to increased rates of anxiety, depression and stress (Rodríguez-Hidalgo et al., 2020; Satıcı et al., 2020). As such, vaccination may be a way to lower fear of COVID-19 and thereby improve mental wellbeing. Therefore, we also investigated whether vaccination status had an impact on fear of COVID-19.

2. Methods

2.1. Participants

Participants for this study were recruited through Prolific platform and were compensated according to the standard rates (i.e., £7.5/h). In the first wave of the study 2000 participants took part. The study was repeated every month between April 2020 and August 2020. One additional follow-up wave was completed in June 2021, in which participants provided information about their vaccination willingness. The data from the participants who completed both the first and the last wave of data collection ($N = 938$) was used for the current study. Table 1 provides an overview of the demographic information of the sample. As shown, most of the participants were equal to or younger than 40 years old (77.1%), highly educated (60.2% had at least a college degree) and reside in Europe or North America (95.9%). The Ethics Review Board of the School of Social and Behavioral Sciences of Tilburg University approved the procedure of this study (RP169).

Table 1. Demographic information of the respondents (total $N = 938$).

	<i>N</i>	%
Age in years		
18-20	99	10.6%
21-30	380	40.5%
31-40	244	26.0%
41-50	120	12.8%
51-60	64	6.8%
61-70	28	3.0%
71-80	3	0.3%
Gender		
Men	451	48.1%
Women	485	51.7%
Prefer not to say	2	0.2%
Education		
Less than high school degree	22	2.3%
High school degree	174	18.6%
Some college, but no degree	178	19.0%
College (2-years)	72	7.7%
College (4-years)	314	33.5%
Master's degree	161	17.2%
Doctorate (PhD or equivalent)	17	1.8%
Region of residence		
Asia	3	0.3%
Europe (incl. Russia)	691	73.7%
Middle East (incl. Israel)	6	0.6%
North America	208	22.2%
Oceania	17	1.8%
South America	5	0.5%
Sub-Sahara Africa	8	0.9%

2.2. Materials & Procedure

2.2.1. Measures

2.2.1.1. Fear of COVID-19

Fear of COVID-19 was measured using the Fear of the Coronavirus Questionnaire (FCQ; Mertens, Gerritsen, et al., 2020). Participants were asked to indicate the extent to which they agree with eight statements on a 5-point Likert scale. Examples of the items are “I am very worried about the coronavirus”, “I am taking precautions to prevent infection (e.g., washing hands, avoiding contact with people, avoiding door handles)”, and “I am constantly following all news updates regarding the virus”. As prior work has demonstrated the reliability and one-dimensionality of the Fear of COVID-19 scale (Mertens, Gerritsen, et al., 2020), here the

construct Fear of COVID-19 was operationalized as the sum of the eight item scores (range = 8 - 40). The internal consistency was 0.74 in the current sample (April 2020 data).

2.2.1.2. Intolerance of uncertainty scale

Intolerance of uncertainty (IU) was measured using the short scale developed and validated by Carleton et al. (2007). IU assesses and individuals' propensity to finding uncertain situations as unpleasant. The short IU scale asks participants to respond to 12 statements on a 5-point Likert scale. Examples of the statements are "Unforeseen events upset me greatly", "It frustrates me not having all the information I need", and "Uncertainty keeps me from living a full life". The internal consistency of this scale was excellent in the current sample (Cronbach's alpha = 0.89; April 2020 data).

2.2.1.3. Penn State Worry Questionnaire

The Penn State Worry Questionnaire (PSWQ) was used to measure participants' tendency to worry. The PSWQ is a well-validated questionnaire often used within clinical setting (Meyer et al., 1990). In this study we used a shortened version asking participants to respond to eight items on a 5-point Likert scale. Examples of items are: "My worries overwhelm me", "Many situations make me worry", and "I know I should not worry about things, but I just cannot help it". The internal consistency of this scale was excellent in the current sample (Cronbach's alpha = 0.95; April 2020 data).

2.2.1.4. Short Health Anxiety Inventory

The Short Health Anxiety Inventory (SHAI) was used to evaluate participants' tendency to worry about their health (Abramowitz et al., 2007; Salkovskis et al., 2002). Participants were

asked to respond to 18 four-choice questions on an ordinal measurement level. An example item is: “1 = I do not worry about my health; 2 = I occasionally worry about my health; 3 = I spend much of my time worrying about my health; 4 = I spend most of my time worrying about my health”. Typically, this scale is subdivided into two subscales: concerns about illness likelihood (14 items) and illness severity (4 items). We have followed this convention in the current study by summing the item scores separately for both constructs. The internal consistency was 0.90 for the illness likelihood subscale and .71 for the illness severity subscale (April 2020 data).

2.2.1.5. Media exposure¹

To measure participants’ voluntary exposure to news about the coronavirus, they were asked to answer the following question: “Have you looked up any extra information regarding the coronavirus outbreak? (not taking into account coincidentally seeing/reading about it in the news)” with yes or no.

2.2.1.6. General health, risk control and risk for loved ones

Participants were also asked to rate their general health, their perceived control to prevent being infected, and the perceived risk for their loved ones, as this may be relevant to predicting their fear for the coronavirus. Particularly, they were asked to answer the following question regarding their health: “Overall, I would rate my general health as:” (options: “Extremely good”, “Somewhat good”, “Neither good nor bad”, “Somewhat bad”, “Extremely bad”). Perceived control of preventing infection was assessed with the following question: “Overall, I believe that

¹ Additional questions were also included about media usage (i.e., which media channels they used and whether they paid attention to the source of the information). However, because these questions were relatively unimportant in previous analyses, we did not include them for the current report (Mertens, Gerritsen, et al., 2020).

I can control or avoid becoming infected by the coronavirus (e.g., by limiting social contact, washing hands, wearing a face mask, etc.):” (options: “Strongly agree”, “Somewhat agree”, “Neither agree nor disagree”, “Somewhat disagree”, “Strongly disagree”). Finally, risk perception for the participants’ loved ones was assessed with the following questions: “Overall, I believe that people that I care about (e.g., grandparents) are at risk of becoming infected and seriously ill due to the coronavirus outbreak:” (options: “Strongly agree”, “Somewhat agree”, “Neither agree nor disagree”, “Somewhat disagree”, “Strongly disagree”).

2.2.1.7. Demographic information

As demographic predictors, participants were asked to indicate their gender they identify with the most (“male”, “female”, “prefer not to say”), their age (in decades), their education level (i.e., their highest degree obtained), whether they work in health care (“yes”, “no”, “unsure (please clarify)”), whether they already got infected by the virus (“yes”, “no”, “unsure”), and their country of residence.

2.2.1.8. Vaccination willingness

Participants were asked to indicate whether they were already vaccinated. The following response options were available: “No, I did not receive an invitation yet”, “No, I don’t want to get vaccinated”, “Other (explain)”, “Yes, the first vaccine”, and “Yes, fully vaccinated”.

2.2.2. Survey administration

The different questionnaires and items were delivered through an online survey using the Qualtrics platform (<https://www.qualtrics.com/>). The online survey could both be completed with

the use of personal computers/laptops and smartphones. The complete survey consisted of 70 self-report items and took approximately 15 minutes to complete.

2.3. Data-analysis approach

In a first step, we recoded participants' answers to the question about their vaccination.

Particularly, the options “No, I did not receive an invitation yet” ($n = 281$; 30%), “Yes, the first vaccine” ($n = 233$; 24.8%), and “Yes, fully vaccinated” ($n = 285$; 30.4%) were recoded as indicating vaccination willingness given that they all indicate that participants do intend to get vaccinated ($n = 799$; 85.2%). Only the response category “No, I don't want to get vaccinated” was coded as an unwillingness to get vaccinated ($n = 101$; 10.8%). The response category “Other (explain)” was omitted from the analyses because this was a small ($n = 38$; 4.1%) and heterogeneous category. Particularly, whereas some of the participants in this category indicated a willingness to get vaccinated (e.g., “I haven't set up an appointment yet as I was infected with the virus recently.”), others were more undecided (e.g., “No, I was waiting to talk to a doctor about my underlying health issues first.”), and still others were more skeptical (e.g.: “No, I am waiting for the side effects to be worked out”). Thus, for the ease of interpretation, this category was not considered.

Next, a binary logistic regression was executed to predict vaccination willingness in June 2021 with fear of COVID-19, anxious personality questionnaires (i.e., intolerance of uncertainty, PSWQ, and SHAI), risk perceptions and factors (i.e., control of infection, risk for family members, and self-rated health), media use (i.e., whether or not they had looked up extra information), and demographic variables measures (i.e., age, gender, education, and previous infection) in April 2020. This logistic regression was followed up with a ROC-curve analysis to

examine the sensitivity and specificity of fear of COVID-19 to predict willingness to get vaccinated. In addition, we conducted the analyses again, but with the predictors measured in June 2021 to investigate whether prediction of vaccination willingness could be improved when the predictors are measured concurrently.

Finally, to examine the potential impact of vaccination on fear of COVID-19, we carried out a repeated-measures ANOVA on fear of COVID-19 scores with the within-subjects factor time (April 2020 and June 2021), the between-subjects factor vaccination status (“No, I did not receive an invitation yet”, “Yes, the first vaccine”, “Yes, fully vaccinated” and “No, I don’t want to get vaccinated”) and their interaction effect to test whether the change in COVID-19 fear between April 2020 and June 2021 differs across vaccination status. All analyses were carried out in SPSS v26 and an alpha level of .05 was used.

3. Results

3.1. Predicting vaccination willingness using data from April 2020

The results of the logistic regression predicting vaccination willingness in June 2021 using variables measured in April 2020 are reported in Table 2. As can be seen in this table, the only significant predictor for vaccination willingness was fear of COVID-19 (OR=1.08, 95%CI = 1.03 to 1.13). This suggests that the odds for vaccination willingness increases with 8% for every one-point increase on the FCQ total score (ranging from 8 to 40). All the other variables did not reliably predict vaccination willingness. The overall model fit was significant, $\chi^2(13) = 29.18, p = .006$, but only explained a limited amount of variance (Nagelkerke $R^2 = 0.07$). To get an impression of the extent to which fear of COVID-19 is a good predictor for vaccination

willingness, we carried out a ROC analysis. These results are shown in Figure 1. The area under the curve was equal to 0.57, indicating that classification of vaccination willingness using fear of COVID-19 measured in April 2020 was better than random, but far from perfect.

Table 2. Results of a logistic regression using predictors from April 2020 to predict COVID-19 vaccination willingness in June 2021. Odds ratios > 1 indicate that higher scores on the predictor increases the odds for willingness to be vaccinated.

Variables	B	SE	Wald	<i>p</i>	OR	95% CI OR
Fear of COVID-19	0.08	0.02	11.10	.001	1.08	[1.03, 1.13]
Age	0.12	0.10	1.39	.232	1.13	[0.93, 1.38]
Male gender	-0.09	0.23	0.14	.709	0.92	[0.59, 1.44]
Highest degree	0.12	0.07	2.75	.097	1.13	[0.98, 1.31]
Previously infected	-0.23	0.21	1.15	.284	0.80	[0.53, 1.21]
General health	0.22	0.15	2.37	.124	1.25	[0.94, 1.66]
Control to prevent infection	-0.08	0.14	0.35	.552	0.92	[0.70, 1.21]
Risk loved ones	-0.10	0.13	0.54	.464	0.91	[0.71, 1.17]
Looked up info	0.23	.25	0.85	.357	1.26	[0.77, 2.04]
IUS	-0.02	0.02	0.75	.387	0.99	[0.95, 1.02]
PSWQ	0.02	0.02	1.17	.280	1.02	[0.98, 1.06]
SHAI likelihood	-0.01	0.02	0.15	.697	0.99	[0.95, 1.03]
SHAI severity	-0.09	0.05	2.99	.084	0.91	[0.82, 1.01]

Note: SE = Standard Error; OR = Odds Ratio; IUS = Intolerance of Uncertainty Scale, PSWQ = Penn State Worry Questionnaire; SHAI = Short Health Anxiety Inventory.

3.2. Predicting vaccination willingness using data from June 2021

To explore whether the prediction of vaccination willingness could be improved when using the same predictors measured concurrently (i.e., in June 2021), we carried out the logistic regression again. The results of this analysis are shown in Table 3. The overall model fit was significant, $\chi^2(13) = 83.46$, $p < .001$ and explained more variance than the previous model (Nagelkerke $R^2 = 0.18$). As can be seen from Table 3, fear of COVID-19 was again a reliable predictor of vaccination willingness (OR=1.11, 95%CI = 1.07 to 1.16). This suggests that the odds for vaccination willingness increases with 11% for every one point increase on the FCQ total score (ranging from 8 to 40). Additionally, perceived control of being able to prevent infection, previous infection with the coronavirus, and IU were also significant predictors in this model. That is, participants who believed that they could avoid getting infected with the coronavirus were more likely to be willing to be vaccinated. Additionally, participants who got previously infected with the coronavirus were less likely to be willing to get vaccinated. Interestingly, the regression coefficient for IU was negative, indicating that participants who scored higher in IU were less willing to be vaccinated against COVID-19. This is somewhat surprising, and possibly reflects concerns and uncertainty regarding the possible side-effects of the vaccines among high IU individuals. Again, we also carried out a ROC analysis to examine the sensitivity and specificity of fear of COVID-19 measured in June 2021 to predict vaccination willingness. The area under the curve was equal to 0.69. Hence, prediction of vaccination willingness was better using data from June 2021. However, there was not cut-off point where both sensitivity and specificity were acceptable (i.e., both $\geq .8$).

Table 3. Results of a logistic regression using predictors from June 2021 to predict COVID-19 vaccination willingness in June 2021. Odds ratios > 1 indicate that higher scores on the predictor increases the odds for willingness to be vaccinated.

Variables	B	SE	Wald	<i>p</i>	<i>OR</i>	<i>95% CI OR</i>
Fear of COVID-19	0.10	0.02	24.65	< .001	1.11	[1.07, 1.16]
Age	0.09	0.10	0.71	.400	1.09	[0.89, 1.34]
Male gender	-0.06	0.23	0.06	.803	0.94	[0.60, 1.49]
Highest degree	0.14	0.08	3.62	.057	1.15	[0.97, 1.34]
Previous infection	-0.65	0.19	12.39	< .001	0.52	[0.36, 0.75]
General health	0.11	0.15	0.54	.463	1.11	[0.84, 1.48]
Control to prevent infection	0.39	0.13	8.32	.004	1.47	[1.13, 1.91]
Risk loved ones	0.06	0.12	0.24	.623	1.06	[0.84, 1.34]
Looked up info	0.42	0.23	3.36	.067	1.53	[0.97, 2.40]
IUS	-0.04	0.02	5.15	.023	0.96	[0.93, 1.00]
PSWQ	0.03	0.02	1.86	.172	1.03	[0.99, 1.07]
SHAI likelihood	-0.01	0.02	0.32	.574	0.99	[0.95, 1.03]
SHAI severity	-0.01	0.05	0.02	.888	0.99	[0.90, 1.10]

Note: SE = Standard Error; OR = Odds Ratio; IUS = Intolerance of Uncertainty Scale, PSWQ = Penn State Worry Questionnaire; SHAI = Short Health Anxiety Inventory.

Figure 1. ROC curves for the prediction of vaccination willingness based on fear of COVID-19 measured in April 2020 (left) and June 2021 (right).

3.3. Robustness analysis

We followed-up the analyses predicting vaccination willingness with an additional robustness test. Particularly, it can be argued that the response option “No, I did not receive an invitation yet” is an ambiguous response that may include both participants who want to get vaccinated and participants who do not want to be vaccinated. As this may have influenced our results, we repeated our earlier analyses without this response option. Instead, the response options “Yes, the first vaccine” and “Yes, fully vaccinated” were combined as clear-cut options indicating vaccination willingness and contrasted with the response option “No, I don’t want to get

vaccinated”. The results of these binary logistic regressions were largely identical to the initial analyses, with the exception that now both age (April 2020: OR = 1.38, 95% CI = [1.12, 1.70]; June 2021: OR = 1.31, 95% CI = [1.06, 1.63]) and education level (April 2020: OR = 1.21, 95% CI = [1.04, 1.41]; June 2021: OR = 1.21, 95% CI = [1.04, 1.42]) also emerged as significant predictors for vaccination willingness. Fear of COVID-19 remained a significant predictor for both the April 2020 (OR = 1.10, 95% CI = [1.05, 1.16]) and June 2021 analyses (OR = 1.10, 95% CI = [1.05, 1.15]).

3.4. The effect of vaccination status on fear of COVID-19

The repeated-measures ANOVA on fear of COVID-19 scores indicated significant main effects of both the factor time (i.e., April 2020 and June 2021), $F(1, 869) = 582.96, p < .001, \eta_p^2 = .40$, and the factor vaccination status (i.e., “No, I did not receive an invitation yet”, “Yes, the first vaccine”, “Yes, fully vaccinated” and “No, I don’t want to get vaccinated”), $F(3, 869) = 15.58, p < .001, \eta_p^2 = .05$. These main effects indicated that fear of COVID-19 decreased substantially over time and that participants who do not want to get vaccinated have lower levels of fear than the other three groups (see Figure 2). Importantly, there was an interaction between time and vaccination status, $F(3, 869) = 9.01, p < .001, \eta_p^2 = .03$. Tukey’s post-hoc tests on the difference scores indicated that this interaction was due to a larger decrease in fear of COVID-19 in the group of participants who do not want to get vaccinated compared to all the other groups (see Figure 2). The other three groups did not differ significantly in their reduction in fear of COVID-19 (p -values $> .05$).

Figure 2. Fear of COVID-19 scores in April 2020 and June 2021 as a function of participants' vaccination status.

4. Discussion

In the current study we find that fear of COVID-19 is a significant predictor for vaccination willingness, even when measured 14 months prior to the assessment of willingness to be vaccinated and after controlling for anxious personality traits, risk perception, self-rated health, media use, and demographic variables. This result suggests that fear of COVID-19 is a relevant construct to consider for explaining vaccination willingness. Additionally, perceived control of being able to prevent infection, IU, age, education level, and previous infection were also significant predictors for vaccination willingness. Finally, participants who did not want to be vaccinated had larger decreases in fear of COVID-19 over time than participants who did get vaccinated or wanted to get vaccinated. These findings have implications for understanding and influencing vaccination willingness.

First, the observation that participants who had higher levels of fear of COVID-19 were more willing to be vaccinated is most likely due to participants with higher levels of fear of COVID-19 perceiving COVID-19 as a threat to their own health or the health of their loved ones. Therefore, these participants are likely more willing to get vaccinated to protect themselves and their loved ones against COVID-19. It is interesting to note that fear of COVID-19 was a very stable predictor because it predicted vaccination willingness with a time lag of 14 months, albeit with reduced sensitivity and specificity compared to when it was measured concurrently. Based on this result, there are several potential approaches to increase vaccination willingness. One approach could be to use threat messages to increase fear. However, the empirical evidence on the usefulness of such fear appeals to increase health behavior is mixed on whether this is a good approach (Peters et al., 2013). We argue for caution when using threat messages to promote

vaccination willingness, because exaggerated or incorrect messaging could fuel vaccine skepticism and erode public trust, which are in turn two other important factors predicting vaccination willingness (Daly et al., 2021; Lindholt et al., 2020). Alternatively, it is possible that people who have low fear of COVID-19 have incorrect ideas about the potential dangers of COVID-19 for their own health and the health of their loved ones (Bono et al., 2021; Gallè et al., 2021; Mahmud et al., 2021). Information campaigns addressing such misconceptions may be a more appropriate way to bring threat perceptions in line with the objective risks and thereby increasing vaccination willingness and compliance with public health measures (Gallè et al., 2021; Huynh et al., 2021; Mahmud et al., 2021).

Second, perceived control of being able to prevent infection, IU, previous infection, age, and education level were also significant predictors of vaccination willingness. Regarding being able to prevent infection, it is possible that participants who were willing to be vaccinated saw vaccination as an effective way to prevent infection with the SARS-CoV-2 virus. This was indeed what initial studies at the time of data collection indicated (e.g., Mallapaty, 2021). However, it is worthwhile to note that more recent data indicates that the effectiveness of the vaccines wanes over time (Tartof et al., 2021) and that vaccination does not fully prevent against infection, especially not for the new strains of the virus (Singanayagam et al., 2021). Furthermore, surprisingly, higher IU was related to a lower willingness to be vaccinated. This is a somewhat unexpected finding because we had anticipated that higher anxious traits would predict higher vaccination willingness. Possibly, this finding reflect uncertainty about possible side-effects of the COVID-19 vaccines. Nonetheless, this is a speculative interpretation and requires further research. Another predictor for vaccination willingness was a previous infection

with the SARS-CoV-2 virus. Particularly, participants were less likely to be willing to be vaccinated if they had had a previous infection with the SARS-CoV-2 virus. This is probably because participants were aware of the fact that immunity against the SARS-CoV-2 virus is also built up after a natural infection (Dan et al., 2021). Finally, both age and education level positively predicted vaccination willingness in the robustness analysis. These findings replicate the findings of earlier cross-sectional studies (Bono et al., 2021; Gagneux-Brunon et al., 2021; Willis et al., 2021). Most likely, age is related to vaccination willingness given that age is one of the main risk factors for serious illness or death due to COVID-19 (Esai Selvan, 2020). Furthermore, education level may be related to COVID-19 vaccination willingness due to higher educated participants being better informed about the dangers of COVID-19 and the benefits of vaccination (Bono et al., 2021).

Third, regarding the effect of vaccination on fear of COVID-19, we found that participants who did not want to be vaccinated generally had lower levels of fear of COVID-19. This corresponds with our main finding that fear of COVID-19 is related to vaccination willingness. In addition, fear reduced more throughout the course of the pandemic for participants who did not want to get vaccinated than for any of the other groups. Hence, differences between the group of participants who did not want to get vaccinated and the rest of the participants became more pronounced throughout the pandemic. This may possibly reflect radicalization and cognitive biases (e.g., confirmation bias) in the group of participants who do not want to be vaccinated, which are possibly driven and maintained by social media channels (Azarpanah et al., 2021; Baines et al., 2021). The other three groups did not differ significantly from one another in changes of fear of COVID-19, suggesting that receiving the vaccine in the

group of participants who were willing to get vaccinated did not significantly reduce fear of COVID-19.

One aspect of our results that warrants some further consideration is that fear of COVID-19 was a reliable, but relatively inaccurate predictor for vaccination willingness. That is, ROC analyses indicated that it was not possible to determine a cut-off point of fear of COVID-19 to increase both the sensitivity and specificity to predict vaccination willingness to acceptable levels (i.e., $\geq .8$). This indicates that other factors besides fear of COVID-19 are also important in determining vaccination willingness. Some of these other factors were identified in the current study (i.e., perceived control of being able to prevent infection, IU, age, and education level). In addition, other factors that were not assessed in the current study are most likely also important, such as perceived risks of side-effects of the vaccines, trust in public health authorities, and prosocial reasons (i.e., getting vaccinated for others who are at risk) (Böhm & Betsch, 2022; Boyon, 2020; Daly et al., 2021). Future studies should consider these different factors, including fear of COVID-19, to get a complete assessment of factors determining vaccination willingness.

There are several limitations to this study. A first limitation is that no nationally representative samples were used, so caution should be used when generalizing these results beyond the predominantly young and highly educated western sample used for this study. Another limitation is that fear of COVID-19 was only measured with one specific scale (i.e., the FCQ developed by Mertens et al., 2020). However, prior research has found that fear of COVID-19 is a multifaceted construct that also includes concerns about the socio-economic consequences of COVID, xenophobia, and stress-related symptoms (e.g., nightmares) (Mertens et al., 2021; Taylor et al., 2020). It is possible that these other facets of fear of COVID-19 are

also related to vaccination willingness. Finally, even though this longitudinal study provides stronger evidence for the causal link between fear of COVID-19 and vaccination willingness than previous cross-sectional studies, experimental studies are needed to draw firm causal conclusions. As strengths, the longitudinal design and the relatively large sample size of this study can be noted.

Taken together, the current study shows that fear of COVID-19 is a relevant predictor for vaccination willingness. As such, fear of COVID-19 may be a relevant target to consider for increasing vaccination rates. Nonetheless, other factors are also important to consider (e.g., concerns about the side-effects of the vaccinations).

Declaration of Interest Statement

The authors declare no conflict of interest regarding the research reported in this article.

Data Availability Statement

The data of the study reported in this article can be obtained at <https://osf.io/ryndg/>.

References

- Abramowitz, J. S., Deacon, B. J., & Valentiner, D. P. (2007). The Short Health Anxiety Inventory: Psychometric Properties and Construct Validity in a Non-clinical Sample. *Cognitive Therapy and Research*, 31(6), 871–883. <https://doi.org/10.1007/s10608-006-9058-1>
- Azarpanah, H., Farhadloo, M., Vahidov, R., & Pilote, L. (2021). Vaccine hesitancy: evidence from an adverse events following immunization database, and the role of cognitive biases. *BMC Public Health*, 21(1), 1686. <https://doi.org/10.1186/s12889-021-11745-1>
- Baines, A., Ittefaq, M., & Abwao, M. (2021). #Scamdemic, #Plandemic, or #Scaredemic: What Parler Social Media Platform Tells Us about COVID-19 Vaccine. *Vaccines*, 9(5), 421. <https://doi.org/10.3390/vaccines9050421>
- Barda, N., Dagan, N., Ben-Shlomo, Y., Kepten, E., Waxman, J., Ohana, R., Hernán, M. A., Lipsitch, M., Kohane, I., Netzer, D., Reis, B. Y., & Balicer, R. D. (2021). Safety of the BNT162b2 mRNA Covid-19 Vaccine in a Nationwide Setting. *New England Journal of Medicine*, 385(12), 1078–1090. <https://doi.org/10.1056/NEJMoa2110475>
- Böhm, R., & Betsch, C. (2022). Prosocial vaccination. *Current Opinion in Psychology*, 43, 307–311. <https://doi.org/10.1016/j.copsyc.2021.08.010>
- Bono, S. A., Faria de Moura Villela, E., Siau, C. S., Chen, W. S., Pengpid, S., Hasan, M. T., Sessou, P., Ditekemena, J. D., Amodan, B. O., Hosseinipour, M. C., Dolo, H., Siewe Fodjo, J. N., Low, W. Y., & Colebunders, R. (2021). Factors Affecting COVID-19 Vaccine Acceptance: An International Survey among Low- and Middle-Income Countries. *Vaccines*, 9(5), 515. <https://doi.org/10.3390/vaccines9050515>
- Boyon, N. (2020). *Three in four adults globally say they would get a vaccine for COVID-19.*

<https://www.ipsos.com/en-us/news-polls/WEF-covid-vaccine-global>

Brewer, N. T. (2021). What Works to Increase Vaccination Uptake. *Academic Pediatrics*, 21(4),

S9–S16. <https://doi.org/10.1016/j.acap.2021.01.017>

Carleton, R. N., Norton, M. A. P. J., & Asmundson, G. J. G. (2007). Fearing the unknown: A

short version of the Intolerance of Uncertainty Scale. *Journal of Anxiety Disorders*, 21(1),

105–117. <https://doi.org/10.1016/j.janxdis.2006.03.014>

Daly, M., Jones, A., & Robinson, E. (2021). Public Trust and Willingness to Vaccinate Against

COVID-19 in the US From October 14, 2020, to March 29, 2021. *JAMA*, 325(23), 2397.

<https://doi.org/10.1001/jama.2021.8246>

Dan, J. M., Mateus, J., Kato, Y., Hastie, K. M., Yu, E. D., Faliti, C. E., Grifoni, A., Ramirez, S.

I., Haupt, S., Frazier, A., Nakao, C., Rayaprolu, V., Rawlings, S. A., Peters, B., Krammer,

F., Simon, V., Saphire, E. O., Smith, D. M., Weiskopf, D., ... Crotty, S. (2021).

Immunological memory to SARS-CoV-2 assessed for up to 8 months after infection.

Science, 371(6529). <https://doi.org/10.1126/science.abf4063>

Dong, E., Du, H., & Gardner, L. (2020). An interactive web-based dashboard to track COVID-19

in real time. *The Lancet Infectious Diseases*, 20(5), 533–534.

[https://doi.org/10.1016/S1473-3099\(20\)30120-1](https://doi.org/10.1016/S1473-3099(20)30120-1)

Esai Selvan, M. (2020). Risk factors for death from COVID-19. *Nature Reviews Immunology*,

20(7), 407–407. <https://doi.org/10.1038/s41577-020-0351-0>

Gagneux-Brunon, A., Detoc, M., Bruel, S., Tardy, B., Rozaire, O., Frappe, P., & Botelho-

Nevers, E. (2021). Intention to get vaccinations against COVID-19 in French healthcare

workers during the first pandemic wave: a cross-sectional survey. *Journal of Hospital*

Infection, 108, 168–173. <https://doi.org/10.1016/j.jhin.2020.11.020>

- Galea, S., Merchant, R. M., & Lurie, N. (2020). The Mental Health Consequences of COVID-19 and Physical Distancing. *JAMA Internal Medicine*.
<https://doi.org/10.1001/jamainternmed.2020.1562>
- Gallè, F., Sabella, E. A., Roma, P., De Giglio, O., Caggiano, G., Tafuri, S., Da Molin, G., Ferracuti, S., Montagna, M. T., Liguori, G., Orsi, G. B., & Napoli, C. (2021). Knowledge and Acceptance of COVID-19 Vaccination among Undergraduate Students from Central and Southern Italy. *Vaccines*, 9(6), 638. <https://doi.org/10.3390/vaccines9060638>
- Hale, T., Angrist, N., Goldszmidt, R., Kira, B., Petherick, A., Phillips, T., Webster, S., Cameron-Blake, E., Hallas, L., Majumdar, S., & Tatlow, H. (2021). A global panel database of pandemic policies (Oxford COVID-19 Government Response Tracker). *Nature Human Behaviour*, 5(4), 529–538. <https://doi.org/10.1038/s41562-021-01079-8>
- Harper, C. A., Satchell, L. P., Fido, D., & Latzman, R. D. (2020). Functional Fear Predicts Public Health Compliance in the COVID-19 Pandemic. *International Journal of Mental Health and Addiction*, 70 3(5), 1. <https://doi.org/10.1007/s11469-020-00281-5>
- Huynh, G., Nguyen, T. Van, Nguyen, D. D., Lam, Q. M., Pham, T. N., & Nguyen, H. T. N. (2021). Knowledge About COVID-19, Beliefs and Vaccination Acceptance Against COVID-19 Among High-Risk People in Ho Chi Minh City, Vietnam. *Infection and Drug Resistance*, Volume 14, 1773–1780. <https://doi.org/10.2147/IDR.S308446>
- Lindholt, M. F., Jørgensen, F., Bor, A., & Petersen, M. B. (2020). Willingness to Use an Approved COVID-19 Vaccine: CrossNational Evidence on Levels and Individual-Level Predictors. *PsyArXiv*. <https://doi.org/https://doi.org/10.31234/osf.io/8kn5f>
- Mahmud, S., Mohsin, M., Khan, I. A., Mian, A. U., & Zaman, M. A. (2021). Knowledge, beliefs, attitudes and perceived risk about COVID-19 vaccine and determinants of COVID-19

- vaccine acceptance in Bangladesh. *PLOS ONE*, 16(9), e0257096.
<https://doi.org/10.1371/journal.pone.0257096>
- Mallapaty, S. (2021). COVID vaccines slash viral spread – but Delta is an unknown. *Nature*, 596(7870), 17–18. <https://doi.org/10.1038/d41586-021-02054-z>
- Mathieu, E., Ritchie, H., Ortiz-Ospina, E., Roser, M., Hasell, J., Appel, C., Giattino, C., & Rods-Guirao, L. (2021). A global database of COVID-19 vaccinations. *Nature Human Behaviour*, 5(7), 947–953. <https://doi.org/10.1038/s41562-021-01122-8>
- Mertens, G., Duijndam, S., Smeets, T., & Lodder, P. (2020). Pandemic panic? Results of a 6-month longitudinal study on fear of COVID-19. *PsyArXiv*.
<https://doi.org/https://doi.org/10.31234/osf.io/xtu3f>
- Mertens, G., Duijndam, S., Smeets, T., & Lodder, P. (2021). The latent and item structure of COVID-19 fear: A comparison of four COVID-19 fear questionnaires using SEM and network analyses. *Journal of Anxiety Disorders*, 81, 102415.
<https://doi.org/10.1016/j.janxdis.2021.102415>
- Mertens, G., Gerritsen, L., Duijndam, S., Salemink, E., & Engelhard, I. M. (2020). Fear of the coronavirus (COVID-19): Predictors in an online study conducted in March 2020. *Journal of Anxiety Disorders*, 102258. <https://doi.org/10.1016/j.janxdis.2020.102258>
- Meyer, T. J., Miller, M. L., Metzger, R. L., & Borkovec, T. D. (1990). Development and validation of the penn state worry questionnaire. *Behaviour Research and Therapy*, 28(6), 487–495. [https://doi.org/10.1016/0005-7967\(90\)90135-6](https://doi.org/10.1016/0005-7967(90)90135-6)
- Miller, I. F., Becker, A. D., Grenfell, B. T., & Metcalf, C. J. E. (2020). Disease and healthcare burden of COVID-19 in the United States. *Nature Medicine*, 26(8), 1212–1217.
<https://doi.org/10.1038/s41591-020-0952-y>

- Milne, S., Sheeran, P., & Orbell, S. (2000). Prediction and Intervention in Health-Related Behavior: A Meta-Analytic Review of Protection Motivation Theory. *Journal of Applied Social Psychology*, 30(1), 106–143. <https://doi.org/10.1111/j.1559-1816.2000.tb02308.x>
- Musche, V., Bäuerle, A., Steinbach, J., Schweda, A., Hetkamp, M., Weismüller, B., Kohler, H., Beckmann, M., Herrmann, K., Tewes, M., Schadendorf, D., Skoda, E.-M., & Teufel, M. (2020). COVID-19-Related Fear and Health-Related Safety Behavior in Oncological Patients. *Frontiers in Psychology*, 11. <https://doi.org/10.3389/fpsyg.2020.01984>
- Park, C.-Y., Villafuerte, J., Abiad, A., Narayanan, B., Banzon, E., Samson, J., Aftab, A., & Tayag, M. C. (2020). *An Updated Assessment of the Economic Impact of COVID-19*. <https://doi.org/10.22617/BRF200144-2>
- Peters, G.-J. Y., Ruiter, R. A. C., & Kok, G. (2013). Threatening communication: a critical re-analysis and a revised meta-analytic test of fear appeal theory. *Health Psychology Review*, 7(sup1), S8–S31. <https://doi.org/10.1080/17437199.2012.703527>
- Rella, S. A., Kulikova, Y. A., Dermitzakis, E. T., & Kondrashov, F. A. (2021). Rates of SARS-CoV-2 transmission and vaccination impact the fate of vaccine-resistant strains. *Scientific Reports*, 11(1), 15729. <https://doi.org/10.1038/s41598-021-95025-3>
- Rodríguez-Hidalgo, A. J., Pantaleón, Y., Dios, I., & Falla, D. (2020). Fear of COVID-19, Stress, and Anxiety in University Undergraduate Students: A Predictive Model for Depression. *Frontiers in Psychology*, 11. <https://doi.org/10.3389/fpsyg.2020.591797>
- Sadoff, J., Gray, G., Vandebosch, A., Cárdenas, V., Shukarev, G., Grinsztejn, B., Goepfert, P. A., Truyers, C., Fennema, H., Spiessens, B., Offergeld, K., Scheper, G., Taylor, K. L., Robb, M. L., Treanor, J., Barouch, D. H., Stoddard, J., Ryser, M. F., Marovich, M. A., ... Douoguih, M. (2021). Safety and Efficacy of Single-Dose Ad26.COV2.S Vaccine against

Covid-19. *New England Journal of Medicine*, 384(23), 2187–2201.

<https://doi.org/10.1056/NEJMoa2101544>

Salkovskis, P. M., Rimes, K. A., Warwick, H. M. C., & Clark, D. M. (2002). The Health Anxiety Inventory: development and validation of scales for the measurement of health anxiety and hypochondriasis. *Psychological Medicine*, 32(05).

<https://doi.org/10.1017/S0033291702005822>

Sallam, M. (2021). COVID-19 Vaccine Hesitancy Worldwide: A Concise Systematic Review of Vaccine Acceptance Rates. *Vaccines*, 9(2), 160. <https://doi.org/10.3390/vaccines9020160>

Satici, B., Saricali, M., Satici, S. A., & Griffiths, M. D. (2020). Intolerance of Uncertainty and Mental Wellbeing: Serial Mediation by Rumination and Fear of COVID-19. *International Journal of Mental Health and Addiction*. <https://doi.org/10.1007/s11469-020-00305-0>

Singanayagam, A., Hakki, S., Dunning, J., Madon, K. J., Crone, M. A., Koycheva, A., Derqui-Fernandez, N., Barnett, J. L., Whitfield, M. G., Varro, R., Charlett, A., Kundu, R., Fenn, J., Cutajar, J., Quinn, V., Conibear, E., Barclay, W., Freemont, P. S., Taylor, G. P., ...

Lackenby, A. (2021). Community transmission and viral load kinetics of the SARS-CoV-2 delta (B.1.617.2) variant in vaccinated and unvaccinated individuals in the UK: a prospective, longitudinal, cohort study. *The Lancet Infectious Diseases*.

[https://doi.org/10.1016/S1473-3099\(21\)00648-4](https://doi.org/10.1016/S1473-3099(21)00648-4)

Tartof, S. Y., Slezak, J. M., Fischer, H., Hong, V., Ackerson, B. K., Ranasinghe, O. N., Frankland, T. B., Ogun, O. A., Zamparo, J. M., Gray, S., Valluri, S. R., Pan, K., Angulo, F. J., Jodar, L., & McLaughlin, J. M. (2021). Effectiveness of mRNA BNT162b2 COVID-19 vaccine up to 6 months in a large integrated health system in the USA: a retrospective cohort study. *The Lancet*, 398(10309), 1407–1416. <https://doi.org/10.1016/S0140->

6736(21)02183-8

Taylor, S., Landry, C. A., Paluszek, M. M., Rachor, G. S., & Asmundson, G. J. G. (2020).

Worry, avoidance, and coping during the COVID-19 pandemic: A comprehensive network analysis. *Journal of Anxiety Disorders*, 76, 102327.

<https://doi.org/10.1016/j.janxdis.2020.102327>

Tenforde, M. W., Olson, S. M., Self, W. H., Talbot, H. K., Lindsell, C. J., Steingrub, J. S.,

Shapiro, N. I., Ginde, A. A., Douin, D. J., Prekker, M. E., Brown, S. M., Peltan, I. D., Gong,

M. N., Mohamed, A., Khan, A., Exline, M. C., Files, D. C., Gibbs, K. W., Stubblefield, W.

B., ... Zunie, T. (2021). Effectiveness of Pfizer-BioNTech and Moderna Vaccines Against

COVID-19 Among Hospitalized Adults Aged ≥ 65 Years — United States, January–March

2021. *MMWR. Morbidity and Mortality Weekly Report*, 70(18), 674–679.

<https://doi.org/10.15585/mmwr.mm7018e1>

Voysey, M., Clemens, S. A. C., Madhi, S. A., Weckx, L. Y., Folegatti, P. M., Aley, P. K.,

Angus, B., Baillie, V. L., Barnabas, S. L., Bhorat, Q. E., Bibi, S., Briner, C., Cicconi, P.,

Collins, A. M., Colin-Jones, R., Cutland, C. L., Darton, T. C., Dheda, K., Duncan, C. J. A.,

... Zuidewind, P. (2021). Safety and efficacy of the ChAdOx1 nCoV-19 vaccine

(AZD1222) against SARS-CoV-2: an interim analysis of four randomised controlled trials

in Brazil, South Africa, and the UK. *The Lancet*, 397(10269), 99–111.

[https://doi.org/10.1016/S0140-6736\(20\)32661-1](https://doi.org/10.1016/S0140-6736(20)32661-1)

Willis, D. E., Andersen, J. A., Bryant-Moore, K., Selig, J. P., Long, C. R., Felix, H. C., Curran,

G. M., & McElfish, P. A. (2021). COVID-19 vaccine hesitancy: Race/ethnicity, trust, and

fear. *Clinical and Translational Science*, cts.13077. <https://doi.org/10.1111/cts.13077>