The role of the amygdala in processing social and affective touch

Katalin M. Gothard and Andrew J. Fuglevand

Departments of Physiology and Neuroscience

College of Medicine

University of Arizona

Tucson, Arizona, USA

Correspondence:

Katalin M. Gothard, MD, PhD

Department of Physiology

College of Medicine

University of Arizona

PO Box 245051

Tucson, AZ 85724-5051

Email: gothard94@gmail.com

Abstract

The amygdala plays a central role in emotion and social behavior, yet its role in processing social and affective touch is not well established. Recent longitudinal studies reveal that touch-deprived infants show later in life exaggerated emotional reactivity that is related to structural and functional changes in multiple brain structures including the amygdala. The internal organization and connectivity of the amygdala is well-suited to process the sensory features of tactile stimuli and also the socio-cognitive dimensions of the received touch. The convergent processing of bottom-up and top-down pathways that carry information about touch results in the elaboration of context appropriate autonomic responses. The positive hedonic value of affective touch in humans and social grooming in non-human primates is correlated with vagal tone and the release of oxytocin and endogenous opioids. Grooming, the non-human primate equivalent of affective touch in humans, also reduces vigilance that has been shown to depend critically on the amygdala. Touch-induced vagal tone and lowered vigilance alter substantially neural activity in the amygdala. Under these circumstances neurons no longer respond phasically to each touch stimulus, rather they appear to signal a sustained functional state in which the amygdala appears decoupled from monitoring the external environment.

Highlights

Touch deprivation during early development causes increased emotional reactivity and impaired emotional regulation in later life most likely through amygdala-dependent mechanisms.

The amygdala is involved in evaluating the hedonic and social aspects of touch and coordinating autonomic changes that reflect the internal state of the body.

Social and affective touch in macaques lowers arousal and vigilance, which is associated with marked changes in amygdala activity.

Introduction

A century ago, the behaviorist school of thought discouraged parents from affectionately holding their babies and hugging their toddlers [1]. Instead of affectionate touch, behaviorists suggested verbal praise, punctuated by occasional handshakes, and pats on the head as rewards for good behavior. Parents judiciously ignored these prescriptions. However, orphanages across the world implemented them with disastrous and irreversible consequences. The “failure to thrive” of institutionalized children has been attributed to emotional neglect and the absence of affective touch [2-4]. Indeed, the lack of affective touch during early development derails normal development of the social brain, leaving individuals unable to establish lasting emotional bonds later in life [5]. These changes appear to be irreversible because affectionate care and extra enrichment provided by the adoptive parents of children neglected and abused in the notorious Romanian orphanages in the 1980's did not reverse the brain abnormalities acquired early in life [6, 7].

It appears, therefore, that physical contact with a caretaker through affective touch during early development lays down the neural foundation for normal emotional reactivity and regulation [8]. Adolescents who were institutionalized during infancy showed increased emotional reactivity associated with persistent hyperactivity of the amygdala and enhanced amygdala connectivity with the prefrontal cortex [9, 10]. The amygdala itself plays a central role in emotional reactivity [11] by setting in register the perceived or expected emotional significance of stimuli with autonomic activation [12, 13]. It is possible that affective touch modifies activity within the amygdala or modifies

neural communication between the amygdala and other brain areas, leading to downregulation of emotional reactivity through vagal mechanisms [14].

Beyond its role in emotional development, the amygdala coordinates emotional behavior throughout adulthood. Affective touch is fully integrated into our adult emotional vocabulary, allowing us to receive and understand a rich variety of emotional signals through our skin [15]. Adults can also experience the negative physical and emotional effects of touch deprivation. Social distancing during the Covid-19 pandemic brought to light our "touch hunger" that shaped the negative experience across age groups [16,17]. Despite the significance of touch in shaping our social and emotional experiences, the specific neural processes in the amygdala that transform particular types of tactile inputs into social and emotional experiences are just beginning to come to light.

The amygdala processes multiple dimensions of touch

It is only recently that subsets of neurons in the non-human primate amygdala have been shown to clearly respond to tactile stimuli [18,19]. It remains unclear, however, what aspects of tactile stimuli these neurons respond to and whether they respond to the positive valence associated with affective touch. Intracranial recordings and neuroimaging studies in humans and animals have converged on the idea that the amygdala responds differentially to the positive or negative valence/value of both non-social and social stimuli [20, 21]. However, neurons in amygdala also respond to neutral

or ambiguous stimuli, to novelty, to alerting stimuli, and even to abstract constructs [reviewed in 22].

Indeed, the majority of the neurons in the amygdala are tuned to multiple dimensions of a stimulus and/or of the ongoing behavior [22]. For example, neurons in the amygdala that respond to touch also respond to visual and auditory stimuli [19] and to the production of facial expressions [18, 23] that activate the mechanoreceptors in the facial skin. It is unclear, however whether these neurons respond to the same viewed and produced facial expressions, as in the canonical mirror neuron system [24].

Furthermore, multidimensional neurons in the amygdala often respond to apparently random combinations of features [25]. For example, in monkeys, the same neurons respond to the magnitude of rewards received and to the social status of familiar conspecifics [26] or to combinations of task-related vigilance, face identity, and reward amount [27]. In humans, the same neurons have been shown to encode both novelty and the identity of visual stimuli [28].

It is also possible that tactile-responsive neurons in the amygdala signal the context in which social touch occurs, or some other aspects of ongoing behavior - a question that we are actively investigating. Based on more extensively explored sensory domains, it is expected that neurons in the amygdala signal the pleasantness/unpleasantness of touch as a function of the relationship between the receiver and the person who delivers the touch, the autonomic state of the receiver, and even the recipient's expectations. Indeed, Ellingsen and colleagues [29] argued that all these factors contribute to the

hedonic value of social touch that emerges through the convergence of bottom up and top-down mechanisms.

The connectivity of the amygdala is ideally suited for processing multiple aspects of touch

The amygdala, by virtue of its nuclear organization and the connectivity of each nucleus, is ideally suited for convergent processing of the physical and the socio-emotional dimensions of tactile signals. The lateral nucleus of the amygdala receives highly processed sensory signals from secondary sensory and multisensory association areas of the temporal and parietal cortices (Fig. 1) [30]. Affective touch that activates C-tactile afferents is processed primarily by the insula [31], which in turn is connected to the amygdala via multiple, bidirectional processing loops [32, 33]. The output of the lateral nucleus projects to the basal nucleus of the amygdala (Fig. 1) where it converges with inputs from the prefrontal cortex [34].

Figure 1. Bottom-up and top-down touch-related signals converge in the basal and accessory basal nuclei of the primate amygdala. Tactile stimuli detected by the canonical skin mechanoreceptors are processed by the tactile subdivision (3b) of the primary somatosensory cortex (SI). Inputs from C-tactile afferents are processed in the insula. The output of these two cortical areas projects to the lateral (L) nucleus of the amygdala, and after local processing, is forwarded to the basal (B) nucleus. In the basal nucleus these bottom-up inputs converge with top-down inputs from limbic cortical areas that include the cingulate cortex (with the dorsal, pre-genual and sub-genual subdivisions) and the orbitofrontal cortex (OFC). The basal nucleus also receives input from the medial segment of the prefrontal cortex (mPFC). The cingulate cortex, the OFC and the mPFC also target the accessory basal nuclei (connections not shown). The connections of the amygdala with limbic cortical areas are reciprocal. The output projections of the basal nucleus broadcast the outcome of the top-down and bottom-up evaluation of touch stimuli to multiple cortical areas. This output is also transferred to the central nucleus (C) that engages autonomic centers in the hypothalamus and brainstem. The insula also has reciprocal connections with the central nucleus. Autonomic centers at each level of brainstem regulate visceral responses to touch and coordinate the relative dominance of sympathetic and parasympathetic responses. Projections to the hypothalamus and brainstem also trigger the release of hormones (e.g., oxytocin) and endogenous opioids that may contribute the subjective experience and the social behaviors elicited by affective touch.

The medial prefrontal cortex and the orbitofrontal cortex (mPFC and OFC, Fig.1), in addition to responding to the interactive aspects of social stimuli [35, 36], also signal social rules and norms [37]. In the tactile domain, strong cultural rules and taboos restrict the area of the body that can be touched by others [38]. Thus, in the basal and accessory basal nuclei (B and AB, Fig. 1) of the amygdala, the physical features of tactile stimuli (processed initially in the lateral nucleus) can be combined with higher-level evaluations of touch in the prefrontal cortex. The outcome of this evaluation is likely broadcast from the amygdala to a large array of targets that include primary and secondary sensory areas, association areas, limbic cortical areas (such as the anterior cingulate cortex, the insula, and the orbitofrontal cortex) and the autonomic centers in the brainstem and hypothalamus via the central nucleus (C, Fig. 1)[39].

There are two main types of cortical projections from the basal (and accessory basal) nucleus. One targets limbic cortical areas, establishing several parallel loops that carry out multiple functions [40]. The other projects to various stages (both association and primary cortices) of sensory processing. Such connectivity may explain the presence of valence-dependent modulation in even early stages of cortical sensory processing. This has been best demonstrated in the visual domain. For example, in macaques, multiple areas of the temporal visual cortex respond differentially to aggressive and friendly facial expressions [41]. Lesions of the amygdala, however, eliminate this valence effect, leaving face detection and face discrimination intact [41]. It is possible, therefore, that the responses in primary somatosensory cortex to the pleasantness of touch or the hedonic value of a caress from an attractive person of the opposite sex [42] is due to the activity

of neurons in the basal nucleus that project to the primary sensory areas [39, 43]. Such neural activity in the basal nucleus may not induce detectable hemodynamic changes in the amygdala itself. Indeed, focal, near-infrared stimulation restricted to a subnuclear region in the basal nucleus of anesthetized monkeys, causes significant increases in the BOLD signal in multiple sensory areas without an equally large activation in the amygdala itself [44].

The basal and accessory basal nuclei also project to the central nuclei of the amygdala either directly or via the intercalated nuclei [45]. The central nuclei are part of an autonomic network that coordinates bodily responses to stimuli of importance [46]. It is likely that the elevated vagal tone in response to affective touch in humans is initiated by the central nucleus of the amygdala [14]. This conjecture gathers momentum if we consider that through the central nuclei, the amygdala is looped into the interoceptive circuit of the brainstem and the insula [47] and may contribute to the coordinated mind-body states elicited by affective touch [48].

The anatomically favorable position of the amygdala for the convergence of pathways that transmit physical and affective dimensions of touch is borne out by empirical data. Despite the challenges of reliably detecting BOLD signal in the amygdala [49, 50], a few neuroimaging studies showed robust activation of the amygdala in response to affective touch. When the amygdala was activated by pleasant tactile stimuli, the BOLD signal was correlated with the perceived pleasantness of touch [51], likely because pleasurable tactile sensations activate the opioid system within the amygdala [52]. A subset of studies

showed affective touch enhanced the functional connectivity between the amygdala and limbic cortical areas. For example, pleasant touch in humans elicited enhanced functional connectivity of the insula and amygdala with medial prefrontal areas [53]. Likewise, pleasant touch, but not painful stimuli, elicited coordinated activity between the anterior cingulate cortex and a region corresponding to the extended amygdala [54].

Context Dependence of Touch Processing in the Amygdala

The subjective value of social touch partially depends on the identity of the toucher. Indeed, the BOLD signal in the amygdala of human subjects differentiates between touch received from a bonded partner or a stranger of opposite sex [55]. This confirms that top-down, higher-level social evaluation of touch is processed by the amygdala, in this case, the relationship between the toucher and the receiver [38]. The pleasantness of social touch also depends on social context. Support for this comes from studies that combine touch with other social signals such as facial expressions. Seeing smiling faces, for example, enhanced the perception of pleasant touch while seeing negative facial expressions potentiated the perception of unpleasant touch [56]. Faces and facial expressions are processed in the amygdala [57], so it is not surprising that patients with amygdala lesions failed to show enhanced detection of tactile stimulation when viewing images of faces being touched (a phenomenon reliably documented in healthy controls) [58]. Such findings suggest a capacity of the amygdala to integrate social context into the processing of tactile information, likely due to the rich connectivity and multidimensionality of its component neurons [22, 27, 59].

Grooming in nonhuman primates is equivalent to affective touch in humans

Many species of non-human primates build and maintain social bonds through reciprocal grooming. Grooming is often traded for coalition support, infant handling, and for tolerance at feeding and drinking sites [60]. Macaque monkeys solicit grooming by "groom presenting" (offering access to body parts) to a trusted social partner. They often lie in front of the groomer, adopt positions that expose vulnerable body parts to the groomer's hands and teeth, close their eyes, and appear fully relaxed [61]. They may fall asleep, relinquishing attention and vigilance to the trusted groomer. Some species of macaques invest 15 - 20% of their daily activity in social grooming, which exceeds hygienic needs, but pays off during social conflict when grooming partners support each other [62]. The hedonic value of grooming is illustrated by the observation that animals readily perform a monotonous and repetitive operant task in exchange for grooming [63]. Indeed, grooming received by macaques from trusted human trainers has comparable physiological effects to grooming from conspecifics, including decreased sympathetic and increased parasympathetic (vagal) activity, indexed by lower heart rate and increased heart rate variability [64]. Many of the beneficial effects of grooming are related to the release of oxytocin that suppresses the effects of cortisol and reduces reactivity to psychosocial stressors [65, 66].

The pleasantness of grooming may be related to the release of endorphins (67 - 70) perhaps triggered by activation of C-tactile fibers in the skin that respond to particular patterns of affective touch [71]. The physiological effects of grooming in macaques match

the effects observed in humans: pleasant touch increases vagal tone, thereby reducing the negative effects of stress [14]. In addition to autonomic changes, stress-reducing effects manifest through measurable behaviors, such as the relaxation of the facial corrugator muscle that is typically contracted during negative affective states [72]. Neural recordings from the amygdala of macaque monkeys that receive social grooming, as we are presently carrying out, are expected to reveal some of the cellular underpinnings of affective touch. The amygdala is expected to signal valence (pleasant or unpleasant), social context, autonomic state, or other, more abstract variables.

Affective touch links the amygdala to attention and vigilance.

We have recorded neural activity simultaneously from the amygdala and somatosensory cortex of macaques in response to alternating blocks of non-social and social tactile stimuli (A Martin et al. *Soc Neurosci Abstr*, 2021, P317.04). Social stimuli were grooming-like finger sweeps to the monkey's face delivered by a trusted human. Gentle airflow delivered to multiple regions of the face that avoided the eyes, nostrils, and the ears served as innocuous non-social stimuli. Grooming induced significant decreases of heart rate confirming that being touched by a human groomer was pleasant. Indeed, the subjects often closed their eyes and appeared to fall asleep. During blocks of non-social tactile stimulation, heart rate was significantly elevated compared to the blocks of grooming. As expected, neurons in the somatosensory cortex responded to both non-social and social stimuli, differentiating between location and contact pressures. Consistent with our previous studies [18, 19], approximately 30% of

neurons in the amygdala responded to non-social tactile stimuli. Contrary to our expectations, however, < 10% of those amygdala neurons that responded to the non-social stimuli responded to social touch, even when the social touch was delivered with comparable pressure and to the same areas of the skin. Furthermore, virtually no amygdala neurons responded selectively to social touch.

The failure of neurons to respond to social touch, suggests a gating mechanism, possibly related to the low sympathetic arousal induced by grooming. Indeed, the small fraction of neurons that responded to both social and non-social touch were recorded in sessions when the heart rate remained high across the blocks. During most grooming blocks, a subset of neurons in the amygdala showed *tonic* changes in firing rate for the entire duration of the block. These neurons increased their activity at the time of switching from the non-social to the social touch, anticipating the period of grooming. During these predictable periods of grooming, neurons in the amygdala did not respond to each grooming sweep. Rather, they appeared to signal an internal state induced by grooming or expected from the predictable sequence of social and non-social stimulus blocks. Taken together these findings suggest that neural activity in the amygdala may be decoupled from monitoring the external environment to monitoring an internal state that has clear autonomic correlates. This phenomenon of gating phasic responses of the amygdala may have not been reported previously because during typical recordings from the amygdala, animals remain vigilant and engage with stimuli that carry emotional or behavioral significance. In response to grooming, however, we observe the opposite, a temporary disengagement of the amygdala from the incoming stimuli.

One explanation for the suppression of episodic neural responses to grooming may be that the animal's sensorium is no longer oriented outwardly, as the responsibility of monitoring the environment for potential danger is transferred to the groomer. Indeed, the amygdala plays a pivotal role in vigilance and attention [73] and shows enhanced, almost automatic facilitation of attention for threat-related stimuli [74, 75]. Accordingly, vigilance and apprehension are correlated with high gamma oscillations in the amygdala that entrain and increase the spiking activity of the resident neurons [76]. It may be that, during periods of positive social contact, the amygdala tonically signals to the rest of the brain the safety of this behavioral context while enhancing activity in the parasympathetic branch of the autonomic nervous system. While the amygdala is broadly known for the "fight or flight" response, it may be just as important for coordinating vagal states. The significant and persistent elevation of heart rate in monkeys with bilateral amygdala lesions [77] suggest the loss of vagal mechanism that slows spontaneous heart rate. In light of these observations, it is not surprising that individuals deprived of affective touch have increased emotional reactivity and poor emotional regulation [6 – 10]. A major challenge for future research will be to determine the cellular mechanisms by which affective touch alters the output of the amygdala to enable positive emotional states.

Acknowledgements

This work was supported by the National Institute of Mental Health R01MH121009

References

1. Watson JB. *Psychological care of infant and child*. Norton & Co; 1928.
2. Bakwin H: **Emotional deprivation in infants**. *J Pediatr* 1949, **35**:512–521.
3. Ardiel EL, Rankin CH: **The importance of touch in development**. *Paediatr Child Health* 2010, **15**:153–156.
4. Johnson DE, Gunnar MR: **IV. Growth Failure in Institutionalized Children**. *Monogr Soc Res Child Dev* 2011, **76**:92–126.
5. O'Connor TG, Rutter M: **Attachment disorder behavior following early severe deprivation: extension and longitudinal follow-up. English and Romanian Adoptees Study Team**. *J Am Acad Child Adolesc Psychiatry* 2000, **39**:703–712.
6. Rutter M: **Developmental catch-up, and deficit, following adoption after severe global early privation. English and Romanian Adoptees (ERA) Study Team**. *J Child Psychol Psychiatry* 1998, **39**:465–476.
7. * Mackes NK, Golm D, Sarkar S, Kumsta R, Rutter M, Fairchild G, Mehta MA, Sonuga-Barke EJS, ERA Young Adult Follow-up team: **Early childhood deprivation is associated with alterations in adult brain structure despite subsequent environmental enrichment**. *Proc. Natl. Acad. Sci. U.S.A.* 2020, **117**:641–649. *Using structural MRI, the authors demonstrate that severe deprivation during early development causes structural alterations in the brain that cannot be reversed by post-hoc compensatory enrichment.
8. ** Cascio CJ, Moore D, McGlone F: **Social touch and human development**. *Developmental Cognitive Neuroscience* 2019, **35**:5–11. **This review discusses the role of affective touch at each stage of human development, from infancy to adulthood.
9. Tottenham N, Hare TA, Millner A, Gilhooly T, Zevin JD, Casey BJ: **Elevated amygdala response to faces following early deprivation**. *Dev Sci* 2011, **14**:190–204.
10. **Silvers JA, Lumian DS, Gabard-Durnam L, Gee DG, Goff B, Fareri DS, Caldera C, Flannery J, Telzer EH, Humphreys KL, et al.: **Previous Institutionalization Is Followed by Broader Amygdala-Hippocampal-PFC Network Connectivity during Aversive Learning in Human Development**. *J. Neurosci.* 2016, **36**:6420–6430. **Using functional MRI, these authors show hyperactivity in the amygdala of previously institutionalized children that alters the functional connectivity between the hippocampus and the prefrontal cortex during aversive learning.
11. * Filippi CA, Sachs JF, Phillips D, Winkler A, Gold AL, Leibenluft E, Pine DS, Fox NA: **Infant behavioral reactivity predicts change in amygdala volume 12 years later**. *Developmental Cognitive Neuroscience* 2020, **42**:100776. Using a longitudinal design, the authors establish a causal relationship between emotional reactivity and the structure and function of the amygdala.
12. LeDoux JE, Iwata J, Cicchetti P, Reis DJ: **Different projections of the central amygdaloid nucleus mediate autonomic and behavioral correlates of conditioned fear**. *Journal of Neuroscience* 1988, **8**:2517–2529.

13. Bohus B, Koolhaas JM, Luiten PG, Korte SM, Roozendaal B, Wiersma A: **The neurobiology of the central nucleus of the amygdala in relation to neuroendocrine and autonomic outflow.** *Prog Brain Res* 1996, **107**:447–460.
14. Triscoli C, Croy I, Steudte-Schmiedgen S, Olausson H, Sailer U: **Heart rate variability is enhanced by long-lasting pleasant touch at CT-optimized velocity.** *Biological Psychology* 2017, **128**:71–81.
15. Hertenstein MJ, Verkamp JM, Kerestes AM, Holmes RM: **The communicative functions of touch in humans, nonhuman primates, and rats: a review and synthesis of the empirical research.** *Genet Soc Gen Psychol Monogr* 2006, **132**:5–94.
16. Field T, Poling S, Mines S, Bendell D, Veazey C: **Touching and Touch Deprivation During a COVID-19 Lockdown.** *International Journal of Psychological Research and Reviews* 2020.
17. Mohr Von M, Kirsch LP, Fotopoulou A: **Social touch deprivation during COVID-19: effects on psychological wellbeing, tolerating isolation and craving interpersonal touch.** 2021.
18. Mosher CP, Zimmerman PE, Fuglevand AJ, Gothard KM: **Tactile Stimulation of the Face and the Production of Facial Expressions Activate Neurons in the Primate Amygdala.** *eNeuro* 2016, **3**.
19. Morrow J, Mosher C, Gothard K: **Multisensory Neurons in the Primate Amygdala.** *J. Neurosci.* 2019, **39**:3663–3675.
20. Morrison SE, Salzman CD: **Re-valuing the amygdala.** *Current Opinion in Neurobiology* 2010, **20**:221–230.
21. Pignatelli M, Beyeler A: **Valence coding in amygdala circuits.** *Curr Opin Behav Sci* 2019, **26**:97–106.
22. **Gothard KM: **Multidimensional processing in the amygdala.** *Nature Reviews Neuroscience* 2020, **21**: 565-575. ** This paper contains a comprehensive review of the social deficits caused by amygdala lesions in non-human primates and presents arguments for high-dimensional representations in the amygdala, based on the multidimensional response properties of its resident neurons.
23. Livneh U, Resnik J, Shohat Y, Paz R: **Self-monitoring of social facial expressions in the primate amygdala and cingulate cortex.** *Proc. Natl. Acad. Sci. U.S.A.* 2012, **109**:18956–18961.
24. Cattaneo L, Rizzolatti G: **The Mirror Neuron System.** *Arch. Neurol.* 2009, **66**:557–560.
25. Kyriazi P, Headley DB, Paré D: **Multi-dimensional Coding by Basolateral Amygdala Neurons.** *Neuron* 2018, **99**:1315–1328.e5.
26. Munuera J, Rigotti M, Salzman CD: **Shared neural coding for social hierarchy and reward value in primate amygdala.** *Nat Neurosci* 2018, doi:10.1038/s41593-018-0082-8.
27. Putnam PT, Gothard KM: **Multidimensional Neural Selectivity in the Primate Amygdala.** *eNeuro* 2019, **6**.
28. Rutishauser U, Ye S, Koroma M, Tudusciuc O, Ross IB, Chung JM, Mamelak AN: **Representation of retrieval confidence by single neurons in the human medial temporal lobe.** *Nat Neurosci* 2015, **18**:1041–1050.

29. *Ellingsen D-M, Leknes S, Løseth G, Wessberg J, Olausson H: **The Neurobiology Shaping Affective Touch: Expectation, Motivation, and Meaning in the Multisensory Context.** *Front Psychol* 2015, **6**:1986. * A review of the bottom-up and top-down factors that contribute to the subjective experiences of the receivers of affective touch.
30. McDonald AJ: **Cortical pathways to the mammalian amygdala.** *Progress in Neurobiology* 1998, **55**:257–332.
31. Olausson H, Lamarre Y, Backlund H, Morin C, Wallin BG, Starck G, Ekholm S, Strigo I, Worsley K, Vallbo ÅB, et al.: **Unmyelinated tactile afferents signal touch and project to insular cortex.** *Nat Neurosci* 2002, **5**:900–904.
32. Jezzini A, Rozzi S, Borra E, Gallese V, Caruana F, Gerbella M: **A shared neural network for emotional expression and perception: an anatomical study in the macaque monkey.** *Front Behav Neurosci* 2015, **9**:243.
33. * Evrard HC: **The Organization of the Primate Insular Cortex.** *Front Neuroanat* 2019, **13**:43. A comprehensive review of the connectivity of different subdivisions of the primate insula including the connectivity with the nuclei of the amygdala.
34. Ghashghaei HT, Barbas H: **Pathways for emotion: interactions of prefrontal and anterior temporal pathways in the amygdala of the rhesus monkey.** *NSC* 2002, **115**:1261–1279.
35. Sliwa J, Freiwald WA: **A dedicated network for social interaction processing in the primate brain.** *Science* 2017, **356**:745–749.
36. Gangopadhyay P, Chawla M, Monte OD, Chang SWC: **Prefrontal–amygdala circuits in social decision-making.** *Nat Neurosci* 2020, doi:10.1038/s41593-020-00738-9.
37. Wallis JD, Anderson KC, Miller EK: **Single neurons in prefrontal cortex encode abstract rules.** *Nature* 2001, **411**:953–956.
38. Suvilehto JT, Glerean E, Dunbar RIM, Hari R, Nummenmaa L: **Topography of social touching depends on emotional bonds between humans.** *Proc. Natl. Acad. Sci. U.S.A.* 2015, **112**:13811–13816.
39. Amaral DG **Anatomical organization of the primate amygdaloid complex.** In: *The amygdala: neurobiological aspects of emotion, memory, and mental dysfunction.* Edited by Aggleton JP, Wiley-Liss, 1992.
40. Bickart KC, Dickerson BC, Barrett LF: **The amygdala as a hub in brain networks that support social life.** *Neuropsychologia* 2014, **63**:235–248.
41. Hadj-Bouziane F, Liu N, Bell AH, Gothard KM, Luh W-M, Tootell RBH, Murray EA, Ungerleider LG: **Amygdala lesions disrupt modulation of functional MRI activity evoked by facial expression in the monkey inferior temporal cortex.** *Proc. Natl. Acad. Sci. U.S.A.* 2012, **109**:E3640–8.
42. Gazzola V, Spezio ML, Etzel JA, Castelli F, Adolphs R, Keysers C: **Primary somatosensory cortex discriminates affective significance in social touch.** *Proc. Natl. Acad. Sci. U.S.A.* 2012, **109**:E1657–66.
43. Freese JL, Amaral DG: **The organization of projections from the amygdala to visual cortical areas TE and V1 in the macaque monkey.** *J. Comp. Neurol.* 2005, **486**:295–317.
44. * Shi S, Xu AG, Rui Y-Y, Zhang X, Romanski LM, Gothard KM, Roe AW: **Infrared neural stimulation with 7T fMRI: A rapid in vivo method for**

- mapping cortical connections of primate amygdala.** *NeuroImage* 2021, **231**:117818. * The functional connectivity of the primate amygdala is mapped using near-infrared focal stimulation of the amygdala in ultra-high field (7T) fMRI. Confirming the known anatomical connectivity of the amygdala, the authors show that the activation of a small number of voxels within the amygdala can cause focal activation in multiple cortical areas that include limbic, sensory and association cortices.
45. Zikopoulos B, John YJ, García-Cabezas MÁ, Bunce JG, Barbas H: **The intercalated nuclear complex of the primate amygdala.** *Neuroscience* 2016, **330**:267–290.
 46. Benarroch E: **Chapter 2 - Central Autonomic Control.** In *Primer on the Autonomic Nervous System (3rd Edition)*. Edited by Robertson D, Biaggioni I, Burnstock G, Low PA, Paton JFR. 2012, Elsevier
 47. ** Berntson GG, Khalsa SS: **Neural Circuits of Interoception.** *Trends in Neurosciences* 2021, **44**:17–28. ** The interoceptive circuits of the brain include the amygdala. In the amygdala interoceptive and exteroceptive pathways intersect contributing to coordinated brain-body states.
 48. **Burleson MH, Quigley KS: **Social interoception and social allostasis through touch: Legacy of the Somatovisceral Afference Model of Emotion.** *Social Neuroscience* 2021, **16**:92–102. ** The most comprehensive to date synthesis of touch processing discussed in the framework of the Theory of Constructed Emotions and the Somatovisceral Afference Model of Emotion.
 49. Robinson SD, Pripfl J, Bauer H, Moser E: **The impact of EPI voxel size on SNR and BOLD sensitivity in the anterior medio-temporal lobe: a comparative group study of deactivation of the Default Mode.** *MAGMA* 2008, **21**:279–290.
 50. Boubela RN, Kalcher K, Huf W, Seidel E-M, Derntl B, Pezawas L, Našel C, Moser E: **fMRI measurements of amygdala activation are confounded by stimulus correlated signal fluctuation in nearby veins draining distant brain regions.** *Sci. Rep.* 2015, **5**:10499–15.
 51. Chen Y, Li Q, Zhang Q, Kou J, Zhang Y, Cui H, Wernicke J, Montag C, Becker B, Kendrick KM, et al.: **The Effects of Intranasal Oxytocin on Neural and Behavioral Responses to Social Touch in the Form of Massage.** *Front Neurosci* 2020, **14**:589878.
 52. Nummenmaa L, Tuominen L, Dunbar R, Hirvonen J, Manninen S, Arponen E, Machin A, Hari R, Jääskeläinen IP, Sams M: **Social touch modulates endogenous μ -opioid system activity in humans.** *NeuroImage* 2016, **138**:242–247.
 53. Gordon I, Voos AC, Bennett RH, Bolling DZ, Pelphrey KA, Kaiser MD: **Brain mechanisms for processing affective touch.** *Hum Brain Mapp* 2013, **34**:914–922.
 54. Rolls ET, O'Doherty J, Kringelbach ML, Francis S, Bowtell R, McGlone F: **Representations of pleasant and painful touch in the human orbitofrontal and cingulate cortices.** *Cereb. Cortex* 2003, **13**:308–317.
 55. ** Suvilehto JT, Renvall V, Nummenmaa L: **Relationship-specific Encoding of Social Touch in Somatosensory and Insular Cortices.** *Neuroscience* 2021, **464**:105–116. ** Using fMRI the authors show that the activation of the amygdala

in response to affective touch depends on the relationship between the toucher and the receiver. They demonstrate that the subjective evaluation of social touch involves amygdala and that the top-down influences, presumably arriving to the amygdala from the prefrontal cortex, modify this evaluation.

56. Ellingsen D-M, Wessberg J, Chelnokova O, Olausson H, Laeng B, Leknes S: **In touch with your emotions: oxytocin and touch change social impressions while others' facial expressions can alter touch.** *Psychoneuroendocrinology* 2014, **39**:11–20.
57. Gothard KM, Battaglia FP, Erickson CA, Spitler KM, Amaral DG: **Neural responses to facial expression and face identity in the monkey amygdala.** *Journal of Neurophysiology* 2007, **97**:1671–1683.
58. * Bertini C, Starita F, Passamonti C, Santoro F, Zamponi N, Michelucci R, Scarpazza C: **Fear-specific enhancement of tactile perception is disrupted after amygdala lesion.** *J Neuropsychol* 2020, **14**:165–182. * Behavioral measures indicate that tactile processing is impaired by amygdala lesions.
59. Ebisch SJH, Salone A, Martinotti G, Carlucci L, Mantini D, Perrucci MG, Saggino A, Romani GL, Di Giannantonio M, Northoff G, et al.: **Integrative Processing of Touch and Affect in Social Perception: An fMRI Study.** *Front. Hum. Neurosci.* 2016, **10**:209.
60. Barrett L, Henzi S, Weingrill T, Lycett J, Hill R: **Female baboons do not raise the stakes but they give as good as they get.** *Animal Behaviour* 2000, **59**:763–770.
61. Boccia ML, Reite M, Laudenslager M: **On the physiology of grooming in a pigtail macaque.** *Physiol. Behav.* 1989, **45**:667–670.
62. Dunbar RIM: **The social role of touch in humans and primates: behavioural function and neurobiological mechanisms.** *Neuroscience & Biobehavioral Reviews* 2010, **34**:260–268.
63. Taira K, Rolls ET: **Receiving grooming as a reinforcer for the monkey.** *Physiol. Behav.* 1996, **59**:1189–1192.
64. * Grandi LC, Ishida H: **The Physiological Effect of Human Grooming on the Heart Rate and the Heart Rate Variability of Laboratory Non-Human Primates: A Pilot Study in Male Rhesus Monkeys.** *Front. Vet. Sci.* 2015, **2**:50. * An important demonstration that affective touch induces vagal tone and a sense of well-being in animals when delivered by a trusted human caretaker.
65. Heinrichs M, Baumgartner T, Kirschbaum C, Ehler U: **Social support and oxytocin interact to suppress cortisol and subjective responses to psychosocial stress.** *Biological Psychiatry* 2003, **54**:1389–1398.
66. Parr LA, Modi M, Siebert E, Young LJ: **Intranasal oxytocin selectively attenuates rhesus monkeys' attention to negative facial expressions.** *Psychoneuroendocrinology* 2013, **38**:1748–1756.
67. Keverne EB, Martensz ND, Tuite B: **Beta-endorphin concentrations in cerebrospinal fluid of monkeys are influenced by grooming relationships.** *Psychoneuroendocrinology* 1989, **14**:155–161.
68. Kalin NH, Shelton SE, Lynn DE: **Opiate systems in mother and infant primates coordinate intimate contact during reunion.** *Psychoneuroendocrinology* 1995, **20**:735–742.

69. Martel FL, Nevison CM, Simpson MJ, Keverne EB: **Effects of opioid receptor blockade on the social behavior of rhesus monkeys living in large family groups.** *Dev Psychobiol* 1995, **28**:71–84.
70. Graves FC, Wallen K, Maestripieri D: **Opioids and attachment in rhesus macaque (*Macaca mulatta*) abusive mothers.** *Behav Neurosci* 2002, **116**:489–493.
71. Olausson H, Wessberg J, Morrison I, McGlone F, Vallbo A: **The neurophysiology of unmyelinated tactile afferents.** *Neuroscience & Biobehavioral Reviews* 2010, **34**:185–191.
72. Mayo LM, Lindé J, Olausson H, Heilig M, Morrison I: **Putting a good face on touch_ Facial expression reflects the affective valence of caress-like touch across modalities.** *Biological Psychology* 2018, **137**:83–90.
73. Davis M, Whalen PJ: **The amygdala: vigilance and emotion.** *Mol Psychiatry* 2001, **6**:13–34.
74. LeDoux J: **The amygdala.** *Current Biology* 2007, **17**:R868–74.
75. Vuilleumier P: **How brains beware: neural mechanisms of emotional attention.** *Trends in Cognitive Sciences* 2005, **9**:585–594.
76. ** Amir A, Headley DB, Lee S-C, Haufler D, Paré D: **Vigilance-Associated Gamma Oscillations Coordinate the Ensemble Activity of Basolateral Amygdala Neurons.** *Neuron* 2018, **97**:656–669.e7. The authors combine behavioral and neurophysiological measures in rats to demonstrate that the output of the amygdala is significantly different at different levels of vigilance.
77. * Mitz AR, Chacko RV, Putnam PT, Rudebeck PH, Murray EA: **Using pupil size and heart rate to infer affective states during behavioral neurophysiology and neuropsychology experiments.** *Journal of Neuroscience Methods* 2017, **279**:1–12. Bilateral amygdala lesions in non-human primates cause significant elevations of heart rate. As heart rate reflects the combination of sympathetic (accelerating) and parasympathetic (decelerating) drive to the heart, the observed increase in heart can be attributed to the loss of parasympathetic drive from the amygdala. These results demonstrate that the amygdala plays a role in coordinating calm, vagal states, not only states of "fight or flight".