Ecological stress, amygdala reactivity, and internalizing problems in preadolescence: Is parenting a buffer?

Michael I. Demidenko¹, Ka I Ip², Dominic P. Kelly¹, Kevin Constante¹, Leigh G. Goetschius¹,

Daniel P. Keating^{1,3}

1. Department of Psychology, University of Michigan.
2. Department of Psychology, Yale University
3. Survey Research Center, Institute for Social Research, University of Michigan

Correspondence concerning this article should be addressed to Michael Demidenko, Department of Psychology, University of Michigan, 530 Church St. 2036, Ann Arbor, MI 48109. E-mail: demidenm@umich.edu

Ecological stress, amygdala reactivity, and internalizing problems

Acknowledgments *A revised version of this manuscript was accepted at Cortex February 2020*

MD & LG are funded by the Eunice Kennedy Shriver National Institute of Child Health and Human Development Developmental Training grant (T32HD007109-34, McLoyd & Monk). KI is funded by the Susan Nolen-Hoeksema Postdoctoral Fellowship.

Data used in the preparation of this article were obtained from the Adolescent Brain Cognitive Development (ABCD) Study (<https://abcdstudy.org>), held in the NIMH Data Archive (NDA). This is a multisite, longitudinal study designed to recruit more than 10,000 children age 9-10 and follow them over 10 years into early adulthood. The ABCD Study is supported by the National Institutes of Health and additional federal partners under award numbers U01DA041022, U01DA041028, U01DA041048, U01DA041089, U01DA041106, U01DA041117, U01DA041120, U01DA041134, U01DA041148, U01DA041156, U01DA041174, U24DA041123, U24DA041147, U01DA041093, and U01DA041025. A full list of supporters is available at <https://abcdstudy.org/federal-partners.html>. A listing of participating sites and a complete listing of the study investigators can be found at <https://abcdstudy.org/scientists/workgroups/>. ABCD consortium investigators designed and implemented the study and/or provided data but did not necessarily participate in analysis or writing of this report. This manuscript reflects the views of the authors and may not reflect the opinions or views of the NIH or ABCD consortium investigators.

Author's Contribution

MD acquired the necessary access and data used agreements with NDA and U-M. MD, KI, DK, KC, LG conceived the study and conducted the literature review. MD & KI, with support from DK, wrote the introduction, KC, LG & KI wrote the methods/analysis section. MD & KI performed analyses. MD, KI, DK & LG reported results. MD & KI wrote the discussion with support from DK, KC & LG. DPK provided in-kind support and assisted MD, KI, DK, KC, LG with study conception and writing. All authors read and approved the manuscript.

Conflict of Interest

The authors declare that they have no conflicts of interest.

Abstract

Ecological stress during adolescent development may increase the sensitivity to negative emotional processes that can contribute to the onset and progression of internalizing behaviors during preadolescence. Although a small number of studies have considered the link among the complex relations in ecological stress, amygdala reactivity and internalizing symptoms in childhood and adolescence, these studies have largely been small, cross-sectional, and often do not consider unique roles of parenting or sex. In the current study, we evaluated the interrelations among ecological stress, amygdala functioning, subsequent internalizing symptoms, and the moderating roles of parenting and sex, 9- and 10-year old preadolescents from the Adolescent Brain Cognitive Development study (ABCD). A subset of participants who met *a priori* quality control criteria for bilateral amygdala during the N-back faces versus places contrast (N = 7,385; Mean Age = 120 months, SD = 7.52; 49.5% Female) were included in the study. A Confirmatory Factor Analysis (CFA) was performed to create a latent variable of ecological stress, and multiple Structural Equation Model (SEM) models were conducted to evaluate the association among baseline ecological stress and internalizing symptoms one year later, the mediating role of amygdala activity, and moderating effects of parental acceptance and sex. The results revealed a significant association among ecological stress and subsequent internalizing symptoms, which was greater in males than females. There was no association between amygdala activity and ecological stress or subsequent internalizing symptoms, and no mediating role of amygdala, or moderating effect of parental acceptance, on the association between ecological stress and internalizing symptoms. An alternative mediation model was tested and revealed that parental acceptance mediated the association between ecological stress and internalizing symptoms, demonstrating lower internalizing symptoms among preadolescents one year later. Given the

Ecological stress, amygdala reactivity, and internalizing problems

lack of association in brain function, ecological stress and internalizing symptoms in this registered report, effects from small studies should be reconsidered in larger samples.

Keywords: Preadolescence, Internalizing, Environment, Amygdala, Sex, Parenting

Introduction

The prevalence of internalizing problems, such as anxiety and depression, increases in preadolescence and becomes more pronounced in adolescence (Costello, Mustillo, Erkanli, Keeler, & Angold, 2003; Cyranowski, Frank, Young, & Shear, 2000; Kessler et al., 2012; Merikangas et al., 2010; Spear, 2000). The increased sensitivity of negative emotional processes during adolescence may have long-term maladaptive health outcomes (Tottenham & Galván, 2016), which can contribute to the onset and progression of chronic disease (Duffy, McLaughlin, & Green, 2018). Identifying sources of ecological stress, defined here as elements of a preadolescent's environment (e.g. family, neighborhood, community) that would be expected to have an adverse effect on the preadolescent (Caspi, Bolger & Eckenrode, 1987), and neural mechanisms in preadolescent development that may contribute to individual differences in emotional sensitivity, and how they both link to the emergence of internalizing problems could provide an important step towards early prognosis and preventive intervention.

Disparate efforts have demonstrated the adverse impacts of early ecological stressors on internalizing problems (Duffy et al., 2018; McEwen, 2012; McEwen & Gianaros, 2011) and amygdala functioning (McEwen & Gianaros, 2011; Swartz, Knodt, Radtke, & Hariri, 2015). Yet, despite recent evidence demonstrating that early adolescence is a behavioral and neurodevelopmental period that is highly sensitive to the influence of ecological stress (Tottenham & Galván, 2016), there is a paucity of studies linking the complex relations among ecological stress, amygdala reactivity and internalizing problems in preadolescence. Swartz and colleagues (2015) investigated how ecological stress, brain, and internalizing symptoms may operate together, whereby amygdala reactivity was a mechanism through which stressful life events were associated with prospective internalizing problems among young adults. This work

Ecological stress, amygdala reactivity, and internalizing problems

offered evidence of how the amygdala reactivity was a mechanism that modified the impacts of ecological stress on internalizing problems, but because it was conducted in a college sample, it is unclear how these effects operate during the critical transitional period of preadolescence.

From the framework of an ecological-transactional model (Lynch & Cicchetti, 1998), the development of internalizing problems may be a dynamic process influenced by ecological factors, by intermediate experiences that are necessary to sustain the progression of vulnerability (Hostinar & Miller, 2019), and by maladaptive brain mechanisms (e.g., amygdala reactivity) that in turn may sustain or amplify the initial vulnerability diathesis (Guyer, 2020). The overarching goal of this study is to examine how ecological stress relates to amygdala function and internalizing symptoms among preadolescents, and whether amygdala function may be a neural mechanism mediating the adverse impact of ecological stress on preadolescents' internalizing symptoms.

According to the allostatic load perspective, cumulative risk factors, as opposed to specific ecological risks, are more predictive of a child's developmental outcomes (Evans, Li, & Whipple, 2013; Gach, Ip, Sameroff, & Olson, 2018). Cumulative stress, however, is a heterogeneous concept (Evans et al., 2013) and various *types* of stress, such as, parental psychopathology (Boecker et al., 2014), family function (Farber, Kim, Knodt, & Hariri, 2019; Vidal-Ribas et al., 2019), socioeconomic status (Noble et al., 2015), childhood neglect (McLaughlin, Sheridan, & Nelson, 2017; Zeanah et al., 2009) and/or abuse (Cohen, Mannarino, Kliethermes, & Murray, 2012; Jaffee, 2017) may have differential effects on symptom development and amygdala functioning (Farah, 2018; McLaughlin et al., 2017), suggesting that it is important to pinpoint how *specific* types of stressors (i.e., ecological stress) may influence neural-behavioral development within a comprehensive model (McLaughlin et al., 2017).

Ecological stress, amygdala reactivity, and internalizing problems

Specifically, our study focuses on understanding the adverse impact of ecological stress on brain-behavioral development for two reasons: 1) ecological stress tends to be a chronic stressor that affects the course of development and poses a high risk for intergenerational transmission, therefore understanding the adverse impact of ecological stress on brain-behavioral functioning may shed light on intervention strategies; and 2) among different types of stressors, ecological stress is one of the most common stressors across diverse populations and geographic locations, thus having strong implications for social policy.

While understanding how amygdala reactivity may mediate the effect of ecological stress and preadolescents' internalizing problems is a fruitful start, increasing work has suggested parenting as one important factor that may moderate the association between stress and amygdala functioning (Farber et al. 2019), and buffer against the development of internalizing symptoms (Gorostiaga, Aliri, Balluerka, & Lameirinhas, 2019). Moreover, internalizing problems affect twice as many girls as boys beginning in late childhood, foreshadowing higher rates of anxiety and depression in females compared to males in adolescence (Pine, Cohen, Gurley, Brook, & Ma, 1998) and into adulthood (Beesdo, Knappe, & Pine, 2009; Wittchen, Nelson, & Lachner, 1998). Greater propensity for internalizing problems in females than in males may stem from differences in neural circuitry (Bangasser & Valentino, 2014). Given these findings, using an ecological-transactional model, we first examine the impact of ecological stress on amygdala-internalizing outcomes. Second, we examine how parenting and sex may moderate the complex associations among ecological stress, amygdala and internalizing problems. We will use a large representative sample from the Adolescent Brain Cognitive Development (ABCD) study to best capture variability in ecological stress and its impact on brain and behavior. A review of ecological stress related to children's internalizing symptoms

Ecological stress, amygdala reactivity, and internalizing problems and amygdala reactivity, and a rationale for how positive parenting and sex may moderate these links, is reviewed next.

The Impact of Multiple Ecological Stressors on Internalizing Problems and Amygdala Reactivity

Increasing evidence has demonstrated that the exposure to multiple ecological risk factors (such as socioeconomic status [SES] and/or disadvantageous communities) has detrimental effects on children's neuro-behavioral development beyond the effect of any singular risk factor (Evans et al., 2016; Gach et al., 2018). Specifically, childhood cumulative risk has been found to predict higher levels of internalizing problems among preadolescents (Evans et al., 2013). The impact of multiple risk factor exposure is often measured using a cumulative risk approach by dichotomizing each risk factor exposure and then summing the dichotomous scores. While this approach is good for parsimony and for preserving statistical power, it is sample- and context-specific. For example, the “cut-off” threshold for high versus low income would be different if the sample was recruited from high versus low SES neighborhoods. This dichotomized approach thus may not be the most informative approach when using the ABCD sample that was recruited across multiple sites and diverse socio-demographic regions. An alternative approach that takes advantage of the characteristics of the ABCD sample is to use latent variable modeling to understand how multiple ecological factors and the underlying latent construct of ecological stress is associated with preadolescent internalizing symptoms and amygdala functioning. Using a latent construct consisting of household financial strain, neighborhood problems and maternal psychological distress, Loukas, Prelow, Suizzo, and Allua (2008) found that exposure to this multiple risk latent construct among 10- to 14- year old youths predicted subsequent internalizing symptoms 16 months later. Our study therefore will first examine whether multiple

Ecological stress, amygdala reactivity, and internalizing problems

ecological risks available in the ABCD study consisting of household income, neighborhood

SES disadvantage, and neighborhood safety comprise a coherent latent construct using

confirmatory factor analysis (CFA), such that its correlates with brain and behavior can then be identified.

The brain plays a pivotal role in efferent and afferent processing of ecological factors, making its architecture a focal point of the effects of stress (McEwen, Gray, & Nasca, 2015).

Recent reviews demonstrate the increasing role that ecological factors play on neural

development (Farah, 2018; McEwen et al., 2015), including its role in emotional processes,

implicating the amygdala as one region whose function is often related to stress (Tottenham et al., 2009). Decreased family income in preadolescence has been shown to be associated with

heightened activation in the amygdala to facial stimuli later in life (Kim et al., 2013). Moreover,

Evans and colleagues (2016) demonstrated that preadolescent cumulative risk (averaged across 9 and 13 years of age) was associated with elevated amygdala response to emotional faces in

adulthood. Likewise, in an epidemiologic cohort examining long-term outcomes of risk factors,

Holz et al. (2017) reported that the accumulated psychosocial risks from 3-months until 11 years of age were inversely associated with amygdala activation in young adulthood. Although earlier

studies evaluated cumulative stress in childhood on the presentation of altered emotional

reactivity in later development, there is limited evidence about neural reactivity during critical

developmental stages. The ABCD sample offers a unique opportunity to evaluate the transitional

window which may have long-term implications for neural functioning in emotional processing,

as well as potentially presenting high-value targets for intervention and prevention.

Parenting as a Moderator of Risk

Parenting is an important factor that may modify the development of emotional reactivity to stress in preadolescence (Fox, Levitt, & Nelson, 2010). Specifically, parental warmth may be one of the protective factors that may reduce the adverse effect of stress on emotional reactivity (as indexed by amygdala structures and functions; Farber et al. 2019; Whittle et al. 2014), and the development of internalizing symptoms (Gorostiaga et al., 2019). For example, Whittle et al. (2014) examined the associations among disadvantaged communities, maternal behavior and amygdala volumes across adolescent development. They found that positive maternal interactions moderated the association between community disadvantages and amygdala volume, whereby youth in a community with higher disadvantages had similar amygdala volumes relative to those in lower SES communities when they experienced positive parenting. With respect to functional activation, Farber et al. (2019) further provided evidence that positive parenting serves an important role in stress and amygdala functioning, reducing amygdala reactivity to interpersonal emotional faces beyond cumulative risk factors. These results offer preliminary evidence that parenting may moderate how the amygdala may react to emotional stimuli in an ecologically risky environment. However, it is unclear how the moderating effect of parenting on stress-amygdala reactivity may contribute to internalizing symptoms. While the mechanism remains unclear, according to a recent review by Gorostiaga and colleagues (2019), parental warmth may be one important factor that modifies the rate of internalizing problems. As evidence for this, Hipwell et al. (2008)'s meta-analysis of 141 studies found that higher parental warmth was associated with fewer internalizing symptoms. It is therefore important to examine whether parental warmth may moderate the link between ecological stress and amygdala reactivity, as well as between amygdala reactivity and internalizing symptoms, in order to better

Ecological stress, amygdala reactivity, and internalizing problems understand the mechanisms that explain the buffering effect of parental warmth. This is especially critical to explore during preadolescence given that it is a time of high neural plasticity (Lillard & Erisir, 2011) when ecological factors may have direct effects on their neuro- and behavioral development (Hackman, Farah, & Meaney, 2010).

Moderating Role of Sex on Mechanisms of Risk

Understanding the relationship between cumulative risk and internalizing symptoms requires understanding differences between sexes. It is well-documented that internalizing problems are more prevalent in females than males (Beesdo et al., 2009; Costello et al., 2003), that this difference first emerges during adolescence (Essau, Lewinsohn, Seeley, & Sasagawa, 2010), and is differentially affected by economic stressors in late childhood/preadolescence (Leventhal & Brooks-Gunn, 2011; Schneider, Waldfogel, & Brooks-Gunn, 2015). While it remains unclear whether sex differences in internalizing symptoms may stem from differences in neural circuitry (i.e., amygdala reactivity and volume), two particularly salient factors could be the effects of parenting and stress. For instance, in the case of income-to-needs ratio and cortical development, Whittle et al. (2017) found that parenting served as a significant buffer in males but not in females. In another study, using a daily diary method, girls' negative family interactions partially explained differences in internalizing symptoms, and positive family interactions reduced sex differences in internalizing symptoms (Telzer & Fuligni, 2013). Collectively, when attempting to model why parenting and stress could affect internalizing symptoms, it is therefore important to consider sex differences in how parenting may moderate stress-amygdala-internalizing associations.

Current Study

The current study examines the effects of an aggregated ecological risk factor on behavioral outcomes, neural mechanisms, and buffers in a large, representative sample of youth (Figure 1). Due to previous studies relying on incomplete definitions of ecological risk (Boecker et al., 2014; Cohen et al., 2012; Farber et al., 2019; Jaffee, 2017; Marshall et al., 2018; McLaughlin et al., 2017; Noble et al., 2015; Odgers et al., 2012; Vidal-Ribas et al., 2019; Zeanah et al., 2009), we use a multi-source latent factor that assesses ecological risk at the individual and neighborhood levels. When creating a comprehensive metric of individual ecological risk, it is critical to account for neighborhood-level resources and characteristics, such as safety, because they provide relevant and distinct information in addition to an individual's report of income (Braveman et al., 2005; Kachmar, Connolly, Wolf, & Curley, 2019). We predict that this latent structure will demonstrate a positive association between higher cumulative risk and higher child internalizing (Hypothesis 1). Subsequently, the mediating role of the amygdala on the association between increased cumulative risk and internalizing will be analyzed, we predict that increased risk will be associated with increased amygdala activity, which in turn will be related to increased internalizing problems (Hypothesis 2-3), and that consequently the direct relationship between cumulative risk and internalizing will be significantly reduced (Hypothesis 4). Based on the evidence of parental warmth functioning as a moderator of internalizing problems (Gorostiaga et al., 2019), the moderating role of parenting will be examined, with the prediction that social-emotionally positive parenting will result in decreased associations between cumulative risk and a) internalizing problems and b) amygdala reactivity, as well as c) the association between amygdala reactivity and internalizing problems. Although prior studies suggest differences in internalizing problems and positive parenting in males and females

Ecological stress, amygdala reactivity, and internalizing problems (Cyranowski et al, 2000), there is little evidence to predict sex differences in the strength of relations between ecological stressors, internalizing problems, the mediating role of the amygdala and the moderating role of parenting, therefore, exploratory analyses will be performed to identify the presence of sex differences for Hypothesis 1 through Hypothesis 4.

Method

Participants

Data were drawn from the release 3.0 (including Baseline and Year 1 data) of the Adolescent Brain Cognitive Development study, a longitudinal national study of 11,878 9- and 10-year old adolescents across 22 sites (<https://nda.nih.gov/study.html?id=901>; Casey et al., 2018). A subset of participants, based on *a priori* criteria (described below), are identified and used in subsequent analyses, N = 7,385 (see inclusion flowchart, supplementary Figure S1). In consideration for the present analysis, participants with missing information on the variable of interest are not included in the final SEM model.

Procedures

For detailed description of study recruitment and procedures, see Garavan et al. (2018) and Casey et al. (2018). Access to the data was established via an unfunded agreement (19-UFA02466). Approval for data access was also granted by the University of Michigan Institutional Review Board (HUM# HUM00159925). Data were accessed by downloading release 3.0 (<https://nda.nih.gov/study.html?id=901>) via the National Institute of Mental health Data Archie (NDA), the pre-packaged data associated with these analyses are NDA ABCD Study #1182451 (complete R-markdown & Mplus files are available at R, <https://osf.io/bq49h/>). This manuscript is part of a Registered Report submission, preregistered on the Open-Science Framework Registry (<https://osf.io/46yru>).

Ecological stress, amygdala reactivity, and internalizing problems

Measures

Income. Parents reported combined household income in the past 12 months by selecting an income category ranging from (1) “less than \$5,000” to (10) “\$200,000 and greater”. Parents were also given the option to select “refuse to answer” or “don’t know”. The variable used in this analysis is based on the baseline items. This item was reverse coded such that higher scores indicated lower income levels, to maintain the same direction on our cumulative risk factor of ecological stress.

Area Deprivation Index. The Area Community Service (ACS) area deprivation index (a single item) uses census block group-level data to approximate neighborhood level of social-economic neighborhood disadvantage (Kind & Buckingham, 2018). Index scores are derived from 17 education, employment, housing-equality, and income measures within a 5-year period at the census block level. Index scores are percentiles converted to decile groups from 1 (groups 1 -10) to 10 (groups 91-100), where high ranked groups indicate more disadvantage. The variables used in this analysis are based on the baseline items.

Neighborhood safety. A PhenX Toolkit measure (3 items) derived from the Neighborhood and Crime Safety scale (Echeverria, Diez-Roux, & Link, 2004; Mujahid, Diez Roux, Morenoff, & Raghunathan, 2007) assessed parents self-reports of neighborhood safety. Parents rated statements such as “My neighborhood is safe ‘from crime,” “Violence is not a problem in my neighborhood”, and “I feel safe walking in my neighborhood, day or night”, on a 5-point Likert scale ranging from (1) “strongly disagree” to (5) “strongly agree”. Items were averaged such that higher scores indicated higher perception of neighborhood safety. The variables used in this analysis are based on the baseline (Wave 1) items. This item was reversed coded, to maintain the same direction on our cumulative risk factor.

Ecological stress, amygdala reactivity, and internalizing problems

Internalizing Symptoms. Parents' reports on the Child Behavior Checklist (Achenbach & Rescorla, 2001), from Wave 1, was used to assess their child's internalizing symptoms using the broadband internalizing subscale (33 items). Parents rated their child's withdrawn, somatic, anxious, and depressive symptoms on a 3-point Likert type scale from (0) "never" to (2) "often". Parents' ratings were used to create an internalizing problem summary score. Raw scores were normalized into t-scores (mean of 50, standard deviation of 10) to provide a common metric, with higher scores indicating more internalizing symptomatology.

Parental Acceptance. The acceptance subscale (5 items) from the Children's Report of Parental Behavior Inventory (Schaefer, 1965) was used to assess parental warmth and acceptance. Preadolescents responded to items regarding the parent or caregiver who completed the ABCD parent protocol. This was typically the child's primary caregiver, but it did not have to be. Statements such as "Makes me feel better after talking about my worries with him/her," were rated on a 3-point Likert type scale ranging from (1) "Not like him/her" to (3) "A lot like him/her". Items were averaged such that higher scores indicated higher perceptions of parental acceptance. The variable used in this analysis is based on the baseline items.

Pubertal Development: Parental reports on child's pubertal stage were assessed using the Pubertal Development Scale (PDS; Petersen et al, 1998). The PDS is a text-based non-invasive measure to assess current pubertal status in females and males. For females, the items assess changes in 1) body hair, 2) breast development and 3) menstruation. For males, the items assess changes in 1) body hair, 2) hair on the face and 3) deepening voice. The pubertal category score provides a 1-5 reference for whether a participant is: 1 = "pre puberty"; 2 = "early puberty"; 3 = "mid puberty"; 4 = "late puberty"; or 5 = "post puberty".

MRI Acquisition

Emotional N-Back (EN-back) Task. The *EN-back* task (Casey et al., 2018; Cohen et al., 2016) taps into both working memory and emotion regulation processes, though in the present study, we focus on the emotion regulation application. Casey et al. (2018) described the *EN-back* task in the ABCD study. The task contains two runs with eight blocks in each run. Stimuli were either emotional faces (happy, fearful, neutral) or places. Emotional faces were taken from the NimStim (Tottenham et al., 2009) and the Racially Diverse Affective Expressions (Conley et al., 2018) stimuli sets.

At the beginning of each block, instructions were given to specify whether it was a 0-back or a 2-back block. Each block consisted of 10 trials. Each trial began with a stimulus being presented for 2 s, during which time the participant would respond with whether the stimulus matched or did not match the target stimulus. Following the stimulus presentation, a fixation cross was presented for 500 ms. There were 80 trials for each of the two memory load conditions and 40 trials of each stimulus type (20 in each of the two memory-load conditions).

Behavioral performance on this task was used as an exclusion criterion. Participants with less than 60% accuracy were considered to have poor performance by the ABCD data analytic core and are thus excluded from the present analyses (Hagler et al., 2019).

MRI acquisition. For additional information regarding MRI acquisition in the ABCD Study, see Casey et al. (2018). Scans were all performed on different 3T scanners from Siemens (Prisma VE11B-C, Siemens Medical Systems, Erlangen, Germany), Philips (Achieva dStream, Ingenia, Philips Medical Systems, Best, the Netherlands), or General Electric (GE) (MR750, DV25-26, General Electric, Milwaukee, WI, USA). At each site, children were desensitized to the scanner environment through either a mock scanner or a play tunnel that was the size of the

Ecological stress, amygdala reactivity, and internalizing problems

scanner bore. To encourage motion compliance in the scanner, behavioral shaping was used to monitor head motion (Epstein et al., 2007) and provide feedback to the child. Additionally, to minimize head motion in the scanner, the head was stabilized with foam padding.

A T1-weighted (T1w) anatomical scan was acquired using the following parameters: Siemens: TR = 2500ms, TE = 2.88ms, TI = 1060 ms, flip angle = 8°, 176 transverse slices; Philips: TR = 6.31ms, TE = 2.9ms, TI = 1060ms, flip angle = 8°, 225 transverse slices; GE: TR = 2500ms, TE = 2ms, TI = 1060ms, flip angle = 8°, 208 transverse slices. Voxel size was 1 x 1 x 1 mm for all three scanner types.

Multi-slice/multiband EPI functional MRI scans for the *EN*-back task were acquired using the following parameters on all scanner types: TR = 800ms, TE = 30ms, flip angle = 52°, 60 transverse slices, multiband acceleration = 6, voxel size = 2.4 x 2.4 x 2.4 mm.

Preprocessing and MRI data analysis.

The data analytic core in the ABCD consortium MRI preprocessing and analyses were performed by the ABCD consortium's data analytic core using their in-house Multi-Modal Preprocessing Stream (v. 248 https://www.nitrc.org/projects/abcd_study) This pipeline uses a combination of MATLAB functions and publicly available neuroimaging packages including Freesurfer v.5.3.0 (Fischl, 2012), AFNI v.2008_02_01_1144 (Cox, 1996) and FSL v.5.0.2.2-centos5_64 (Jenkinson, Beckmann, Behrens, Woolrich, & Smith, 2012; Smith et al., 2004). For more details, see Hagler et al. (2019).

Structural images. T1w structural images were used during the preprocessing of functional MRI data and the parcellations used for the ROI time course extractions. T1w images were corrected for gradient non-linearity distortions (Jovicich et al., 2006) and for intensity inhomogeneity using B₁-bias fields which are estimated using sparse spatial smoothing and white

Ecological stress, amygdala reactivity, and internalizing problems

matter segmentation. Images were then resampled with 1mm isotropic voxels and registered to standard reference brain for subsequent subcortical segmentation using an automated, atlas-based volumetric segmentation procedure (Fischl et al., 2002) in Freesurfer.

Functional images. Functional images were corrected for head motion by registering each frame to the first using AFNI's 3dvolreg (Cox, 1996) and B₀ field inhomogeneity, which causes spatial and intensity distortions, using a reverse gradient method (Holland, Kuperman, & Dale, 2010). Images were then corrected for gradient non-linearity distortions (Jovicich et al., 2006) and between-scan motion was corrected using alignment to a reference scan. Following this realignment, a registration matrix is calculated to allow for rigid body transformation between fMRI and T1W anatomical images. The fMRI images remain in “native-space” and have a 2.4mm isotropic resolution.

Following image preprocessing, the first 16 frames of the scan were removed to ensure equilibration of the T1W signal and then voxel time series were normalized by dividing by the mean across time of each voxel. Baseline and quadratic trends in the time-series data were removed using linear regression, as were motion estimates and their derivatives (Power et al., 2014). Volumes with greater than 0.9mm framewise displacement were censored (Siegel et al., 2014). Time course data were then bandpass filtered to remove signal between 0.31-0.43 Hz using an infinite impulse response notch filter.

The *EN*-back task was modeled in the data at the individual subject level using a general linear model implemented in AFNI's 3dDeconolve (Cox, 1996). The HRF was modelled using a gamma basis function variate and its temporal derivative. The duration of cues in the *EN*-back task (2.5 seconds) were modeled as square waves that were convolved with the HRF. Incorrect trials were modeled as separate conditions.

Ecological stress, amygdala reactivity, and internalizing problems

ROI data. The present study made use of tabulated data sheets resulting from the ABCD consortium's data analytic core (Hagler et al., 2019). The average signal for the left and right amygdala were calculated based on atlas-based subcortical parcellations in Freesurfer (Fischl et al., 2002) for each run of the *EN*-back task separately across both the 0-back and 2-back conditions for the emotional faces vs places. The average signal across both runs was weighted by the number of frames remaining after motion censoring minus the number of model parameters (not accounting for temporal autocorrelation). Conditions with no valid trials were marked as undefined and had empty cells in the ABCD tabulated data. Conditions with few trials and extremely high standard error of the mean for the beta estimates were also censored following the ABCD protocol for exclusion. We did not have hypotheses regarding laterality, so we averaged the extracted values for the left and right amygdalae; however, lateral (left & right amygdala models) exploratory results will be provided in supplementary materials. Although we cannot use a traditional method to get at mean activation in the amygdala for the aforementioned condition, beta values will be used to provide a distribution of mean-level signal and overall effect of the task condition for the amygdala in supplementary materials. Further, participants were excluded if their fMRI data quality is labeled as 'Reject' based on the FreeSurfer Quality Control variable (*fsqc_qc*) and exceed 0.9 for mean framewise displacement, specific to our analyses.

Statistical Analysis

Pearson correlation, descriptive statistics and missing data pattern analysis will be conducted first for exclusions relating to fMRI data. Data transformation (e.g., log) will be performed if variables (e.g., CBCL internalizing T-score) are deemed to be significantly skewed. Mplus 7.2 will be used to test the hypothesized model (see Figure 1). First, confirmatory factor

Ecological stress, amygdala reactivity, and internalizing problems analysis (CFA) of household income, neighborhood socioeconomic disadvantage level, and neighborhood safety will be performed to examine whether various ecological risk factors should be loaded as one or multiple latent constructs. The CFA will be run on the individual variables (e.g. reverse coded income, area deprivation index, and the reverse coded neighborhood safety). Second, a SEM model will be conducted to examine the associations among latent construct(s) from CFA at Step 1, amygdala reactivity and internalizing symptoms. Mediation/indirect effects will be tested using MODEL INDIRECT in Mplus. Third, to examine the moderating role of parental warmth in the hypothesized stress-amygdala-internalizing mediation model, we will examine the interaction effect of ecological stress by parental warmth on amygdala and internalizing problems. If the interaction term is significant, simple slope analysis will be performed and partial residuals will be plotted using the Visreg package in R to visualize the simple slope analysis. To examine the moderating effect of sex, a multi-group model within Mplus will be conducted to examine whether the hypothesized associations will vary between males and females. To account for the complex sampling effect of siblings within a family cluster, all CFA and path models will be estimated using Taylor-series linearization using Type = Complex in Mplus. To account for the multiple site design of the ABCD study, we specified Stratification = ABCD site for all models in Mplus. Covariates including puberty, race and scanner type will be included in all models.

Multiple-fit statistics are reported and interpreted as outlined (Kline, 2015): (a) Pearson X^2 for which nonsignificant values signify good fit and a X^2/df ratio <3 is acceptable; (b) Comparative Fit Index (CFI) for which a value .90 is considered a good fit; and (3) root mean square error of approximation (RMSEA), for which a value $<.08$ is considered acceptable and $<.05$ is considered good. While missing data will be handled using the Mplus Full Information

Ecological stress, amygdala reactivity, and internalizing problems

Maximum Likelihood (FIML) feature (Muthén & Muthén, 2012), additional analysis will be conducted with and without participants with missing data to ensure that the results are not altered by missingness.

Results

Descriptive Statistics

Based on four exclusion criteria relating to amygdala ROI beta, a participant had to have a) a beta estimate present, b) the FreeSurfer Quality Control variable (fsqc_qc) had to have received “Accept”, c) mean framewise displacement did not exceed .9, and d) behavioral performance during the *EN*-back task had to be greater than 60% accuracy, a subset of participants ($N = 7,385$) from the full sample were retained for subsequent analyses (flowchart, supplementary Figure S1).

For ease of interpreting subsequent SEM models and to maintain continuity throughout, as mentioned in the measures section, parental self-report of a) neighborhood safety and b) baseline income have been reverse coded. Table 1 shows the relevant descriptive statistics for the key variables in the study, both for the participants included in this study ($N = 7385$) and those excluded from the study for reasons outlined in the Methods section ($N = 4493$). There were some notable differences between those included and excluded from these analyses. On average, the included sample were 2 months older than the excluded sample and were in a higher family income decile compared to the excluded sample (decile #3, \$75,000 through \$99,999 and decile #4, \$50,000 through \$74,999, respectively). These effects can both be considered as small in magnitude (Cohen, 1988). The included sample was more likely to be White. In addition, compared to the excluded sample, the included sample had higher neighborhood safety ($d = .19$)

Ecological stress, amygdala reactivity, and internalizing problems

and associated with a lower area deprivation index ($d = -.14$) – however, the size of these minimal-to-small (Cohen, 1988).

Sex differences in pubertal development at baseline were expected given the known differences in the trajectories of pubertal development by age between males and females. On a scale between 1 (e.g., prepuberty) and 5 (e.g., post puberty), females ($M = 2.2$, $SD = 0.9$) showed significantly more pubertal development than males ($M = 1.3$, $SD = 0.6$), $t(5965.4) = 45.9$, $p < .001$, $d = 1.1$, as expected. There was also a significant sex difference for Year 1 internalizing between Males ($M = 49.1$, $SD = 10.5$) and Females ($M = 47.7$, $SD = 10.3$), $t(7100) = 18.9$, $p < .001$; however, this effect was small ($d = -0.13$).

All distributions for variables used in these analyses are reported in supplementary materials, see Figures in Supplemental Section 2.2 to Section 2.3.

Correlations

Correlations were conducted between key demographic variables, (sex, age, pubertal development, income), three indicators of ecological stress (area deprivation index and reverse coded: neighborhood safety, parental neighborhood safety), internalizing symptoms, amygdala activations (left, right and bilateral) and *EN*-back mean framewise displacement. Figure 2 shows the relations between all aforementioned variables; select key correlations are reviewed here.

As expected, pubertal development had a moderate positive correlation with age and income (both r 's = .19) and a large negative correlation with being male ($r = .48$). Lower income was correlated with higher (implying more disadvantage) scores for the area deprivation index ($r = .36$) and lower neighborhood safety ($r = .35$), both at moderate effect sizes. Area deprivation index and neighborhood safety had a small correlation with each other ($r = .20$). As expected,

Ecological stress, amygdala reactivity, and internalizing problems bilateral, left and right amygdala activations (see mean beta estimates in supplemental section 2.2, Figure S6 and S7) were highly correlated with each other, $r = .88 - .90$.

Confirmatory Factor Analysis: Ecological Stress

CFA revealed that household income (factor loading = .81, $p < .001$), area deprivation (factor loading = .45, $p < .001$) and neighborhood safety (factor loading = .44, $p < .001$) are loaded as a single latent construct.

Structural Equation Models (SEM)

Hypothesis 1: Ecological stress latent construct will positively associate with preadolescents' internalizing symptoms.

Supporting our hypothesis, SEM revealed that higher levels of ecological stress (latent construct at baseline) was significantly associated with higher levels of internalizing symptoms at 1 -year follow-up (Figure 3A). Model fit indices indicated adequate fit in the model: Chi-square (17) = 174, $p < .001$, REMSEA = .036, CFI = .93, SRMR = .02.

Multi-group SEM model (and Wald Test of Parameter Constraints for path coefficient differences) was conducted to test whether the effect of ecological stress on internalizing symptoms differed by sex. Results indicated that the effect of ecological stress on preadolescents' internalizing symptoms was stronger in males than females (Wald (1) = 5.43, $p = .02$; Figure 3B).

Hypothesis 2: Increased ecological stress will be associated with increased amygdala activity

No association was found between ecological stress and bilateral amygdala reactivity (Figure 3A), nor left or right amygdala activity (see Supplementary Figure S10). No sex difference was found (Figure 3B).

Ecological stress, amygdala reactivity, and internalizing problems

Hypothesis 3: Increased amygdala activity at baseline will be related to increased internalizing problems at 1-year follow.

No association was found between any amygdala activity and internalizing symptoms. (Figure 3A; Supplementary **Figure S10**). No sex difference was found (Figure 3B).

Hypothesis 4: The direct relationship between ecological stress and internalizing will be significantly reduced when including amygdala reactivity as a mediator

Bilateral (or left or right) amygdala activity at baseline did not mediate the link between ecological stress and internalizing symptoms at 1-year follow-up (Figure 3A). No sex difference was found (Figure 3B). Which is to be expected given that hypothesis 2 and 3, mentioned above, were not supported.

Hypothesis 5: The moderating role of parental acceptance

High parental acceptance at baseline was negatively associated with preadolescent's internalizing symptoms at 1-year follow-up. However, parental acceptance did not moderate any links (ecological stress- internalizing problems, ecological stress- amygdala, and amygdala - internalizing symptoms) in our model (Figure 3A). No sex difference was found (Figure 3B).

Post Hoc Analyses

Given that parental acceptance significantly associated with preadolescents' internalizing symptoms and we did not find a moderating effect of parental acceptance, we also tested for potential mediation effects. Specifically, we conducted a mediation analysis using MODEL INDIRECT in Mplus to test for whether parental acceptance mediated the associations between ecological stress at baseline and preadolescents' internalizing symptoms at 1-year follow-up, with the same covariates (puberty, race and scanner type), cluster (family) and stratification (ABCD site) in the model. Results indicated that there was a significant indirect effect (est =

Ecological stress, amygdala reactivity, and internalizing problems

.002, $p = .025$), such that lower parental acceptance mediated the associations between ecological stress and internalizing symptoms, indicating that the effect of ecological stress on internalizing symptomatology was higher for preadolescents reporting lower parental acceptance (Supplementary **Figure S11**). Model fit indices indicated an adequate fit Chi-square (43) = 161, $p < .001$, REMSEA = .036, CFI = .94, SRMR = .02.

Multi-group analysis of sex effect was also conducted, and the indirect effect of parental acceptance did not reveal significant mediation when examining the model separately by sex.

Sensitivity Analyses

In our original registration of this project, we had proposed four exclusion criteria for the amygdala ROI beta estimates that were central to the model: a) a participant had to have a beta estimate present, b) the FreeSurfer Quality Control variable (fsqc_qc) had to have been labeled as “Accept”, c) mean framewise displacement did not exceed .9, and d) behavioral performance during the *EN*-back task had to be greater than 60% accuracy. Upon downloading the Release 3.0 (NDA ABCD Study #1182451), we examined the ABCD Consortium recommendations for inclusion/exclusions criteria related to the *EN*-back task data. Their recommendations were more stringent than those we had originally proposed. As such, it was prudent to check the results of our models with these stricter exclusion criteria, as artifacts in the neural data may contribute additional noise into our models and impact the results related to amygdala activation.

In addition to our original criteria (a-c), there were eight additional criteria used for inclusion/exclusion to perform our sensitivity analysis. We ensured that both the Raw 1) Functional *EN*-back (iqc_nback_ok_ser) and 2) T1 images (iqc_t1_ok_ser) passed quality control; 3) >200 degrees of freedom were available for the *EN*-back task (tfmri_nback_all_beta_dof); 4) each participant had > 100 trials of *EN*-back data

Ecological stress, amygdala reactivity, and internalizing problems

(fmri_nb_all_beh_total_nt) ; 5) there was a match with the e-prime data

(iqc_nback_ep_t_series_match); 6) the B0 fieldmaps were available

(apqc_fmri_bounwarp_flag); 7) there were no discrepancies between manual ratings in fMRI QC

(fmri_postqc_qc); and 8) the ventral FOV cutoff score was <60 (apqc_fmri_fov_cutoff_ventral).

This resulted in a sensitivity sample of $N = 5,807$ (see flowchart in supplementary Figure S12).

The reduction in eligible participants was largely driven by the exclusion due to rater

discrepancy (fMRI Manual QC), whereby $N = 1,476$ were excluded due to discrepancy between raters.

Our results with respect to the links between amygdala, ecological stress, internalizing and parental warmth did not change when applying this sensitivity check. Furthermore, the Pearson correlations among the variables between the full ($N = 7,385$) and constrained sample ($N = 5,807$) remained relatively stable. Specifically, with the exception of two bivariate correlations that had a change of $r = |0.02|$, a small fraction of bivariate associations reflects a change in $r < |.015|$. The SEM model and Pearson correlations for the constrained sample are reported in supplementary materials, Section 4: Sensitivity Analyses.

Discussion

In light of the paucity of the extant literature on the mediating role of brain function in ecological stress and behavioral outcomes, the overarching goal of this study was to evaluate the neurodevelopmental link between ecological stress, amygdala reactivity and internalizing problems, and the buffering effect of parental acceptance, in a large, geographically and socio-economically diverse sample of preadolescents. In this well-powered registered report, we created a latent construct of ecological stress based on household income, self-reported neighborhood safety, and neighborhood disadvantage. Using this construct, we confirmed, in

Ecological stress, amygdala reactivity, and internalizing problems

part, our hypothesis that there would be a significant association between ecological stress at baseline and subsequent internalizing symptoms at 1-year follow-up; however, this association varied by sex. We did not find evidence to support our hypothesis that there would be a link between ecological stress and amygdala activation during the *Emotional N-back (EN-back)* task. Likewise, we found no support for the hypothesis that amygdala activation mediated the relationship between ecological stress and internalizing symptoms at the 1-year follow-up. Finally, although we did not find a moderating effect of parental acceptance on either of the relationships in the model, post hoc analyses did reveal a mediating role of parental acceptance on the association between ecological stress and internalizing symptoms.

It is well documented that cumulative ecological stress has adverse effects on psychological development (Duffy et al., 2018; McEwen, 2012), such as internalizing problems (Evan et al., 2013). Consistent with this broad literature, in this large longitudinal sample of preadolescents, we find a significant association between baseline ecological stress, measured at 9-10 years of age, and internalizing symptoms 1 year later. This finding is consistent with work reporting that multiple risk factors during early adolescence predicted higher rates of internalizing symptoms at a 16-month follow-up (Loukas et al., 2008). Our findings extend the adolescent literature to show that ecological stress impacts internalizing symptoms during preadolescent development in a large sample.

Previous work on internalizing symptoms has predominantly focused on changes during adolescence, reporting that rates of internalizing problems are higher in females than males (Beesdo et al., 2009; Costello et al., 2003). In contrast to this past work, the preadolescent males in our sample reported significantly more internalizing symptoms than females, and the association between ecological stress and internalizing symptoms was stronger in males than

Ecological stress, amygdala reactivity, and internalizing problems

females. Previous work suggests that sex differences in internalizing symptoms become more pronounced in adolescence with pubertal development (Angold et al., 1998; Salk et al., 2017).

The preadolescents in our sample were in relatively early stages of puberty, based on parent self-reported Pubertal Development Scale ($M = 1.73$ on a scale of 1-5), and thus it may be too early to observe the post-pubertal increase in internalizing symptoms. Additionally, the effect size of the mean difference in internalizing symptoms was very small, explaining <1% of variance (issues of effect sizes in ABCD are discussed below). As the ABCD sample gets older, it will be essential to consider how sex differences in internalizing symptoms change and would be expected to become more pronounced over time.

A central hypothesis of the present study considered the role of amygdala activation in the relation to ecological stress and internalizing symptoms. Previous work has found evidence for an association between childhood poverty and amygdala reactivity to emotional faces in early adulthood (Evans et al., 2016; Kim et al., 2013). This link was not supported, however, in the transitional window of preadolescence when looking at the association between childhood ecological stress and concurrent amygdala activation in faces versus places during the *EN*-back task. We considered the association between ecological stress and an average of bilateral amygdala activation, as well as amygdala activation in left or right hemispheres during the *N*-back contrast to assess for possible hemispheric differences. Estimates were not significant in either approach, even when the analyses were constrained to a smaller group who met rigorous quality control standards for the fMRI data. It is possible that the influence of ecological stress on amygdala reactivity is stronger when these ecological factors are assessed in early childhood development when the amygdala is more sensitive to affective stimuli (Silvers et al., 2017), relative to adolescents. This highlights the need to understand how developmental timing of

Ecological stress, amygdala reactivity, and internalizing problems

stress exposure (Cohodes et al., 2020) may moderate the effect of ecological stress on amygdala activation and its relation with internalizing symptoms. Nevertheless, ecological stress tends to be stable across time and the extant literature has found evidence that amygdala activation mediates the association between cumulative stress and internalizing symptoms in a college sample (Swartz et al., 2015), but this was also not found in our preadolescent sample. Future work in the ABCD sample could examine whether *changes* of amygdala reactivity across development may mediate the associations between ecological risk and internalizing symptoms using the ABCD longitudinal fMRI design.

Moreover, our study found no association between parental acceptance and amygdala activation (to emotional faces versus places). Additionally, although it does have a significant direct effect on internalizing symptoms, parental acceptance is not a buffer against the adverse impact of ecological stress on amygdala and subsequent internalizing symptoms. This differs from work that has reported the important role of positive parenting on amygdala-medial PFC functional connectivity in young adults (Farber et al., 2019). These results may differ as a function of age and units of analyses. While it is beyond the scope of our study, future work in the ABCD study may consider how parental acceptance is associated with fronto-amygdala connectivity, and whether or not the association between parenting and amygdala reactivity changes with development.

Although parental acceptance was not found to be an important moderator, in post hoc analyses parental acceptance was found to mediate the relationship between ecological stress and internalizing symptoms. Parenting behavior is thought to impact the effect of stress exposure on youth (Gorostiaga et al., 2019). Thus, parental acceptance could be changing the magnitude of the relationship between ecological stress and internalizing symptoms (moderator), or

Ecological stress, amygdala reactivity, and internalizing problems

alternatively, it could be the mechanism (mediator) explaining the relationship between ecological stress and internalizing symptoms. Similar to previous work showing that higher rates of parental warmth are associated with lower depressive symptoms in a longitudinal adolescent sample (Garthe et al., 2015), our results show that higher parental acceptance at baseline is associated with lower internalizing symptoms at 1-year follow-up. Likewise, our results suggest that parental acceptance mediates the relationship between ecological stress and follow-up internalizing symptoms. However, this effect was small and other aspects of parenting may confer different results (Gorostiaga et al., 2019). Future analyses in the ABCD study should consider the possible mechanistic role that parental acceptance plays in the link between ecological stress and negative developmental outcomes.

While these analyses are the first to examine the mediating role of amygdala function in the association between baseline cumulative stress and 1-year follow-up internalizing symptoms in the ABCD data, several cross-sectional analyses have considered associations between socioeconomic variables and brain structure and cognitive function. In a cross-sectional analysis of the baseline ABCD study data, Assari (2020) reported that parental socioeconomic status (SES), as measured by neighborhood income (ADI) and family income, mediated the relationship between ethnoracial differences and amygdala volume. Meanwhile, in their cross-sectional analysis, Taylor and colleagues (2020) probed the relationship between neighborhood poverty, as measured by 9 of 17 ADI variables, hippocampal and prefrontal cortex (PFC) volumes, and cognitive assessments across seven cognitive domains from the NIH toolbox. In their results, Taylor and colleagues (2020) found that higher household income was positively associated with hippocampal and PFC volumes, and that neighborhood poverty was negatively related to the volume of some PFC regions and the right hippocampus. They also found that

Ecological stress, amygdala reactivity, and internalizing problems

higher neighborhood poverty was negatively associated with scores across all seven cognitive domains. Likewise, Vargas et al., (2020) reported that neighborhood deprivation, as measured by ADI quantiles (1-5; five being the highest deprivation), was negatively related to cognitive function and that this association was partially mediated by PFC surface area. Although all of these analyses considered cross-sectional associations, it will be important to consider how SES and brain associations change across time -- a strength of the ABCD design. Furthermore, although the brain was identified as mediator in cross-sectional analyses of the ABCD sample (Assari, 2020; Assari et al., 2020; Vargas et al., 2020), it will be important to reexamine the mechanistic role of specific brain regions using the longitudinal ABCD data to distinguish between cross-sectional and longitudinal effects. Finally, to describe how SES manifests in the brain (Farah, 2017) and relates to cognitive and behavioral outcomes, future work should examine differences in the analytic pipeline that contribute to convergence and differences in findings.

Study Considerations

This study's findings are novel and provide insight into the relations between ecological stress, internalizing symptoms and amygdala function. The major strengths of this study include its large sample size that is far greater than the median sample size, $N < 50$, in MRI studies (Szucs & Ioannidis, 2020), and our attempt to reduce concerns relating to questionable research practices (i.e., p-hacking, publication bias, HARKing, etc.) (Chambers et al., 2014) and researcher degrees of freedom (Simmons et al., 2011) by using a registered report format. Nevertheless, this study must be understood within the context of several important factors that affect its generalizability.

Ecological stress, amygdala reactivity, and internalizing problems

First, there continue to be important differences in how ecological stress and early adversity is defined and measured. Competing theoretical frameworks exist which support examining neural and behavior correlates of early life stress by modeling adversity as a multidimensional (McLaughlin et al., 2014) and unidimensional construct (Smith & Pollak, 2020). Additionally, there is variability in the items included in early life stress constructs (e.g., home, community). These approaches all have merit though the heterogeneity in constructs can make it difficult to compare results across studies. Therefore, it is worth noting that our *a priori* ecological stress construct may differ from others and thus may influence the generalizability of our results. Specifically, the latent factor derived from the *a priori* set of variables in our confirmatory factor analysis was weighted towards self-reported household income, which is a family level stress and may be different from nuanced community level factors that employ cost-of-living adjustments.

Importantly, a substantial portion (approximately 38%) of the overall participants who had at least partially complete data available needed to be excluded from the present analyses based on MRI exclusion criteria. Analyses of the differences in demographic variables between included and excluded participants suggest that the included sample 1) is less ethnically diverse and 2) has a higher average SES than the excluded sample, although the differences are small in magnitude. While the decision to exclude participants with particularly low accuracy on the working memory measure (Hagler et al., 2019) and with comparatively high amounts of motion in the scanner in order to reduce noise are justified (Caballero-Gaudes & Reynolds, 2017), it could be argued that such exclusions risk the models in these analyses not reflecting children with lower working memory skills or with less ability to remain still during imaging. This issue

Ecological stress, amygdala reactivity, and internalizing problems

is not new in the field and requires continued efforts to create developmentally appropriate cognitive measures and correct for in-scanner motion while excluding as little data as possible.

The effect sizes in this study are comparably smaller than what might be expected based on the extant literature; however, they may be closer to the “true effects” for this population. Given that it has been established that smaller samples tend to artificially inflate effects due to their comparatively increased sample variability (Button et al., 2013; Marek et al., 2020; Szucs & Ioannidis, 2017), and that the estimated median sample in fMRI is less than 50 (Szucs & Ioannidis, 2020), former studies may be underpowered to find small effects. Furthermore, Owens et al. (2020) argues that given that the median correlation between questionnaire and task variables in the ABCD study is only $r = .03$, which by the standards set in Cohen (1988) would be considered very small, it is likely that the size of effects in behavioral psychology has been overestimated due to inflation from underpowered studies. Therefore, although the effects are small compared to the previous literature, there should be relative confidence in the magnitude of the effects found in these analyses.

Finally, to probe amygdala activation in these analyses, we focused on the contrast of faces versus places during the *EN*-back task. Although there is a basis for behavioral and neural effects of emotional faces in development (Somerville et al., 2011), there are likely differences in explicit, implicit and passive-viewing designs that elicit different activation (García-García et al., 2016). Thus, this analytic choice may contribute to differences in the results, as this would change the mean beta estimates of amygdala activity. While keeping in mind the limitations in the subtraction process of task-based fMRI (Cacioppo et al., 2003; Price & Friston, 2005), future work should consider how brain function in alternative contrasts relates to ecological stress.

Conclusion

The ABCD study presents a large-scale opportunity to learn about adolescent brain development and the role of ecological stress. The present study investigated how ecological stress, defined as household and community economic resources and neighborhood safety, could increase risk for preadolescents' development of internalizing symptoms. Ecological stress was indeed associated with developing more internalizing symptoms one year later and to a slightly larger extent for males compared to females. In contrast to what has been reported among adolescent and young adult samples, amygdala reactivity to emotional stimuli did not mediate the relation between ecological stress and preadolescent's internalizing symptoms. In addition, parental acceptance did not lessen the association between ecological risk, amygdala reactivity, and internalizing symptoms. However, parental acceptance was meaningfully related to ecological stress and internalizing symptoms with higher rates of acceptance relating to lower internalizing symptoms one year later. This finding has strong implication in the design of parenting intervention that aims to reduce internalizing problems among preadolescents. These results also suggest that the amygdala activity alone does not play a meaningful role in the association between ecological stress and internalizing symptoms in this age group and in our sample. Additionally, they shed light on alternative mechanisms for the role of parenting in stressful environments in a preadolescent sample.

Figure 1: Conceptual model testing the moderation-mediation model of baseline ecological risk, amygdala reactivity (mediator), parental warmth (moderator) and Wave 1 internalizing problems. Income = baseline parental self-reported income; ADI = baseline area deprivation index; P-NB = baseline parental self-report of neighborhood safety. Amygdala = baseline bilateral activation of amygdala on *EN*-back task contrasting faces versus places.

Ecological stress, amygdala reactivity, and internalizing problems

Table 1. Table showing differences in key demographic and cognitive variables between the included and excluded samples at baseline.

	Included Sample (Baseline) N=7385	Excluded Sample (Baseline) N=4493	Effect size
Sex			$\Phi = .04$
Female	3656 (49.5%)	2026 (45.1%)	
Male	3729 (50.5%)	2467 (54.9%)	
Race			$\Phi = .18$
White	4259 (57.7%)	1925 (42.8%)	
Black	787 (10.7%)	998 (22.2%)	
Hispanic	1414 (19.1%)	997 (22.2%)	
Asian	165 (2.2%)	89 (2.0%)	
Other	760 (10.3%)	487 (10.8%)	
	<i>M(SD)</i>	<i>M(SD)</i>	
Age (Months)	120 (7.52)	118 (7.33)	$d = .24$
Family Income Deciles	7.6 (2.2)	6.6 (2.7)	$d = .38$
1. \$200,000 and greater	885 (12.0%)	365 (8.1%)	
2. \$100,000 through \$199,999	2325 (31.5%)	990 (22.0%)	
3. \$75,000 through \$99,999	1061 (14.4%)	511 (11.4%)	
4. \$50,000 through \$74,999	944 (12.8%)	555 (12.3%)	
5. \$35,000 through \$49,999	541 (7.3%)	393 (8.7%)	
6. \$25,000 through \$34,999	355 (4.8%)	299 (6.7%)	
7. \$16,000 through \$24,999	273 (3.7%)	251 (5.6%)	
8. \$12,000 through \$15,999	138 (1.9%)	136 (3.0%)	
9. \$5,000 through \$11,999	194 (2.6%)	227 (5.1%)	
10. Less than \$5,000	151 (2.0%)	266 (5.9%)	
Refuse to answer	278 (3.8%)	234 (5.2%)	
Don't Know	240 (3.2%)	264 (5.9%)	
Pubertal Development	1.73 (0.86)	1.79 (0.88)	$d = -.07$
ADI	91.4 (24.4)	94.9 (25.4)	$d = -.14$
Internalizing Symptoms	48.20 (10.4)	48.90 (11.0)	$d = -.06$
Neighborhood Safety	3.96 (0.93)	3.77 (1.04)	$d = .19$
Parental Acceptance	2.79 (0.29)	2.77 (0.32)	$d = .06$

Note: Unless specified otherwise, each cell refers to the N of that group and the percentage that group composes of the given variable within either the included or excluded sample. Positive effects refer to higher mean scores for the included sample; negative effects refer to higher mean scores for the excluded sample.

Figure 2. Correlation table showing the direction and magnitude of effects between key study variables. As indicated in the key, the larger the colored slice of the pie chart, the larger the effect. Positive correlations are shown in blue; negative correlations are shown in red. Pie slices denote Pearson's r values, and the associated magnitude is represented by the color in the colorbar.

ADI = Area Deprivation Index; NSafe = Parental Report Neighborhood Safety; Accept = Parental Acceptance; Bilat Amyg = Bilateral Amygdala; meanFD = Mean Framewise Displacement during *EM-Nback*. Sex: Male = 1, Female = 0. CBCL(t) = Internalizing Symptoms

Ecological stress, amygdala reactivity, and internalizing problems

Figure 3: Results from the models testing the main model proposed in Figure 1. A) Full model evaluating the role of ecological risk, bilateral amygdala activation, moderating effect of parental acceptance on follow-up internalizing symptoms; B) Model freely estimating ecological risk to follow-up internalizing symptoms, indicating a difference across sex. Both models controlled for age, puberty, race, scanner type, clustered by family, stratified by site.

*Note: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$*

References

- Achenbach, T. M., & Rescorla, L. (2001). *Manual for the ASEBA school-age forms & profiles: An integrated system of multi-informant assessment*. Retrieved from <https://trove.nla.gov.au/version/11167240>
- Angold, A., Costello, E. J., & Worthman, C. M. (1998). Puberty and depression: The roles of age, pubertal status and pubertal timing. *Psychological Medicine*, 28(1), 51–61. <https://doi.org/10.1017/S003329179700593X>
- Assari, S. (2020). Socioeconomic Status Inequalities Partially Mediate Racial and Ethnic Differences in Children’s Amygdala Volume. *Studies in Social Science Research*, 1(2), 62–79. <https://doi.org/10.22158/sssr.v1n2p62>
- Assari, S., Boyce, S., Bazargan, M., & Caldwell, C. H. (2020). Family Income Mediates the Effect of Parental Education on Adolescents’ Hippocampus Activation During an N-Back Memory Task. *Brain Sciences*, 10(8), 520. <https://doi.org/10.3390/brainsci10080520>
- Bangasser, D. A., & Valentino, R. J. (2014). Sex differences in stress-related psychiatric disorders: Neurobiological perspectives. *Frontiers in Neuroendocrinology*, 35(3), 303–319. <https://doi.org/10.1016/j.yfrne.2014.03.008>
- Beesdo, K., Knappe, S., & Pine, D. S. (2009). Anxiety and anxiety disorders in children and adolescents: Developmental issues and implications for DSM-V. *The Psychiatric Clinics of North America*, 32(3), 483–524. <https://doi.org/10.1016/j.psc.2009.06.002>
- Boecker, R., Holz, N. E., Buchmann, A. F., Blomeyer, D., Plichta, M. M., Wolf, I., ... Laucht, M. (2014). Impact of early life adversity on reward processing in young adults: EEG-fMRI results from a prospective study over 25 years. *PloS One*, 9(8), e104185. <https://doi.org/10.1371/journal.pone.0104185>

Ecological stress, amygdala reactivity, and internalizing problems

Braveman, P. A., Cubbin, C., Egerter, S., Chideya, S., Marchi, K. S., Metzler, M., & Posner, S.

(2005). Socioeconomic Status in Health Research: One Size Does Not Fit All. *JAMA*, 294(22), 2879–2888. <https://doi.org/10.1001/jama.294.22.2879>

Button, K. S., Ioannidis, J. P. A., Mokrysz, C., Nosek, B. A., Flint, J., Robinson, E. S. J., &

Munafò, M. R. (2013). Power failure: Why small sample size undermines the reliability of neuroscience. *Nature Reviews Neuroscience*, 14(5), 365–376.

<https://doi.org/10.1038/nrn3475>

Caballero-Gaudes, C., & Reynolds, R. C. (2017). Methods for cleaning the BOLD fMRI signal.

NeuroImage, 154, 128–149. <https://doi.org/10.1016/j.neuroimage.2016.12.018>

Cacioppo, J. T., Berntson, G. G., Lorig, T. S., Norris, C. J., Rickett, E., & Nusbaum, H. (2003).

Just because you're imaging the brain doesn't mean you can stop using your head: A primer and set of first principles. *Journal of Personality and Social Psychology*, 85(4), 650–661. <https://doi.org/10.1037/0022-3514.85.4.650>

Casey, B. J., Cannonier, T., Conley, M. I., Cohen, A. O., Barch, D. M., Heitzeg, M. M., ...

ABCD Imaging Acquisition Workgroup. (2018). The Adolescent Brain Cognitive Development (ABCD) study: Imaging acquisition across 21 sites. *Developmental Cognitive Neuroscience*, 32, 43–54. <https://doi.org/10.1016/j.dcn.2018.03.001>

Chambers, C. D., Feredoes, E., Muthukumaraswamy, S. D., & Etchells, P. (2014). Instead of

“playing the game” it is time to change the rules: Registered Reports at AIMS

Neuroscience and beyond. *AIMS Neuroscience*, 1(1), 4–17.

<https://doi.org/10.3934/Neuroscience.2014.1.4>

Cohen, A. O., Breiner, K., Steinberg, L., Bonnie, R. J., Scott, E. S., Taylor-Thompson, K. A., ...

Casey, B. J. (2016). When Is an Adolescent an Adult? Assessing Cognitive Control in

Ecological stress, amygdala reactivity, and internalizing problems

Emotional and Nonemotional Contexts. *Psychological Science*, 27(4), 549–562.

<https://doi.org/10.1177/0956797615627625>

Cohen, J. (1988). *Statistical Power Analysis for the Behavioral Sciences*. Academic Press.

Cohen, J. A., Mannarino, A. P., Kliethermes, M., & Murray, L. A. (2012). Trauma-focused CBT for youth with complex trauma. *Child Abuse & Neglect*, 36(6), 528–541.

<https://doi.org/10.1016/j.chiabu.2012.03.007>

Cohodes, E. M., Kitt, E. R., Baskin-Sommers, A., & Gee, D. G. (2020). Influences of early-life stress on frontolimbic circuitry: Harnessing a dimensional approach to elucidate the effects of heterogeneity in stress exposure. *Developmental Psychobiology*, n/a(n/a).

<https://doi.org/10.1002/dev.21969>

Conley, M. I., Dellarco, D. V., Rubien-Thomas, E., Cohen, A. O., Cervera, A., Tottenham, N., & Casey, B. (2018). The racially diverse affective expression (RADIATE) face stimulus set. *Psychiatry Research*, 270, 1059–1067. <https://doi.org/10.1016/j.psychres.2018.04.066>

Costello, E. J., Mustillo, S., Erkanli, A., Keeler, G., & Angold, A. (2003). Prevalence and development of psychiatric disorders in childhood and adolescence. *Archives of General Psychiatry*, 60(8), 837–844. <https://doi.org/10.1001/archpsyc.60.8.837>

Cox, R. W. (1996). AFNI: Software for analysis and visualization of functional magnetic resonance neuroimages. *Computers and Biomedical Research, an International Journal*, 29(3), 162–173. <https://doi.org/10.1006/cbmr.1996.0014>

Cyranowski, J. M., Frank, E., Young, E., & Shear, M. K. (2000). Adolescent onset of the gender difference in lifetime rates of major depression: A theoretical model. *Archives of General Psychiatry*, 57(1), 21–27. <https://doi.org/10.1001/archpsyc.57.1.21>

Ecological stress, amygdala reactivity, and internalizing problems

Duffy, K. A., McLaughlin, K. A., & Green, P. A. (2018). Early life adversity and health-risk behaviors: Proposed psychological and neural mechanisms. *Annals of the New York Academy of Sciences*, 1428(1), 151–169. <https://doi.org/10.1111/nyas.13928>

Echeverria, S. E., Diez-Roux, A. V., & Link, B. G. (2004). Reliability of self-reported neighborhood characteristics. *Journal of Urban Health: Bulletin of the New York Academy of Medicine*, 81(4), 682–701. <https://doi.org/10.1093/jurban/jth151>

Epstein, J. N., Casey, B. J., Tonev, S. T., Davidson, M., Reiss, A. L., Garrett, A., ... Spicer, J. (2007). Assessment and Prevention of Head Motion During Imaging of Patients with Attention Deficit Hyperactivity Disorder. *Psychiatry Research*, 155(1), 75–82. <https://doi.org/10.1016/j.psychresns.2006.12.009>

Essau, C. A., Lewinsohn, P. M., Seeley, J. R., & Sasagawa, S. (2010). Gender differences in the developmental course of depression. *Journal of Affective Disorders*, 127(1–3), 185–190. <https://doi.org/10.1016/j.jad.2010.05.016>

Evans, G. W., Li, D., & Whipple, S. S. (2013). Cumulative risk and child development. *Psychological Bulletin*, 139(6), 1342–1396. <https://doi.org/10.1037/a0031808>

Evans, G. W., Swain, J. E., King, A. P., Wang, X., Javanbakht, A., Ho, S. S., ... Liberzon, I. (2016). Childhood Cumulative Risk Exposure and Adult Amygdala Volume and Function. *Journal of Neuroscience Research*, 94(6), 535–543. <https://doi.org/10.1002/jnr.23681>

Farah, M. J. (2017). The Neuroscience of Socioeconomic Status: Correlates, Causes, and Consequences. *Neuron*, 96(1), 56–71. <https://doi.org/10.1016/j.neuron.2017.08.034>

Ecological stress, amygdala reactivity, and internalizing problems

Farah, M. J. (2018). Socioeconomic status and the brain: Prospects for neuroscience-informed policy. *Nature Reviews. Neuroscience*, 19(7), 428–438. <https://doi.org/10.1038/s41583-018-0023-2>

Farber, M. J., Kim, M. J., Knodt, A. R., & Hariri, A. R. (2019). Maternal overprotection in childhood is associated with amygdala reactivity and structural connectivity in adulthood. *Developmental Cognitive Neuroscience*, 40, 100711. <https://doi.org/10.1016/j.dcn.2019.100711>

Fischl, B. (2012). FreeSurfer. *NeuroImage*, 62(2), 774–781. <https://doi.org/10.1016/j.neuroimage.2012.01.021>

Fischl, B., Salat, D. H., Busa, E., Albert, M., Dieterich, M., Haselgrove, C., ... Dale, A. M. (2002). Whole Brain Segmentation: Automated Labeling of Neuroanatomical Structures in the Human Brain. *Neuron*, 33(3), 341–355. [https://doi.org/10.1016/S0896-6273\(02\)00569-X](https://doi.org/10.1016/S0896-6273(02)00569-X)

Fox, S. E., Levitt, P., & Nelson, C. A. (2010). How the timing and quality of early experiences influence the development of brain architecture. *Child Development*, 81(1), 28–40. <https://doi.org/10.1111/j.1467-8624.2009.01380.x>

Gach, E. J., Ip, K. I., Sameroff, A. J., & Olson, S. L. (2018). Early cumulative risk predicts externalizing behavior at age 10: The mediating role of adverse parenting. *Journal of Family Psychology: JFP: Journal of the Division of Family Psychology of the American Psychological Association (Division 43)*, 32(1), 92–102. <https://doi.org/10.1037/fam0000360>

Garavan, H., Bartsch, H., Conway, K., Decastro, A., Goldstein, R. Z., Heeringa, S., ... Zahs, D. (2018). Recruiting the ABCD sample: Design considerations and procedures.

Ecological stress, amygdala reactivity, and internalizing problems

Developmental Cognitive Neuroscience, 32, 16–22.

<https://doi.org/10.1016/j.dcn.2018.04.004>

García-García, I., Kube, J., Gaebler, M., Horstmann, A., Villringer, A., & Neumann, J. (2016).

Neural processing of negative emotional stimuli and the influence of age, sex and task-related characteristics. *Neuroscience & Biobehavioral Reviews*, 68, 773–793.

<https://doi.org/10.1016/j.neubiorev.2016.04.020>

Garthe, R. C., Sullivan, T., & Kliwer, W. (2015). Longitudinal Relations Between Adolescent and Parental Behaviors, Parental Knowledge, and Internalizing Behaviors Among Urban Adolescents. *Journal of Youth and Adolescence*, 44(4), 819–832.

<https://doi.org/10.1007/s10964-014-0112-0>

Gorostiaga, A., Aliri, J., Balluerka, N., & Lameirinhas, J. (2019). Parenting Styles and

Internalizing Symptoms in Adolescence: A Systematic Literature Review. *International Journal of Environmental Research and Public Health*, 16(17).

<https://doi.org/10.3390/ijerph16173192>

Gunnar, M., & Quevedo, K. (2007). The neurobiology of stress and development. *Annual Review of Psychology*, 58, 145–173. <https://doi.org/10.1146/annurev.psych.58.110405.085605>

Hackman, D. A., Farah, M. J., & Meaney, M. J. (2010). Socioeconomic status and the brain:

Mechanistic insights from human and animal research. *Nature Reviews. Neuroscience*, 11(9), 651–659. <https://doi.org/10.1038/nrn2897>

Hagler, D. J., Hatton, S., Cornejo, M. D., Makowski, C., Fair, D. A., Dick, A. S., ... Dale, A. M.

(2019). Image processing and analysis methods for the Adolescent Brain Cognitive Development Study. *NeuroImage*, 202, 116091.

<https://doi.org/10.1016/j.neuroimage.2019.116091>

Ecological stress, amygdala reactivity, and internalizing problems

Hipwell, A., Keenan, K., Kasza, K., Loeber, R., Stouthamer-Loeber, M., & Bean, T. (2008).

Reciprocal influences between girls' conduct problems and depression, and parental punishment and warmth: A six year prospective analysis. *Journal of Abnormal Child Psychology*, 36(5), 663–677. <https://doi.org/10.1007/s10802-007-9206-4>

Holland, D., Kuperman, J. M., & Dale, A. M. (2010). Efficient correction of inhomogeneous static magnetic field-induced distortion in Echo Planar Imaging. *NeuroImage*, 50(1), 175–183. <https://doi.org/10.1016/j.neuroimage.2009.11.044>

Holz, N. E., Boecker-Schlier, R., Buchmann, A. F., Blomeyer, D., Jennen-Steinmetz, C., Baumeister, S., ... Laucht, M. (2017). Ventral striatum and amygdala activity as convergence sites for early adversity and conduct disorder. *Social Cognitive and Affective Neuroscience*, 12(2), 261–272. <https://doi.org/10.1093/scan/nsw120>

Hostinar, C. E., & Miller, G. E. (2019). Protective factors for youth confronting economic hardship: Current challenges and future avenues in resilience research. *American Psychologist*, 74(6), 641–652. <https://doi.org/10.1037/amp0000520>

Jaffee, S. R. (2017). Child Maltreatment and Risk for Psychopathology in Childhood and Adulthood. *Annual Review of Clinical Psychology*, 13, 525–551. <https://doi.org/10.1146/annurev-clinpsy-032816-045005>

Jenkinson, M., Beckmann, C. F., Behrens, T. E. J., Woolrich, M. W., & Smith, S. M. (2012). FSL. *NeuroImage*, 62(2), 782–790. <https://doi.org/10.1016/j.neuroimage.2011.09.015>

Jovicich, J., Czanner, S., Greve, D., Haley, E., van der Kouwe, A., Gollub, R., ... Dale, A. (2006). Reliability in multi-site structural MRI studies: Effects of gradient non-linearity correction on phantom and human data. *NeuroImage*, 30(2), 436–443. <https://doi.org/10.1016/j.neuroimage.2005.09.046>

Ecological stress, amygdala reactivity, and internalizing problems

Kachmar, A. G., Connolly, C. A., Wolf, S., & Curley, M. A. Q. (2019). Socioeconomic Status in Pediatric Health Research: A Scoping Review. *The Journal of Pediatrics*, 213, 163–170.

<https://doi.org/10.1016/j.jpeds.2019.06.005>

Kessler, R. C., Avenevoli, S., Costello, E. J., Georgiades, K., Green, J. G., Gruber, M. J., ...

Merikangas, K. R. (2012). Prevalence, persistence, and sociodemographic correlates of DSM-IV disorders in the National Comorbidity Survey Replication Adolescent Supplement. *Archives of General Psychiatry*, 69(4), 372–380.

<https://doi.org/10.1001/archgenpsychiatry.2011.160>

Kim, P., Evans, G. W., Angstadt, M., Ho, S. S., Sripada, C. S., Swain, J. E., ... Phan, K. L.

(2013). Effects of childhood poverty and chronic stress on emotion regulatory brain function in adulthood. *Proceedings of the National Academy of Sciences of the United States of America*, 110(46), 18442–18447. <https://doi.org/10.1073/pnas.1308240110>

Kind, A. J. H., & Buckingham, W. R. (2018). Making Neighborhood-Disadvantage Metrics Accessible—The Neighborhood Atlas. *The New England Journal of Medicine*, 378(26), 2456–2458. <https://doi.org/10.1056/NEJMp1802313>

Kline, R. B. (2015). *Principles and Practice of Structural Equation Modeling, Fourth Edition*. Guilford Publications.

Leventhal, T., & Brooks-Gunn, J. (2011). Changes in neighborhood poverty from 1990 to 2000 and youth's problem behaviors. *Developmental Psychology*, 47(6), 1680–1698.

<https://doi.org/10.1037/a0025314>

Lillard, A. S., & Erisir, A. (2011). Old Dogs Learning New Tricks: Neuroplasticity Beyond the Juvenile Period. *Developmental Review: DR*, 31(4), 207–239.

<https://doi.org/10.1016/j.dr.2011.07.008>

Ecological stress, amygdala reactivity, and internalizing problems

Loukas, A., Prelow, H. M., Suizzo, M.-A., & Allua, S. (2008). Mothering and Peer Associations

Mediate Cumulative Risk Effects for Latino Youth. *Journal of Marriage and Family*,

70(1), 76–85. <https://doi.org/10.1111/j.1741-3737.2007.00462.x>

Lynch, M., & Cicchetti, D. (1998). An ecological-transactional analysis of children and contexts:

The longitudinal interplay among child maltreatment, community violence, and

children's symptomatology. *Development and Psychopathology*, 10(2), 235–257.

Marek, S., Tervo-Clemmens, B., Calabro, F. J., Montez, D. F., Kay, B. P., Hatoum, A. S.,

Donohue, M. R., Foran, W., Miller, R. L., Feczko, E., Miranda-Dominguez, O., Graham,

A. M., Earl, E. A., Perrone, A. J., Cordova, M., Doyle, O., Moore, L. A., Conan, G.,

Uriarte, J., ... Dosenbach, N. U. F. (2020). Towards Reproducible Brain-Wide

Association Studies. *BioRxiv*, 2020.08.21.257758.

<https://doi.org/10.1101/2020.08.21.257758>

Marshall, N. A., Marusak, H. A., Sala-Hamrick, K. J., Crespo, L. M., Rabinak, C. A., &

Thomason, M. E. (2018). Socioeconomic disadvantage and altered corticostriatal

circuitry in urban youth. *Human Brain Mapping*, 39(5), 1982–1994.

<https://doi.org/10.1002/hbm.23978>

McEwen, B. S. (2012). Brain on stress: How the social environment gets under the skin.

Proceedings of the National Academy of Sciences, 109(Supplement 2), 17180–17185.

<https://doi.org/10.1073/pnas.1121254109>

McEwen, B. S., & Gianaros, P. J. (2011). Stress- and Allostasis-Induced Brain Plasticity. *Annual*

Review of Medicine, 62(1), 431–445. <https://doi.org/10.1146/annurev-med-052209->

100430

Ecological stress, amygdala reactivity, and internalizing problems

McEwen, B. S., Gray, J., & Nasca, C. (2015). Recognizing Resilience: Learning from the Effects of Stress on the Brain. *Neurobiology of Stress*, 1, 1–11.

<https://doi.org/10.1016/j.ynstr.2014.09.001>

McLaughlin, K. A., Sheridan, M. A., & Lambert, H. K. (2014). Childhood Adversity and Neural Development: Deprivation and Threat as Distinct Dimensions of Early Experience.

Neuroscience and Biobehavioral Reviews, 47, 578–591.

<https://doi.org/10.1016/j.neubiorev.2014.10.012>

McLaughlin, K. A., Sheridan, M. A., & Nelson, C. A. (2017). Neglect as a Violation of Species-Expectant Experience: Neurodevelopmental Consequences. *Biological Psychiatry*, 82(7), 462–471. <https://doi.org/10.1016/j.biopsych.2017.02.1096>

Merikangas, K. R., He, J., Burstein, M., Swanson, S. A., Avenevoli, S., Cui, L., ... Swendsen, J. (2010). Lifetime Prevalence of Mental Disorders in US Adolescents: Results from the National Comorbidity Study-Adolescent Supplement (NCS-A). *Journal of the American Academy of Child and Adolescent Psychiatry*, 49(10), 980–989.

<https://doi.org/10.1016/j.jaac.2010.05.017>

Mujahid, M. S., Diez Roux, A. V., Morenoff, J. D., & Raghunathan, T. (2007). Assessing the measurement properties of neighborhood scales: From psychometrics to econometrics.

American Journal of Epidemiology, 165(8), 858–867.

<https://doi.org/10.1093/aje/kwm040>

Muthén, L. K., & Muthén, B. O. (2012). *Mplus: Statistical Analysis with Latent Variables – User’s Guide*.

Ecological stress, amygdala reactivity, and internalizing problems

Noble, K. G., Houston, S. M., Brito, N. H., Bartsch, H., Kan, E., Kuperman, J. M., ... Sowell, E.

R. (2015). Family income, parental education and brain structure in children and adolescents. *Nature Neuroscience*, 18(5), 773–778. <https://doi.org/10.1038/nn.3983>

Odgers, C. L., Caspi, A., Russell, M. A., Sampson, R. J., Arseneault, L., & Moffitt, T. E. (2012).

Supportive parenting mediates neighborhood socioeconomic disparities in children's antisocial behavior from ages 5 to 12. *Development and Psychopathology*, 24(3), 705–721. <https://doi.org/10.1017/S0954579412000326>

Owens, M. M., Potter, A., Hyatt, C., Albaugh, M., Thompson, W. K., Jernigan, T., Yuan, D.,

Hahn, S., Allgaier, N., & Garavan, H. (2020). Recalibrating expectations about effect size: A multi-method survey of effect sizes in the ABCD study. PsyArXiv.

<https://doi.org/10.31234/osf.io/tn9u4>

Pine, D. S., Cohen, P., Gurley, D., Brook, J., & Ma, Y. (1998). The risk for early-adulthood

anxiety and depressive disorders in adolescents with anxiety and depressive disorders.

Archives of General Psychiatry, 55(1), 56–64. <https://doi.org/10.1001/archpsyc.55.1.56>

Power, J. D., Mitra, A., Laumann, T. O., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E.

(2014). Methods to detect, characterize, and remove motion artifact in resting state fMRI.

NeuroImage, 84, 320–341. <https://doi.org/10.1016/j.neuroimage.2013.08.048>

Price, C. J., & Friston, K. J. (2005). Functional ontologies for cognition: The systematic

definition of structure and function. *Cognitive Neuropsychology*, 22(3), 262–275.

<https://doi.org/10.1080/02643290442000095>

Salk, R. H., Hyde, J. S., & Abramson, L. Y. (2017). Gender differences in depression in

representative national samples: Meta-analyses of diagnoses and symptoms.

Psychological Bulletin, 143(8), 783–822. <https://doi.org/10.1037/bul0000102>

Ecological stress, amygdala reactivity, and internalizing problems

Schaefer, E. S. (1965). Children's reports of parental behavior: An inventory. *Child Development*, 36(2), 413–424. <https://doi.org/10.2307/1126465>

Schneider, W., Waldfogel, J., & Brooks-Gunn, J. (2015). The Great Recession and Behavior Problems in 9-Year Old Children. *Developmental Psychology*, 51(11), 1615–1629. <https://doi.org/10.1037/dev0000038>

Siegel, J. S., Power, J. D., Dubis, J. W., Vogel, A. C., Church, J. A., Schlaggar, B. L., & Petersen, S. E. (2014). Statistical improvements in functional magnetic resonance imaging analyses produced by censoring high-motion data points. *Human Brain Mapping*, 35(5), 1981–1996. <https://doi.org/10.1002/hbm.22307>

Silvers, J. A., Insel, C., Powers, A., Franz, P., Helion, C., Martin, R., Weber, J., Mischel, W., Casey, B. J., & Ochsner, K. N. (2017). The transition from childhood to adolescence is marked by a general decrease in amygdala reactivity and an affect-specific ventral-to-dorsal shift in medial prefrontal recruitment. *Developmental Cognitive Neuroscience*, 25, 128–137. <https://doi.org/10.1016/j.dcn.2016.06.005>

Simmons, J. P., Nelson, L. D., & Simonsohn, U. (2011). False-positive psychology: Undisclosed flexibility in data collection and analysis allows presenting anything as significant. *Psychological Science*, 22(11), 1359–1366. <https://doi.org/10.1177/0956797611417632>

Smith, K. E., & Pollak, S. D. (2020). Rethinking Concepts and Categories for Understanding the Neurodevelopmental Effects of Childhood Adversity. *Perspectives on Psychological Science*, 1745691620920725. <https://doi.org/10.1177/1745691620920725>

Smith, S. M., Jenkinson, M., Woolrich, M. W., Beckmann, C. F., Behrens, T. E. J., Johansen-Berg, H., ... Matthews, P. M. (2004). Advances in functional and structural MR image

Ecological stress, amygdala reactivity, and internalizing problems

analysis and implementation as FSL. *NeuroImage*, 23 Suppl 1, S208-219.

<https://doi.org/10.1016/j.neuroimage.2004.07.051>

Somerville, L. H., Fani, N., & McClure-Tone, E. B. (2011). Behavioral and neural representation of emotional facial expressions across the lifespan. *Developmental Neuropsychology*, 36(4), 408–428. <https://doi.org/10.1080/87565641.2010.549865>

Spear, L. P. (2000). The adolescent brain and age-related behavioral manifestations.

Neuroscience and Biobehavioral Reviews, 24(4), 417–463.

Swartz, J. R., Knodt, A. R., Radtke, S. R., & Hariri, A. R. (2015). A neural biomarker of psychological vulnerability to future life stress. *Neuron*, 85(3), 505–511.

<https://doi.org/10.1016/j.neuron.2014.12.055>

Szucs, D., & Ioannidis, J. P. A. (2017). Empirical assessment of published effect sizes and power in the recent cognitive neuroscience and psychology literature. *PLOS Biology*, 15(3), e2000797. <https://doi.org/10.1371/journal.pbio.2000797>

Szucs, D., & Ioannidis, J. PA. (2020). Sample size evolution in neuroimaging research: An evaluation of highly-cited studies (1990–2012) and of latest practices (2017–2018) in high-impact journals. *NeuroImage*, 221, 117164. <https://doi.org/10.1016/j.neuroimage.2020.117164>

Taylor, R. L., Cooper, S. R., Jackson, J. J., & Barch, D. M. (2020). Assessment of Neighborhood Poverty, Cognitive Function, and Prefrontal and Hippocampal Volumes in Children. *JAMA Network Open*, 3(11). <https://doi.org/10.1001/jamanetworkopen.2020.23774>

Telzer, E. H., & Fuligni, A. J. (2013). Positive daily family interactions eliminate gender differences in internalizing symptoms among adolescents. *Journal of Youth and Adolescence*, 42(10), 1498–1511. <https://doi.org/10.1007/s10964-013-9964-y>

Ecological stress, amygdala reactivity, and internalizing problems

Tottenham, N., & Galván, A. (2016). Stress and the adolescent brain: Amygdala-prefrontal cortex circuitry and ventral striatum as developmental targets. *Neuroscience and Biobehavioral Reviews*, 70, 217–227. <https://doi.org/10.1016/j.neubiorev.2016.07.030>

Tottenham, N., Tanaka, J. W., Leon, A. C., McCarry, T., Nurse, M., Hare, T. A., ... Nelson, C. (2009). The NimStim set of facial expressions: Judgments from untrained research participants. *Psychiatry Research*, 168(3), 242–249. <https://doi.org/10.1016/j.psychres.2008.05.006>

Vargas, T., Damme, K. S. F., & Mittal, V. A. (2020). Neighborhood deprivation, prefrontal morphology and neurocognition in late childhood to early adolescence. *NeuroImage*, 220, 117086. <https://doi.org/10.1016/j.neuroimage.2020.117086>

Vidal-Ribas, P., Benson, B., Vitale, A. D., Keren, H., Harrewijn, A., Fox, N. A., ... Stringaris, A. (2019). Bidirectional Associations Between Stress and Reward Processing in Children and Adolescents: A Longitudinal Neuroimaging Study. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, 4(10), 893–901. <https://doi.org/10.1016/j.bpsc.2019.05.012>

Whittle, S., Simmons, J. G., Dennison, M., Vijayakumar, N., Schwartz, O., Yap, M. B. H., ... Allen, N. B. (2014). Positive parenting predicts the development of adolescent brain structure: A longitudinal study. *Developmental Cognitive Neuroscience*, 8, 7–17. <https://doi.org/10.1016/j.dcn.2013.10.006>

Whittle, S., Vijayakumar, N., Simmons, J. G., Dennison, M., Schwartz, O., Pantelis, C., ... Allen, N. B. (2017). Role of Positive Parenting in the Association Between Neighborhood Social Disadvantage and Brain Development Across Adolescence. *JAMA Psychiatry*, 74(8), 824–832. <https://doi.org/10.1001/jamapsychiatry.2017.1558>

Ecological stress, amygdala reactivity, and internalizing problems

Wittchen, H. U., Nelson, C. B., & Lachner, G. (1998). Prevalence of mental disorders and psychosocial impairments in adolescents and young adults. *Psychological Medicine*,

28(1), 109–126. <https://doi.org/10.1017/s0033291797005928>

Zeanah, C. H., Egger, H. L., Smyke, A. T., Nelson, C. A., Fox, N. A., Marshall, P. J., & Guthrie, D. (2009). Institutional rearing and psychiatric disorders in Romanian preschool children. *The American Journal of Psychiatry*, 166(7), 777–785.

<https://doi.org/10.1176/appi.ajp.2009.08091438>