On the relationship of arousal and attentional distraction by emotional novel sounds

Authors

Carolina Bonmassar¹⁾, Florian Scharf²⁾, Andreas Widmann¹⁾³⁾, Nicole Wetzel¹⁾⁴⁾⁵⁾

Affiliations

¹⁾Leibniz Institute for Neurobiology, Magdeburg, Germany

²⁾University of Kassel, Germany

³⁾Leipzig University, Germany

⁴⁾Center for Behavioral Brain Sciences Magdeburg, Germany

⁵⁾University of Applied Sciences Magdeburg-Stendal, Germany

Short Title

Distraction effects and PDRs to emotional sounds

Corresponding Author

Carolina Bonmassar <https://orcid.org/0000-0002-8412-6502>, carolina.bonmassar@lin-magdeburg.de, +49-391-6263-93431, Brenneckestraße 6, 39118 Magdeburg

Abstract

Unexpected and task-irrelevant sounds can impair performance in a task. It has been shown that highly arousing emotional distractor sounds impaired performance less compared to moderately arousing neutral distractor sounds. The present study tests whether these differential emotion-related distraction effects are directly related to an enhancement of arousal evoked by processing of emotional distractor sounds.

We disentangled costs of orienting of attention and benefits of increased arousal levels during the presentation of highly arousing emotional and moderately arousing neutral novel sounds that were embedded in a sequence of repeated standard sounds. We used sound-related pupil dilation responses as a marker of arousal and RTs as a marker of distraction in a visual categorization task in 57 healthy young adults. Multilevel analyses revealed increased RT and increased pupil dilation in response to novel vs. standard sounds. Emotional novel sounds reduced distraction effects on the behavioral level and increased pupil dilation responses compared to neutral novel sounds. Bayes Factors revealed strong evidence against an inverse proportional relationship between behavioral distraction effects and sound-related pupil dilation responses for emotional sounds. Given that the activity of the locus coeruleus has been linked to both changes in pupil diameter and arousal, it may embody an indirect relationship as a common antecedent by the release of norepinephrine into brain networks involved in attention control and control of the pupil. The present study provides new insights into the relation of changes in arousal and attentional distraction during the processing of emotional task-irrelevant novel sounds.

Keywords: attention, emotion, novel, arousal, pupil, Locus coeruleus, norepinephrine, distraction effects

47

48 **1.Introduction**

49 A sudden cry can capture our attention, impair performance in a task at hand and increase the
50 level of arousal to prepare us for a fight or flight reaction. The orienting of attention and the
51 increase in arousal are two aspects of the orienting response, reflecting costs of attention
52 distraction and benefits of arousal increase (Näätänen, 1992; Sokolov, 1963). The present
53 study investigates the direct relation of distraction costs and arousal benefits by the co-
54 registration of performance and pupil size in a well-established distraction paradigm.

55

56 The involuntary capture of attention by unexpected stimuli occurring outside the current focus
57 of attention enables the detection of potentially relevant events in the environment (e.g., a
58 ringing smartphone). The involved orienting and evaluation mechanisms include capacity
59 limited processes (Näätänen, 1992) that can result in impaired performance in a task
60 (distraction effect, e.g., Escera et al., 1998; Schröger & Wolff, 1998). The underlying
61 mechanisms have been described by a three-stage model of involuntary attention (e.g., Escera
62 & Corral, 2007; Schröger, 1997). In the first stage a predictive model of the acoustic
63 environment is created automatically. Unexpected sounds (e.g., new sounds) violate the
64 prediction (Winkler et al., 2009; Winkler & Schröger, 2015; but see, May & Tiitinen, 2010).
65 This can trigger an orienting of attention and further evaluation of the unexpected sound. If no
66 adaptation of behavior is required, attention is reoriented to the task at hand. When applying
67 an oddball paradigm including frequently repeated standard sounds and rare, randomly
68 presented distractor or oddball sounds (also termed novel or deviant sounds), distractor
69 sounds frequently cause prolonged reaction times (RTs) in a task not related to the sound
70 sequence or to the deviant feature (auditory task, Berti et al., 2004; Horváth et al., 2008;
71 Muller-Gass & Schröger, 2007; visual task, Escera et al., 1998; Parmentier et al., 2008).

72

Nonetheless, some studies reported reduced distraction effects or even improved performance when task-irrelevant emotional information was presented (Lindström & Bohlin, 2011; Lorenzino & Caudek, 2015; Max et al., 2015). These reduced distraction or facilitation effects have been thought to be caused by an increased level of arousal caused by the emotional content of the distractor event (Max et al., 2015). This explanation is in line with the models by Näätänen (1992) and Sokolov (1963) postulating that the orienting response includes costs of orienting and benefits of enhanced arousal (see also Hoyer et al., 2021). However, a direct relation between distraction effects and distractor-related changes in the arousal level was not yet evidenced in the context of emotional and novel sounds and will be investigated in the present study. Nonetheless, previous studies in the visual domain investigated this relationship already by means of emotional attentional blink tasks (McHugo et al., 2013) and demonstrated that attention capture by distractors are modulated by state levels of arousal, in particular highly emotional stimuli facilitated the task performance (e.g., threat of shock, (Kim & Anderson, 2020; Kim et al., 2021; Lee et al., 2012; Sutherland & Mather, 2015).

We used changes in pupil diameter as a marker of novel and emotional sound-related increase in arousal. Several studies confirmed that novel oddball sounds (Bonmassar et al., 2020; Liao, Kidani, et al., 2016; Liao, Yoneya, et al., 2016; Wetzel et al., 2016; Widmann et al., 2018) and emotional events (Bradley et al., 2008, 2017; Hess & Polt, 1960) cause a transient dilation of the pupil. In recent oddball studies, emotionally negative novel sounds, that were interspersed in a sequence of repeated standard sounds, evoked stronger pupil dilation than emotionally neutral novel sounds (Bonmassar et al., 2020; Widmann et al., 2018). Additional evidence for emotional arousal being linked to an increase in pupil dilation is the simultaneous use of skin conductance responses and heart rate as a marker of emotional arousal. Recent studies observed a simultaneous increase in skin conductance responses and

heart rate as well as pupil dilation responses to emotional stimuli (Bradley et al., 2008, 2017; Wang et al., 2018).

In sum, cumulative evidence supports a model postulating that distraction effects include costs of orienting and benefits of arousal. To test this hypothesis, we applied a well-established auditory-visual oddball paradigm, including emotional highly arousing and neutral moderately arousing environmental novel sounds while participants focused on a visual categorization task. We expected first task-irrelevant novel sounds to prolong RTs compared to standard sounds (distraction effect; Escera, 1998; Schröger & Wolff, 1998) and increase PDRs (Murphy et al., 2011; Widmann et al., 2018). Second, we expected reduced distraction effects in response to emotional novel sounds compared to neutral novel sounds (Max et al., 2015) but increased amplitudes of the PDRs to emotional vs. neutral novel sounds (Bonmassar et al., 2020). Third, we hypothesized a direct relationship between emotion-related distraction effects and emotion-related PDRs. That is, we expected concomitant shorter RTs and higher PDRs in trials with emotional but not for neutral novel sounds. Importantly, the relationship between RT and PDR needs to be analyzed at both within- and between-participant level to disentangle effects of the average PDR (e.g., participants with higher average PDR show smaller distraction effects) from effects at single trial level (e.g., smaller distraction effects occur in *trials* with larger PDR). This was achieved by means of adequate centering strategies in a linear mixed-effects models (for an example of different levels of analysis see LoTemplio et al., 2021). That is, we expected shorter RTs in trials with higher PDR for emotional but not for neutral novel sounds. Moreover, the present experimental approach includes ethically harmless stimuli and has no potential to cause anxiety, fear or threat. Thus, this paradigm can be applied in developmental studies even with young children as well as with patients, enabling insights in the relation between attentional distraction and arousal in developmental and clinical samples.

2. Method

2.1 Participants

61 participants took part in the experiment. Four participants were excluded for the following reasons: pupil data available from one eye only, reaction times deviating more than two standard deviations from the average (two participants) and an accidental double participation in the experiment. The data of 57 healthy adults ($M_{\text{age}}=25$ years, range 18-36, 31 females, 5 left-handed) were used in the study. Participation was rewarded by money (10€/hour). All participants gave written informed consent. Participants confirmed a normal or corrected-to-normal vision, normal hearing, no medication with effects on the nervous system, and no history of attention-related disorders. Handedness was measured with an abbreviated German version of the Oldfield Handedness Inventory (Oldfield, 1971). The project was approved by the local ethics committee.

2.2 Stimuli

Auditory stimuli. A total of 48 environmental novel sounds¹ were collected from the database of a previous study (Max et al., 2015). Max and colleagues selected a set of 210 auditory stimuli, collected from the International Affective Digitized Sounds study (IADS, Bradley & Lang, 2007), and from other data bases as described by Max et al. (2015).

In the present study, sounds were allocated to three categories: 24 highly arousing emotionally negative sounds (for example an ambulance siren), 24 moderately arousing neutral sounds (for example toasting glasses), and 3 moderately arousing neutral sounds used as standard sounds (for example a musical instrument). Descriptive statistic and independent samples *t*-test are reported in Table 1 and Table 2. Sounds had a duration of 500 ms including faded ends of 5 ms. They were presented at a loudness of 54.5 dB SPL (measured with PAA3

PHONIC Handheld audio analyzer, Phonic Corporation, Taipei, Taiwan). Loudness of sounds was equalized with root mean square normalization.

Visual stimuli. Three different target categories were presented in separate blocks: (a) princesses vs. knights (Fig.1), (b) cats vs. hens, (c) butterflies vs. fish. For each target figure two versions were presented (slightly differing in shape, color and direction). All versions of the target figures were presented with equal probability (e.g. 25% princess with a pink dress, 25% princess with a blue dress, 25% knights with gray armor, 25% knight with blue armor) in a pseudorandomized order, that is, we implemented some constrictions to avoid biases which would appear in a complete randomized order. For example, the first two trials of each block were standard trials and we implemented two standard trials following a novel sound. For each target category, a different scene was used as a background. Princesses and knights were presented in front of a palace (left side) and a fortress (right side), cats and hens were presented in front of a basket (left side) and a hen-roost (right side), and butterflies and fishes were presented with a flowering shrub (left side) and a pond (right side). The background landscapes' pictures were displayed at the center of a screen with a size of 960 x 720 px, 267 x 200 mm, (24.3° x 18.3° visual angle from a viewing distance of 620 mm). Picture mean luminance without targets or feedback was 51.2 cd/m² on a grey background screen with a mean luminance of 2.9 cd/m² (princess/knights 50.3 cd/m², cat/hen 55.1 cd/m², butterfly/fish 48.2 cd/m²). The different versions of the targets and background scenes were presented to apply exactly the same paradigm to children in a future study (not reported here).

Apparatus and Software. The auditory stimuli were presented via loudspeakers (Bose Companion 2 series III Multimedia speaker system) located at the left and the right of the screen. The visual stimuli were presented on a VIEWPixx/EEG display (VPixx Technologies Inc.) with a resolution of 1920 x 1080 (23,6-inch diagonal display size) and a refresh rate of 120 Hz. Responses to the target were given pressing a button on a response box (RTbox)

located in the front of the screen (Li et al., 2010). The experimental stimulation was presented via Psychtoolbox (Version 3.0.15, Kleiner et al., 2007) using Octave (Linux, Version 4.0.0).

Table 1

Descriptive statistics of the categories of sounds used in the experiment.

Sound	Number	Valence		Arousal	
		Mean	SD	Mean	SD
Standard	3	5.66	0.21	4.43	0.40
Emotional	24	2.59	0.57	6.68	0.45
Neutral	24	5.21	0.46	4.81	0.37

Note. Sounds had been rated on a 9-point scale for valence (1 = unpleasant — 5 = neutral — 9 = pleasant) and arousal (1 = calm — 9 = arousing).

Table 2

Independent t-tests of the categories of sounds used in the experiment.

Condition	Valence					Arousal				
	<i>t</i>	<i>df</i>	<i>p</i>	<i>d</i>	<i>BF</i> ₁₀	<i>t</i>	<i>df</i>	<i>p</i>	<i>d</i>	<i>BF</i> ₁₀
emotional vs. neutral	-17.36	46	< .001	-5.012	2.845×10^{18}	15.56	46	< .001	4.493	4.358×10^{16}
emotional vs. standard	-9.003	25	< .001	-5.513	1.747×10^6	8.103	25	< .001	4.962	282390.711
neutral vs. standard	-1.666	25	.108	-1.020	1.098	1.631	25	.115	0.999	1.061

Note. Statistically significant results are marked in bold.

186

187 Fig.1. Trial structure. In every trial, a sound was presented for 500 ms. 100 ms after sound
 188 offset, each sound was followed by the target (e.g. princess). The target was presented for 500

ms. Participants were instructed to press the left button when a princess appeared and the right button when a knight appeared. The response time window was 2,000 ms after target onset. Correct responses within the response time window were directly followed by a feedback motion, which consisted of two images with a duration of 150 and 450 ms (total feedback duration 600ms). The visual background was presented during the entire trial.

2.3 Procedure

The experiment was conducted in an acoustically attenuated and electromagnetically shielded cabin. Illuminance of the cabin was held constant at a level of 48.9 lx (measured with MAVOLUX 5032B USB, GOSSEN Foto- and Lichtmesstechnik GmbH, Nürnberg, Germany). Participants sat in front of a screen, having their right and left index finger on the RTbox buttons. Each experimental block started with a five-point eye-tracker calibration and validation procedure.

Task and Feedback. Participants were instructed to press the left button when a princess (or cat or butterfly) appeared on the screen and the right button when a knight (or hen or fish) appeared (see Fig.1). They were asked to respond to the target stimuli as fast and correctly as possible and to ignore the sounds. Correct responses were followed by feedback, that is, the target moved towards the left or right side. For example, the princess moved to the palace on the left side and the knight moved to the fortress on the right side (Fig. 1). The feedback motion consisted of two images with a duration of 150 and 450 ms.

Trial and block structure. Each sound was presented with a fixed stimulus onset asynchrony (SOA) of 3300 ms (Fig. 1). 100 ms after sound offset, each sound was followed by a visual target. The target was presented for 500 ms. After target onset, participants had a 2 s time window to respond. The feedback was presented with a duration of 600 ms directly after the response, but not earlier than 200 ms after target offset. A total of six blocks were presented, each consisting of 40 trials. Two blocks included princesses and knights as target figure, two

blocks of cats and hens and two blocks of butterflies and fish. The order of blocks containing different scenes were balanced across participants. Blocks containing the same scene were always presented one after another. Each block lasted about 2 minutes.

Sound sequence. The sound sequence included standard sounds (80%), emotional (10%) and neutral (10%) novel sounds. These probabilities of sound type presentation were equal over each block. That is, each of the six blocks included 32 standard sounds, 4 emotional novel sounds and 4 neutral novel sounds. In total, 192 standard sounds, 24 emotional novel sounds and 24 neutral novel sounds were presented. The sound sequence was unique for each participant. This ensured that potential changes in brightness were *not* systematically related to the occurrence of different sound types. For each scene (princesses vs. knights, cats vs. hens, butterflies vs. fish) a different standard sound was presented. This prevented potential effects of specific stimulus features of a single standard sound on performance. The assignment of standard sounds to the scene was counter-balanced across participants. The sound sequence was pseudo-randomized so that each novel was preceded by at least two standard sounds. Each novel was presented only once in total.

Training blocks. To familiarize participants with each of the three different scenes in the experimental block, three short training blocks including 8 trials each (6 standard sounds, 1 emotional and 1 neutral novel sounds) were performed. Sounds presented in the training blocks were not presented in the experimental blocks. If more than 50% of the trials was answered incorrectly, the training was repeated. Because the experiment was designed to be suitable for children, all participants of the present study understood the task promptly and no repetition was needed.

2.4 Data Analysis

The first two standard trials per block are required for the formation of a predictive model of the upcoming stimuli (Bendixen et al., 2007). Because the two standard trials immediately

following an oddball sound can be affected by previous distractor sound processing (Wetzel, 2015), these were removed from all analyses. Only corresponding identical trials from the behavioral and pupil data, including a correct response, were used for analysis. Trials with incorrect or missing responses were excluded from pupil data analysis and trials with missing pupil data or blinks which could not be interpolated (see below) were also excluded from RT analysis.

2.5 Pupil data processing

The pupil diameter of both eyes was recorded with an infrared EyeLink Portable Duo eye-tracker (SR Research Ltd., Mississauga, Ontario, Canada). The eye tracking was set up in remote mode at a sampling rate of 500 Hz.

The eye-tracker automatically reports the number of pixels below a specific threshold as belonging to the pupil (in case area is recorded or in case diameter is recorded, as here, a transformation of area to diameter by: $(256 * \sqrt{\text{area in pixel} \div \Pi})$). By maintaining constant distance between the participant and the eye-tracker, the number of pixels actually reflects a meaningful and valid physical unit which can be converted to other meaningful units by simple linear transformations (e.g., mm; as described in several publications, for example Hayes & Petrov, 2016; Klingner et al., 2008). We converted the eye tracker pupil diameter digital counts to mm as suggested by Steinhauer and colleagues (2022). Pupil size analysis was implemented with MATLAB software. Eye saccade and blink information were provided by the eye tracker. Partial blinks were detected during post-processing from the smoothed velocity times series by an additional custom function, i.e., pupil diameter changes exceeding 20 mm/s including a 50 ms pre-blink and a 100 ms post-blink interval (Merritt et al., 1994). We applied Kret et al.'s (2019) dynamic offset algorithm to average data from both eyes. Isolated data segments between blinks or missing data shorter than 10 ms were considered as missing data. Subsequently, segments with blinks or missing data shorter than 1 s were

interpolated with linear interpolation, longer segments were removed from the continuous data. Data were segmented in epochs of 2 s of duration (including a -0.2 to 0 s pre-stimulus interval), baseline corrected by subtracting the mean amplitude of the baseline period (-0.2 to 0.2 s) from each epoch. Typically, the pupil is not able to contract or dilate any earlier than 200 ms after stimulus onset (Mathôt et al., 2018). Thus, baseline correction was extended to range from -0.2 to 0.2 s, which allows for a wider span of baseline activity. The mean PDRs were computed in a time window around the peak between 1.3 and 1.5 s for each trial and each participant. In addition, for each trial, the average pupil size in the baseline period was computed.

2.6 Behavioral data (reaction times, RTs)

Incorrect responses, responses faster than 100 ms after target onset and missing responses (or responses given later than 2 s after target onset) were excluded from RT and pupil analysis. Participants deviating more than 2 standard deviations from the average reaction times were excluded from the analysis. A total mean of 132 trials per participant (SD= 5.19; max= 140; min= 109) were used for analysis.

2.7 Statistical analysis

Analysis of Condition Effects

A paired samples t-test and Bayesian paired samples t-test were performed to compare PDR amplitudes in response to standard sounds, emotional novel and neutral novel sounds in the selected analysis mean amplitude time window (1.3-1.5 s). The same analysis was performed for the RTs in trials including standard sounds, emotional novel, and neutral novel sounds. All t-tests and Bayesian t-tests were performed using the R packages stats (v4.0.3, R Core Team, 2019) and BayesFactor (v0.9.12-4.2, R. D. Morey et al., 2011; Morey & Rouder, 2011; Rouder et al., 2009).

Analysis of Statistical Associations

We analyzed the relationship between RT and PDR both at trial and participant level with Linear mixed effect models (LMMs) to account for the dependencies between trials within participants. Trials were treated as primary unit of investigation (level 1) nested within participants (level 2). All models were estimated with the Maximum Likelihood method using the R packages lme4 (v1.1-27, Bates et al., 2015), and lmerTest (v3.1-3, Kuznetsova et al., 2017). As measures of goodness-of-fit model, we computed marginal and conditional R^2 (Nakagawa & Schielzeth, 2013), that is, the proportion of the total variability explained by the fixed effects and by all fixed and random effects together, respectively. Please note that relatively low values for R^2 are not uncommon due to the considerable variability of RTs between trials. Degrees of freedom for statistical tests were approximated using Satterthwaite's approximation. Bayes Factors were approximated from differences in the Bayesian Information Criterion, that is, $BF_{01} = \exp\left(\frac{BIC_0 - BIC_1}{2}\right)$ (Raftery, 1995). Specifically, we followed the logic of a "Type III" analysis of variance and computed the Bayes Factors from the comparison of the full model versus the full model excluding the respective effect.

We explored a range of conceivable models in which *RT* was modeled as a function of the various candidate predictors. To systemize the search for the best model, we applied a best subset selection and selected the best-fitting model using the Bayesian information criterion (BIC; Burnham & Anderson, 2004; Schwarz, 1978, Table 3). The set of candidate predictors contained various predictors at trial- and participant-level. Following from the experimental design, we always included Novelty (Standard vs. Distractor) and Emotionality (Neutral novel sound vs. Emotional novel sound) of the presented sounds as predictors. We applied a

contrast coding such that the coefficient of Novelty (0 for standard, 1 for novels *irrespective* of the emotional content) is an estimate of the predicted difference in RT between standards and neutral novels whereas the coefficient of Emotionality (0 for standard sounds and neutral novels, 1 for emotional novels) reflects the predicted difference in RT between emotional and neutral novels. We included a random intercept (i.e., varying average RT) and a random slope for the predictor Novelty across participants (i.e., varying distraction effects). A random slope for Emotionality was not supported by the data and resulted in a singular fit².

We considered several potential relationships between pupil diameter and RTs: Both pupil diameter within the baseline period (baseline PD) and during the pupil dilation response (PDR) were used as potential predictors (LoTemplio et al., 2021; Murphy et al., 2011). The baseline PD was included in the selection process of the best model only to improve the model estimates (Alday, 2019) by controlling for potential confounding due to differences in baseline PD. This approach is comparable to an ANCOVA approach where covariates are included – although they are not of substantive interest – to control for confounding. We did not interpret the resulting baseline effects as this would have gone beyond the scope of the manuscript (however, for possible interpretation see Supplementary material).

We considered that baseline PD and PDR can vary from trial to trial but there may also be systematic differences between participants and both these sources of variation could affect response times differentially. To give an intuition why this can happen: When the raw PDR takes a “large” value (e.g., relative to the grand mean) it remains unclear what “large” exactly implies, because large values could be due to the respective participant generally showing large PDRs or due to the specific trial showing a large PDR. If the raw baseline PD or PDR values were used as predictors, the trial and participant level effects of these predictors would be confounded and uninterpretable.

An established way to disentangle trial level from participant level variance is to create two variables: a trial level variable which is centered around the mean *within* participants and a participant level variable which represents the mean of each participant centered around the grand mean of all participants (Enders & Tofighi, 2007). The former variable represents the effect of fast fluctuations on trial level (e.g., do participants respond faster in trials with a larger PDR *relative* to the participant's individual average?). The later variable represents the effect of interindividual differences which are stable over the course of the experimental session (e.g., do participants with a generally larger PDR have larger behavioral distraction effects?). Both baseline PD and PDR were treated this way to separate the two sources of variation. We refer to the trial-level variables by the index "trial" (e.g., PDR_{trial}) and to the participant-level variables by the index "participant" (e.g., PDR_{participant}).

The least complex model under consideration contained Novelty, Emotionality, PDR_{trial} and Baseline_{trial} as simple effects. The most complex model could include Novelty, Emotionality, linear, and quadratic effects of PDR and baseline PDR both at trial level and participant³ level as well as their interactions with Novelty and Emotionality. Between these models, all possible alternative models were considered in the model space with two restrictions: (1) Any model containing a quadratic effect or interaction should also include the respective lower order ("simple") effect. (2) Any interaction including either Novelty or Emotionality should always be accompanied by an interaction with the respective other predictor because potential differences between the sound types are of genuine substantive interest to our study. All model effects specified based on the BIC selection are listed in Table 4. Except for a moderate skewness (2.29) in the level-1 residuals due to very slow responses in some single trials, all model assumptions were respected. We decided to keep these rare trials in the dataset because their removal would not have changed the results in any meaningful way given the large number of trials available for the model (7554)⁴.

369 Table 3

370 *Overview of the 10 best BIC-selected models.*

371

BIC model selection

	1 st best Model	2 nd best Model	3 rd best Model	4 th best Model	5 th best Model	6 th best Model	7 th best Model	8 th best Model	9 th best Model	10 th best Model
Intercept	X	X	X	X	X	X	X	X	X	X
Baseline _{trial}	X	X	X	X	X	X	X	X	X	X
(Baseline _{trial}) ²	X		X	X	X	X	X	X	X	X
Baseline _{participant}										X
(Baseline _{participant}) ²							X			
Novelty	X	X	X	X	X	X	X	X	X	X
Emotionality	X	X	X	X	X	X	X	X	X	X
PDR _{trial}	X	X	X	X	X	X	X	X	X	X
PDR _{participant}	X	X		X	X	X	X	X	X	X
Baseline _{trial} × Novelty	X	X		X	X	X	X	X	X	X
Baseline _{trial} × Emotionality										
Baseline _{participant} × Novelty										
Baseline _{participant} × Emotionality										
Novelty × (Baseline _{trial}) ²									X	
Novelty × (Baseline _{participant}) ²										
Novelty × PDR _{trial}				X						
Novelty × PDR _{participant}								X		
Emotionality × (Baseline _{trial}) ²					X					
Emotionality × (Baseline _{participant}) ²										
Emotionality × PDR _{trial}						X				
Emotionality × PDR _{participant}										
Df	12.00	11.00	11.00	13.00	13.00	13.00	13.00	13.00	13.00	13.00
loglik	-43851.65	-43858.42	-43858.55	-43849.99	-43850.11	-43850.19	-43850.34	-43850.69	-43851.20	-43851.25
BIC	87810.46	87815.06	87815.33	87816.08	87816.30	87816.46	87816.76	87817.47	87818.48	87818.59
delta		4.61	4.87	5.62	5.84	6.00	6.31	7.01	8.02	8.13

weight	0.62	0.06	0.05	0.04	0.03	0.03	0.03	0.02	0.01	0.01

Note. The parameters or predictors that are included in the model are marked with a “X”.

372

373

374

375 Table 4
 376 *Coefficients for the mixed-effects model.*
 RT-PDR-model

Dependent variable	Predictor/Fixed effect	Estimate	SE	df	t-value	P	BF	Results ^a description
Reaction times	Intercept	406.040	5.144	57.683	78.941	<0.001	–	mean (predicted) standard PDRs
	Novelty	22.859	2.963	109.884	7.714	<0.001	8.7 · 10⁷	novel RTs minus standard RTs
	Emotionality	-10.947	3.077	7444.833	-3.558	<0.001	6.4	emotional RTs minus neutral RTs
	Baseline _{trial}	-2.954	4.140	7465.697	-0.713	0.475	1	
	Baseline _{trial} × Novelty	35.670	8.335	7468.218	4.280	<0.001	107.9	<i>fastest</i> RTs to novel sounds when the baseline is <i>smaller</i> than average
	Baseline _{trial} × Emotionality	9.101	10.525	7492.214	0.865	0.387	0.02	
	(Baseline _{trial}) ²	17.239	4.609	7500.540	3.740	<0.001	12.4	<i>fastest</i> RTs when the baseline is <i>at average</i> within each participant
	PDR _{trial}	12.506	5.000	7477.175	2.501	0.013	2.6	at a trial level, <i>slower</i> RTs for trials with <i>larger</i> PDRs
	PDR _{participant}	-189.637	47.221	57.174	-4.016	<0.001	11.3	participants with <i>generally larger</i> PDRs showed <i>faster</i> RTs
Random effects		Variance	SD	Corr				
Participant	Intercept	1424.1	37.74					
	Novelty	153.4	12.39	0.37				
Residuals		6261.0	79.13					

Note. Statistically significant results are marked in bold.

RT = reaction time; SE= standard error; df = degree of freedom; SD = standard deviation; Corr = correlation; PDR = pupil dilation response; BF = Bayes Factor.

3. Results

PDR

Tests of the three a-priori hypotheses were conducted using Bonferroni adjusted alpha levels of .016 per test (.05/3). The analysis of the pupil dilation responses in the time window selected showed an effect of novel sounds (emotional vs. standard sounds: $t(56)= 11.2$, $p <.001$, $BF_{10}= 1.01 \times 10^{13}$; neutral vs. standard sounds: $t(56)= 7.90$, $p <.001$, $BF_{10}= 9.04 \times 10^7$, see Fig. 2 and Fig. 3, Panel B), resulting from larger amplitudes for novel sounds compared to standard sounds. An effect of emotional sounds (emotional vs. neutral sounds: $t(56)= 5.86$, $p <.001$, $BF_{10}= 5.77 \times 10^4$) resulted from larger amplitudes for emotional novel sounds compared to neutral novel sounds.

RTs

Tests of the three a-priori hypotheses were conducted using Bonferroni adjusted alpha levels of .016 per test (.05/3). The analysis of the reaction times showed an effect of novel sounds (emotional vs. standard sounds: $t(56)= 4.79$, $p <.001$, $BF_{10}= 1.5 \times 10^3$; neutral vs. standard sounds: $t(56)= 7.58$, $p <.001$, $BF_{10}= 2.8 \times 10^7$, see Fig. 3, Panel A), resulting from slower reaction times for novel sounds compared to standard sounds. An effect of emotional sounds (emotional vs. neutral sounds: $t(56)= -3.92$, $p <.001$, $BF_{10}= 98.24$) resulted from faster reaction times for emotional novel sounds compared to neutral novel sounds.

RT-PDR-model

The best-fitting BIC-selected model contained the predictors Novelty, Emotionality, $\text{Baseline}_{\text{trial}}^*$, $\text{Baseline}_{\text{trial}} \times \text{Novelty}$, $\text{Baseline}_{\text{trial}} \times \text{Emotionality}^*$, $(\text{Baseline}_{\text{trial}})^2$, $\text{PDR}_{\text{trial}}$ and $\text{PDR}_{\text{participant}}$. The effects of the predictors Novelty and Emotionality showed that participants responded significantly slower to novel than to standard sounds but significantly faster to

emotional novels compared to neutral novel sounds (Fig. 3, Panel A; Table 4, Novelty and Emotionality effects) – resembling the results of the confirmatory analyses above. In addition, we found dissociable relationships between PDR and RT at trial and participant level. At trial level, slower RTs were predicted for trials with larger PDR (Fig. 4, Panel A; Table 4, effect PDR_{trial}), but participants with generally larger PDRs tended to respond *faster* (Fig. 4, Panel B; Table 4, effect $PDR_{\text{participant}}$). The model also revealed effects of the baseline PD on RTs and distraction effects that are described in detail in the Supplement material.

With respect to our research questions, the existence of an interaction of pupil dilation and behavioral distraction effects (either at trial or participant level) was of major interest. Therefore, we computed additional Bayes Factors comparing the BIC-selected model with models in which we added such interactions. At trial level, there was strong evidence against the inclusion of the terms $PDR_{\text{trial}} \times \text{Novelty}$ and $PDR_{\text{trial}} \times \text{Emotionality}$ into the BIC-model ($BF = 0.002$). At participant level, there was strong evidence against the inclusion of the terms $PDR_{\text{participant}} \times \text{Novelty}$ and $PDR_{\text{participant}} \times \text{Emotionality}$ ($BF = 0.006$). That is, the model did not support an interaction of these factors neither on the trial level nor on the participants level.

Fig. 2. Grand-average pupil dilation responses (PDRs) for emotional novel sounds, neutral novel sounds, and standard sounds. Sound onset is at time point zero. Shading indicates the 95% confidence interval. The gray window indicates the time window used for analysis. Novel sounds evoked statistically significantly increased PDRs compared to standard sounds. Emotional novel sounds evoked statistically significantly increased PDRs compared to neutral novel sounds.

Fig. 3. Panel A: Mean reaction time (RT) for standard, neutral novel and emotional novel sounds. Novel sounds evoked increased RTs compared to standard sounds, demonstrating a distraction effect. Emotional novel sounds caused reduced RTs compared to neutral novel sounds, indicating a facilitation effect. Panel B: Mean pupil dilation response for standard, neutral novel and emotional novel sounds. Novel sounds evoked larger PDRs compared to standard sounds. Emotional novel sounds evoked larger PDR compared to neutral novel sounds, indicating an increase in arousal. Panel C: Mean distraction effects (RT novel minus

RT standard sound) and pupil dilation differences between PDR to novel and standard sound. This plot displays the hypothesized relationship between faster reaction times and larger pupil dilations to emotional novel sounds. This relation has been disconfirmed by the multilevel analysis. The plots show 95% confidence interval.

Fig. 4. Panel A: The relationship between performance (reaction times, RT) and pupil dilation response (PDR) at trial level. Larger pupil dilations in a trial were associated with slower reaction times. Panel B: The relationship between performance (standard reaction times, RT) and pupil dilation response (PDR) at participant level. Participants with larger pupil dilations showed faster average response times.

4. Discussion

This study investigated the direct relations of emotion-related distraction effects on performance in a primary task and increased levels of arousal evoked by processing of such emotional distractor sounds. Novel sounds, compared to standard sounds, prolonged RTs in a visual categorization task and evoked a transient dilation of the pupil. On the behavioral level, distraction effects were reduced in response to emotional compared to neutral novel sounds while the pupil dilated even more in response to emotional novel sounds vs. neutral novel

sounds. However, mixed-model effects could not provide any evidence for a correlation between performance and transient changes in pupil diameter that was specific to a sound's novelty or emotional content. This result was confirmed by Bayes Factors.

Novel sounds impaired performance in a subsequent categorization task compared to standard sounds. This result is consistent with current models of distraction of attention (Corbetta & Shulman, 2002; Escera & Corral, 2007; Näätänen, 1992; Posner, 1980, 2016; Sokolov, 1963). New, salient, and task-irrelevant events can involuntarily capture attention and can impair performance. This distraction effect (difference between RTs to distractor and RTs to standard sounds) has been observed in the auditory, visual, and tactile modality (Akatsuka et al., 2007; Escera, 1998; Schröger & Wolff, 1998) and has been replicated many times (Berti & Schröger, 2001, 2004; Hoyer et al., 2021; Parmentier, 2014; Wetzell et al., 2019). Task-irrelevant emotional novel sounds significantly decreased distraction effects compared to neutral novel sounds. Our results indicate that task-irrelevant emotional stimuli facilitated processing and improved behavioral performance in a subsequent task (Lindström & Bohlin, 2011; Lorenzino & Caudek, 2015; Max et al., 2015; Phelps et al., 2006; Zeelenberg & Bocanegra, 2010). Similar effects have been observed in the visual modality as well (e.g., Kim & Anderson, 2020; Kim et al., 2021; Lee et al., 2012; Sutherland & Mather, 2015). Emotional stimuli innately achieve prioritized processing due to their high motivational relevance (Mather et al., 2016; Schupp et al., 2003). However, previous studies also observed that emotional information can capture higher attentional resources than less salient stimuli resulting in impaired performance (for example, Kanske, 2012; Most et al., 2005; Pereira et al., 2006; Pessoa, 2008; for a review on visual emotional stimuli, see Bradley et al., 2012). Opposite results on modulations of emotional information on performance and attentional capacities may be explained due to differences between experimental designs, sensory stimulus presentation, task relevance and task assignment. For example, stimuli presented in

the same modality interfere more with each other than stimuli presented in different modalities (e.g., Duncan et al., 1997; Schupp et al., 2008; Soto-Faraco & Spence, 2002). Thus, emotional distraction effects may be reduced by presenting the task-irrelevant emotional stimulus in a different modality compared to the target (reduced response times De Houwer et al., 2002; Jiang et al., 2007; Scott et al., 2009; improved identification of visual targets Brosch et al., 2007, 2008; Zeelenberg & Bocanegra, 2010).

On a psychophysiological level, we observed a larger transient pupil dilation response (PDR) to novel sounds compared to standard sounds. This finding is in line with our hypothesis and with previous studies (Gilzenrat et al., 2010; Marois et al., 2020; Murphy et al., 2011) and has been discussed to reflect a transient increase in arousal (Joshi et al., 2016; Joshi & Gold, 2020; Krebs et al., 2018; Murphy et al., 2014; for a review see, Eckstein et al., 2017; Poe et al., 2020; Zekveld et al., 2018). Moreover, larger pupil dilation was observed in response to emotional novel sounds compared to neutral novel sounds (as e.g., Bonmassar et al., 2020; Partala & Surakka, 2003; Widmann et al., 2018). Thus, our study could replicate effects of unexpected, task-irrelevant, and emotionally arousing sounds on both performance and pupil size from previous studies. A dilation of the pupil is likely to be modulated by the increased activity of LC-NE system (Poe et al., 2020), which primarily modulates the level of arousal (Aston-Jones & Cohen, 2005; Corbetta et al., 2008; Poe et al., 2020) and is discussed to be involved in the processing of emotion (for review see Nieuwenhuis et al., 2011; Ranganath & Rainer, 2003; Sara & Bouret, 2012). The bias in favor of infrequent, salient and high-priority information is modulated by the norepinephrine release of the locus coeruleus (Joshi & Gold, 2020; Mather et al., 2016; Poe et al., 2020) and is represented by the pupil dilation (Megemont, 2022). In fact, animal studies evidenced a causal relation between stimulations of LC neurons and the subsequent pupil dilation for both tonic and phasic LC activity (for tonic see Gilzenrat et al., 2010; Hayat et al., 2020; Joshi & Gold, 2020; Liu et al., 2017, for phasic

see Gilzenrat et al., 2010; Joshi et al., 2016; Joshi & Gold, 2020; Reimer et al., 2016; Sara & Bouret, 2012).

It is worth mentioning that the pupil can also partly reflect the activity of structures other than the LC such as the superior colliculus (SC, for review see Wang & Munoz, 2015) and the anterior cingulate cortex (ACC). Activity in such brain areas is coordinated with pupil fluctuations that are associated with certain aspects of cognitive processing, including attention and orienting to salient stimuli (Joshi et al., 2016; Wang & Munoz, 2015). However, LC and SC might subserve complementary functions (for review see Einhäuser, 2017), for example, the SC is part of a coeruleo-cortical pathway that modulates attentional functions (Wang & Munoz, 2015). All these multiple pathways may help to explain the more variable timing of pupil-related modulations of neuronal activity.

Even though emotional novel sounds evoked larger PDRs and reduced distraction effects separately, the applied multilevel model did not support a correlation between both effects neither on the trial nor on the participant level. The lack of a correlation was confirmed by the computation of Bayes Factors, which showed that the data provide strong evidence against such interactions. That is, emotional novel sounds evoking larger PDRs did not show systematically larger behavioral facilitation effects and participants showing larger average PDRs in response to emotional distractor sounds did not show correspondingly larger behavioral facilitation effects. Based on these results, we suggest that the emotion-related facilitation effect on the behavioral level and the increase in arousal reflected by the PDR do most likely not reflect the operation of identical processes. They are presumably caused by at least partly independent mechanisms. This does not exclude common precursor processes. It can be speculated that one of the involved processes does not show proportional behavior, for example due to all-or-nothing effects or ceiling or floor effects. Taken together, we propose

that our behavioral and psychophysiological results indicate the operation of possibly related, but not identical mechanisms contributing to emotion-related decreased effects of distraction.

Even though we did not find an emotion-specific correlation between reduced distraction effects and increased PDR, our exploratory analysis showed two opposite relationships between RTs and PDRs at the trial and the participant level, independent of the sound type presented. On a participant level, participants with larger mean PDR, responded faster to target stimuli. Behaviorally relevant stimuli can dilate the pupil (e.g., Beatty, 1982; Murphy et al., 2014). The negative correlation could indicate that participants with increased PDR have continuously and more effectively used the sounds as a temporal cue for both the occurrence and the timing of the upcoming target (Hackley, 2009; Hackley & Valle-Inclán, 2003) and effectively prepared for the onset of the to-be-categorized stimulus (Volosin et al., 2016; Wetzel et al., 2013). This can result in faster responses compared to participants who were less engaged in the task. Strauch and colleagues suggested that pupil dilation might be interpreted as a readout of all three attentional subsystems, alerting, orienting and executive attention as suggested by Petersen & Posner (2012). Following this suggestion, the negative correlation at participant level could also reflect higher-level attentional factors related to the executive functions: Participants with larger PDR employed more attentional resources because they were more engaged in performing the task. In both cases, the negative correlation between RTs and PDRs might reflect participant-level aspects of task engagement (see e.g. also Hopstaken et al., 2015).

At the trial level, we observed a positive correlation between RTs and PDRs, that is, in trials with slower reaction times we observed larger PDRs in the same trial irrespective of sound type (as previously reported by Murphy et al., 2011). Again, following the suggestion by Strauch et al. (2022), this positive correlation could reflect intermediate-level factors related to alerting and orienting of attention (Petersen & Posner, 2012): at a trial level, pupil dilation

indicates orienting and distraction of attention in response to stimuli occurring in the surrounding. Interestingly, as we did not observe an interaction effect of trial-level PDR and novelty, the relation slower RT-larger PDR apparently might also hold for standard trials. We suggest that attentional orienting and enhanced stimulus processing observable in distracting novel trial at larger scales might also occur in standard trials at smaller scales, for example in relation to increased phasic NE release, potentially due to attentional orienting toward sound stimulation and spontaneous fluctuations of the LC activity (Jepma & Nieuwenhuis, 2011), resulting in enhanced processing of the current sound (and vice versa; Aston-Jones & Cohen, 2005) also in standard trials. The enhanced processing of the task-irrelevant standard and novel sounds can impair subsequent target stimulus-related processes, resulting in increased RT at trial-level.

Since PDR and RT on a trial level consider deviations relative to the participants' individual averages whereas participant level RT and PDR consider deviations of the participants individual averages from the grand average, these correlations represent differentiable sources of variance and the relationships can point in different directions (Enders & Tofighi, 2007; see also LoTempio et al., 2021 for the relation of P3b and RT in an oddball task). More generally, the trial-to-trial fluctuation of activity could reflect brain processes specific to that stimulus-driven behavior, whereas a difference between participants could reflect a general individual response bias to incoming stimuli. Our results demonstrate that the centering strategies, common in multilevel models, can also be applied effectively to disentangle enduring and transient effects in experimental settings.

6. Conclusion

Our findings indicate that task-irrelevant and unexpected novel sounds impair performance in a categorization task and distraction effects are reduced in response to emotional compared to

584 neutral novel sounds. Transient changes in pupil size are larger in response to novel sounds
585 compared to standard sounds and this increase is larger for emotional than for neutral novel
586 sounds. Our frequentist and Bayesian results disconfirm our hypothesis of a direct relation
587 between reduced distraction effects on the behavioral level and increased arousal reflected by
588 larger PDR to emotional novel sounds. We suggest that both performance and pupil diameter
589 reflect partly distinct processes. Given that the PDR has been discussed to indirectly reflect
590 the activity of the locus coeruleus, the LC-NE system may embody a common antecedent for
591 both effects, spreading norepinephrine to cortical areas involved in attention control and
592 control of the pupil. In addition, the observed emotion-unspecific correlations between
593 performance (RT) and levels of arousal (PDR), that differ on the trial and the participants
594 level, provide new insights into the underlying mechanisms of potential fluctuation of the LC-
595 NE system, aspects of individual task engagement and their effects on performance.

596 At last, the present paradigm makes use of ethically harmless stimuli which do not cause
597 anxiety, fear or threat. Thus, it can be applied in the research of a relation between attentional
598 distraction and arousal in developmental and clinical studies.

599

600 **Author Notes**

601 We thank Dunja Kunke and Tjerk Dercksen for proofreading and Carolin Albrecht for her
602 assistance in investigating the cause for the singular model fits. We are grateful for support in
603 data acquisition by Dunja Kunke and Gabriele Schoeps. We thank all participants for support.
604 The project was funded by the DFG (WE 5026/1-2), the Center for Behavioral Brain Sciences
605 Magdeburg funded by the European Regional Development Fund (ZS/ 2016/04/78120), and
606 the Leibniz Association (P58/2017).

607

608 **Declarations of Interest**

609 None.

610 **For reprints:** Carolina Bonmassar, Carolina.Bonmassar@lin-magdeburg.de

611

Footnotes

¹All sounds used in the present study were environmental sounds. In the following we will omit the specification “environmental” and we will term the sounds as “standard”, “emotional novel” and “neutral novel” in order to improve readability.

² Initially, we intended to use the predictor Condition (Standard vs. Emotional novel sound vs. Neutral novel sound). However, a random slope of Condition results in 2 separate random slopes for the differences between emotional novels vs. standards and neutral novels vs. standards, respectively. Both for RT and PDR as dependent variable, this resulted in a singular model due to the high correlation of these random slopes – indicating that a single random slope across both novel types was more appropriate. Re-parameterization into the separate predictors Novelty and Emotionality enabled a model with a common random slope (i.e., individual distraction effects) across both types of novel sounds.

³ Theoretically, the participant-level PDR/Baseline could be computed from the average across all sound types or separately for each sound Type. The correlations between the values from these approaches were very high (all $r_s > .80$) and the choice between these methods did therefore not change the results in any meaningful way. For the sake of comprehensibility, we only report the results when averaging PDR/Baseline across all sound types.

⁴ We investigated the potential impact of the misspecified level-1-residual distribution by comparing our model with normal level-1 distribution with a model with an exgaussian distribution which can account for the considerable skewness of RTs at trial-level using brms (Bürkner, 2017, 2018; Carpenter et al., 2016) which utilizes a Bayesian estimation algorithm.

637 The exgaussian model fit the data substantially better, but this did not affect any substantive
638 conclusion, because none of the parameters changed its sign or effect size fundamentally.

639

640 *These terms were added manually following the substantive restrictions outlined above. This
641 change did not affect size or hypothesis test of any other effect in the model.

642

643

References

- Akatsuka, K., Wasaka, T., Nakata, H., Kida, T., Hoshiyama, M., Tamura, Y., & Kakigi, R. (2007). Objective examination for two-point stimulation using a somatosensory oddball paradigm: An MEG study. *Clinical Neurophysiology*, 118(2), 403–411. <https://doi.org/https://doi.org/10.1016/j.clinph.2006.09.030>
- Alday, P. M. (2019). How much baseline correction do we need in ERP research? Extended GLM model can replace baseline correction while lifting its limits. *Psychophysiology*, 56(12), e13451. <https://doi.org/https://doi.org/10.1111/psyp.13451>
- Aston-Jones, G., & Cohen, J. D. (2005). An integrative theory of locus coeruleus-norepinephrine function: Adaptive gain and optimal performance. *Annual Review of Neuroscience*, 28, 403–450. <https://doi.org/10.1146/annurev.neuro.28.061604.135709>
- Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, 67(1), 1–48. <https://doi.org/10.18637/jss.v067.i01>
- Beatty, J. (1982). Task-evoked pupillary responses, processing load, and the structure of processing resources. In *Psychological Bulletin* (Vol. 91, Issue 2, pp. 276–292). American Psychological Association. <https://doi.org/10.1037/0033-2909.91.2.276>
- Bendixen, A., Roeber, U., & Schröger, E. (2007). Regularity extraction and application in dynamic auditory stimulus sequences. *Journal of Cognitive Neuroscience*, 19(10), 1664–1677. <https://doi.org/10.1162/jocn.2007.19.10.1664>
- Berti, S., Roeber, U., & Schröger, E. (2004). Bottom-Up Influences on Working Memory: Behavioral and Electrophysiological Distraction Varies with Distractor Strength. *Experimental Psychology*, 51(4), 249–257. <https://doi.org/10.1027/1618-3169.51.4.249>
- Berti, S., & Schröger, E. (2001). A comparison of auditory and visual distraction effects: behavioral and event-related indices. *Cognitive Brain Research*, 10(3), 265–273. [https://doi.org/https://doi.org/10.1016/S0926-6410\(00\)00044-6](https://doi.org/https://doi.org/10.1016/S0926-6410(00)00044-6)

670 Berti, S., & Schröger, E. (2004). Distraction effects in vision: Behavioral and event-related
671 potential indices. *NeuroReport*, 15(4), 665–669. [https://doi.org/10.1097/00001756-](https://doi.org/10.1097/00001756-200403220-00018)
672 200403220-00018

673 Bonmassar, C., Widmann, A., & Wetzel, N. (2020). The impact of novelty and emotion on
674 attention-related neuronal and pupil responses in children. *Developmental Cognitive*
675 *Neuroscience*, 42(August 2019), 100766. <https://doi.org/10.1016/j.dcn.2020.100766>

676 Bradley, M. M., Keil, A., & Lang, P. J. (2012). Orienting and emotional perception:
677 Facilitation, attenuation, and interference. *Frontiers in Psychology*, 3(NOV), 1–6.
678 <https://doi.org/10.3389/fpsyg.2012.00493>

679 Bradley, M. M., & Lang, P. J. (2007). The International Affective Digitized Sounds (; IADS-
680 2): Affective ratings of sounds and instruction manual. *University of Florida,*
681 *Gainesville, FL, Tech. Rep. B-3.*

682 Bradley, M. M., Miccoli, L., Escrig, M. A., & Lang, P. J. (2008). The pupil as a measure of
683 emotional arousal and autonomic activation. *Psychophysiology*, 45(4), 602–607.
684 <https://doi.org/10.1111/j.1469-8986.2008.00654.x>

685 Bradley, M. M., Sapigao, R. G., & Lang, P. J. (2017). Sympathetic ANS modulation of pupil
686 diameter in emotional scene perception: Effects of hedonic content, brightness, and
687 contrast. *Psychophysiology*, 54(10), 1419–1435. <https://doi.org/10.1111/psyp.12890>

688 Brosch, T., Grandjean, D., Sander, D., & Scherer, K. R. (2008). Behold the voice of wrath:
689 Cross-modal modulation of visual attention by anger prosody. *Cognition*, 106(3), 1497–
690 1503. <https://doi.org/10.1016/j.cognition.2007.05.011>

691 Brosch, T., Sander, D., & Scherer, K. R. (2007). That baby caught my eye... attention capture
692 by infant faces. *Emotion*, 7 3, 685–689.

693 Burnham, K. P., & Anderson, D. R. (2004). Multimodel inference: Understanding AIC and
694 BIC in model selection. *Sociological Methods and Research*, 33(2), 261–304.
695 <https://doi.org/10.1177/0049124104268644>

696 Corbetta, M., Patel, G., & Shulman, G. L. (2008). The Reorienting System of the Human
697 Brain: From Environment to Theory of Mind. *Neuron*, 58(3), 306–324.
698 <https://doi.org/10.1016/j.neuron.2008.04.017>

699 Corbetta, M., & Shulman, G. L. (2002). Control of goal-directed and stimulus-driven
700 attention in the brain. *Nature Reviews Neuroscience*, 3(3), 201–215.
701 <https://doi.org/10.1038/nrn755>

702 De Houwer, J., Hermans, D., Rothermund, K., & Wentura, D. (2002). Affective priming of
703 semantic categorisation responses. *Cognition and Emotion*, 16(5), 643–666.
704 <https://doi.org/10.1080/02699930143000419>

705 Duncan, J., Martens, S., & Ward, R. (1997). Restricted attentional capacity within but not
706 between sensory modalities. *Nature*, 387(6635), 808–810. <https://doi.org/10.1038/42947>

707 Eckstein, M. K., Guerra-Carrillo, B., Miller Singley, A. T., & Bunge, S. A. (2017). Beyond
708 eye gaze: What else can eyetracking reveal about cognition and cognitive development?
709 *Developmental Cognitive Neuroscience*, 25, 69–91.
710 <https://doi.org/10.1016/j.dcn.2016.11.001>

711 Einhäuser, W. (2017). The pupil as marker of cognitive processes. *Cognitive Science and*
712 *Technology*, 141–169. https://doi.org/10.1007/978-981-10-0213-7_7

713 Enders, C. K., & Tofighi, D. (2007). Centering predictor variables in cross-sectional
714 multilevel models: a new look at an old issue. *Psychological Methods*, 12(2), 121–138.
715 <https://doi.org/10.1037/1082-989X.12.2.121>

716 Escera, C. (1998). *Neural Mechanisms of Involuntary Attention*. 590–604.

717 Escera, C., Alho, K., Winkler, I., & Näätänen, R. (1998). Neural Mechanisms of Involuntary
718 Attention to Acoustic Novelty and Change. *Journal of Cognitive Neuroscience*, 10(5),
719 590–604. <https://doi.org/10.1162/089892998562997>

720 Escera, C., & Corral, M. J. (2007). Role of mismatch negativity and novelty-P3 in involuntary
721 auditory attention. *Journal of Psychophysiology*, 21(3–4), 251–264.

722 <https://doi.org/10.1027/0269-8803.21.34.251>

723 Gilzenrat, M. S., Nieuwenhuis, S., Jepma, M., & Cohen, J. D. (2010). Pupil diameter tracks
 724 changes in control state predicted by the adaptive gain theory of locus coeruleus
 725 function. *Cogn Affect Behav Neurosci*, 10(2), 252–269.
 726 <https://doi.org/10.3758/CABN.10.2.252>.Pupil

727 Hackley, S. A. (2009). The speeding of voluntary reaction by a warning signal: Presidential
 728 Address, 2006. *Psychophysiology*, 46(2), 225–233. [https://doi.org/10.1111/j.1469-](https://doi.org/10.1111/j.1469-8986.2008.00716.x)
 729 [8986.2008.00716.x](https://doi.org/10.1111/j.1469-8986.2008.00716.x)

730 Hackley, S. A., & Valle-Inclán, F. (2003). Which stages of processing are speeded by a
 731 warning signal? *Biological Psychology*, 64(1), 27–45.
 732 [https://doi.org/https://doi.org/10.1016/S0301-0511\(03\)00101-7](https://doi.org/10.1016/S0301-0511(03)00101-7)

733 Hayat, H., Regev, N., Matosevich, N., Sales, A., Paredes-Rodriguez, E., Krom, A. J.,
 734 Bergman, L., Li, Y., Lavigne, M., Kremer, E. J., Yizhar, O., Pickering, A. E., & Nir, Y.
 735 (2020). *Locus coeruleus norepinephrine activity mediates sensory-evoked awakenings*
 736 *from sleep*. <https://www.science.org>

737 Hayes, T. R., & Petrov, A. A. (2016). Mapping and correcting the influence of gaze position
 738 on pupil size measurements. *Behavior Research Methods*, 48(2), 510–527.
 739 <https://doi.org/10.3758/s13428-015-0588-x>

740 Hess, E. H., & Polt, J. M. (1960). Pupil Size as Related to Interest Value of Visual Stimuli.
 741 *Science*, 132(3423), 349 LP – 350. <https://doi.org/10.1126/science.132.3423.349>

742 Hopstaken, J. F., van der Linden, D., Bakker, A. B., & Kompier, M. A. J. (2015). The
 743 window of my eyes: Task disengagement and mental fatigue covary with pupil
 744 dynamics. *Biological Psychology*, 110, 100–106.
 745 <https://doi.org/10.1016/j.biopsycho.2015.06.013>

746 Horváth, J., Winkler, I., & Bendixen, A. (2008). Do N1/MMN, P3a, and RON form a strongly
 747 coupled chain reflecting the three stages of auditory distraction? *Biological Psychology*,

748 79(2), 139–147. <https://doi.org/10.1016/j.biopsycho.2008.04.001>

749 Hoyer, R. S., Elshafei, H., Hemmerlin, J., Bouet, R., & Bidet-Caulet, A. (2021). Why Are
750 Children So Distractible? Development of Attention and Motor Control From Childhood
751 to Adulthood. *Child Development*, 00(0), 1–22. <https://doi.org/10.1111/cdev.13561>

752 Jepma, M., & Nieuwenhuis, S. (2011). Pupil Diameter Predicts Changes in the Exploration–
753 Exploitation Trade-off: Evidence for the Adaptive Gain Theory. *Journal of Cognitive*
754 *Neuroscience*, 23(7), 1587–1596. <https://doi.org/10.1162/jocn.2010.21548>

755 Jiang, Y., Vagnini, V., Clark, J., & Zhang, Q. (2007). Reduced sensitivity of older adults to
756 affective mismatches. *TheScientificWorldJournal*, 7(February), 641–648.
757 <https://doi.org/10.1100/tsw.2007.115>

758 Joshi, S., & Gold, J. I. (2020). Pupil size as a window on neural substrates of cognition.
759 *Trends in Cognitive Sciences*, 24(6), 466–480. <https://doi.org/10.1016/j.tics.2020.03.005>

760 Joshi, S., Li, Y., Kalwani, R. M., & Gold, J. I. (2016). Relationships between Pupil Diameter
761 and Neuronal Activity in the Locus Coeruleus, Colliculi, and Cingulate Cortex. *Neuron*,
762 89(1), 221–234. <https://doi.org/10.1016/j.neuron.2015.11.028>

763 Kanske, P. (2012). On the influence of emotion on conflict processing. *Frontiers in*
764 *Integrative Neuroscience*, 6(JULY 2012), 2010–2013.
765 <https://doi.org/10.3389/fnint.2012.00042>

766 Kim, Andy J., & Anderson, B. A. (2020). Arousal-biased competition explains reduced
767 distraction by reward cues under threat. *ENeuro*, 7(4), 1–12.
768 <https://doi.org/10.1523/ENEURO.0099-20.2020>

769 Kim, Andy Jeesu, Lee, D. S., & Anderson, B. A. (2021). The influence of threat on the
770 efficiency of goal-directed attentional control. *Psychological Research*, 85(3), 980–986.
771 <https://doi.org/10.1007/s00426-020-01321-4>

772 Kleiner, M., Brainard, D., & Pelli, D. (2007). What’s new in psychtoolbox-3? *Perception*
773 *(ECVP Abstract Supplement)* 14.

774 Klingner, J., Kumar, R., & Hanrahan, P. (2008). Measuring the Task-Evoked Pupillary
775 Response with a Remote Eye Tracker. *Proceedings of the 2008 Symposium on Eye*
776 *Tracking Research & Applications*, 69–72.
777 <https://doi.org/10.1145/1344471.1344489>

778 Krebs, R. M., Park, H. R. P., Bombeke, K., & Boehler, C. N. (2018). Modulation of locus
779 coeruleus activity by novel oddball stimuli. *Brain Imaging and Behavior*, 12(2), 577–
780 584. <https://doi.org/10.1007/s11682-017-9700-4>

781 Kret, M. E., & Sjak-Shie, E. E. (2019). Preprocessing pupil size data: Guidelines and code.
782 *Behavior Research Methods*. <https://doi.org/10.3758/s13428-018-1075-y>

783 Kuznetsova, A., Brockhoff, P., & Christensen, R. (2017). lmerTest Package: Tests in Linear
784 Mixed Effects Models. *Journal of Statistical Software*, 82, 1–26.

785 Lee, T. H., Itti, L., & Mather, M. (2012). Evidence for arousal-biased competition in
786 perceptual learning. *Frontiers in Psychology*, 3(JUL), 1–9.
787 <https://doi.org/10.3389/fpsyg.2012.00241>

788 Li, X., Liang, Z., Kleiner, M., & Lu, Z.-L. (2010). RTbox: A device for highly accurate
789 response time measurements. *Behavior Research Methods*, 42(1), 212–225.
790 <https://doi.org/10.3758/BRM.42.1.212>

791 Liao, H. I., Kidani, S., Yoneya, M., Kashino, M., & Furukawa, S. (2016). Correspondences
792 among pupillary dilation response, subjective salience of sounds, and loudness.
793 *Psychonomic Bulletin and Review*, 23(2), 412–425. [https://doi.org/10.3758/s13423-015-](https://doi.org/10.3758/s13423-015-0898-0)
794 0898-0

795 Liao, H. I., Yoneya, M., Kidani, S., Kashino, M., & Furukawa, S. (2016). Human pupillary
796 dilation response to deviant auditory stimuli: Effects of stimulus properties and voluntary
797 attention. *Frontiers in Neuroscience*, 10(FEB). <https://doi.org/10.3389/fnins.2016.00043>

798 Lindström, B. R., & Bohlin, G. (2011). Emotion processing facilitates working memory
799 performance. *Cognition and Emotion*, 25(7), 1196–1204.

800   <https://doi.org/10.1080/02699931.2010.527703>

801   Liu, Y., Rodenkirch, C., Moskowitz, N., Schriver, B., & Wang, Q. (2017). Dynamic
802   Lateralization of Pupil Dilation Evoked by Locus Coeruleus Activation Results from
803   Sympathetic, Not Parasympathetic, Contributions. *Cell Reports*, 20(13), 3099–3112.
804   <https://doi.org/10.1016/j.celrep.2017.08.094>

805   Lorenzino, M., & Caudek, C. (2015). Task-irrelevant emotion facilitates face discrimination
806   learning. *Vision Research*, 108, 56–66. <https://doi.org/10.1016/j.visres.2015.01.007>

807   LoTempio, S., Silcox, J., Federmeier, K. D., & Payne, B. R. (2021). Inter- and intra-
808   individual coupling between pupillary, electrophysiological, and behavioral responses in
809   a visual oddball task. *Psychophysiology*, 58(4), 1–14. <https://doi.org/10.1111/psyp.13758>

810   Marois, A., Pozzi, A., & Vachon, F. (2020). Assessing the Role of Stimulus Novelty in the
811   Elicitation of the Pupillary Dilation Response to Irrelevant Sound. *Auditory Perception*
812   & *Cognition*, 00(00), 1–17. <https://doi.org/10.1080/25742442.2020.1820290>

813   Mather, M., Clewett, D., Sakaki, M., & Harley, C. W. (2016). Norepinephrine ignites local
814   hotspots of neuronal excitation: How arousal amplifies selectivity in perception and
815   memory. *Behavioral and Brain Sciences*, 39.
816   <https://doi.org/10.1017/S0140525X15000667>

817   Mathôt, S., Fabius, J., Van Heusden, E., & Van der Stigchel, S. (2018). Safe and sensible
818   preprocessing and baseline correction of pupil-size data. *Behavior Research Methods*,
819   50(1), 94–106. <https://doi.org/10.3758/s13428-017-1007-2>

820   Max, C., Widmann, A., Kotz, S. A., Schröger, E., & Wetzels, N. (2015). Distraction by
821   emotional sounds: Disentangling arousal benefits and orienting costs. *Emotion*, 15(4),
822   428–437. <https://doi.org/10.1037/a0039041>

823   May, P. J. C., & Tiitinen, H. (2010). Mismatch negativity (MMN), the deviance-elicited
824   auditory deflection, explained. *Psychophysiology*, 47(1), 66–122.
825   <https://doi.org/10.1111/j.1469-8986.2009.00856.x>

826 McHugo, M., Olatunji, B. O., & Zald, D. H. (2013). The emotional attentional blink: What we
827 know so far. Running title: The emotional attentional blink. *Frontiers in Human*
828 *Neuroscience*, 7(APR 2013), 1–9. <https://doi.org/10.3389/fnhum.2013.00151>

829 Merritt, S. L., Keegan, A. P., & Mercer, P. W. (1994). Artifact Management in Pupillometry.
830 *Nursing Research*, 43(1).
831 [https://journals.lww.com/nursingresearchonline/Fulltext/1994/01000/Artifact_Managem](https://journals.lww.com/nursingresearchonline/Fulltext/1994/01000/Artifact_Management_in_Pupillometry.12.aspx)
832 [ent_in_Pupillometry.12.aspx](https://journals.lww.com/nursingresearchonline/Fulltext/1994/01000/Artifact_Management_in_Pupillometry.12.aspx)

833 Morey, R. D., & Rouder, J. N. (2011). Bayes factor approaches for testing interval null
834 hypotheses. In *Psychological Methods* (Vol. 16, Issue 4, pp. 406–419). American
835 Psychological Association. <https://doi.org/10.1037/a0024377>

836 Morey, R. D., Rouder, J. N., Pratte, M. S., & Speckman, P. L. (2011). Using MCMC chain
837 outputs to efficiently estimate Bayes factors. *Journal of Mathematical Psychology*, 55,
838 368–378.

839 Most, S. B., Chun, M. M., Widders, D. M., Haven, N., & Zald, D. H. (2005). *Attentional*
840 *rubbernecking: Cognitive control and personality in emotion-induced blindness*. 12(4),
841 654–661.

842 Muller-Gass, A., & Schröger, E. (2007). Perceptual and cognitive task difficulty has
843 differential effects on auditory distraction. *Brain Research*, 1136, 169–177.
844 <https://doi.org/https://doi.org/10.1016/j.brainres.2006.12.020>

845 Murphy, P. R., O’Connell, R. G., O’Sullivan, M., Robertson, I. H., & Balsters, J. H. (2014).
846 Pupil diameter covaries with BOLD activity in human locus coeruleus. *Human Brain*
847 *Mapping*, 35(8), 4140–4154. <https://doi.org/10.1002/hbm.22466>

848 Murphy, P. R., Robertson, I. H., Balsters, J. H., & O’connell, R. G. (2011). Pupillometry and
849 P3 index the locus coeruleus-noradrenergic arousal function in humans.
850 *Psychophysiology*, 48(11), 1532–1543. [https://doi.org/10.1111/j.1469-](https://doi.org/10.1111/j.1469-8986.2011.01226.x)
851 [8986.2011.01226.x](https://doi.org/10.1111/j.1469-8986.2011.01226.x)

852 Näätänen, R. (1992). *Attention and Brain Function*.

853 Nakagawa, S., & Schielzeth, H. (2013). A general and simple method for obtaining R² from
854 generalized linear mixed-effects models. *Methods in Ecology and Evolution*, 4(2), 133–
855 142. <https://doi.org/10.1111/j.2041-210x.2012.00261.x>

856 Nieuwenhuis, S., De Geus, E. J., & Aston-Jones, G. (2011). The anatomical and functional
857 relationship between the P3 and autonomic components of the orienting response.
858 *Psychophysiology*, 48(2), 162–175. <https://doi.org/10.1111/j.1469-8986.2010.01057.x>

859 Oldfield, R. C. (1971). The assessment and analysis of handedness: The Edinburgh inventory.
860 *Neuropsychologia*, 9(1), 97–113. [https://doi.org/https://doi.org/10.1016/0028-](https://doi.org/https://doi.org/10.1016/0028-3932(71)90067-4)
861 3932(71)90067-4

862 Parmentier, F. B. R. (2014). The cognitive determinants of behavioral distraction by deviant
863 auditory stimuli: A review. *Psychological Research*, 78(3), 321–338.
864 <https://doi.org/10.1007/s00426-013-0534-4>

865 Parmentier, F. B. R., Elford, G., Escera, C., Andrés, P., & San Miguel, I. (2008). The
866 cognitive locus of distraction by acoustic novelty in the cross-modal oddball task.
867 *Cognition*, 106(1), 408–432. <https://doi.org/10.1016/j.cognition.2007.03.008>

868 Partala, T., & Surakka, V. (2003). Pupil size variation as an indication of affective processing.
869 *International Journal of Human Computer Studies*, 59(1–2), 185–198.
870 [https://doi.org/10.1016/S1071-5819\(03\)00017-X](https://doi.org/10.1016/S1071-5819(03)00017-X)

871 Pereira, M. G., Volchan, E., de Souza, G. G. L., Oliveira, L., Campagnoli, R. R., Pinheiro, W.
872 M., & Pessoa, L. (2006). Sustained and transient modulation of performance induced by
873 emotional picture viewing. *Emotion*, 6(4), 622–634.

874 Pessoa, L. (2008). *On the relationship between emotion and cognition*. 9(2).
875 www.nature.com/reviews/neuro

876 Petersen, S. E., & Posner, M. I. (2012). The attention system of the human brain: 20 years
877 after. *Annual Review of Neuroscience*, 35, 73–89. <https://doi.org/10.1146/annurev-neuro->

878 062111-150525

879 Phelps, E. A., Ling, S., & Carrasco, M. (2006). Emotion facilitates perception and potentiates
880 the perceptual benefits of attention. *Psychological Science*, 17(4), 292–299.
881 <https://doi.org/10.1111/j.1467-9280.2006.01701.x>

882 Poe, G. R., Foote, S., Eschenko, O., Johansen, J. P., Bouret, S., Aston-Jones, G., Harley, C.
883 W., Manahan-Vaughan, D., Weinshenker, D., Valentino, R., Berridge, C., Chandler, D.
884 J., Waterhouse, B., & Sara, S. J. (2020). Locus coeruleus: a new look at the blue spot.
885 *Nature Reviews Neuroscience*, 21(11), 644–659. [https://doi.org/10.1038/s41583-020-](https://doi.org/10.1038/s41583-020-0360-9)
886 0360-9

887 Posner, M. I. (1980). Orienting of Attention. *Quarterly Journal of Experimental Psychology*,
888 32(1), 3–25. <https://doi.org/10.1080/00335558008248231>

889 Posner, M. I. (2016). Orienting of Attention: Then and Now. *Quarterly Journal of*
890 *Experimental Psychology*, 69(10), 1864–1875.
891 <https://doi.org/10.1080/17470218.2014.937446>

892 R Core, T. (2019). *R: a language and environment for statistical computing*. R Foundation for
893 *Statistical Computing, Austria*.

894 Raftery, A. E. (1995). Bayesian Model Selection in Social Research. *Sociological*
895 *Methodology*, 25, 111–163. <https://doi.org/10.2307/271063>

896 Ranganath, C., & Rainer, G. (2003). Cognitive neuroscience: Neural mechanisms for
897 detecting and remembering novel events. *Nature Reviews Neuroscience*, 4(3), 193–202.
898 <https://doi.org/10.1038/nrn1052>

899 Reimer, J., McGinley, M. J., Liu, Y., Rodenkirch, C., Wang, Q., McCormick, D. A., & Tolias,
900 A. S. (2016). Pupil fluctuations track rapid changes in adrenergic and cholinergic activity
901 in cortex. *Nature Communications*, 7(May), 1–7. <https://doi.org/10.1038/ncomms13289>

902 Rouder, J. N., Speckman, P. L., Sun, D., Morey, R. D., & Iverson, G. J. (2009). Bayesian t
903 tests for accepting and rejecting the null hypothesis. *Psychonomic Bulletin & Review*,

904 16, 225–237.

905 Sara, S. J., & Bouret, S. (2012). Orienting and Reorienting: The Locus Coeruleus Mediates
 906 Cognition through Arousal. *Neuron*, 76(1), 130–141.
 907 <https://doi.org/10.1016/j.neuron.2012.09.011>

908 Schröger, E. (1997). On the detection of auditory deviations: A pre-attentive activation model.
 909 *Psychophysiology*, 34(3), 245–257. [https://doi.org/https://doi.org/10.1111/j.1469-](https://doi.org/10.1111/j.1469-8986.1997.tb02395.x)
 910 [8986.1997.tb02395.x](https://doi.org/10.1111/j.1469-8986.1997.tb02395.x)

911 Schröger, E., & Wolff, C. (1998). Attentional orienting and reorienting is indicated by human
 912 event-related brain potentials. *Cognitive Brain Research*, 7(1), 71–87.
 913 [https://doi.org/10.1016/S0926-6410\(98\)00013-5](https://doi.org/10.1016/S0926-6410(98)00013-5)

914 Schupp, H. T., Junghöfer, M., Weike, A. I., & Hamm, A. O. (2003). Attention and emotion:
 915 An ERP analysis of facilitated emotional stimulus processing. *NeuroReport*, 14(8),
 916 1107–1110. <https://doi.org/10.1097/00001756-200306110-00002>

917 Schupp, H. T., Stockburger, J., Bublatzky, F., Junghöfer, M., Weike, A. I., & Hamm, A. O.
 918 (2008). The selective processing of emotional visual stimuli while detecting auditory
 919 targets: An ERP analysis. *Brain Research*, 1230, 168–176.
 920 [https://doi.org/https://doi.org/10.1016/j.brainres.2008.07.024](https://doi.org/10.1016/j.brainres.2008.07.024)

921 Schwarz, G. (1978). Estimating the Dimension of a Model. *The Annals of Statistics*, 6(2),
 922 461–464. <https://doi.org/10.1214/aos/1176344136>

923 Scott, G. G., O'Donnell, P. J., Leuthold, H., & Sereno, S. C. (2009). Early emotion word
 924 processing: Evidence from event-related potentials. *Biological Psychology*, 80(1), 95–
 925 104. <https://doi.org/10.1016/j.biopsycho.2008.03.010>

926 Sokolov, E. N. (1963). Higher Nervous Functions: The Orienting Reflex. *Annual Review of*
 927 *Physiology*, 25(1), 545–580. <https://doi.org/10.1146/annurev.ph.25.030163.002553>

928 Soto-Faraco, S., & Spence, C. (2002). Modality-specific auditory and visual temporal
 929 processing deficits. *The Quarterly Journal of Experimental Psychology Section A*, 55(1),

930 23–40. <https://doi.org/10.1080/02724980143000136>

931 Steinhauer, S. R., Bradley, M. M., Siegle, G. J., Roecklein, K. A., & Dix, A. (2022).
 932 Publication guidelines and recommendations for pupillary measurement in
 933 psychophysiological studies. *Psychophysiology*, 59(4), 1–36.
 934 <https://doi.org/10.1111/psyp.14035>

935 Strauch, C., Wang, C.-A., Einhäuser, W., Van der Stigchel, S., & Naber, M. (2022).
 936 Pupillometry as an integrated readout of distinct attentional networks. *Trends in*
 937 *Neurosciences*, 1–13. <https://doi.org/10.1016/j.tins.2022.05.003>

938 Sutherland, M., & Mather, M. (2015). Negative Arousal Increases the Effects of Stimulus
 939 Salience in Older Adults. *Exp Aging Res*, 41(3).
 940 <https://doi.org/10.1080/0361073X.2015.1021644>.

941 Volosin, M., Grimm, S., & Horváth, J. (2016). Exploiting temporal predictability: Event-
 942 related potential correlates of task-supportive temporal cue processing in auditory
 943 distraction. *Brain Research*, 1639, 120–131.
 944 <https://doi.org/10.1016/j.brainres.2016.02.044>

945 Wang, C. A., Baird, T., Huang, J., Coutinho, J. D., Brien, D. C., & Munoz, D. P. (2018).
 946 Arousal Effects on Pupil Size, Heart Rate, and Skin Conductance in an Emotional Face
 947 Task. *Frontiers in Neurology*, 9(December), 1–13.
 948 <https://doi.org/10.3389/fneur.2018.01029>

949 Wang, C. A., & Munoz, D. P. (2015). A circuit for pupil orienting responses: Implications for
 950 cognitive modulation of pupil size. *Current Opinion in Neurobiology*, 33, 134–140.
 951 <https://doi.org/10.1016/j.conb.2015.03.018>

952 Wetzel, N. (2015). Effects of the short-term learned significance of task-irrelevant sounds on
 953 involuntary attention in children and adults. *International Journal of Psychophysiology*,
 954 98(1), 17–26. <https://doi.org/https://doi.org/10.1016/j.ijpsycho.2015.06.003>

955 Wetzel, N., Buttelmann, D., Schieler, A., & Widmann, A. (2016). Infant and adult pupil

- dilation in response to unexpected sounds. *Developmental Psychobiology*, 58(3), 382–392. <https://doi.org/10.1002/dev.21377>
- Wetzel, N., Scharf, F., & Widmann, A. (2019). Can't Ignore—Distraction by Task-Irrelevant Sounds in Early and Middle Childhood. *Child Development*, 90(6), e819–e830. <https://doi.org/10.1111/cdev.13109>
- Wetzel, N., Schröger, E., & Widmann, A. (2013). The dissociation between the P3a event-related potential and behavioral distraction. *Psychophysiology*, 50(9), 920–930. <https://doi.org/10.1111/psyp.12072>
- Widmann, A., Schröger, E., & Wetzel, N. (2018). Emotion lies in the eye of the listener: Emotional arousal to novel sounds is reflected in the sympathetic contribution to the pupil dilation response and the P3. *Biological Psychology*, 133(January 2017), 10–17. <https://doi.org/10.1016/j.biopsycho.2018.01.010>
- Winkler, I., Denham, S. L., & Nelken, I. (2009). Modeling the auditory scene: predictive regularity representations and perceptual objects. *Trends in Cognitive Sciences*, 13(12), 532–540. <https://doi.org/https://doi.org/10.1016/j.tics.2009.09.003>
- Winkler, I., & Schröger, E. (2015). Auditory perceptual objects as generative models: Setting the stage for communication by sound. *Brain and Language*, 148, 1–22. <https://doi.org/10.1016/j.bandl.2015.05.003>
- Zeelenberg, R., & Bocanegra, B. R. (2010). Auditory emotional cues enhance visual perception. *Cognition*, 115(1), 202–206. <https://doi.org/10.1016/j.cognition.2009.12.004>
- Zekveld, A. A., Koelewijn, T., & Kramer, S. E. (2018). The Pupil Dilation Response to Auditory Stimuli: Current State of Knowledge. *Trends in Hearing*, 22, 1–25. <https://doi.org/10.1177/2331216518777174>