Advancing the Study of Resilience to Daily Stressors

Anthony D. Ong^{1,2*} and Kate Leger³

¹Department of Psychology, Cornell University

²Center for Integrative Developmental Science, Cornell University

³Department of Psychology, University of Kentucky

Author Note

Anthony D. Ong <https://orcid.org/0000-0002-5032-667X>

The authors declare no conflict of interest with respect to the authorship or publication of this article. This article was prepared while the first author was on sabbatical leave at the Department of Medical Social Science, Feinberg School of Medicine, Northwestern University.

Correspondence concerning this article should be addressed to Anthony Ong, Department of Psychology, Cornell University, Ithaca, NY 14853-4401. Contact: anthony.ong@cornell.edu

Accepted manuscript in press at *Perspectives on Psychological Science*

Abstract

Historically, studies of childhood and adult resilience have typically focused on adaptation to chronic life adversities, such as poverty and maltreatment, or isolated and potentially traumatic events, such as bereavement and serious illness. Here, we present a complementary view, suggesting that stressors experienced in daily life may also forecast individual health and well-being. We argue that daily process approaches that incorporate intensive sampling of individuals in natural settings can provide powerful insights into unfolding adaptational processes. In making this argument, we review studies that link within-person dynamics with diverse health-related phenomena. Findings from this research provide support for a multiple-levels-analysis perspective that embraces greater unity in pivotal resilience constructs invoked across childhood and adult literatures. Drawing on insights and principles derived from life-span theory, we conclude by outlining promising directions for future work and considering their broader implications for the field of resilience.

Keywords: daily stress, resilience, dampened reactivity, accelerated recovery, toughening, richness

Advancing the Study of Resilience to Daily Stressors

Resilience has numerous meanings in prior research, but generally refers to the capacity of a dynamic system to adaptively respond to environmental adversity. Inherent in the construct of resilience are two distinct dimensions: exposure to significant risks and evidence of positive adjustment (Luthar & Cicchetti, 2000; Luthar, Cicchetti, & Becker, 2000; Zautra, Hall, & Murray, 2008). A key implication is that resilience is best understood as an active dynamic adaptation to stressors rather than as an inert trait or predisposition. Although indicators of adaptational processes have varied across developmental and ecological contexts, predominant conceptualizations of resilience emphasize three key elements: sustainability, recovery, and steeling. *Sustainability* refers to the maintenance of health and well-being in the face of major life stressors (Bonanno, 2004; Masten, Best, & Garmezy, 1990). *Recovery* refers to how quickly and effectively people bounce back or regain equilibrium following challenge and adversity (Curtis & Cicchetti, 2003; Davidson, 2000). *Steeling* refers to the propensity for prior stressor exposure to increase coping capacity in the face of future stressors (Rutter, 2012; Seery, 2011). Evidence of these core elements of resilience in the context of toxic environmental circumstances (Luthar, 2006) and potentially traumatic events (Bonanno, Westphal, & Mancini, 2011; Seery, 2011) have been well-documented, but the propensity for resilience in the face of naturally occurring day-to-day stressors is not well understood.

In this article, we review current research demonstrating that the phenomenon of resilience is not limited to major life adversities but applies to relatively minor events encountered in daily life. Importantly, this research illustrates how dynamic daily processes can be conceived as resilient factors that describe individuals' inherent capacity for change (Kalisch, Müller, & Tüscher, 2014; Ram & Gerstorf, 2009) and, in turn, are linked to physical health and psychological functioning. Drawing on principles from life-span theory (Baltes, 1987;

Staudinger, Marsiske, & Baltes, 1993), we present a multiple-levels-analysis perspective that takes into account resilience processes operating across different timescales (Fig. 1), including both short-term, *intraindividual variability* and long-term, *intraindividual change* (for definitions of terms in italicized type, see Table 1). We highlight the benefits of *measurement-burst-designs* (Nesselroade, 1991b; Sliwinski, 2008) and point to the unfulfilled potential of existing time series tools (Brose, Neubauer, & Schmiedek, 2021; Hamaker, Asparouhov, Brose, Schmiedek, & Muthen, 2018) for investigating and modeling *interindividual differences* in intraindividual variability (Wang, Hamaker, & Bergeman, 2012). We conclude by discussing remaining questions and future directions, including how daily process inquiries hold great promise for elucidating resilience processes that can inform new targets for intervention research and practice.

Resilience as Dynamic Daily Process

The idea that resilience reflects a dynamic process is not new. Developmental researchers and theorists have long noted that the temporal unfolding nature of resilience implies that it is not a trait or personality characteristic (Luthar, 2006; Luthar et al., 2000; Rutter, 2006). Commenting on this issue, Luthar and colleagues (2000) maintained that a core distinction between resilience and traits that purport to capture resilience is that only the former presupposes a dynamic process and exposure to adversity. Similar concerns have been raised about the use of single-administration trait questionnaires in studies of adult resilience (Bonanno et al., 2011; Kalisch et al., 2017). The overarching message emanating from these influential programs of research is clear: To better understand the adaptational processes that underpin resilience, we need to examine them as they unfold.

The conception of resilience as an unfolding dynamic process necessitates research designs that combine *idiographic* (patterns of variability and change within individuals) and *nomothetic* (patterns of differences across individuals) methods. This hybrid approach is exemplified in the daily process paradigm (Bolger & Zuckerman, 1995; Tennen & Affleck, 2002), which uses *intensive longitudinal methods* (e.g., experience sampling, daily diary assessments) to examine individual differences in the patterning of temporal events and behavior. These methods enhance ecological validity, strengthen causal inference, and minimize recall error. Importantly, process-oriented designs permit the assessment of resilience processes closer to their real-time moments of change.

Applications of Daily Process Formulations of Resilience to Health

While the theoretical significance of daily process designs for resilience research has been recognized (Almeida, 2005; Ong, Bergeman, & Boker, 2009), empirical studies to date have primarily explored how individual differences in personal vulnerabilities and resources predict exposure and responses to daily stressful events (Bolger & Zuckerman, 1995; Zautra, Affleck, Tennen, Reich, & Davis, 2005). More recently, researchers have turned to daily process designs to probe the dynamic processes that give rise to interindividual differences (Charles, Piazza, Mogle, Sliwinski, & Almeida, 2013; Leger, Charles, & Almeida, 2018; Smyth et al., 2018). In this section, we review research demonstrating the value of conceptualizing specific daily *resilience processes* as interindividual difference characteristics (Ram & Gerstorf, 2009). Working at the interface between substantive theory and methodological implementation, we delineate four dynamic processes underlying individual differences in resilience to everyday stressors: *dampened reactivity*, *accelerated recovery*, *toughening* and *inoculation*, and *richness* and *balance*. Each resilience process is described along with illustrative examples of how it is

operationalized and measured in daily life. Studies that explore how these dynamic processes are associated with psychological functioning and physical health are then reviewed (for details of each study, see Table 2) followed by recommendations for future research. Findings from this work demonstrate how the use of within-person, process-oriented methods can serve as powerful tools for illuminating context-specific protective processes (Luthar et al., 2000), thereby helping to identify individuals who are most at risk for maladaptive adjustment and poor health.

Dampened Reactivity

Leading models of stress and health posit that heightened stress reactivity plays a prominent role in the development of psychiatric disorders and disease risk (Epel et al., 2018). While individual differences in physiological responses to standardized laboratory stressors have been widely reported, growing research suggests that individuals may also differ in their reactivity and recovery from naturally occurring stressors. Here we focus on studies of affective responses to daily stressors which characterize the bulk of existing intensive longitudinal studies. Operationally, affective reactivity has been conceptualized as interindividual differences in the degree of within-person coupling of daily stress and affect (Sin, Graham-Engeland, Ong, & Almeida, 2015). From a resilience perspective, these intraindividual parameters measure a continuum of interindividual differences in affective reactivity to daily stressors, ranging from resilience (i.e., *dampened reactivity*) on one end to vulnerability on the other (i.e., *heightened reactivity*).

Measurement. Affective reactivity is typically estimated as the regression coefficient (β_{1j}) in a within-person regression model,

$$\text{Affect}_{ij} = \beta_{0j} + \beta_{1j} (\text{stressor}_{ij}) + r_{ij}$$

where the $stressor_{ij}$ variable is a binary indicator that distinguishes event and non-event occasions. The regression coefficient β_{1j} captures the expected change in affect for person j in response to a same-day stressor event (Ong et al., 2013; Sin et al., 2015).

Associations with mental and physical health. Growing evidence indicates that dampened negative affect (NA) reactivity to daily stressors may be protective against subsequent mental health problems (Bai, Robles, Reynolds, & Repetti, 2020; Charles et al., 2013; Cohen, Gunthert, Butler, O'Neill, & Tolpin, 2005), chronic health conditions (Piazza, Charles, Sliwinski, Mogle, & Almeida, 2013), marital risk (Ong, Gardner, Urganci, Günaydin, & Selçuk, 2020), allostatic load (Piazza, Stawski, & Sheffler, 2019), and even mortality (Chiang, Turiano, Mroczek, & Miller, 2018; Stanton, Selcuk, Farrell, Slatcher, & Ong, 2019). Likewise, the maintenance of positive affect (PA) in the face of daily stressors is associated with more favorable profiles of sleep (Ong et al., 2013), health-related biomarkers (Sin et al., 2015), mental health (Ong & Burrow, 2018; Zhaoyang, Scott, Smyth, Kang, & Sliwinski, 2019), and longevity (Mroczek et al., 2015).

Recommendations. Assessing dynamic resilience processes as stable individual differences requires measures that are reliable and sensitive to within-person change. A challenge in assessing the reliability of person-specific estimates (random slopes) reflecting daily stress reactivity is that it is unknown how many measurement occasions are needed for the individual slope estimates to be accurate and valid measures of interindividual differences. These problems can be addressed by applying a dynamic structural equation modeling approach, where the random effects are treated as latent variables in a general latent variable modeling framework (Asparouhov, Hamaker, & Muthen, 2018; Hamaker et al., 2018). Using simulated data on daily stress reactivity and change in affective distress, Brose et al. (2021) demonstrated that parameter

estimates became closer to the true parameter estimates when using a one-step multilevel structural equation modeling (MSEM) approach compared a two-step approach.

In addition to linking dampened reactivity to changes in well-being, MSEM approaches that incorporate measurement-burst-designs (Nesselroade, 1991b; Sliwinski, 2008) can be used to establish the temporal stability of interindividual differences in daily resilience processes. Lifespan developmentalists have long recognized the importance of longitudinal designs for understanding temporal aspects of development, including intraindividual variability and intraindividual change (Baltes & Nesselroade, 1979; Nesselroade, 1991a; Wohlwill, 1973). Figure 1 illustrates a measurement-burst design that consists of intensive ‘bursts’ of measurements obtained over microtime scales (e.g., hours, days, weeks) from a single individual and that are repeated over macrotime scales (e.g., years, decades). The magnified circles A, B, C, and D depict four dynamic attributes of resilience (i.e., dampened reactivity, accelerated recovery, toughness/inoculation, and richness/balance) that theoretically capture interindividual differences in capacity or potential for change. By comparison, the solid line connecting the bursts represent long-term intraindividual change processes (e.g., development) that accrue with advancing age. Employing two-wave measurement burst data from the National Study of Daily Experiences (NSDE), Rush, Rast, Almeida, and Hofer (2019) reported a significant average within-person association in stress reactivity across bursts, albeit with considerable variability in the strength of the association within each burst. Using a MSEM approach, future work could benefit from assessing the timing of effects or temporal specificity of associations between dampened reactivity and well-being and the extent to which they exert reciprocal effects on each other (Brose et al., 2021; Rush et al., 2019).

Accelerated Recovery

Beyond reactivity, recent theory and research suggest that interindividual differences in the rate of affective recovery from daily stressors may also have implications for long-term health. Whereas *dampened reactivity* reflects the magnitude of responses to stress, *accelerated recovery* reflects the speed with which stress responses return to baseline (Epel et al., 2018).

Measurement. Affective recovery can be estimated as the regression coefficient (β_{1j}) in a within-person regression model,

$$\text{Affect}_{ij} = \beta_{0j} + \beta_{1j} (\text{stressor}_{ij-1}) + r_{ij}$$

where the stressor_{ij} variable is a binary indicator that distinguishes event and non-event occasions. The regression coefficient β_{1j} captures the expected change in affect for person j in response to a stressor event experienced on the previous day (Leger et al., 2018).

Associations with mental and physical health. Like differences in stressor reactivity, individuals differ in the rate or speed with which they recover from daily stressors. Using a daily-burst design (i.e., 8-day diary study nested within a 10-year longitudinal study design), Leger, Charles, and Almeida (2018) demonstrated that temporary or short-lived NA in response to daily stressors was associated with fewer numbers of chronic conditions and lower functional impairment 10 years later. Similarly, Bergeman and Deboeck (2014) found the interindividual differences in the rate of stress reduction or dissipation was inversely associated with depressive symptoms over a 5-year period. These findings demonstrate the unique contribution of daily process studies to elucidating dynamic resilience processes (e.g., dampened reactivity and accelerated recovery). Collectively, these studies show how daily study designs can be incorporated into longitudinal studies to make inferences about within-person dynamics, thereby generating more highly predictive models of stress and health (Epel et al., 2018). Critically, the

scope of these investigations offers insights into daily processes that simply could not have been ascertained from traditional trait reports of resilience assessed at a single point in time (Kalisch et al., 2017).

Recommendations. Dynamic operationalizations of stress recovery depend crucially on the length of time between measurements. Linking retrospective reports of daily stress and affect over an entire day may thus obscure recovery processes that manifest across relatively faster timescales (e.g., minutes, hours). More frequent measurement bursts assessed at shorter intervals would therefore permit a better understanding of stress recovery processes as they naturally occur in daily life (Hamaker, Ceulemans, Grasman, & Tuerlinckx, 2015). More generally, daily process studies of stress reactivity and recovery should be broadened to include measures of health that go beyond self-report (Gordon & Mendes, 2021; Leger et al., 2018). Experimental studies have assessed physiological recovery from laboratory-based stressors and their links to physical health (e.g., Panaite, Salomon, Jin, & Rottenberg, 2015; Steptoe & Marmot, 2005). Future studies that combine experimental manipulations, intensive longitudinal designs, and objective indices of physical health may yield new insights into the dynamic mechanisms involved in accelerated recovery from daily stressors. Finally, from the perspective of psychological traits, consistency in idiographic structure (i.e., intraindividual variability patterns defined across time and situations) is fundamental to understanding individual differences in personality (Beck & Jackson, 2020; Shoda, Mischel, & Wright, 1994). Yet evidence for temporal stability and cross-situational consistency in stressor reactivity and recovery have not been established, and therefore constitute an important future research direction. Further, the potential contaminating influence of personality traits, such as neuroticism, on the relation between affective reactivity/recovery from daily stressors and health has received scant attention (but see Sin et al.,

2015; Stanton et al., 2019). Hence, an important methodological issue for future studies of interindividual differences in resilience processes and health is whether associations are independent of neuroticism and allied personality traits.

Toughening and Inoculation

Although much work has focused on resilience as the capacity to absorb and recover from stressful events, there is growing evidence that stressful experiences themselves may also contribute to the capacity for resilience (Seery, Holman, & Silver, 2010). This conceptualization of resilience holds that stressors that are challenging, but manageable, can play an adaptive role in preparing individuals for coping with later stressors, a protective phenomenon referred to as *toughening* or *inoculation* (Dienstbier, 1989; Meichenbaum, 1993). Most importantly, toughening and inoculation effects are not limited to major life adversities, but may also influence adaptation to minor daily stressors. DiCorcia and Tronick (2011) reviewed developmental research showing that successful regulation of everyday stressors, scaffolded by caregiver reparatory sensitivity, prepares infants for coping with subsequent stressors. Seery and Quinton (2016) reviewed social psychological evidence demonstrating U-shaped relationships between daily stressor exposure and well-being. While not directly employing intensive longitudinal designs, this research underscores the potential of daily process studies to advance understanding of the protective benefits or toughening qualities of daily stressor exposures that until now have been ascribed to cumulative lifetime adversities (Seery et al., 2010).

Measurement. One component of everyday stressor exposure that may be of particular importance for health is the effect of *stressor pile-up* in daily life (Schilling & Diehl, 2014; Smyth et al., 2018). Researchers interested in stressor pile-up have used various indices to quantify patterns of stressor accumulation (e.g., frequency counts of daily stressors, number of

stressor reactivity-recover cycles). Following Schilling et al. (2014), stressor pile-up can be expressed with the following equation:

$$A_{(k)ti} = \frac{\sum_{j=1}^k (k - j + 1) S_{(t-j)i}}{\sum_{j=1}^k j}$$

where $A_{(k)ti}$ is an index of stressor pile-up across k days for individual i , and k specifies the number of days that precede day t (Schilling & Diehl, 2014).

Associations with mental and physical health. Research suggest that stressor pile-up is a common daily phenomenon (Almeida, Wethington, & Kessler, 2002; Bolger, DeLongis, Kessler, & Schilling, 1989) that can have adverse consequences for mental health and well-being, especially in the short term (Bolger & Schilling, 1991; Grzywacz & Almeida, 2008; Serido, Almeida, & Wethington, 2004). Using data from a 30-day diary study, Schilling et al. (2014) found that stressor pile-up over the course of a week had an independent effect on daily NA above and beyond the effect of concurrent daily stress. In a coordinated analysis of data from two Ecological Momentary Assessment (EMA) studies, Almeida and colleagues (2020) reported that greater stressor pile-up was more strongly associated with physical activity compared to reactivity and recovery. Extending this research to a clinical sample, Smith et al. (2021) found that the cumulative build-up of stressors over recent hours predicted greater subsequent binge-eating symptoms among adults with binge-eating disorder.

Recommendations. Although exposure to mild everyday stressors has been theorized to foster resilience (Seery & Quinton, 2016), this has yet to be tested empirically using intensive longitudinal data. To the extent that exposure to daily stressors builds resilience through toughness and inoculation, individual differences in daily stressor accumulation and pile-up should demonstrate U-shape relationships with well-being, with moderate levels of stressor pile-

up (relative to no or high levels) contributing to improved health over time. Charles et al. (2021) provide some evidence consistent with this premise, finding among a sample of adults that leading a stress-free life, while associated with higher emotion well-being, may be linked to lower cognitive functioning. A potential fruitful direction for future work would be to explicitly test curvilinear relationships between daily stressor pile-up and subsequent health.

To date, almost all investigations of daily stress processes have created aggregate measures of individual differences and then drawn inferences about more dynamic processes underlying psychological (mal)adjustment. In contrast, the study by Almeida et al. (2020) used a within-person approach to capture everyday stress processes and demonstrated substantial variation in the temporal patterning of stressor pile-up both within and across days. This approach represents a significant advance in the assessment of stressor pile-up as it allows researchers to begin to explicitly test resilience processes (e.g., toughening and inoculation) across different timescales, as they unfold in real-time and in individuals' natural environments. Do low-to-moderate levels of stressor pile-up reflect adaptive flexibility, such that exposure to some stress in daily life is more likely to provide opportunities to develop toughness than exposure to either no stress or high stress? Do toughening mechanisms (e.g., mastery, perceived control and belief in the ability to cope) that have been theorized to account for resilience in the face of major life adversities (Seery et al., 2010) also explain how and why exposure to minor daily hassles may be beneficial to overall mental health and well-being? To date, no studies have systematically examined these questions.

Richness and Balance

Distinguished from cumulative stressor counts are indices that capture *stressor diversity* or the *richness* and *balance* of “hassles” (e.g., home chores, work deadlines, interpersonal

tensions) across multiple domains of daily life. Consistent with a conservation model of stress (Hobfoll, 1989), high stressor diversity theoretically functions as a resource that confers differential well-being (Koffer, Ram, Conroy, Pincus, & Almeida, 2016). Following techniques used in the natural sciences to assess the biodiversity of ecosystems (Magurran, 2004; Morin, 1999), measures of diversity have been used to assess a variety of social and psychological phenomena, including racial and ethnic diversity (Budescu & Budescu, 2012), behavioral flexibility (Ram, Conroy, Pincus, Hyde, & Molloy, 2012), population genetics (Sherwin, 2010), community social networks (Li, Zhang, Feng, & Wu, 2015), emotional diversity (Ong, Benson, Zautra, & Ram, 2018; Quoidbach et al., 2014; Urban-Wojcik et al., 2020), and activity diversity (Lee et al., 2018; Lee, Urban-Wojcik, Charles, & Almeida, 2021).

Measurement. Stressor diversity can be estimated using Shannon's (1948) entropy:

$$SD_i = -\left(\frac{1}{\ln(m)}\right) \sum_{j=1}^m p_{ij} \ln p_{ij}$$

where SD_i is an index that quantifies the relative variety or *richness* and evenness or *balance* in stressor experiences (j) across all study days for individual i (Koffer et al., 2016).

Associations with mental and physical health. Employing data from two independent diary studies, Koffer and colleagues (Koffer, Ram, & Almeida, 2018; Koffer et al., 2016) used an entropy index to quantify the dispersion of daily stressors across multiple domains (e.g., health, financial, work, interpersonal) and found that higher daily stressor diversity was associated with lower NA and weaker links between daily stressor exposure and NA. Using EMA data, Koffer, Dickman, and Kamarck (2021) reported that lower stressor diversity (i.e., higher number of stressors concentrated in one domain) coupled with higher stressor exposure was associated with higher diastolic blood pressure in a sample of middle-aged adults.

Recommendations. The number of stressor events sampled may affect the interpretation of stressor diversity and its association with health and well-being. Coarse assessments of individuals' overall stressor ecosystems could restrict the degree to which richness and balance of stressful experiences are adequately captured (see Brown & Coyne, 2017). Thus, future research should examine whether the number of stressors assessed influences the rank order of stressor diversity scores (see Benson, Ram, Almeida, Zautra, & Ong, 2018). Further, while extant work has focused on the immediate consequences of stressor richness and balance, it is plausible that exposure to many types of daily stressors that are appraised as manageable rather than overwhelming may also contribute to the propensity for resilience to future stressors, be they major life adversities or minor daily hassles (Seery & Quinton, 2016). Finally, methods used to operationalize stressor diversity, such as Shannon's entropy, assume that stressor events are independent and identically distributed across time (Ram & Gerstorf, 2009). This assumption is untenable, however, when stressors are conceptualized as continuous phenomena. One such class of stressors are chronic strains which represent unresolved, recurrent demands that people face in their daily lives (Pearlin & Skaff, 1996). Here, alternative diversity indices that incorporate heterogeneity in time-series data, such as measures of *turbulence* (Koffer et al., 2016), may be used to quantify the amount and distribution of individuals' daily stressor exposure.

Examples of turbulent variation abound in nature. Unchecked by natural controls, invasive species, for example, can spread quickly and displace native plants, animals, and other organisms, causing dramatic biodiversity loss and ecosystem degradation. In a similar vein, differential exposure to turbulent (stochastic) daily stressors may disrupt the continuity of everyday life and, over time, threaten the health and functioning of the emotional ecosystem

(Quoidbach et al., 2014). By contrast, low turbulent stressor ecologies may be those characterized by greater homogeneity (predictability) in patterns of daily stressor exposure (Ram, Brinberg, Pincus, & Conroy, 2017).

Integrating Life-Span Principles in the Study of Resilience

As the preceding discussions suggest, resilience is a heterogeneous construct that encompasses an array of dynamic processes including dampened reactivity, rapid recovery, moderate stressor exposure, and high stressor diversity. We have argued that a daily process conceptualizations of resilience may reveal the adaptive ways in which individuals respond to stressors in everyday life, and thereby complement traditional formulations of resilience that primarily focus on major life adversities (Luthar et al., 2000; Zautra et al., 2008). In this final section, we summarize key principles and concepts from life-span theory and their implications for advancing the study of resilience in daily life. Although some of these concepts are inherent in previous models of resilience, a life-span perspective provides an interpretive framework for understanding variation in resilient functioning (Infurna, 2021; Staudinger et al., 1993).

Multidirectionality and multidimensionality

A key insight from lifespan theory is that the development of adaptive capacities throughout life are characterized by the simultaneous unfolding of increases, decreases, and maintenance in functioning. Furthermore, development is a process that spans multiple domains, such that decrements in one domain may co-exist with stability or even increments in other domains. Taken together, these *multidirectionality* and *multidimensionality* perspectives may help to explain the heterogeneity in functioning across domains observed among those labeled as resilient (Luthar, 2006). Indeed, evidence of discordance between behavior and physiological functioning have led some scholars to question whether resilience is a veridical construct or one

that is only “skin-deep” (Brody et al., 2013). Yet as developmentalists have pointed out, resilience is not an “across-the-board phenomenon” (Infurna & Luthar, 2017; Luthar, 2006), and adaptation across diverse spheres of functioning is never uniform but manifest in co-occurring profiles of successive gains and losses (Staudinger et al., 1993).

Recommendations. Translating multidirectional and multidimensional conceptions of adaptation to empirically tractable questions is critical for advancing the study of resilience in daily life. To what extent do daily manifestations of dampened reactivity, accelerated recovery, toughening, and stressor richness and balance change across the life span? How does the timing, direction, and rate of change in these resilience processes differ before and after adversity? Do affective and non-affective domains of resilience in daily life constitute related, but distinct, processes (Nezlek, 2005)? While there is growing recognition of the multidirectional and multidimensional nature of resilience to major life stressors (Infurna & Luthar, 2017), little is known about how resilience processes cohere across multiple domains (e.g., affective, cognitive, physiological) in daily life. Furthermore, to date, the resilience literature has largely focused on single outcomes, which prevents a comparison of trajectories of adaptation within and across different domains of functioning (Infurna & Luthar, 2017). How to relate resilience processes that manifest on microtime scales (e.g., hours, days) to “varieties” in resilient outcomes (Ryff et al., 2012) that unfold over macrotime scales (e.g., years, decades)? The answers to these questions await further investigation.

Plasticity and reserve capacity

From a lifespan perspective, whether individuals are able to sustain, recover, or benefit from adversity depends critically on the degree of adaptive potential or *plasticity* (Staudinger et al., 1993). Further, the extent to which such adaptive plasticity protects against future stressors is

reflected in the level and accumulation of latent capabilities or *reserve capacity* (Cullati, Kliegel, & Widmer, 2018). Although the concepts of plasticity and reserve capacity have been recognized in life course models of resilience (Gallo, de los Monteros, & Shivpuri, 2009; Ryff & Singer, 2008), their explicit connections to resilience in daily life have not been addressed.

Recommendations. Incorporating life-span principles more systematically into daily process research can enhance understanding of resilience to everyday stressors. Is plasticity or within-person variability in short-term resilience processes (e.g., dampened reactivity, toughening) modifiable through intervention (Baltes, 1987; Staudinger et al., 1993)? When considering the timing of potential interventions, is midlife a critical, more malleable period in the life course for examining resilience in the face of challenge or the potential for change and plasticity (Infurna, 2021)? If so, what effect will preventive interventions in midlife have on catalyzing future benefits in the form of greater reserve capacity in later life? Pursuit of these questions may lead to important insights into how intensive longitudinal designs can be incorporated into randomized controlled studies of resilience to determine causal within-person mechanisms (Hamaker & Wichers, 2017).

Conclusion

The study of resilience in everyday life offers critically needed complements to existing research on potential trauma and extreme adversity (Bonanno et al., 2011). In this review, we have discussed the utility of four idiographic indices (i.e., dampened reactivity, accelerated recovery, toughness/inoculation, and richness/balance), as well as key principles of lifespan development (i.e., multidirectionality, multidimensionality, plasticity, and reserve capacity) that warrant greater attention in daily process studies of resilience. Continued research in this area will deepen understanding of the mechanisms by which individuals' inherent capacity for

change, conceived as dynamic daily processes, exert their health-promoting effects. Beyond dynamic models that depict life as it is lived, daily process research holds great promise for unifying diverse formulations of resilience across the child and adult literatures (e.g., sustainability, recovery, steeling), thereby affording greater insight into what it means to be well in the face of adversity.

References

- Almeida, D. M. (2005). Resilience and vulnerability to daily stressors assessed via diary methods. *Current Directions in Psychological Science*, 14, 64-68.
- Almeida, D. M., Marcusson-Clavertz, D., Conroy, D. E., Kim, J., Zawadzki, M. J., Sliwinski, M., & Smyth, J. M. (2020). Everyday stress components and physical activity: examining reactivity, recovery and pileup. *Journal of Behavioral Medicine*, 43, 108-120.
- Almeida, D. M., Wethington, E., & Kessler, R. C. (2002). The Daily Inventory of Stressful Events: An interview-based approach for measuring daily stressors. *Assessment*, 9, 41-55.
- Asparouhov, T., Hamaker, E. L., & Muthen, B. (2018). Dynamic structural equation models. *Structural Equation Modeling*, 25, 359–388.
- Bai, S., Robles, T. F., Reynolds, B. M., & Repetti, R. L. (2020). Daily mood reactivity to stress during childhood predicts internalizing problems three years later. *Journal of Abnormal Child Psychology*, 48, 1063-1075.
- Baltes, P. B. (1987). Theoretical propositions of life-span developmental psychology: On the dynamics between growth and decline. *Developmental Psychology*, 23(5), 611-626.
doi:10.1037/0012-1649.23.5.611
- Baltes, P. B., & Nesselroade, J. R. (1979). History and rationale of longitudinal research. In J. R. Nesselroade & P. B. Baltes (Eds.), *Longitudinal research in the study of behavior and development* (pp. 1–39). New York: Academic Press.
- Beck, E. D., & Jackson, J. J. (2020). Idiographic traits: A return to Allportian approaches to personality. *Current Directions in Psychological Science*, 29, 301-308.
- Benson, L., & Ram, N. (2018). Biodiversity metrics in lifespan developmental methodology. In *Oxford Research Encyclopedia of Psychology*: Oxford University Press.

- Benson, L., Ram, N., Almeida, D., Zautra, A., & Ong, A. D. (2018). Fusing biodiversity metrics into investigations of daily life and healthy aging: Recommendations and illustrations with emodiversity. *The Journals of Gerontology, Series B: Psychological Sciences and Social Sciences*, 73, 75-86.
- Bergeman, C. S., & Deboeck, P. R. (2014). Trait stress resistance and dynamic stress dissipation on health and well-being: The reservoir model. *Research in Human Development*, 11, 108–125.
- Bolger, N., DeLongis, A., Kessler, R. C., & Schilling, E. A. (1989). Effects of daily stress on negative mood. *Journal of Personality and Social Psychology*, 57(5), 808-818.
- Bolger, N., & Schilling, E. A. (1991). Personality and the problems of everyday life: The role of neuroticism in exposure and reactivity to daily stressors. *Journal of Personality*, 59(3), 355-386.
- Bolger, N., & Zuckerman, A. (1995). A framework for studying personality in the stress process. *Journal of Personality and Social Psychology*, 69(5), 890-902.
- Bonanno, G. A. (2004). Loss, trauma, and human resilience: Have we underestimated the human capacity to thrive after extremely aversive events? *American Psychologist*, 59, 20-28.
- Bonanno, G. A., Westphal, M., & Mancini, A. D. (2011). Resilience to loss and potential trauma. *Annual Review of Clinical Psychology*, 7, 511-535.
- Brody, G. H., Yu, T., Chen, E., Miller, G. E., Kogan, S. M., & Beach, S. R. H. (2013). Is resilience only skin deep?: Rural African Americans' socioeconomic status–related risk and competence in preadolescence and psychological adjustment and allostatic load at age 19. *Psychological Science*, 24, 1285-1293.

- Brose, A., Neubauer, A. B., & Schmiedek, F. (2021). Integrating state dynamics and trait change: A tutorial using the example of stress reactivity and change in well-being. *European Journal of Personality*.
- Brown, N., & Coyne, J. C. (2017). Emodiversity: Robust predictor of outcomes or statistical artifact? *Journal of Experimental Psychology: General*, 146, 1372–1377.
- Budescu, D. V., & Budescu, M. (2012). How to measure diversity when you must. *Psychological Methods*, 17, 215–227.
- Charles, S. T., Mogle, J., Chai, H. W., & Almeida, D. M. (2021). The mixed benefits of a stressor-free life. *Emotion*.
- Charles, S. T., Piazza, J. R., Mogle, J., Sliwinski, M. J., & Almeida, D. M. (2013). The wear-and-tear of daily stressors on mental health. *Psychological Science*, 24, 733–741.
- Chiang, J. J., Turiano, N. A., Mroczek, D. K., & Miller, G. E. (2018). Affective reactivity to daily stress predicts 20-year mortality risk in adults with chronic illness: Findings from the national study of daily experience. *Health Psychology*, 37, 170-178.
- Cohen, J. H., Gunthert, K. C., Butler, A. C., O'Neill, S. C., & Tolpin, L. H. (2005). Daily reactivity as a prospective predictor of depressive symptoms. *Journal of Personality*, 73, 1687-1713.
- Cullati, S., Kliegel, M., & Widmer, E. (2018). Development of reserves over the life course and onset of vulnerability in later life. *Nature Human Behavior*, 2, 551-558.
- Curtis, W., & Cicchetti, D. (2003). Moving research on resilience into the 21st century: Theoretical and methodological considerations in examining the biological contributors to resilience. *Development and Psychopathology*, 15, 773-810.

- Davidson, R. J. (2000). Affective style, psychopathology, and resilience: Brain mechanisms and plasticity. *American Psychologist*, 55, 1196-1214.
- DiCorcia, J. A., & Tronick, E. (2011). Quotidian resilience: Exploring mechanisms that drive resilience from a perspective of everyday stress and coping. *Neuroscience & Biobehavioral Reviews*, 35, 1593–1602.
- Dienstbier, R. A. (1989). Arousal and physiological toughness: Implications for mental and physical health. *Psychological Review*, 96, 84–100.
- Epel, E. S., Crosswell, A. D., Mayer, S. E., Prather, A. A., Slavich, G. M., Puterman, E., & Mendes, W. B. (2018). More than a feeling: A unified view of stress measurement for population science. *Frontiers in Neuroendocrinology*, 49, 146–169.
- Gallo, L. C., de los Monteros, K. E., & Shivpuri, S. (2009). Socioeconomic status and health: what is the role of reserve capacity? . *Current Directions in Psychological Science*, 18, 269–274.
- Gordon, A. M., & Mendes, W. B. (2021). A large-scale study of stress, emotions, and blood pressure in daily life using a digital platform. *Proceedings of the National Academy of Sciences*.
- Grzywacz, J. G., & Almeida, D. M. (2008). Stress and binge drinking: A daily process examination of stressor pile-up and socioeconomic status in affect regulation. *International Journal of Stress Management*, 15, 364–380.
- Hamaker, E. L., Asparouhov, T., Brose, A., Schmiedek, F., & Muthen, B. (2018). At the frontiers of modeling intensive longitudinal data: Dynamic structural equation models for the affective measurements from the COGITO study. *Multivariate Behavioral Research*, 53, 820–841.

- Hamaker, E. L., Ceulemans, E., Grasman, R., & Tuerlinckx, F. (2015). Modeling affect dynamics: State of the art and future challenges. *Emotion Review*, 7, 316-322.
- Hamaker, E. L., & Wichers, M. (2017). No time like the present: Discovering the hidden dynamics in intensive longitudinal data. *Current Directions in Psychological Science*, 26, 10-15.
- Hobfoll, S. E. (1989). Conservation of resources: A new attempt at conceptualizing stress. *American Psychologist*, 44, 513-524.
- Infurna, F. J. (2021). Utilizing principles of life-span developmental psychology to study the complexities of resilience across the life span. *The Gerontologist*, 61, 807-818.
- Infurna, F. J., & Luthar, S. S. (2017). The multidimensional nature of resilience to spousal loss. *Journal of Personality and Social Psychology*, 112, 926-947.
- Kalisch, R., Baker, D. G., Basten, U., Boks, M. P., Bonanno, G. A., Brummelman, E., . . . Kleim, B. (2017). The resilience framework as a strategy to combat stress-related disorders. *Nature Human Behavior*, 1, 784-790
- Kalisch, R., Müller, M. B., & Tüscher, O. (2014). A conceptual framework for the neurobiological study of resilience. *Behavioral and Brain Sciences* 1-79.
- Koffer, R. E., Dickman, K., & Kamarck, T. (2021). High stressor exposure and low stressor diversity are linked to high blood pressure across age. *Unpublished manuscript*.
- Koffer, R. E., Ram, N., & Almeida, D. M. (2018). More than counting: An intraindividual approach to categorical repeated measures. *Journal of Gerontology: Psychological Sciences and Social Sciences*, 73, 87-99.

- Koffer, R. E., Ram, N., Conroy, D. E., Pincus, A. L., & Almeida, D. M. (2016). Stressor diversity: Introduction and empirical integration into the daily stress model. *Psychology and Aging, 31*, 301-320.
- Lee, S., Koffer, R. E., Sprague, B. N., Charles, S. T., Ram, N., & Almeida, D. M. (2018). Activity diversity and its associations with psychological well-being across adulthood. *Journal of Gerontology: Psychological Sciences, 73*, 985-995.
- Lee, S., Urban-Wojcik, E. J., Charles, S. T., & Almeida, D. M. (2021). Rich and balanced experiences of daily emotions are associated with activity diversity across adulthood. *Journal of Gerontology: Psychological Sciences and Social Sciences*.
- Leger, K. A., Charles, S. T., & Almeida, D. M. (2018). Letting it go: Lingering negative affect in response to daily stressors is associated with physical health in later years. *Psychological Science, 29*, 1283-1290.
- Li, Y., Zhang, G., Feng, Y., & Wu, C. (2015). An entropy-based social network community detecting method and its application to scientometrics. *Scientometrics 102*, 1003-1017.
- Luthar, S. S. (2006). Resilience in development: A synthesis of research across five decades. In D. Cicchetti & D. Cohen (Eds.), *Developmental psychopathology: Risk, disorder, and adaptation* (2 ed., Vol. 3). Hoboken: John Wiley & Sons.
- Luthar, S. S., & Cicchetti, D. (2000). The construct of resilience: Implications for interventions and social policies. *Development and Psychopathology, 12*, 857-885.
- Luthar, S. S., Cicchetti, D., & Becker, B. (2000). The construct of resilience: A critical evaluation and guidelines for future work. *Child Development, 71*, 543-562.
- Magurran, A. E. (2004). *Measuring biology diversity*. Malden: Blackwell Publishing.

- Masten, A. S., Best, K. M., & Garmezy, N. (1990). Resilience and development: Contributions from the study of children who overcome adversity. *Development and Psychopathology*, 2, 425-444.
- Meichenbaum, D. (1993). Stress inoculation training: A twenty year update. In R. L. Woolfolk & P. M. Lehrer (Eds.), *Principles and practices of stress management* (2nd ed., pp. 373–406). New York: Guilford Press.
- Morin, P. J. (1999). *Community ecology*. Malden: Blackwell Publishing.
- Mroczek, D. K., Stawski, R. S., Turiano, N. A., Chan, W., Almeida, D. A., Neupert, S. D., & Spiro III, A. (2015). Emotional reactivity predicts mortality: Longitudinal findings from the VA Normative Aging Study. *Journal of Gerontology: Psychological Sciences and Social Sciences*, 70, 398–406.
- Nesselroade, J. R. (1991a). Interindividual differences in intraindividual change. In L. M. Collins & J. L. Horn (Eds.), *Best methods for the analysis of change: Recent advances, unanswered questions, future directions* (pp. 92-105). Washington, DC: American Psychological Association.
- Nesselroade, J. R. (1991b). The warp and woof of the developmental fabric. In R. M. Downs, L. S. Liben, & D. S. Palermo (Eds.), *Visions of aesthetics, the environment & development: The legacy of Joachim F. Wohlwill* (pp. 214-242). Hillsdale: Erlbaum.
- Nezlek, J. B. (2005). Distinguishing affective and non-affective reactions to daily events. *Journal of Personality*, 73, 1539–1568.
- Ong, A. D., Benson, L., Zautra, A., & Ram, N. (2018). Emodiversity and biomarkers of inflammation. *Emotion*, 18, 3-14.

- Ong, A. D., Bergeman, C. S., & Boker, S. M. (2009). Resilience comes of age: Defining features in later adulthood. *Journal of Personality*, 77(6), 1777-1804.
- Ong, A. D., & Burrow, A. L. (2018). Affective reactivity to daily racial discrimination as a prospective predictor of depressive symptoms in African American graduate and post-graduate students. *Development and Psychopathology*, 30, 1649-1659.
- Ong, A. D., Exner-Cortens, D., Riffin, C., Steptoe, A., Zautra, A. J., & Almeida, D. M. (2013). Linking stable and dynamic features of positive affect to sleep. *Annals of Behavioral Medicine*, 46, 52-61.
- Ong, A. D., Gardner, S., Urganci, B., Günaydın, G., & Selçuk, E. (2020). Affective reactivity, resting heart rate variability, and marital quality: A 10-year longitudinal study of U.S. adults. *Journal of Family Psychology*, 34, 375-882.
- Panaite, V., Salomon, K., Jin, A., & Rottenberg, J. (2015). Cardiovascular recovery from psychological and physiological challenge and risk for adverse cardiovascular outcomes and all-cause mortality. *Psychosomatic Medicine*, 77, 215–226.
- Pearlin, L. I., & Skaff, M. M. (1996). Stress and the life course: A paradigmatic alliance. *Gerontologist*, 36(2), 239-247.
- Piazza, J. R., Charles, S. T., Sliwinski, M. J., Mogle, J., & Almeida, D. M. (2013). Affective reactivity to daily stressors and long-term risk of reporting a chronic physical health condition. *Annals of Behavioral Medicine*, 45, 110-120.
- Piazza, J. R., Stawski, R. S., & Sheffler, J. L. (2019). Age, daily stress processes, and allostatic load: A longitudinal study. *Journal of Aging and Health*, 31, 1671-1691.

Quoidbach, J., Gruber, J., Mikolajczak, M., Kogan, A., Kotsou, I., & Norton, M. (2014).

Emodiversity and the emotional ecosystem. *Journal of Experimental Psychology: General*, *143*, 2057-2066.

Ram, N., Brinberg, M., Pincus, A. L., & Conroy, D. E. (2017). The questionable ecological validity of ecological momentary assessment: Considerations for design and analysis. *Research in Human Development*, *14*, 253-270.

Ram, N., Conroy, D. E., Pincus, A. L., Hyde, A. L., & Molloy, L. (2012). Tethering theory to method: Using measures of intraindividual variability to operationalize individuals' dynamic characteristics. In G. Hancock & J. Harring (Eds.), *Advances in longitudinal modeling in the social and behavioral sciences* (pp. 81–110). Charlotte: Information Age Publishing.

Ram, N., & Gerstorf, D. (2009). Time-structured and net intraindividual variability: Tools for examining the development of dynamic characteristics and processes. *Psychology and Aging*, *24*, 778-791. doi:10.1037/a0017915

Rush, J., Rast, P., Almeida, D. M., & Hofer, S. M. (2019). Modeling long-term changes in daily within-person associations: An application of multilevel SEM. *Psychology and Aging*, *34*, 163-176.

Rutter, M. (2006). Implications of resilience concepts for scientific understanding. *Annals of the New York Academy of Sciences*, *1094*, 1-12.

Rutter, M. (2012). Resilience as a dynamic concept. *Development and Psychopathology*, *24*, 335-344.

Ryff, C. D., Friedman, E., Fuller-Rowell, T., Love, G., Miyamoto, Y., Morozink, J., . . .

Tsenkova, V. (2012). Varieties of resilience in midus. *Social and Personality Psychology Compass*, 6, 792-806.

Ryff, C. D., & Singer, B. (2008). Thriving in the face of challenge: The integrative science of human resilience. In F. Kessel, P. L. Rosenfield, & N. B. Anderson (Eds.),

Interdisciplinary research: Case studies from health and social science. (pp. 198-227).

New York, NY, US: Oxford University Press.

Schilling, O. K., & Diehl, M. (2014). Reactivity to stressor pile-up in adulthood: Effects on daily negative and positive affect. *Psychology and Aging*, 29, 72-83.

Seery, M. D. (2011). Resilience: A silver lining to experiencing adverse life events. *Current Directions in Psychological Science*, 20, 390-394.

Seery, M. D., Holman, E. A., & Silver, R. C. (2010). Whatever does not kill us: Cumulative lifetime adversity, vulnerability, and resilience. *Journal of Personality and Social Psychology*, 99, 1025–1041.

Seery, M. D., & Quinton, W. J. (2016). Understanding resilience: From negative life events to everyday stressors. *Advances in Experimental Social Psychology*, 54, 181-245.

Serido, J., Almeida, D. M., & Wethington, E. (2004). Chronic stressors and daily hassles: Unique and interactive relationships with psychological distress. *Journal of Health and Social Behavior*, 45, 17-33.

Shannon, C. (1948). A mathematical theory of communication. *The Bell System Technical Journal*, 27, 379 – 423.

Sherwin, W. B. (2010). Entropy and information approaches to genetic diversity and its expression: Genomic geography. *Entropy*, 12, 1765–1798.

- Shoda, Y., Mischel, W., & Wright, J. C. (1994). Intraindividual stability in the organization and patterning of behavior: Incorporating psychological situations into the idiographic analysis of personality. *Journal of Personality and Social Psychology*, 67, 674-687.
- Sin, N. L., Graham-Engeland, J. E., Ong, A. D., & Almeida, D. M. (2015). Positive and negative affective responses to daily stressors are associated with inflammation. *Health Psychology*, 34, 1154-1165.
- Sliwinski, M. (2008). Measurement-burst designs for social health research. *Social and Personality Psychology Compass*, 2, 245–261.
- Smith, K. E., Mason, T. B., Schaefer, L. M., Anderson, L. M., Critchley, K., Crosby, R. D., . . . Peterson, C. B. (2021). Dynamic stress responses and real-time symptoms in binge-eating disorder. *Annals of Behavioral Medicine*, 55, 758-768.
- Smyth, J. M., Sliwinski, M. J., Zawadzki, M. J., Scott, S. B., Conroy, D. E., Lanza, S. T., & Almeida, D. M. (2018). Everyday stress response targets in the science of behavior change. *Behaviour Research and Therapy*, 101, 20 –29.
- Stanton, S. C., Selcuk, E., Farrell, A. K., Slatcher, R. B., & Ong, A. D. (2019). Perceived partner responsiveness, daily negative affect reactivity, and all-cause mortality: A 20-year longitudinal study. *Psychosomatic Medicine*, 81, 7-15.
- Staudinger, U. M., Marsiske, M., & Baltes, P. B. (1993). Resilience and levels of reserve capacity in later adulthood: Perspectives from life-span theory. *Development and Psychopathology*, 5(4), 541-566. doi:10.1017/S0954579400006155
- Steptoe, A., & Marmot, M. G. (2005). Impaired cardiovascular recovery following stress predicts 3-year increases in blood pressure. *Journal of Hypertension*, 23, 529-536.

- Tennen, H., & Affleck, G. (2002). The challenge of capturing daily processes at the interface of social and clinical psychology. *Journal of Social and Clinical Psychology, 21*(6), 610-627.
- Urban-Wojcik, E. J., Mumford, J. A., Almeida, D. M., Lachman, M. E., Ryff, C. D., Davidson, R. J., & Schaefer, S. M. (2020). Emodiversity, health, and well-being in the Midlife in the United States (MIDUS) daily diary study. *Emotion*. doi:10.1037/emo0000753
- Wang, L., Hamaker, E. L., & Bergeman, C. S. (2012). Investigating inter-individual differences in short-term intra-individual variability. *Psychological Methods, 17*, 567-581.
- Wohlwill, J. F. (1973). *The study of behavioral development*. New York: Academic Press.
- Zautra, A. J., Affleck, G. G., Tennen, H., Reich, J. W., & Davis, M. C. (2005). Dynamic approaches to emotions and stress in everyday life: Bolger and Zuckerman reloaded with positive as well as negative affects. *Journal of Personality, 76*, 1511-1538.
- Zautra, A. J., Hall, J. S., & Murray, K. E. (2008). Resilience: a new integrative approach to health and mental health research. *Health Psychology Review, 2*, 41-64.
- Zhaoyang, R., Scott, S. B., Smyth, J. M., Kang, J., & Sliwinski, M. J. (2019). Emotional responses to stressors in everyday life predict long-term trajectories of depressive symptoms. *Annals of Behavioral Medicine, 54*, 402-412.

Table 1*Glossary of Terms*

Construct	Definition
Inoculation/toughening	Psychological and physiological changes that promote effective coping, including appraising stressors as more manageable (rather than overwhelming).
Intensive longitudinal methods	Methods that employ repeated assessments of individuals (e.g., experience sampling, daily diary) that allow researchers to study people's thoughts, emotions and behavior
Interindividual differences	Between-person characteristics that are relatively stable over time
Intraindividual change	Enduring, long-term change over macro timescale (e.g., years, decades)) due to developmental or aging effects
Intraindividual variability	Short-term fluctuations over micro timescales (e.g., days, weeks) due to individual adaptation or contextual influences
Measurement burst designs	A design that incorporates bursts of intensive repeated assessment within a relatively short period of time that are repeated longitudinally over more widely spaced intervals
Multidimensionality	The complex interplay of factors influencing development across the lifespan, including biological, cognitive, and socioemotional changes.
Multidirectionality	Adaptive capacities throughout life are characterized by the simultaneous unfolding of increases, decreases, and maintenance in functioning
Plasticity	The capacity to change in response to experience or environmental stimulation throughout the lifespan.
Reactivity	Changes in affect, cognitive, or physiology in response to daily stressors
Recovery	Reduced activation levels in affection, cognition, and physiology following a stressor
Reserve capacity	Level and accumulation of latent capabilities or individual resources for responding effectively to challenging conditions
Resilience Processes	Adaptive responses to daily stressors (i.e., dampened reactivity, accelerated recovery, toughness/inoculation, richness/balance) the increase the likelihood of subsequent health and well-being

Table 1, (continued)

Construct	Definition
Resiliency	The capacity of a dynamic system to adaptively respond to environmental adversity
Richness and balance, or diversity	The spread of stressors across multiple domains of daily life
Steeling	The propensity for prior stressor exposure to increase coping capacity in the face of future stressors
Stressor pile-up	Cumulative stressors that build up over a period of time
Sustainability	The maintenance of health and well-being in the face of major life stressors
Turbulence	A type of diversity index that incorporates contiguity in time-series data

Table 2*Overview of Studies Examining Links between Resilience Processes and Health Outcomes*

Study	Predictor	Outcome	Summary of Findings
Almeida et al., 2020	Reactivity, recovery, and pileup	Physical activity	Greater stressor pileup, but not NA reactivity or recovery, predicted lower subsequent levels of physical activity
Anderson et al., 2021	NA reactivity	Depressive and anxiety symptoms	Dampened NA reactivity in Latinx high school students predicted decreased depressive symptoms, but not anxiety symptoms, during the first year of college.
Bai et al., 2020	PA and NA reactivity	Internalizing problems	Dampened NA and PA reactivity in response to school stressors predicted fewer internalizing problems across three years in children age 8-13
Bergeman & Deboeck, 2014	Rate of stress dissipation	Depressive symptoms	Higher rates of stress dissipation predicted lower depressive symptoms, particularly in individuals low in trait resistance
Booij et al., 2018	PA and NA reactivity	Psychotic and depressive symptoms	Attenuated PA and NA reactivity predicted lower depressive symptoms, but not psychotic symptoms, in a general population
Charles et al., 2013	NA reactivity	General affective distress/an affective disorder	Dampened NA reactivity predicted lower general affective distress and decreased likelihood of an affective disorder
Charles et al., 2021	Stressor exposure	Affective well-being, chronic conditions, and cognitive functioning	Experiencing no daily stressors was related to better physical and emotional well-being but lower levels of cognitive functioning
Chiang et al., 2018	NA reactivity	Mortality	Dampened NA reactivity predicted lower mortality in adults with chronic illness
Cohen et al., 2005	NA reactivity	Depression reduction and depressive symptoms	Attenuated NA reactivity predicted lower depressive symptoms

Table 2, (continued)

Study	Predictor	Outcome	Summary of Findings
Drake et al., 2021	NA reactivity	Flourishing	Dampened NA reactivity was associated with higher levels of flourishing
Koffer et al., (unpublished)	Stressor diversity	Blood pressure	Lower stressor diversity was related to higher diastolic blood pressure in older adulthood
Koffer et al., 2016	Stressor diversity	Affective well-being	Higher stressor diversity was linked with better affective well-being
Leger et al., 2018	NA the day following a stressor	Chronic conditions and functional limitations	Lower NA the day following a stressful event predicted fewer chronic conditions and functional limitations over a period of 10 years
Mrozcek et al., 2015	NA and PA reactivity	Mortality	Dampened PA reactivity predicted lower mortality risk, but NA reactivity did not
Ong et al., 2013	PA reactivity	Sleep outcomes - rest and quality	Attenuated PA reactivity was associated with impaired sleep, especially in those high in trait PA
Ong et al., 2018	PA and NA reactivity	Depressive symptoms	Dampened NA and PA reactivity to daily discrimination predicted lower depressive symptoms among African Americans
Ong et al., 2020	NA reactivity	Marital quality	Dampened NA reactivity predicted better marital quality 10 years later. This was moderated by HRV
Piazza et al., 2013	NA reactivity	Chronic conditions	Dampened NA reactivity predicted decreased risk of chronic health conditions
Piazza et al., 2019	NA reactivity	Allostatic load	Dampened NA reactivity predicted lower allostatic load but only among older adults with high levels of stressor exposure
Rook et al., 2016	NA reactivity	Blood glucose levels	Decreased NA reactivity predicted lower fasting glucose levels in patients with diabetes
Rush et al., 2019	NA Reactivity	Life satisfaction, PWB	Decreased NA reactivity over time predicted higher levels of life satisfaction and PWB

Table 2, (continued)

Study	Predictor	Outcome	Summary of Findings
Schilling & Diehl, 2014	Pileup	Affective well-being	Stressor pile-up over the week was associated with increases in NA, but not PA
Selcuk et al., 2016	NA Reactivity	Changes in PWB	NA reactivity mediated the relationship between partner responsiveness and PWB, with decreased NA reactivity predicting higher PWB
Sin et al., 2015	NA and PA reactivity	Inflammation (Il-6, CRP)	Attenuated NA reactivity predicted greater levels of subsequent PWB Decreased PA reactivity was associated with lower Il-6; Decreased NA reactivity was associated with higher CRP among women
Sin et al., 2016	NA reactivity	HRV (SDRR, HF-HRV, RMSSD)	Decreased NA reactivity was associated with higher heart-rate variability
Smith et al., 2021	Reactivity, recovery, and pileup	Binge Eating Disorder	Stressor pileup, but not reactivity or recovery, was associated with more binge-eating symptoms in adults with binge eating disorder
Stanton et al., 2019	NA and PA reactivity	Mortality	NA reactivity, but not PA reactivity, mediated the relationship between partner responsiveness and mortality, with dampened NA reactivity predicting lower mortality risk
Zhaoyang et al., 2020	PA and NA reactivity	Depressive symptoms	Dampened PA reactivity predicted decreased depressive symptoms over 18 months - NA reactivity did not

Note. NA = negative affect; PA = positive affect; PWB = psychological well-being. For studies utilizing indices of reactivity and recovery, results are summarized on a continuum, ranging from resilience to vulnerability.

Intraindividual Variability and Change *Descriptive Framework*

Figure 1. A visual representation of resilience operating across different timescales, including both short-term dynamics (*intraindividual variability*) and long-term changes (*intraindividual change*). Dynamic resilience processes are characterized as intensive ‘bursts’ of measurements and depicted in magnified circles A (*dampened reactivity*), B (*accelerated recovery*), C (*toughness/inoculation*), and D (*richness/balance*). The solid line connecting the bursts represent long-term intraindividual change processes (e.g., development) that accrue with advancing age. Figure adapted from Nesselroade (1991b), Ram and Gerstorf (2009), and Benson and Ram (2018).