Personality Traits in the International Classification of Diseases 11th Revision (ICD-11)

Joshua R. Oltmanns, Ph.D.

Stony Brook University

Citation:

Oltmanns, J. R. (2021). Personality Traits in the International Classification of Diseases 11th Revision (ICD-11). *Current Opinion in Psychiatry*, 34, 48-53. <https://doi.org/10.1097/YCO.0000000000000656>

Author's note:

Correspondence should be addressed to Joshua R. Oltmanns, Department of Psychiatry and Behavioral Sciences, 10-060E Health Sciences Center, Stony Brook University, Stony Brook, NY, 11794. Email: jroltmanns@gmail.com

Abstract

Purpose of review: The ICD-11 officially adopted a dimensional system of personality disorder that was a paradigm shift for the classification of personality disorders. The purpose of this article is to review the growing amount of research on one component of that system—the personality trait domain model. Importantly, several self-report measures have been developed to measure the ICD-11 domains and have been subjected to initial validation through examination of their factor structure, multi-method use, convergent and discriminant validity with other prominent dimensional personality models (such as the Five-Factor Model), and criterion validity for important life outcomes. *Recent Findings:* Studies indicate the ICD-11 domains align with the Five-Factor Model and prior influential models of dimensional personality traits, as expected, and thus rest on an impressive body of empirical research. They also capture large amounts of variance included in the ICD-10/DSM-5 Section II personality disorders. *Summary:* Together these findings support the construct validity of the ICD-11 trait domains. However, continued validation research is necessary, as well as research on how to implement these domains into clinical practice, and research on the more specific facet-level of the trait domains—although the ICD-11 model is only officially at the domain-level.

Keywords: ICD-11, personality disorder, personality traits, dimensional model, questionnaire

Key Points

- The ICD-11 trait domains align with prior well-established and validated dimensional personality models such as the Five-Factor Model of personality, providing them with a strong base of pre-existing research support.
- The ICD-11 trait domains capture variance in important life outcomes such as depression, life satisfaction, and insomnia, as well as the ICD-10/DSM-5-Section II personality disorder categories.
- Several self-report measures of the ICD-11 trait domains are available, as well as an informant-report measure, all are freely available for use, and have a growing amount of research support.

Personality Traits in the International Classification of Diseases 11th Revision (ICD-11)

Introduction

The recent shift to a dimensional system of personality disorder classification in ICD-11 is a leap forward for psychiatry. Within the classification system is a five-domain trait model, for which several measures have recently been developed and for which initial validation has been provided (see Table 1). The present review examines recent validation research on the ICD-11 traits.

The ICD-11 trait model eliminates the diagnostic personality disorder categories and replaces them with a three-component system: (a) a general severity rating, (b) five prominent personality trait domains, and (c) a borderline pattern specifier, which assesses DSM-5 Section II/ICD-10 borderline personality disorder (1). The five domains of the ICD-11 maladaptive personality trait model are directly aligned with the FFM: negative affectivity aligns with FFM neuroticism, detachment with low FFM extraversion, dissociality with low FFM agreeableness, anankastia with high FFM conscientiousness, and disinhibition with low FFM conscientiousness (2,3). Of note is that anankastia and disinhibition align with opposite ends of FFM conscientiousness, and thus are theoretically bipolar opposites. More specific *facets* of each domain are described in ICD-11, but only the domain-level is used for clinical description. This decision was made in order to preserve parsimony (4).

Recent Findings

Personality Inventory for ICD-11 (PiCD).

The PiCD was developed as the first direct self-report measure of the ICD-11 trait domains according to descriptions of the ICD-11 trait domains made by the ICD-11 Workgroup for the Revision of Personality Disorders (5). The measure consists of 60 items, with 12

measuring each of the five domains, and is freely available for use. In the initial validation study, the scales showed good convergent and discriminant validity and joint factor structure with measures of FFM-aligned trait domains of normal and maladaptive personality, including the Personality Inventory for DSM-5 (PID-5) (5). The PiCD showed good factor structure in an item-level analysis of all 60 items, with 90% of the items loading above $\lambda = .40$ on their expected domain factor. A four-factor solution consisting of negative affectivity, detachment, and dissociality factors, and a bipolar anankastia-versus-disinhibition factor fit the data better than a five-factor solution with separate factors for anankastia and disinhibition. In a five-factor item-level solution, the bipolar anankastia-versus-disinhibition factor divided into two bipolar anankastia-versus-disinhibition factors, a finding that has since been replicated several times (6–8). Notably the four-factor structure with a bipolar factor is somewhat confusing, but the factor structure has no bearing on clinical assessment, because a clinician would score and interpret the PiCD for five separate domains regardless of the factor structure.

Oltmanns and Widiger (2019) examined the relations of the PiCD with the other components of the ICD-11 model, as well as replicated the alignment of the PiCD scales with the FFM and the four-factor structure of the PiCD in an online sample of 269 adults with mental health treatment histories (3). We found that all domains of the PiCD except for anankastia correlated with measures of the ICD-11 and DSM-5 Section III general severity ratings and the borderline specifier. Gutierrez and colleagues replicated the four-factor structure and the PiCD scales' relations to ICD-11 general severity in large samples of 2,522 Spanish adults and 797 inpatients (8). McCabe and Widiger (9) as well identified the four-factor structure of the PiCD, this time in a joint analysis with the facets of the PID-5. Further, they found that the PiCD scales accounted for over 60% of the variance in measures of borderline personality disorder.

Carnovale and colleagues (7) found the four-factor structure and that the PiCD scales accounted for a moderate amount of variance in the MMPI-2-RF scales (median $R^2 = .30$) in a sample of 518 Canadian college students.

Crego and Widiger (10) examined the relations of the PiCD domains to foundational dimensional models of personality disorder traits: the Schedule for Nonadaptive and Adaptive Personality (SNAP) (11), which measures three maladaptive trait domains, and the Dimensional Assessment of Personality Pathology (DAPP) (12), which measures four maladaptive trait domains. They found convergent, discriminant, and structural validity for the PiCD domains with the SNAP and DAPP, including support for the four-factor structure. Of note is that the DAPP includes a compulsivity domain—analogue to ICD-11 anankastia—and the SNAP includes three facet scales of compulsivity (Propriety and Workaholism). The PiCD Anankastia scale correlated $r = .69$ with DAPP Compulsivity, $r = .55$ with SNAP Propriety, and $r = .42$ with SNAP Workaholism.

Somma and colleagues (13) provided validation evidence for the Italian PiCD in a community sample of 1,122 adults. They replicated the convergent and discriminant validity of the PiCD with the FFM and the PID-5 (shorter form), and the four-factor structure. In a subset of the sample ($n = 262$), they found two-week test-retest reliability, with r values ranging from .81 (anankastia) to .89 (negative affectivity and detachment). PiCD negative affectivity, detachment, dissociality, and disinhibition domains correlated with two forms of personality impairment.

Oltmanns and Widiger (14)—in developing and validating an informant-version of the PiCD—also examined the criterion validity of the PiCD for several criteria variables such as depressive symptoms, relationship satisfaction, insomnia symptoms, social adjustment, satisfaction with life, health perceptions, and cognitive functioning in a sample of 714

community older adults. Specifically, PiCD Negative Affectivity, Disinhibition, and Detachment were correlated with depressive symptoms, dissatisfaction with life, and worse perceptions of mental health. Negative affectivity had a uniquely strong relationship with insomnia symptoms and detachment had a more unique relationship with social problems. These findings are an important step forward for the criterion validation of the ICD-11 and PiCD domains.

Informant-Personality Inventory for ICD-11.

Oltmanns and Widiger (14) developed the Informant-Personality Inventory for ICD-11 (IPiC), an informant-version of the PiCD. The IPiC contains the same 60 items as the PiCD, with pronoun changes and minor adjustments to facilitate reporting from the other-perspective about a target person, and is freely available. Bach and colleagues (6) first examined the factor structure of the IPiC as completed by 133 clinicians describing 238 different patients. The IPiC yielded a four-factor solution at the item-level, again with the bipolar anankastia-disinhibition factor, and a five-factor solution with two separate bipolar anankastia-versus-disinhibition factors (along with negative affectivity, detachment, and dissociality factors).

In a more recent study, the IPiC and PiCD were administered to 714 community adults and 569 informants. The IPiC again demonstrated a four-factor solution at the item-level. Self–other agreement correlations on the domains ranged from $r = .28$ (for anankastia and disinhibition) to $r = .44$ (detachment), which are consistent with prior levels of self–other agreement on maladaptive trait domains (15). The IPiC correlated with self-reported external criteria such as depressive symptoms, insomnia symptoms, lack of social support, dissatisfaction with life, and worse mental health perceptions. IPiC scales were more strongly associated with informant-reported criteria: correlating even higher with informant-reported relationship dissatisfaction, worse health status, and cognitive problems.

FFiCD.

The ICD-11 Workgroup decided to limit the trait model to the broader domain-level. Yet within each of the ICD-11 trait domain descriptors, there are clear indicators of more specific, facet-level traits that contribute to each domain. Further, it is clear from research that the facet-level provides a clearer and more specific description of personality (16–18). For example, both paranoid and avoidant personality disorders are defined by high levels of domain-level neuroticism. However, paranoid is defined at the facet-level—within neuroticism—by high angry hostility, while avoidant is defined by high anxiousness, depressiveness, and self-consciousness facets (16).

Taking this into consideration, Oltmanns and Widiger (19) developed a facet-level self-report measure of the ICD-11 trait domains called the FFiCD. The measure was developed from items of over 100 scales of the dimensional Five-Factor Model of Personality Disorder (FFMPD) (20). The FFiCD consists of 121 items that score the five domains, as well as 20 facets, and it is freely available for use. The initial validation article provided evidence of convergent, discriminant, and structural validity of the FFiCD facets with the PiCD, the FFM, and the PID-5. The domains and facets of the FFiCD also display a four-factor structure including a bipolar anankastia-versus-disinhibition factor. Further, the FFiCD includes 47 “nuances,” which are 2-3 item scales below the facets that contribute to an even more specific diagnostic picture of the individual.

PID-5.

After the PiCD was published online, Bach and colleagues published a scoring algorithm for the ICD-11 domains via the PID-5 assessment of the DSM-5 Section III Criterion B trait domains (21). Creating a “cross-walk” between the ICD-11 and DSM-5 Section III Criterion B

trait domains was a relatively straightforward task for the domains of negative affectivity, detachment, dissociality (i.e., antagonism), and disinhibition, as those four domains are essentially analogous across ICD-11 and DSM-5 Section III. The tougher part has been assessing anankastia with the PID-5. Anankastia is not included in the DSM-5 trait model. PID-5 ICD-11 Anankastia is scored with the PID-5 Rigid Perfectionism and PID-5 Perseveration facets. Including Perseveration in this scoring has proven problematic, as this scale was developed to measure negative affectivity.

Bach and colleagues (21) originally hypothesized PID-5 facet/ICD-11 domain alignments and tested the algorithm in a sample of 1,541 Danish outpatients and community adults and replicated it in a sample of 637 US undergraduates. Exploratory analyses demonstrated structural validity for a five-factor solution, including separate anankastia and disinhibition factors along with negative affectivity, detachment, and dissociality factors. However, in the Danish sample, the anankastia factor—defined by PID-5 Perseveration—correlated $r = .63$ with the negative affectivity factor. Bach and colleagues (22) did find that the anankastia scale correlated $r = .60$ with obsessive-compulsive personality disorder, providing evidence that it captures content related to compulsivity/anankastia. Lotfi and colleagues (23) and Lugo and colleagues (24) both showed that the general structure of the PID-5—as scored for the ICD-11 trait domains—was supported in Iranian and Brazilian adults, respectively. However, they both again found that the anankastia factor was highly correlated with negative affectivity.

Sellbom and colleagues (25) examined the PID-5 ICD-11 scoring with the MMPI-2-RF and measures of DSM-IV personality disorders and FFM traits in a sample of 343 psychiatric outpatients. They found support for the five-factor structure and associations with the FFM and the MMPI-2-RF Personality Psychopathology 5 (PSY-5). The ICD-11 domains also captured

large amounts of variance in each of the DSM-IV personality disorder scores. However, the PID-5 ICD-11 anankastia factor again correlated highly with the negative affectivity factor ($r = .67$), and it correlated $r = .56$ with the PSY-5 Negative Emotionality scale—which was even higher than all other hypothesized convergent correlations between corresponding scales, except for negative affectivity. PID-5 ICD-11 Anankastia did again, however, correlate $r = .54$ with obsessive-compulsive personality pathology.

McCabe and Widiger (9) directly compared PID-5 ICD-11 Anankastia with PiCD Anankastia. The correlation was only $r = .34$. This was likely because the PID-5 Perseveration facet correlated $r = -.17$ with PiCD Anankastia. In contrast, PID-5 Rigid Perfectionism correlated $r = .54$ with PiCD Anankastia. In a four-factor solution of the PiCD scales and the facets of the PID-5, Perseveration loaded $\lambda = .69$ on a negative affectivity factor and not on a bipolar anankastia-versus-disinhibition factor that included PiCD Anankastia and PiCD Disinhibition, as well as PID-5 disinhibition facets. Together these studies indicated no problems in the PID-5 assessment of ICD-11 negative affectivity, detachment, disinhibition, or dissociality, but indicated problems in the assessment of anankastia, probably due to the inclusion of PID-5 Perseveration in the scoring of PID-5 ICD-11 Anankastia.

PID-5, Brief Form Plus.

Kerber and colleagues (26) derived the Personality Inventory for DSM-5, Brief Form Plus (PID5BF+) as a brief measure scorable for both the DSM-5 Section III Criterion B traits and the ICD-11 traits in data from English and German-speaking samples ($N = 2,927$). They used ant colony optimization to select 34 items that assess 17 of the original 25 PID-5 facets. Internal consistency and convergent, discriminant, and criterion validity support was provided for negative affectivity, detachment, antagonism, and disinhibition domains. However, PID-5 ICD-

11 anankastia “had the lowest reliability among all six domains and showed remarkable differences between the constructs of perseveration and rigid perfectionism” (26, p. 25).

Bach and colleagues (27) provided an interesting solution to the problematic PID-5 assessment of anankastia. In its development, the PID-5 at one point included more specific facets of compulsivity/anankastia within Rigid Perfectionism (28). However, these facets were eventually combined into PID-5 Rigid Perfectionism. In light of the problems with the Perseveration facet in the assessment of anankastia, Bach and colleagues altered the PID5BF+ to delete the perseveration items and divide PID-5 Rigid Perfectionism into three separate two-item facets: orderliness, rigidity, and perfectionism. This led to a modified PID5BF+ that includes 36 items assessing six domains and 18 facets, with two items per facet. Support was found in archival data of 16 samples from Europe, the US, and Brazil for a six-factor structure including negative affectivity, detachment, disinhibition, anankastia, antagonism (i.e., dissociality), and psychoticism factors. Notably, the factor correlations between anankastia and negative affectivity were much lower, suggesting improved discriminant validity for PID5BF+ Anankastia. However, the PID-5 Rigid Perfectionism facet, from which these two-item facets derive, was developed by retaining only items that loaded highly on a single latent factor (i.e., $\lambda > .50$), suggesting high intercorrelations of the items. It is possible that these three facets may be relatively homogeneous.

Personality Assessment Questionnaire-11 (PAQ-11).

Kim and colleagues (29) developed a brief 17-item domain-level self-report measure for the ICD-11 trait domains in samples of Korean adult students and psychiatric patients. The PAQ-11 was derived from the Personality Assessment Schedule interview (30), which was used previously in field trials for initial validation of the ICD-11 domains (31). The PAQ-11 includes

five items to assess negative affectivity, four items for anankastia, four items for detachment, two items for disinhibition, and two items for dissociality. The PAQ-11 domains showed convergent and discriminant validity with and against with the FFM and PID-5 trait measures. Negative affectivity also showed convergent relations with three measures assessing depression, anxiety, anger, and emotional dysregulation. The PAQ-11 may be a useful brief measure of the ICD-11 domains.

Language.

Finally, Barroilhet and colleagues (32,33) have conducted interesting work scoring the ICD-11 traits via transcribed language in electronic health records. In 3,623 psychiatric patients, disinhibition and negative affectivity predicted longer hospital stays. In 12,274 medical patients, they found that disinhibition predicted greater mortality risk and detachment increased the likelihood of readmission. The use of electronic health records to assess the ICD-11 traits is a promising avenue for future research.

Conclusion

The ICD-11 officially shifted the classification of personality disorder from a categorical system to a dimensional one. This occurred because of the extensive research support on the validity of dimensions for assessing personality (34). Research on the five ICD-11 trait domains has been developing rapidly in the past two years, providing empirical support for the ICD-11 dimensions specifically. The ultimate goal is that—through this improvement in the classification of personality disorders—we will be able to develop and refine treatments that may improve both mental and physical health outcomes.

Acknowledgements

None.

Financial support and sponsorship

None.

Conflicts of interest

None.

References

1. World Health Organization. International statistical classification of diseases and related health problems (11th ed.) [Internet]. 2019. Available from: <https://icd.who.int/>
2. Mulder RT, Horwood J, Tyrer P, Carter J, Joyce PR. Validating the proposed ICD-11 domains: ICD-11 PD domains. *Personal Ment Health*. 2016 May;10(2):84–95.
3. Oltmanns JR, Widiger TA. Evaluating the assessment of the ICD-11 personality disorder diagnostic system. *Psychol Assess*. 2019;31(5):674–84.
4. Tyrer P, Crawford M, Mulder R, Blashfield R, Farnam A, Fossati A, et al. The rationale for the reclassification of personality disorder in the 11th revision of the International Classification of Diseases (ICD-11): ICD-11 reclassification. *Personal Ment Health*. 2011 Nov;5(4):246–59.
5. Oltmanns JR, Widiger TA. A self-report measure for the ICD-11 dimensional trait model proposal: The Personality Inventory for ICD-11. *Psychol Assess*. 2018;30(2):154–69.
6. Bach B, Christensen S, Kongerslev MT, Sellbom M, Simonsen E. Structure of clinician-reported ICD-11 personality disorder trait qualifiers. *Psychol Assess*. 2020 Jan;32(1):50–9.
7. Carnovale M, Sellbom M, Bagby RM. The Personality Inventory for ICD-11: Investigating Reliability, Structural and Concurrent Validity, and Method Variance. 2020;10.
8. Gutierrez F, Aluja A, Ruiz-Rodriguez J, Garcia LF, Garriz M, Gutierrez-Zotes JA, et al. Personality Disorders in the ICD-11: Spanish Validation of the PiCD and the SASPD in a Mixed Community and Clinical Sample. *Assessment*. in press;
9. McCabe GA, Widiger TA. A comprehensive comparison of the ICD-11 and DSM–5 section III personality disorder models. *Psychol Assess*. 2020 Jan;32(1):72–84. *

A comparison of the ICD-11 and DSM-5 Section III models of personality disorders. Relations between general severity measures and trait measures from both models, as well as relations with borderline personality disorder, are examined.

10. Crego C, Widiger TA. The convergent, discriminant, and structural relationship of the DAPP-BQ and SNAP with the ICD-11, DSM–5, and FFM trait models. *Psychol Assess*. 2020 Jan;32(1):18–28. *

A comparison of the ICD-11 and DSM-5 Section III trait models with prior prominent dimensional trait models of personality and personality disorder. Nuances regarding similarities and differences of the histories and relations between the models are discussed.

11. Clark LA. Manual for the schedule for nonadaptive and adaptive personality (SNAP). Minneapolis: University of Minnesota Press; 1993.

12. Livesley WJ, Jackson DN. Manual for the dimensional assessment of personality pathology-basic questionnaire. Port Huron, MI: Sigma Press; 2009.
13. Somma A, Gialdi G, Fossati A. Reliability and construct validity of the Personality Inventory for ICD-11 (PiCD) in Italian adult participants. *Psychol Assess.* 2020 Jan;32(1):29–39.
14. Oltmanns JR, Widiger TA. Consensual, Structural, and Criterion Validation of the Informant-Personality Inventory for ICD-11 in Older Adults. under review.
15. Oltmanns JR, Oltmanns TF. Self–other agreement on ratings of personality disorder symptoms and traits: Three meta-analyses. In: Letzring TD, Spain JS, editors. *The Handbook of Accurate Personality Judgment: Theory and Empirical Findings.* Oxford University Press; in press.
16. Samuel D, Widiger T. A meta-analytic review of the relationships between the five-factor model and DSM-IV-TR personality disorders: A facet level analysis☆. *Clin Psychol Rev.* 2008 Dec;28(8):1326–42.
17. Reynolds SK, Clark LAnna. Predicting Dimensions of Personality Disorder From Domains and Facets of the Five Factor Model. *J Pers.* 2001 Apr;69(2):199–222.
18. Sprock J. A Comparative Study of the Dimensions and Facets of the Five-Factor Model in the Diagnosis of Cases of Personality Disorder. *J Personal Disord.* 2002 Oct;16(5):402–23.
19. Oltmanns JR, Widiger TA. The Five-Factor Personality Inventory for ICD-11: A facet-level assessment of the ICD-11 trait model. *Psychol Assess.* 2020;32(1):60–71. *

Development and initial validation of a facet-level measure for the ICD-11 trait model. The measure was developed from the Five-Factor Model of Personality Disorder (FFMPD) measures and includes 121 items assessing the five ICD-11 domains, 20 facets, and 47 nuances.

20. Widiger TA, Lynam DR, Miller JD, Oltmanns TF. Measures to Assess Maladaptive Variants of the Five-Factor Model. *J Pers Assess.* 2012;94(5):450–5.
21. Bach B, Sellbom M, Kongerslev M, Simonsen E, Krueger RF, Mulder R. Deriving ICD-11 personality disorder domains from dsm-5 traits: initial attempt to harmonize two diagnostic systems. *Acta Psychiatr Scand.* 2017 Jul;136(1):108–17. *

The initial development of a scoring system for ICD-11 personality traits via the Personality Inventory for DSM-5. The process of translating the ICD-11 trait model to the PID-5 measure of the DSM-5 Section III trait model is described and initially evaluated.

22. Bach B, Sellbom M, Skjernov M, Simonsen E. ICD-11 and DSM-5 personality trait domains capture categorical personality disorders: Finding a common ground. *Aust N Z J Psychiatry.* 2018 May;52(5):425–34.

23. Lotfi M, Bach B, Amini M, Simonsen E. Structure of DSM-5 and ICD-11 personality domains in Iranian community sample: Structure of DSM-5 and ICD-11 traits in Iran. *Personal Ment Health*. 2018 May;12(2):155–69.
24. Lugo V, de Oliveira SES, Hessel CR, Monteiro RT, Pasche NL, Pavan G, et al. Evaluation of DSM-5 and ICD-11 personality traits using the Personality Inventory for DSM-5 (PID-5) in a Brazilian sample of psychiatric inpatients: PID-5 personality traits in Brazilian inpatients. *Personal Ment Health*. 2019 Feb;13(1):24–39.
25. Sellbom M, Solomon-Krakus S, Bach B, Bagby RM. Validation of Personality Inventory for DSM-5 (PID-5) algorithms to assess ICD-11 personality trait domains in a psychiatric sample. *Psychol Assess*. 2020 Jan;32(1):40–9. *

A comparison of the ICD-11 scoring from the PID-5 in relation to important validation criteria including the DSM-IV personality disorders, the MMPI-2-RF, and the FFM. The scores showed convergent and discriminant validity, with problems in the PID-5 assessment of anankastia.

26. Kerber A, Schultze M, Müller S, Wright AGC, Spitzer C, Krueger R, et al. Development of a Short and Reliable Measure for DSM-5 and ICD-11 Maladaptive Personality Traits Using Ant Colony Optimization Algorithms [Internet]. *PsyArXiv*; [cited 2020 Jan 16]. Available from: <https://osf.io/rsw54>
27. Bach B, Kerber A, Aluja A, Bastiaens T, Keeley JW, Claes L, et al. International Assessment of DSM-5 and ICD-11 Personality Disorder Traits: Toward a Common Nosology in DSM-5.1. *Psychopathology*. 2020 May 5;1–10. *

A scoring algorithm is developed to briefly assess both ICD-11 and DSM-5 Section III trait models using the PID-5. Problems with the PID-5 assessment of ICD-11 anankastia are addressed by splitting the Rigid Perfectionism facet into three separate two-item facets.

28. Krueger RF, Derringer J, Markon KE, Watson D, Skodol AE. Initial construction of a maladaptive personality trait model and inventory for DSM-5. *Psychol Med*. 2012 Sep;42(9):1879–90.
29. Kim Y, Tyrer P, Hwang S. Personality Assessment Questionnaire for ICD-11 personality trait domains: Development and testing. *Personal Ment Health*. 2020 Jul 7;pmh.1493. *

Initial development and validation of the PAQ-11, a brief measure of the five ICD-11 domains. The measure was developed from the PAS and consists of 17 items. The scales were validated with the important external criteria such as the FFM and measures of emotion dysregulation.

30. Tyrer P, Alexander MS, Cicchetti D, Cohen MS, Remington M. Reliability of a Schedule for Rating Personality Disorders. *Br J Psychiatry*. 1979 Aug;135(2):168–74.
31. Tyrer P, Mulder R, Kim Y-R, Crawford MJ. The Development of the ICD-11 Classification of Personality Disorders: An Amalgam of Science, Pragmatism, and Politics. *Annu Rev Clin Psychol*. 2019 May 7;15(1):481–502. *

The original chair of the ICD-11 Working Group for the Revision of Personality Disorders, Dr. Tyrer, and colleagues describe an in-depth history of the development of the ICD-11 personality disorder diagnostic system.

32. Barroilhet SA, Pellegrini AM, McCoy TH, Perlis RH. Characterizing DSM-5 and ICD-11 personality disorder features in psychiatric inpatients at scale using electronic health records. *Psychol Med*. 2019 Sep 23;1–9.
33. Barroilhet SA, Bieling A, McCoy TH, Perlis RH. Association between DSM-5 and ICD-11 personality dimensional traits in a general medical cohort and readmission and mortality. *Gen Hosp Psychiatry* [Internet]. 2020; Available from: <http://www.sciencedirect.com/science/article/pii/S0163834320300190>
34. Clark LA. Assessment and Diagnosis of Personality Disorder: Perennial Issues and an Emerging Reconceptualization. *Annu Rev Psychol*. 2007 Jan;58(1):227–57.

Table 1

Questionnaire Measures of the ICD-11 Trait Domains

Measure	Number of scales	Number of items and items-per-scale (on average)
PiCD	5 domain scales	60 items, 12 items per domain
IPiC	5 domain scales	60 items, 12 items per domain
FFiCD	5 domain scales, 20 facet scales, 47 nuance scales	121 items, 24 items per domain, 6 items per facet, 2-3 items per nuance
PID-5 for ICD-11	5 domain scales, 16 facet scales	143 items, 28 items per domain, 9 items per facet
PID-5BF+ (original)	6 domain sales, 17 facet scales	34 items, 6 items per domain, 2 items per facet
PID-5BF+ (revised)	6 domain scales, 18 facet scales	36 items, 6 items per domain, 2 items per facet
PAQ-11	17 items, 5 domain scales	17 items, 3-4 items per domain

Note. PiCD = Personality Inventory for ICD-11, IPiC = Informant-Personality Inventory for ICD-11, FFiCD = Five-Factor Inventory for ICD-11, PID-5 for ICD-11 = Personality Inventory for DSM-5 scored for ICD-11, PID-5-BF+ = Personality Inventory for DSM-5-Brief Form-Plus, PAQ-11 = Personality Assessment Questionnaire for ICD-11.