©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

Connections during Crisis: Adolescents' social dynamics and mental health during COVID-19

Reuma Gadassi Polack^{1a}, Haran Sened², Samantha Aubé³, Adam Zhang¹, Jutta Joormann^{1*}, Hedy Kober^{1*}

¹**Yale University**

²**Bar-Ilan University**

³**Boston College**

*Authors contributed equally to this work

^a Correspondence concerning this article should be addressed to Reuma Gadassi Polack, Department of Psychology, Yale University, 2 Hillhouse Avenue, New Haven, CT, 06520. Email: reuma.gadassipolack@yale.edu.

Author's Note

Our hypothesis regarding changes in spillover processes during COVID-19 was pre-registered at https://osf.io/eq62h/?view_only=742c1d2916454517b6416302d6211926. This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 786064, the National Institute of Mental Health Translational Developmental Neuroscience Training Grant (T32 #MH18268), The Israeli Council for Higher Education Postdoctoral Research Fellowship for Women awarded to Dr. Gadassi Polack, the Azrieli Fellowship awarded to Dr. Sened, and the National Institute of Mental Health R21 MH119552 awarded to Dr. Joormann and Dr. Kober. The authors thank Itay Polack Gadassi for his help setting up the study, and to Ralitz Gueorguiva for advice regarding data analysis. The authors also thank the research assistants who helped with data collection and the families who participated.

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

Connections during Crisis: Adolescents' social dynamics and mental health during COVID-19

Abstract

Adolescence is a critical period for social development, which COVID-19 has dramatically altered. Quarantined youths had limited in-person interactions with peers. The present study used an intensive longitudinal design to investigate changes in interpersonal dynamics and mental health during COVID-19. Specifically, we investigated whether the associations between different social contexts – i.e., “spillover” – changed during COVID-19, and whether changes in social interactions during COVID-19 was associated with changes in depressive symptoms. Approximately one year prior to the onset of COVID-19, 139 youths reported depressive symptoms and daily interactions with parents, siblings, and friends, every day for 21 days via online questionnaires. Shortly after schools closed due to COVID-19, 115 of these youths completed a similar 28-day diary. Analyses included 112 youths (62 girls, 73% Caucasian; $M_{\text{age}}=11.77$, range 8-15 in Wave 1) who completed at least 13 diary days in each data wave. Our results show that younger adolescents experienced significant decreases in negative and positive interactions with friends, whereas older adolescents showed significant decreases in negative interactions with friends and significant increases in positive interactions with siblings. As predicted, within-day spillover of positive interactions and person-level association of negative interactions increased within the family during COVID-19, whereas within-day spillover of positive interactions between family and friends decreased. We also found a dramatic increase in depressive symptoms. More negative interactions and fewer positive interactions with family members were associated with changes in depressive symptoms. Our study sheds light on how youths' social development may be impacted by COVID-19.

Keywords: depression, adolescents, family processes, peer relationship, COVID-19, spillover

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

Connections during Crisis: Adolescents' social dynamics and mental health during COVID-19

The COVID-19 pandemic has affected many individuals, families, and communities around the world (Choi et al., 2020; Wang et al., 2020; WHO, 2020). Many governments implemented quarantines and restricted social gatherings to decrease infection concerns (CDC, 2020). As a result, many individuals markedly diminished in-person contact. Previous quarantines have shown that such dramatic social changes can lead to feelings of social isolation and a greater likelihood of developing mental health issues (Brooks et al., 2020; Ghrebreyesus et al., 2020). Studies have begun to delineate the mental-health consequences of COVID-19 and related quarantine and already noted increases in psychopathology, including depression (e.g., Chanal et al., 2020; Ghebreyesus et al., 2020).

Children and adolescents may be at a particularly high risk for deleterious mental health outcomes during COVID-19 given their position at a critical developmental stage in which social interactions, particularly with peers, are central (Ellis et al., 2020; Orben, Tomova, Blakemore, 2020). Indeed, adolescence is a time during which youth are at increasingly high risk of developing mental health disorders (e.g., Petersen et al., 2018; Salk, Peterson, Abramson & Hyde, 2016). In addition, adolescents can be particularly vulnerable to stress (Cohodes, Kitt, Baskin-Sommers, & Gee, 2020), which may render them susceptible to mental health ramifications from COVID-19. However, most studies thus far on the mental health impact of COVID-19 were based on cross-sectional designs (c.f., Chanal et al., 2020); thus, their ability to deduce causality is limited. Moreover, such studies have typically speculated about changes in social relationships during quarantine, and have not examined it directly.

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

During typical transition to adolescence, youths begin spending an increasing amount of time with their peers, and less time with their parents (e.g., Lam et al., 2012; Larson et al., 1991; Larson et al., 1996). Additionally, peer groups become more influential, while parental impact is lessened (e.g., De Goede et al., 2006; Hadiwijaya et al., 2017). These processes of becoming closer to peers is thought to support youths' social-emotional development and increased independence from their family (Nelson et al., 2005). In this context, COVID-19 and the social distancing measures it required – especially during early stages – may have been particularly harmful to youths' social-emotional development. Indeed, recent cross-sectional evidence shows that youth have experienced distress over the maintenance of friendships during quarantine (Ellis et al., 2020). Although online social interactions may partially compensate for a lack of in-person meetings (Ellis et al., 2020; Orben et al., 2020), in-person interactions are likely to be more influential and foster more connectedness. Importantly, the current investigation focused on the initial stages of the pandemic, when “stay at home” recommendations were widespread.

One way to assess relationship connectedness is derived from theories that view the social world as comprised of different subsystems (i.e., relationships) that are interconnected (e.g., the mother-adolescent dyad is a subsystem, as is the mother-father dyad; Bronfenbrenner & Morris, 2007; Minuchin, 1985). Although these systems are interconnected, they ideally also have boundaries between them: thus, if there is a problem in a specific subsystem, it would not proliferate to other connected subsystems (e.g., mother-father conflict would not lead to mother-child conflict; Davies & Sturge-Apple, 2014). However, boundaries between relationships are permeable, and events or emotions from one subsystem/relational context are often associated with similar events or emotions in other relational contexts, via a process dubbed “spillover” (e.g., Chung et al., 2011; Flook & Fuligni, 2008; Kaufman et al., 2020; Krishnakumar &

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: [10.1037/dev0001211](https://doi.org/10.1037/dev0001211)

Buehler, 2000; Mastrotheodoros et al., 2020). The process of spillover has predominately been studied in the context of how parental conflict affects parent-child relationships (e.g., Chung et al., 2011; Kaufman et al., 2020; Mastrotheodoros et al., 2020; Timmons & Margolin, 2015). Excessive spillover can reflect “enmeshed” family boundaries (Minuchin, 1985), which may be detrimental to youth’s mental health, especially when the spillover is of negative interactions or emotions (e.g., Koçak et al., 2017; Lindblom et al., 2017). Spillover can also occur between family and peer relationships, which can create vicious cycles of relational negativity. For instance, higher parental rejection predicted increases in peer victimization a year later (Kaufman et al., 2020). In the current investigation, we used spillover processes to assess connectedness within the family system, as well as between family subsystems and peer relationships.

Our study involved two waves of data collection – one a year prior to COVID-19 and one during the beginning of the pandemic. Each wave was comprised of an intensive longitudinal design (daily diaries), which allowed us to examine the impact of COVID-19 using two levels of analyses: the day-level and the person-level. Separating day-level from person-level allowed us to separately analyze state (i.e., a given day) and trait (i.e., a specific person) effects (Bolger & Laurenceau, 2013; Trull & Ebner-Priemer, 2020). The distinction between trait and state is of theoretical importance because trait-level effects reflect individual differences in strength of association between relational contexts, whereas state-level effects shed light on daily processes across individuals. Moreover, associations found on the state-level do not necessarily indicate trait-level associations, and vice-versa, highlighting the need to examine them separately.

Taken together, the present study contributes to the existing COVID-19 literature in several ways. First, we recruited participants who already provided daily-diary data a year prior to COVID-19 (Gadassi Polack, Chertkof, Kober & Joormann, 2021), which served as a baseline.

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

Then, we collected diary data from the same participants shortly after quarantine began. Using both waves, we assessed how COVID-19 impacted youths' mental health as well as their social interactions with family and friends. Importantly, we examined whether these social interactions accounted for changes in mental health during quarantine. Second, by examining spillover effects we assessed *how* each of these youths' relational contexts were connected to the other. If youths are meant to become more independent from their parents, closer to their friends, and more influenced by them during normative development, we expected that the quarantine and social distancing would disrupt such typical development, resulting in youths becoming closer to and more influenced by their family (vs. friends). Finally, we included siblings as a social context separately from parents. Evaluating sibling relationships was particularly critical during COVID-19, when siblings were the closest thing to a peer group available to most youths.

Our study also contributes to the literature on spillover processes. First, we assessed a wide range of both positive (e.g., support) and negative (e.g., conflict) social interactions reported by youths. This broader approach is necessary given that the literature has historically focused on conflict and negative interactions only (e.g., Chung et al., 2011; Davies & Sturge-Apple, 2014; Kaufman et al., 2020; Mastrotheodoros et al., 2020; Timmons & Margolin, 2015), despite substantial findings showing that positive interactions are important and can proliferate via self-enhancing cycles (Fredrickson, 2013). Secondly, this study's nested (two-wave) intensive longitudinal design and daily diary approach treats COVID-19 as an "experiment in nature" (Gleason et al., 2008), thus allowing us to examine how spillover processes change under stressful situations, and how physical proximity contributes to these processes.

Finally, our study contributes to the literature on social deprivation during adolescence, which has scarcely been investigated in humans due to ethical considerations (Orben et al.,

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

2020). The preclinical literature shows that interactions with peers are particularly important for adolescent animals, and that isolation has a lasting detrimental impact on mood, behavioral, and cognitive outcomes (Panksepp, 1981), including increases in anxiety and depression-like behaviors (Orben et al., 2020). Even with partial isolation, adolescent animals exhibit changes in the prefrontal cortex, a brain region associated with executive function (Bell et al., 2010).

Although no comparable studies were conducted on human adolescents, a recent study that examined the influence of the COVID-19 lockdown on college students found that their social networks were more sparse compared to prior cohorts, that they had more mental-health difficulties, and that having fewer social interactions was associated with worse mental health outcomes (Elmer et al., 2020).

The following questions guided our analyses:

- (1) **How did social interactions change due to COVID-19?** We hypothesized that during (vs. prior to) COVID-19, youths would report more interactions with family members and fewer interactions with friends. We examined age as a moderator of these changes: we expected older (vs. younger) youths to have more interactions with friends and fewer interactions with parents both prior to and during COVID-19. We had no hypothesis regarding siblings.
- (2) **How did within-day spillover and person-level associations of social interactions change during COVID-19?** We hypothesized that within-day spillover and person-level associations between family members would increase in strength during COVID-19 (vs. before), whereas within-day spillover and person-level associations between family and friends would decrease. We examined these hypotheses separately for positive and negative interactions. This hypothesis was pre-registered at

https://osf.io/eq62h/?view_only=742c1d2916454517b6416302d6211926.

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

(3) Did youth depressive symptoms increase during COVID-19? And were these increases

moderated by age? We hypothesized that depressive symptoms would increase during COVID-19 and that this increase would be particularly prominent among girls, as gender differences in depression typically emerge during adolescence (e.g., Salk et al., 2016).

(4) Do social interactions during COVID-19 account for increases in depressive symptoms

from prior to and during COVID-19? Specifically, we hypothesized that a lower number of positive interactions and a higher number of negative interactions would account for increases in depressive symptoms from prior to during COVID-19.

Method

All procedures have been approved by the Yale University Institutional Review Board (protocol number: 2000022492, title: “Emotion regulation in risk for depression”). Study materials can be accessed online (Gadassi Polack et al., 2021

https://osf.io/eq62h/?view_only=742c1d2916454517b6416302d6211926)

Participants

Wave 1: One hundred forty-eight youths were recruited via flyers in the University area, on Craigslist, and on social media. Advertisements invited youths 9-15 years old to a diary study about emotions and social interactions. Inclusion criteria were daily access to an internet-enabled device and consent from a legal guardian. Siblings were allowed to participate. Participants received \$40 if they completed 60% of surveys and \$60 if they completed 90%. Those who completed <60% received \$10 for participation. One hundred and thirty-nine youths (94%) completed at least 13 diary entries. Since we included children who turned 9 during the diary period, the age range was 8-15.

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

Wave 2: We contacted participants who (1) completed Wave 1, and (2) indicated interest in additional studies. They were invited to participate in a one-time background questionnaire session and a 28-day diary study similar to the one they had completed during Wave 1. Due to the increased length of the study, compensation was \$50 if they completed 60% of surveys and \$70 if they completed 90%; those who completed <60% received \$10 for participation. One hundred and fifteen out of the 117 participants who started Wave 2 completed at least 13 entries (98%). Supplementary table S1 presents demographics for both data waves.

Final sample characteristics. We included 112 participants (62 girls) who completed at least 13 diary entries in each data wave. Mean age of participants in Wave 1 was 11.77 ($SD=2.13$, median=12), and 12.64 ($SD=2.12$, median=13) in Wave 2. The sample was mainly Caucasian (73%). Average time between the two waves was 310 days ($SD=58.81$, range=222–436). Youths included in the final sample were similar to those who did not complete Wave 2 in terms of gender ($\chi^2[1]=1.30$, $p=.254$), race ($\chi^2[6]=7.80$, $p=.253$), and age at Wave 1 ($t[146]=1.60$, $p=.112$), and all other research variables (vs. excluded) in Wave 2 (all $ps \geq .072$).

Power analysis. The present study is part of a larger project on emotions and social interactions; sample size was determined for Wave 1. To offset attrition, we increased the diary period in the Wave 2 to 28 days.

Procedure

Wave 1. Data were collected between 1/31/2019 and 9/23/2019. Participants came to the lab for an initial visit, during which a researcher reviewed the daily diary questionnaire to ensure clarity. Youths signed assent forms and their parents signed consent forms. Participants also completed a practice survey and a demographics questionnaire. Subsequently, every evening for 21 days, participants received a link via email to the daily survey, which they completed on a

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

secure website (Qualtrics). Participants were instructed to complete the survey before going to bed. The link expired after 14 hours.

Wave 2. Data were collected between 3/30/2020 and 6/8/20. On 3/10/20, the governor of Connecticut declared a state public health emergency. Public schools were closed on 3/13/20, public amusement places on 3/18/20, and non-essential businesses on 3/23/20. “Stay at home” and mandatory face-covering orders were issued on 3/28/20. Connecticut started phase 1 of reopening 5/20/20; phase 2 started after the end of data collection. According to the CT Department of Public Health (<https://data.ct.gov/Health-and-Human-Services/COVID-19-Tests-Cases-Hospitalizations-and-Deaths-S/rf3k-f8fg>), during data collection there were 44,179 confirmed cases of COVID-19, between 293-1,972 individuals were hospitalized per day due to COVID-19, and 4,097 COVID-related deaths in Connecticut, and 1,961,781 confirmed cases and 111,774 deaths reported across the U.S.

Participants were contacted by e-mail or phone. If they expressed interest, an online Zoom session was scheduled for them and their parents, during which they gave assent and consent respectively (through Qualtrics) and received explanations about the diary. Then, youths filled out a demographic questionnaire as well as some additional questionnaires not used in the current investigation. Subsequently, every evening for 28 days, participants received a link via email to the daily survey. Participants were instructed to complete the survey before going to bed. The link expired after 16 hours.

Background COVID-19-related questions

In the initial session for Wave 2, we asked youths background questions, including if their school had been moved to an online format, if their usual activities had been canceled, how many people were living in their household, and if their parents are working from home.

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

Diary measures

The present study is part of a larger study; only relevant measures are described.

Interpersonal interactions. Every day, participants were asked to report whether or not the following events occurred, and if they did, with whom (mom/dad/sibling/friend/romantic partner in Wave 1; mother/father/sibling/friend/other kid/other adult in Wave 2). The list of interpersonal interactions included positive (e.g., *was included by, was complimented by, was supported by*) and negative (e.g., *had a fight with, was left out by, was criticized by, was insulted by, was made fun of by, and was let down by*) interpersonal interactions. Each event within a given social context was scored “1” (*event occurred*) or “0” (*event did not occur*). Number of events were summed every day within social context (e.g., if participant indicated that they had a fight with their mother and were criticized by her on a certain day, they received a “2” in the daily score of negative events with mother).

Depressive Symptoms. A self-report Children’s Depression Inventory – short version (CDI-S; Kovacs, 1985), which consists of 10 items assessed the severity of depressive symptoms. The short form is similar to the full measure in its specificity and sensitivity (Allgaier et al., 2012). Each item consists of three sentences representing different degrees of symptom severity (from 0 to 2), from which participants choose those that best describe them. Instructions were adapted for the daily diary by asking participants to choose the sentence that best describes them at the time they were answering the survey. The score suggested as the clinical cutoff for the short version is ≥ 3 (Allgaier et al., 2012). We calculated the between- and within-subject reliabilities using procedures outlined in Shrout and Lane (2012). For a given measure, the between-subject reliability coefficient is the expected between-subject reliability estimate for a single typical day. The within-subject reliability coefficient is the expected within-subject

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

reliability of change within individuals over the daily-diary entries. The between-person and within-person reliabilities were .91 and .75 in Wave 1 and .92 and .74 in Wave 2. These reliabilities are considered good for within-individual measures (Nezlek, 2017).

Data Analytic Plan

Day-level analyses. Data within each wave were hierarchically nested: days within individuals to account for the non-independence of day-level data, and to prevent inflation of effects (Krull & MacKinnon, 2001). Data were analyzed using the nlme package (Pinheiro, Bates, DebRoy, Sarkar, & R Core Team, 2014) of R Studio software (R Core Team, 2013). Level 1 was the day-level and Level 2 was the person-level. We centered day-level predictors at the person-mean to make interpretation of intercepts clearer, and to separate Level 1 and Level 2 effects (see Zhang, Zyphur, & Preacher, 2009). We used a compound symmetry structure across the daily errors. As covariates, we entered into the analyses (1) the lagged mean-centered outcome score (i.e., the previous day's outcome variable, entered as a deviation from the mean) and (2) the person's mean outcome score (averaged across the entire diary period). Thus, the outcome (interpersonal interactions) became a residualized change score.

For example, to test our hypothesis that daily variations in positive interactions with mother predicted daily variations in interactions with father and that this association was moderated by data wave, we ran a model in which data wave (dummy coded such that 0=Wave 1 and 1=Wave 2), day-level positive interactions with mother, and the interaction between day-level positive interactions with mother and data wave were the predictors of variations in positive interactions with father. In addition, we entered yesterday's positive interactions with father into the model, along with the participant's mean level of positive interactions with father. Including lagged positive interactions with father means that whatever effect we find for positive

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

interactions with mother would not include variance that is due to yesterday's positive interactions with father and its effects on positive interactions with mother (or directly on today's positive interactions with father). We also entered the participant's mean score of the predictors (e.g., their mean level of positive interactions with mother). Including the person-mean variables allows estimation of both person-level and day-level effects (Bolger & Laurenceau, 2013), and also allows us to rule out static spurious "third variables" as alternative explanations.

Y_{ijkl} (Positive interactions with dad during wave l on day k for person j) =

$(\beta_0 + b_{0j}) +$

$(\beta_1 + b_{1j}) * (\text{lagged positive interactions with dad [day } k-1]) +$

$(\beta_2 + b_{2j}) * (\text{Positive interactions with mom on day } k) +$

$\beta_3 * (\text{Data Wave}) +$

$\beta_4 * (\text{Data Wave} * \text{Positive interactions with mom on day } k)$

$\beta_5 * (\text{mean Positive interactions with dad}) +$

$\beta_6 * (\text{mean Positive interactions with mom}) +$

ϵ_{ijkl}

Person-level analyses. Thirty (26.1%) of our participants were siblings, which entered non-independence to the data. Thus, we added a variable identifying familial membership to the model's random statement. However, since variance of the random effect of family membership was zero, person-level analyses were evaluated using linear models. To do this, we averaged daily scores of our variables across the entire diary period within individual. For example, depressive symptoms before and during COVID-19 were summed within day, and then averaged across the diary period (21 days for Wave 1, 28 days for Wave 2) within individual. Examining

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

person-level effect using summed scores across the diary period follows our prior work (e.g., Gadassi, Mor, & Rafaeli, 2011) and recommendations in the field (Bolger & Laurenceau, 2013).

To examine if person-level associations between social interactions changed during COVID-19, we examined multilevel models in which social interactions (e.g., positive interactions with mother) were associated with social interactions from a different social context (e.g., positive interaction with father). To examine whether these associations were different in size during COVID-19, we added a dummy-coded variable to indicate “Wave” (coded “0” for Wave 1 and “1” for Wave 2).

$$Y_i \text{ (Mean number of positive interactions with father for person } j) = \\ \beta_0 + \beta_1 \text{ (mean number of positive interactions with mother)} + \\ \beta_2 \text{ (Data Wave)} + \\ \beta_3 \text{ (mean number of positive interactions with mother *Data Wave)} + \\ \epsilon_{jk}$$

All analyses were repeated adjusting for gender and number of days between data waves; these are reported below only when this changed the effects.

Moderation by age. To examine for possible moderation by age of hypotheses 1 & 3, we conducted all analyses with participants’ z-transformed age at Wave 2 as a moderator. Supplementary Table S4 presents correlations between participants’ age and study variables.

Results

Youths’ lives during COVID-19

The vast majority of our sample (97.3%) reported school moving to a full or partial online format (104 participants reported attending school completely online and four reported partially online). Similarly, most (91.9%) reported that after-school activities were partially or

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

fully cancelled (93 reported all activities cancelled, 9 reported some were cancelled). The vast majority of our sample, 107 participants (95.5%), reported living with three or more people in their household. Finally, most participants reported that at least one parent stayed home (70 participants reported both parents at home, 35 reported at least one parent; together 94.6%).

Changes in social interactions during COVID-19, and their moderation by age.

To examine our hypothesis that the number of social interactions with family members will increase and the number of social interactions with friends will decrease, and whether these changes will be moderated by youths' age, we conducted a 4-way mixed ANOVA with Valence (positive vs. negative interactions), Time (before vs. during COVID-19) and Context (mother, father, sibling or friend) as within-participant factors, and a Z-score of participant age at Wave 2 as a covariate. Given the significant Valence by Context interaction ($F[1, 110]=98.08, p<.001, \eta^2=.47$) indicating that positive interactions ($M=1.43, SE=0.10$) were much more frequent than negative interactions ($M=0.37, SE=0.04$), and to simplify reporting, we report 3-way mixed ANOVAs separately for positive and negative interactions.

Positive social interactions. To examine how positive interactions change during COVID-19, we conducted a 3-way (Time by Context by Age) mixed ANOVA. The ANOVA did not reveal a significant main effect for Time ($F[1, 110]=0.17, p=.677$). However, the Time by Context ($F[3, 330]=10.70, p<.001, \eta^2=.09$) and Time by Context by Age ($F[3, 330]=3.52, p=.015, \eta^2=.03$) interactions were significant. In addition, the main effects of Context ($F[3, 330]=30.68, p<.001, \eta^2=.22$), Age ($F[1, 110]=5.47, p=.021, \eta^2=.05$), and Context by Age interaction ($F[3, 330]=20.72, p<.001, \eta^2=.16$) were significant. The Time by Age interaction was not ($F[1, 110]=1.87, p=.666$).

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

To better understand the 3-way Time by Context by Age interaction, we first conducted paired-sample *t*-tests for each social context within each age group. For younger participants, there was a significant decrease in number of positive interactions with friends from prior ($M=1.44$, $SD=1.16$) to during COVID-19 ($M=0.74$, $SD=1.15$; $t[51]=4.86$, $p<.001$) and, for older participants, there was a significant increase in number of positive interactions with siblings from prior ($M=0.67$, $SD=0.93$) to during COVID-19 ($M=0.87$, $SD=1.12$; $t[59]=2.03$, $p=.047$; See Table 1). All other comparisons were not significant ($ps\geq.069$).

To better understand the interaction of Context and Age, we averaged positive interactions across the two data waves. For participants 12 years of age or younger, positive interactions with mothers were the most frequent, followed by fathers ($t[51]=3.24$, $p=.002$) and siblings ($t[51]=5.77$, $p<.001$; Table 1). Least frequent were positive interactions with friends, although they did not significantly differ from siblings ($t[51]=1.48$, $p=.146$). For participants 13 years of age or older, positive interactions with mothers and friends were the most frequent, with no significant difference ($t[59]=0.49$, $p=.629$). Next were positive interactions with fathers, which were significantly less frequent than with friends ($t[59]=2.34$, $p=.023$). Least frequent were positive interactions with siblings ($t[59]=4.88$, $p<.001$).

Negative social interactions. To examine how negative interactions change during COVID-19, we conducted a 3-way (Time by Context by Age) mixed ANOVA. The ANOVA did not reveal a significant main effect for Time ($F[1, 110]=1.38$, $p=.244$). However there was a significant Time by Context interaction ($F[3, 330]=4.52$, $p=.004$, $\eta^2=.04$). In addition, there was a main effect for relational Context ($F[3, 330]=22.11$, $p<.001$, $\eta^2=.17$), and a significant Context by Age interaction ($F[3, 330]=7.04$, $p<.001$, $\eta^2=.06$). To better understand the source of the Time by Context interaction, we conducted a series of paired-sample *t*-tests comparing the frequency

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

of negative interactions within Context. The number of negative interactions with friends significantly decreased during COVID-19, but there were no changes in frequency of negative interactions in other relational contexts (Table 1).

To explore the Context by Age interaction, we averaged the number of negative interactions across data waves, and split participants by median age ($\text{Median}_{\text{Wave2}} = 13$). For participants who were 12 years of age or younger, negative interactions with siblings were the most frequent, followed by negative interactions with mother ($t[51]=4.35, p<.001$; Table 1). Negative interactions with mothers were similar in frequency to fathers ($t[51]=1.48, p=.146$) and friends ($t[51]=1.34, p=.185$). For participants 13 years of age or older, negative interactions with siblings were the most frequent, together with friends ($t[59]=1.31, p=.194$). Next were negative interactions with mothers, which were less frequent than with siblings ($t[59]=2.28, p=.026$) but similar in frequency to friends ($t[59]=1.28, p=.205$). Negative interactions with fathers were the least frequent, significantly lower than mothers ($t[59]=2.98, p=.004$). The main effect of age ($F[1, 110]=0.20, p=.659$), as well as its interactions with Time ($F[1, 110]=0.49, p=.485$), and with Time and context ($F[3, 330]=0.92, p=.433$) were not significant.

Positive and negative spillover & associations

Day-level. For simplicity, we denote spillover of positive social interactions from one context to the other as *positive spillover* and spillover of negative social interactions *negative spillover*. Significant within-day positive and negative spillover was found between almost every couple of interactions examined – mother-father, mother-siblings, mother-friends, father-siblings, father-friends, and siblings-friends. Supplementary Tables S2 and S3 present all day-level spillover analyses. The only interactions that did not spill-over were negative interactions with friends; father-friends and sibling-friends negative spillovers in both directions were not

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

significant across data waves. Similarly, negative interactions with friends did not spill-over to negative interactions with mothers. Moderation by Wave was found for positive spillover between family and friends: positive spillover from mothers, fathers, and siblings to friends was weaker during COVID-19 versus prior (Table 2b-d, and Figure 1B-D). Negative spillover from friends to fathers was stronger during COVID-19 (Table 3, Figure 1E). The only moderation of within-family spillover was the positive spillover from fathers to mothers (Table 2a, Figure 1A), which intensified during COVID-19. Adjusting for gender and time altered the results in one case -- sibling-friend negative spillover became significant.

Person-level.

Positive person-level associations. The associations between positive interactions within family and from family to friends were positive and significant (Table 4). Specifically, individuals who had more positive interactions with one family member had more positive interactions with other family members as well as with friends. However, our hypothesis that the strength of the associations would increase during COVID-19 was not supported, as can be seen by the non-significant Context x Wave interaction.

Negative person-level associations. The associations between negative interactions within family and between family and friends were positive and significant (Table 5). Specifically, individuals who had more negative interactions with one family member had more negative interactions with other family members as well as with friends. Importantly, the strength of most of these associations changed significantly during COVID-19 in the predicted direction. The association between interactions with mothers and interactions with fathers (Table 4a, Figure 2A), as well as between fathers and siblings (Table 4b, Figure 2B) increased in strength, whereas the person-level associations between family members and friends decreased

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: [10.1037/dev0001211](https://doi.org/10.1037/dev0001211)

in strength (Table 4d, e, & f, Figure 2C-E), as can be seen by the significant Context x Wave interactions. The only association that remained unchanged was the person-level associations of negative interactions between mothers and siblings.

Changes in depressive symptoms during COVID-19

To examine our hypothesis that depressive symptoms would increase during COVID-19, and that this increase would be moderated by age and gender, we conducted a mixed model ANOVA with Time (prior to vs. during COVID-19) as the within-participant variable, and participants' Gender and Age (z-transformed) as between-participant factors. As predicted, there was a main effect of Time: there was a significant increase in depressive symptoms from prior to during COVID-19 ($F[1, 109]=19.82, p<.001, \eta^2=.15$; Table 1). This increase was not moderated by Age ($F[1, 109]=0.31, p=.579$) or Gender ($F[1, 109]=0.51, p=.476$). The main effect of Age was not significant ($F[1, 109]=2.81, p=.097$). However, the main effect of Gender was significant ($F[1, 109]=14.62, p<.001, \eta^2=.12$), indicating that, across data waves, girls had higher levels of depressive symptoms ($M=4.38, SE=0.38$) compared to boys ($M=2.20, SE=0.43$).

Do social interactions during COVID-19 explain increases in depressive symptoms?

To examine this, we conducted a hierarchical regression analysis predicting depressive symptoms during COVID-19. In the first step, depressive symptoms before COVID-19 were the predictor. In the second step of the regression, we added positive and negative interactions with friends and with family members. Due to the high associations between interactions within the family, we created a compound variable that summarized number of interactions across family members within individuals. In line with our hypotheses, positive and negative interactions with family members significantly predicted depressive symptoms during COVID-19 above and

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

beyond depressive symptoms before COVID-19 (Table 6). Number of interactions with friends did not significantly contribute to the model¹.

Discussion

The current study investigated how social dynamics of children and adolescents (youths) and their mental health changed during the emergence of COVID-19. Specifically, we explored changes in the frequency of these interactions as well as in the associations between them during COVID-19, and how these interactions subsequently related to increases in depressive symptoms. This is a unique investigation utilizing two waves of a daily diary design: one prior to COVID-19 and the other shortly after pandemic-induced school closures. The results largely support our hypotheses as they point to significant changes in social dynamics and depression during the pandemic and shed light on the association between the two.

One key finding was that the number of daily negative social interactions with friends decreased during COVID-19 for all participants. Interestingly, the decrease in positive interactions was observed only for younger participants (9-12 year old during COVID-19) – and was not evident at all for participants 13 years or older. Additionally, for older participants only, there was a significant increase in the number of positive interactions with siblings during COVID-19. Surprisingly, although the vast majority of our participants reported being at home with their parents during COVID-19, there was no change in the number of interactions with parents. Overall, it appears that younger participants were impacted from the sequela of COVID-19 via greater changes in social interactions. A possible explanation for this may be that older (vs. younger) adolescents have more advanced social skills (Bialecka-Pikul et al., 2020; Dumontheil et al., 2010), which they are able to use to preserve friendships and improve their relationships with their siblings. Importantly, given data on typical development, we would have

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

expected all our participants to show an increase in the number of interactions with friends, and decreases in interactions with parents (e.g., De Goede et al., 2009; Hadiwijaya et al., 2017; Lam et al., 2012; Larson & Richards, 1991; Larson, Richards, Moneta, Holmbeck, & Duckett, 1996). This suggests that the COVID-19 pandemic has significantly impacted the developmental trajectory of adolescents' interpersonal relationships.

Further, the current results show that it is not only the *number* of interactions that changed but also the *magnitude* of their reciprocal influence, evaluated by spillover processes and person-level associations. In general, across interaction valence and both data waves, results highlighted within-family as well as family-to-friends (and vice-versa) daily spillover processes as well as person-level associations. Importantly, the current results show that it is not only the number of social interactions that changed during COVID-19 but also the magnitude of their reciprocal influence. Prior to COVID-19, social interactions in one social context were significantly and positively associated with social interactions in other social contexts (e.g., positive interactions with mother were associated with positive interactions with father). These associations were found both at day-level and person level. As predicted, the strength of the associations changed during COVID-19. Specifically, on the day level, we detected significant decreases in positive spillover (i.e., within-day association) from family members to friends during COVID-19, and an intensification of fathers-to-mothers positive spillover. Contrary to predictions, friends-to-father negative spillover was stronger (vs. weaker) during COVID-19.

Person-level associations further supported our hypotheses: the association of mother-father and father-sibling negative interactions increased, whereas family-friends associations of negative interactions decreased during COVID-19. Since what was assessed here was the change in person-level correlation, these results reflect an intensification of individual differences –

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: [10.1037/dev0001211](https://doi.org/10.1037/dev0001211)

youths who had, for example, a high correlation between negative interactions with father and negative interactions with mother, had an even higher correlation now. Taken together, these results suggest that social dynamics during COVID-19 changed in ways that rendered the family more of a “closed system”, decreasing spillover/associations outside it and increasing daily spillover and person-level associations within it. It should be noted that the pattern of decreases in the family-to-friend association is in line with normative developmental patterns (e.g., Larson et al., 1996; Lam et al., 2012), and therefore we cannot conclude that they are facilitated by COVID-19 (vs. peers). Nevertheless, the combination of reductions in day-level positive spillover with increases in within-person association of negative interactions suggests that COVID-19 has had a detrimental effect on social dynamics, but more so to youth who had more negative family relationships prior to the pandemic.

In addition to exploring changing social dynamics due to COVID-19, we also examined its mental health consequences. As predicted, youth experienced significant increases – almost 40% – in depressive symptoms over just under one year. Importantly, a higher frequency of negative interactions and a lower frequency of positive interactions with family members partially accounted for these increases. These findings are in line with prior longitudinal studies showing that adolescents experience increases in depressive symptoms during times of stress and uncertainty (Jenness et al., 2019), and specifically during COVID-19 (Chahal et al., 2020). Moreover, the findings are in line with prior work on the social context of depression (Schwartz et al., 2012), and on the impact of social deprivation (Orben et al., 2020). Importantly, these findings add to the accumulating literature showing the mental health impact of COVID-19 (Choi et al., 2020; Wang et al., 2020; WHO, 2020).

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

It is important to note that – unlike most of the findings regarding changes in social interactions, which were opposite to the expected course of typical development, and therefore likely caused by COVID-19 – results regarding depression are more challenging to interpret. Studies have long established typical increases in depressive symptoms from middle childhood to adolescence (e.g., Salk et al., 2016), with steeper increases in girls (e.g., Marcotte et al., 2002; Salk et al., 2016). Although we found increases in depression symptoms from Wave 1 to Wave 2, this increase was not associated with the number of days between Wave 1 and Wave 2, or with age. Furthermore, we found that, although girls had higher levels of depressive symptoms in both data waves, gender did not moderate the increases in symptoms, as would be expected in typical development (e.g., Salk et al., 2016). This suggests that COVID-19 acted as a stressor for both boys and girls (versus other stressors that emerge during adolescence, such as body image; Marcotte et al., 2002). Nevertheless, we cannot completely separate typical developmental changes in depression from the impact of COVID-19. Another important caveat is the issue of directionality. Although changes in social interactions were associated with changes in depressive symptoms, we also believe that this association is likely significant in the opposite direction as well; in fact, prior literature has shown that the association between depression and social interactions is bi-directional (e.g., Hankin et al., 2010).

Collectively, our results suggest that, during the early stages of COVID-19, the family unit's importance increased, whereas peer-group influence decreased – a pattern that opposes the expected developmental trajectory (e.g., De Goede et al., 2009; Hadiwijaya et al., 2017; Lam et al., 2012; Larson & Richards, 1991; Larson et al., 1996). These findings have at least two possible explanations. The first and more obvious explanation is that, despite youths having grown up in an electronic age with ubiquitous online communication (Jensen et al., 2019),

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

physical presence and encounters with peers *still matters*. Indeed, similar concerns have arisen for children hospitalized for extended periods and who are thus away from school for a long time (Wadley et al., 2014). Considering that almost all the youths in our sample no longer had in-person school or after-school activities, this interpretation has important public policy implications, as it suggests that conducting school online poses unique challenges for adolescents, especially those whose home life is more disrupted. Further, increased youth reliance on the family unit, especially since COVID-19 is ongoing, may exert a high price on parents. The current study focused on the children's perspective; but social dynamics within family include the parents. Future research should examine parents' needs and test ways of providing them with the support they may require.

Another possible explanation for the increased importance and influence of the family during COVID-19 is less specific. In times of more extreme stress and threat from the outside world, youths may regress back into the safety of their immediate family (Pfefferbaum, Jacobs, Jones, Reyes, & Wyche, 2017). Indeed, research on youths exposed to natural disasters acknowledges that the parents' role in offering support and safety becomes amplified due to youths' greater need for this reassurance post-disaster (Costa, Weems, & Pina, 2009). Although the family unit may indeed serve as a protective factor at such times (Banks & Weems, 2014), this phenomenon of safety-seeking within the family may thwart youths' developmental goals, especially as the pandemic persists. Thus, although physical health is – and should be – our utmost priority, it is important to try and find creative ways for youths to interact with their peers amidst this crisis. For example, schools may encourage students to work on assignments in person, in pairs (or small “pods”), or via virtual one-on-one meetings, which may create more

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

socialization opportunities. Families could try and ensure youths have the means and opportunities to communicate with peers privately, whenever possible.

Although these findings may appear grim, we are encouraged by two hopeful observations regarding within-family bonds. First, even during the pandemic, positive interactions were significantly more frequent compared to negative interactions. Moreover, more positive experiences with family members were associated with less steep increases in depressive symptoms during COVID-19. Second, the steep increase in positive (but not negative) interactions with siblings found for older adolescents during the pandemic also offers some hope. This is particularly interesting as when surveyed pre-pandemic, older adolescents had the least amount of social interactions with their siblings. However, during the early stages of COVID-19, they had as many positive interactions with siblings as with peers; indeed, the only significant increase in positive interactions (almost 30%) was with siblings. Together with the absence of change in negative interactions with siblings, this finding suggests that the sibling relationship may serve, at least for older adolescents, as a natural compensation for the decrease in peer relationships. Consistently, previous studies have shown that siblings can have a beneficial influence on youth's mental health (McHale, Updegraff, & Whiteman, 2012); this may even be intensified by COVID-19. Pre-COVID research showed that adolescents without siblings experience greater levels of solitude than adolescents with siblings, and that those without siblings also derive lower levels of satisfaction from both time with friends and solitude (Wikle, Ackert, & Jenson, 2019). Given the social isolation induced by the pandemic, it is likely that the impact of being an only child may exacerbate the experience of loneliness in youths.

One of the main contributions of the current investigation is our focus on a wide assessment of youths' social networks. Although the developmental psychology literature widely

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

recognizes the important role of the social context during adolescence (Flynn, Felmlee, & Conger, 2017; Orben et al., 2020) and acknowledges the influence of parents (Brinberg, Fosco, & Ram, 2017), siblings (McHale et al., 2012), and peers (Schacter & Margolin, 2019), the majority of the literature examines the family separately from the peer group (Schacter & Margolin, 2019; though see Kaufman et al., 2020), rather than exploring these contexts together. The current study is one of the first to investigate the reciprocal associations between youths' interactions with family members and friends (Kaufman et al., 2020; Chung et al., 2011) and adds to past work in several ways. First, we investigated how an external stressor alters the strength of these associations. Second, whereas previous studies on spillover focused on a single positive (e.g., support; Schacter & Margolin, 2019) or negative (e.g., conflict; Ehrlich et al., 2012) interaction, we included a wide range of both positive and negative interactions. Thus, our study reveals that the spillover process is relevant to a wide range of behaviours.

Clinical implications

The results of the present study have direct implications for clinical practice and public health. First and foremost, our data describe changes in the social dynamics of youths and suggest that their normative developmental trajectory may be thwarted. In typical development, adolescence is a time of transition from reliance on immediate family to seeking support from peer groups, thus establishing autonomy and self-identity (Savard, Joussemet, Emond Pelletier, & Mageau, 2013). The opposite pattern of development appeared in our sample, whereby youths became more emotionally removed from peers and instead, more intimately involved in family relationships. This finding – in conjunction with the increased rates of depressive symptoms in our sample – highlights the importance of creating a normative environment in which youth can interact with peers, preferably in person. More broadly, this topic has far-reaching implications

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

even beyond COVID-19, as it can generalize to youths who may be isolated from their peers in other circumstances such as illness, war, or immigration, to name a few.

Limitations and future research

The present research has several limitations. First, COVID-19 is a “natural experiment.” This means that we could not control intervening variables or examine dose-response effects. Therefore, although the analyses ruled out the alternative explanation that time between Waves explains the results, our ability to deduce causality is limited. Future studies manipulating spillover processes or examining other types of stressors are needed to better understand the nature of changes in spillover processes. Similarly, our assessment of day-level spillover is constrained to same-day processes, limiting our understanding of temporal precedence in these effects; future studies examining different time lags are needed. Secondly, while daily diaries are far better than single timepoint retrospective data, they nevertheless also rely on self-reports (Stone, Shiffman, & Atienza, 2007). Thus, future research using, for example, reports from other informants (e.g., parents) are required to further validate the findings. Third, from a statistical perspective, we tested each hypothesis multiple times for several types of spillover. This is mitigated by the fact that findings followed consistent patterns (e.g., multiple types of spillover to friends were weaker during COVID-19), and largely remained significant when controlling for gender and time that passed between Waves. Still, future studies could focus on specific types of spillover and replicate our findings. Another limitation of our statistical analyses is the use of multilevel models, rather than more advanced statistical methods that may be better suited for our data. Like other data sets that combine multiple waves of intensive longitudinal designs, our data combined two different questions: (1) how do youth change over a year (from before to during COVID) – specifically how did the mean levels change (e.g., mean level of depressive

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

symptoms), and (2) how daily variability around the mean in one variable explains daily variability around the mean in another variable. Growth models are best suited to examine the first question, whereas multilevel models are best suited to examine the second question (McNeish & Hamaker, 2020). Recently developed methods combine these models of change (e.g., Dynamic Structural Equation Modelling) and could be a better fit for similar datasets that combine panel designs with intensive longitudinal designs as we did here. Fourth, the current study captured youths' initial reactions to COVID-19 and school closures. It is quite possible that, as youths and their families start adapting to the "new normal," social dynamics will revert back to their normal developmental course and depressive symptoms will subside. However, the animal/preclinical literature suggests that social deprivation from peers during adolescence may have long-term effects on mental health and brain development (Orben et al., 2020). Conducting additional follow-ups is necessary to characterize the long-term sequelae of COVID-19 on youths' social-emotional development. Finally, since we aimed to explore the social context in all of its complexity (i.e., four different relational contexts, with positive and negative interactions in each), we conducted a high number of statistical tests. Although we believe that there are theoretical benefits to understanding the complexity of the social context, we recognize that conducting multiple analyses decreased the power of the study. Future studies using larger samples are needed to replicate these findings.

Summary

The present study is an innovative investigation of social dynamics and mental health symptoms in children and adolescents, and how they changed during the COVID-19 pandemic and its associated quarantine. To examine these factors, we used a unique design in which youths filled out daily diaries twice: approximately a year prior to the onset of COVID-19 as well as

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: [10.1037/dev0001211](https://doi.org/10.1037/dev0001211)

during the early stages of the pandemic. Our results suggest that quarantine and the restriction of in-person interactions to the immediate family thwarted normative youths' social-emotional development. Rather than gaining more independence from their family and affiliating more with friends – as is foundational in typical developmental trajectories – youths became more dependent on their family members. The present investigation demonstrated the relevance of social interactions to mental health by showing that social interactions contributed to increases in depressive symptoms observed in youth during COVID-19.

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

References

- Allgaier, A. K., Pietsch, K., Frühe, B., Sigl-Glöckner, J., & Schulte-Körne, G. (2012). Screening for depression in adolescents: validity of the patient health questionnaire in pediatric care. *Depression and Anxiety*, 29, 906-913. <https://doi.org/10.1002/da.21971>
- Banks, D. M., & Weems, C. F. (2014). Family and peer social support and their links to psychological distress among hurricane-exposed minority youth. *American Journal of Orthopsychiatry*, 84, 341–352. <https://doi.org/10.1037/ort0000006>
- Bialecka-Pikul, M., Szpak, M., Zubek, J., Bosacki, S., & Kołodziejczyk, A. (2020). The psychological self and advanced theory of mind in adolescence. *Self and Identity*, 19, 85-104.
- Bell, H. C., Pellis, S. M., & Kolb, B. (2010). Juvenile peer play experience and the development of the orbitofrontal and medial prefrontal cortices. *Behavioural brain research*, 207, 7–13. <https://doi.org/10.1016/j.bbr.2009.09.029>
- Bolger, N., & Laurenceau, J. P. (2013). *Intensive longitudinal methods: An introduction to diary and experience sampling research*. Guilford Press.
- Brinberg, M., Fosco, G. M., & Ram, N. (2017). Examining inter-family differences in intra-family (parent–adolescent) dynamics using grid-sequence analysis. *Journal of Family Psychology*, 31, 994 –1004. <https://doi.org/10.1037/fam0000371>
- Bronfenbrenner, U., & Morris, P. A. (2006). The bioecological model of human development. In W. Damon & R. M. Lerner (Eds.), *Handbook of child psychology, Vol. 1: Theoretical models of human development* (6th ed., pp. 793 – 828). New York: Wiley. <https://doi.org/10.1002/9780470147658.chpsy0114>

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

Brooks, S. K., Webster, R. K., Smith, L. E., Woodland, L., Wessely, S., Greenberg, N., & Rubin,

G. J. (2020). The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *The Lancet*, 395, 912-920. [https://doi.org/10.1016/S0140-6736\(20\)30460-8](https://doi.org/10.1016/S0140-6736(20)30460-8)

Centers for Disease Control and Prevention. (2020). Interim guidance on management of coronavirus disease 2019 (COVID-19) in correctional and detention facilities.

Chahal, R., Kirshenbaum, J. S., Miller, J. G., Ho, T. C., & Gotlib, I. H. (2020). Higher executive control network coherence buffers against puberty-related increases in internalizing symptoms during the COVID-19 pandemic. *Biological Psychiatry*. <https://doi.org/10.1016/j.bpsc.2020.08.010>

Choi, K. R., Heilemann, M. V., Fauer, A., & Mead, M. (2020). A Second Pandemic: Mental Health Spillover From the Novel Coronavirus (COVID-19). *Journal of the American Psychiatric Nurses Association*, 26, 340–343. <https://doi.org/10.1177/1078390320919803>

Chung, G. H., Flook, L., & Fuligni, A. J. (2011). Reciprocal associations between family and peer conflict in adolescents' daily lives. *Child Development*, 82, 1390–1396. <https://doi.org/10.1111/j.1467-8624.2011.01625.x>

Cohodes, E. M., Kitt, E. R., Baskin-Sommers, A., & Gee, D. G. (2020). Influences of early-life stress on frontolimbic circuitry: Harnessing a dimensional approach to elucidate the effects of heterogeneity in stress exposure. *Developmental Psychobiology*. doi: 10.1002/dev.21969.

Costa, N. M., Weems, C. F., & Pina, A. A. (2009). Hurricane Katrina and youth anxiety: The role of perceived attachment beliefs and parenting behaviors. *Journal of Anxiety Disorders*, 23, 935-941. <https://doi.org/10.1016/j.janxdis.2009.06.002>

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

Davies, P.T., Sturge-Apple, M.L. (2014). Family context in the development of psychopathology.

In: Lewis, M., Rudolph, K.D., editors. *Handbook of developmental psychopathology*.

Springer; US: 2014. p. 143-161.

De Goede, I. H., Branje, S. J., Delsing, M. J., & Meeus, W. H. (2009). Linkages over time

between adolescents' relationships with parents and friends. *Journal of Youth and*

Adolescence, 38, 1304-1315. doi: 10.1007/s10964-009-9403-2

Dumontheil, I., Apperly, I. A., & Blakemore, S. J. (2010). Online usage of theory of mind

continues to develop in late adolescence. *Developmental Science*, 13, 331-338. DOI:

10.1111/j.1467-7687.2009.00888.x

Ehrlich, K. B., Dykas, M. J., & Cassidy, J. (2012). Tipping points in adolescent adjustment:

predicting social functioning from adolescents' conflict with parents and friends. *Journal*

of Family Psychology, 26, 776–783. <https://doi.org/10.1037/a0029868>

Ellis, W. E., Dumas, T. M., & Forbes, L. M. (2020). Physically isolated but socially connected:

Psychological adjustment and stress among adolescents during the initial COVID-19

crisis. *Canadian Journal of Behavioural Science*, 52, 177-187.

<http://dx.doi.org/10.1037/cbs0000215>

Elmer T, Mepham K, Stadtfeld C (2020). Students under lockdown: Comparisons of students'

social networks and mental health before and during the COVID-19 crisis in Switzerland.

PLoS ONE, 15: e0236337. <https://doi.org/10.1371/journal.pone.0236337>

Flook, L., & Fuligni, A. J. (2008). Family and school spillover in adolescents' daily lives. *Child*

Development, 79, 776-787. <https://doi.org/10.1111/j.1467-8624.2008.01157.x>

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

- Flynn, H. K., Felmlee, D. H., & Conger, R. D. (2017). The social context of adolescent friendships: Parents, peers, and romantic partners. *Youth & Society*, 49, 679-705.
<https://doi.org/10.1177/0044118X14559900>
- Fredrickson, B. L. (2013). Positive emotions broaden and build. In *Advances in experimental social psychology* (Vol. 47, pp. 1-53). Academic Press.
- Gadassi, R., Mor, N., & Rafaeli, E. (2011). Depression and empathic accuracy in couples: An interpersonal model of gender differences in depression. *Psychological Science*, 22, 1033-1041. <https://doi.org/10.1177/0956797611414728>
- Gadassi Polack, R., Chertkof, J., Kober, H., & Joormann, J. (in press). Maternal depression history moderates the association between criticism (but not praise) and depressive symptoms in youth. *Research on Child and Adolescent Psychopathology*.
- Gadassi Polack, R., Sened, H., Joormann, J., & Kober, H. (2021, April 21). Spillover. Retrieved from osf.io/eq62h
- Ghebreyesus, T.A. (2020). Addressing mental health needs: an integral part of COVID-19 response. *World Psychiatry*, 19, 129–130. <https://doi.org/10.1002/wps.20768>
- Gleason, M. E., Iida, M., Shrout, P. E., & Bolger, N. (2008). Receiving support as a mixed blessing: evidence for dual effects of support on psychological outcomes. *Journal of Personality and Social Psychology*, 94, 824–838. <https://doi.org/10.1037/0022-3514.94.5.824>
- Hadiwijaya, H., Klimstra, T. A., Vermunt, J. K., Branje, S. J., & Meeus, W. H. (2017). On the development of harmony, turbulence, and independence in parent–adolescent relationships: A five-wave longitudinal study. *Journal of Youth and Adolescence*, 46, 1772-1788. DOI 10.1007/s10964-016-0627-7

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

Hankin, B. L., Stone, L., & Wright, P. A. (2010). Corumination, interpersonal stress generation, and internalizing symptoms: accumulating effects and transactional influences in a multiwave study of adolescents. *Development and Psychopathology*, 22, 217–235.
<https://doi.org/10.1017/S0954579409990368>

Jenness, J. L., Peverill, M., King, K. M., Hankin, B. L., & McLaughlin, K. A. (2019). Dynamic associations between stressful life events and adolescent internalizing psychopathology in a multiwave longitudinal study. *Journal of Abnormal Psychology*, 128, 596–609. <https://doi.org/10.1037/abn0000450>

Jensen, M., George, M. J., Russell, M. R., & Odgers, C. L. (2019). Young adolescents' digital technology use and mental health symptoms: Little evidence of longitudinal or daily linkages. *Clinical Psychological Science*, 7, 1416-1433. doi: 10.1177/2167702619859336.

Kaufman, T. M., Kretschmer, T., Huitsing, G., & Veenstra, R. (2020). Caught in a vicious cycle? Explaining bidirectional spillover between parent-child relationships and peer victimization. *Development and Psychopathology*, 32, 11-20. doi: 10.1017/S0954579418001360.

Krishnakumar, A., & Buehler, C. (2000). Interparental conflict and parenting behaviors: A meta-analytic review. *Family Relations*, 49, 25-44. <https://doi.org/10.1111/j.1741-3729.2000.00025.x>

Koçak, A., Mouratidis, A., Sayıl, M., Kindap-Tepe, Y., & Uçanok, Z. (2017). Interparental conflict and adolescents' relational aggression and loneliness: The mediating role of maternal psychological control. *Journal of Child and Family Studies*, 26, 3546-3558.
DOI 10.1007/s10826-017-0854-x

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

Kovacs, M. (1985). The Children's Depression Inventory. *Psychopharmacology Bulletin*, 21, 995–998.

Krull, J. L., & MacKinnon, D. P. (2001). Multilevel modeling of individual and group level mediated effects. *Multivariate Behavioral Research*, 36, 249-277. doi: 10.1207/S15327906MBR3602_06.

Lam, C. B., McHale, S. M., & Crouter, A. C. (2012). Parent–child shared time from middle childhood to late adolescence: Developmental course and adjustment correlates. *Child Development*, 83, 2089-2103. DOI: 10.1111/j.1467-8624.2012.01826.x

Larson, R., & Richards, M. H. (1991). Daily companionship in late childhood and early adolescence: Changing developmental contexts. *Child Development*, 62, 284-300. <https://doi.org/10.2307/1131003>

Larson, R. W., Richards, M. H., Moneta, G., Holmbeck, G., & Duckett, E. (1996). Changes in adolescents' daily interactions with their families from ages 10 to 18: Disengagement and transformation. *Developmental Psychology*, 32, 744–754. <https://doi.org/10.1037/0012-1649.32.4.744>

Lindblom, J., Vänskä, M., Flykt, M., Tolvanen, A., Tiitinen, A., Tulppala, M., & Punamäki, R. L. (2017). From early family systems to internalizing symptoms: The role of emotion regulation and peer relations. *Journal of Family Psychology*, 31, 316-326. <http://dx.doi.org/10.1037/fam0000260>

Marcotte, D., Fortin, L., Potvin, P., & Papillon, M. (2002). Gender differences in depressive symptoms during adolescence: Role of gender-typed characteristics, self-esteem, body image, stressful life events, and pubertal status. *Journal of Emotional and Behavioral Disorders*, 10, 29-42. <https://doi.org/10.1177/106342660201000104>

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

Mastrotheodoros, S., Van Lissa, C. J., Van der Graaff, J., Deković, M., Meeus, W. H., & Branje,

S. J. (2020). Day-to-day spillover and long-term transmission of interparental conflict to adolescent–mother conflict: The role of mood. *Journal of Family Psychology*. doi: 10.1037/fam0000649.

McHale, S. M., Updegraff, K. A., & Whiteman, S. D. (2012). Sibling Relationships and Influences in Childhood and Adolescence. *Journal of Marriage and the Family*, 74, 913–930. <https://doi.org/10.1111/j.1741-3737.2012.01011.x>

[McNeish, D., & Hamaker, E. L. \(2020\). A primer on two-level dynamic structural equation models for intensive longitudinal data in Mplus. *Psychological Methods*, 25, 610–635. <https://doi.org/10.1037/met0000250>](https://doi.org/10.1037/met0000250)

Minuchin, P. (1985). Families and individual development: Provocations from the field of family therapy. *Child Development*, 56, 289–302. DOI: 10.2307/1129720

Nelson, E. E., Leibenluft, E., McClure, E. B., & Pine, D. S. (2005). The social re-orientation of adolescence: a neuroscience perspective on the process and its relation to psychopathology. *Psychological Medicine*, 35, 163174. DOI: 10.1017/S0033291704003915

Nezlek, J. B. (2017). A practical guide to understanding reliability in studies of within-person variability. *Journal of Research in Personality*, 69, 149–155. <https://doi.org/10.1016/j.jrp.2016.06.020>

Orben, A., Tomova, L., & Blakemore, S. J. (2020). The effects of social deprivation on adolescent development and mental health. *The Lancet Child & Adolescent Health*. [https://doi.org/10.1016/S2352-4642\(20\)30186-3](https://doi.org/10.1016/S2352-4642(20)30186-3)

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: [10.1037/dev0001211](https://doi.org/10.1037/dev0001211)

Panksepp J. (1981). The ontogeny of play in rats. *Developmental Psychobiology*, 14, 327–332.

<https://doi.org/10.1002/dev.420140405>

Petersen, I. T., Lindhiem, O., LeBeau, B., Bates, J. E., Pettit, G. S., Lansford, J. E., & Dodge, K.

A. (2018). Development of internalizing problems from adolescence to emerging adulthood: Accounting for heterotypic continuity with vertical scaling. *Developmental Psychology*, 54, 586–599. <https://doi.org/10.1037/dev0000449>

Pfefferbaum, B., Jacobs, A. K., Jones, R. T., Reyes, G., & Wyche, K. F. (2017). A skill set for supporting displaced children in psychological recovery after disasters. *Current Psychiatry Reports*, 19, 60. <https://doi.org/10.1007/s11920-017-0814-6>

Pinheiro, J., Bates, D., DebRoy, S., & Sarkar, D. (2014). R Core Team (2014) nlme: linear and nonlinear mixed effects models. R package version 3.1-117. Available at <http://CRAN.R-project.org/package=nlme>.

R Core Team (2013). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>

Salk, R. H., Petersen, J. L., Abramson, L. Y., & Hyde, J. S. (2016). The contemporary face of gender differences and similarities in depression throughout adolescence: Development and chronicity. *Journal of Affective Disorders*, 205, 28-35.
<http://dx.doi.org/10.1016/j.jad.2016.03.071>

Savard, A., Joussemet, M., Pelletier, J. E., & Mageau, G. A. (2013). The benefits of autonomy support for adolescents with severe emotional and behavioral problems. *Motivation and Emotion*, 37, 688-700. doi: [10.1007/s11031-013-9351-8](https://doi.org/10.1007/s11031-013-9351-8)

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: [10.1037/dev0001211](https://doi.org/10.1037/dev0001211)

- Schacter, H. L., & Margolin, G. (2019). The Interplay of Friends and Parents in Adolescents' Daily Lives: Towards A Dynamic View of Social Support. *Social development*, 28, 708–724. <https://doi.org/10.1111/sode.12363>
- Schwartz, O. S., Dudgeon, P., Sheeber, L. B., Yap, M. B., Simmons, J. G., & Allen, N. B. (2012). Parental behaviors during family interactions predict changes in depression and anxiety symptoms during adolescence. *Journal of Abnormal Child Psychology*, 40, 59–71. doi: 10.1007/s10802-011-9542-2.
- Shrout, P. E., & Lane, S. P. (2012). *Psychometrics*. In M. R. Mehl & T. S. Conner (Eds.), *Handbook of research methods for studying daily life* (p. 302–320). The Guilford Press.
- Stone, A., Shiffman, S., Atienza, A., & Nebeling, L. (2007). *The science of real-time data capture: Self-reports in health research*. Oxford University Press.
- Timmons, A. C., & Margolin, G. (2015). Family conflict, mood, and adolescents' daily school problems: moderating roles of internalizing and externalizing symptoms. *Child Development*, 86, 241–258. <https://doi.org/10.1111/cdev.12300>
- Trull, T. J., & Ebner-Priemer, U. W. (2020). Ambulatory assessment in psychopathology research: A review of recommended reporting guidelines and current practices. *Journal of Abnormal Psychology*, 129, 56–63. <https://doi.org/10.1037/abn0000473>
- Wadley, G., Vetere, F., Hopkins, L., Green, J., & Kulik, L. (2014). Exploring ambient technology for connecting hospitalised children with school and home. *International Journal of Human-Computer Studies*, 72, 640–653. <https://doi.org/10.1016/j.ijhcs.2014.04.003>

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

Wang, C., Pan, R., Wan, X., Tan, Y., Xu, L., Ho, C. S., & Ho, R. C. (2020). Immediate Psychological Responses and Associated Factors during the Initial Stage of the 2019 Coronavirus Disease (COVID-19) Epidemic among the General Population in China. *International Journal of Environmental Research and Public Health*, 17, 1729. <https://doi.org/10.3390/ijerph17051729>

World Health Organization. Facing Mental Health Fallout from the Coronavirus Pandemic. <https://www.who.int/news-room/feature-stories/detail/facing-mental-health-fallout-from-the-coronavirus-pandemic> (29 September 2020, date last accessed).

Wikle, J. S., Ackert, E., & Jensen, A. C. (2019). Companionship patterns and emotional states during social interactions for adolescents with and without siblings. *Journal of Youth and Adolescence*, 48, 2190–2206. <https://doi.org/10.1007/s10964-019-01121-z>

Zhang, Z., Zyphur, M. J., & Preacher, K. J. (2009). Testing multilevel mediation using hierarchical linear models: Problems and solutions. *Organizational Research Methods*, 12, 695–719. <https://doi.org/10.1177/1094428108327450>

Tables

Table 1. *Depressive symptoms and social interactions before and during COVID-2019.*

		Wave 1	Wave 2	% change	<i>t</i>	<i>p</i> -value
Depressive symptoms	Younger ¹	1.99 (2.54)	3.36 (3.87)	+69%	3.50	.001
	Older	3.60 (3.42)	4.48 (3.68)	+24%	2.74	.008
	All	2.85 (3.14)	3.96 (3.79)	+39%	4.24	<.001
Positive interactions						
Mother	Younger	2.46 (1.45)	2.56 (1.72)	+4%	0.44	.658
	Older	1.47 (1.41)	1.53 (1.61)	+4%	0.38	.702
	All	1.93 (1.50)	2.01 (1.73)	+4%	0.59	.556
Father	Younger	1.97 (1.38)	2.05 (1.70)	+4%	0.44	.665

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

	Older	1.05 (1.38)	0.94 (1.43)	-11%	-1.00	.322
	All	1.48 (1.45)	1.45 (1.66)	-2%	-0.22	.829
Sibling	Younger	1.17 (1.24)	1.46 (1.52)	+25%	1.86	.069
	Older	0.67 (0.93)	0.87 (1.12)	+30%	2.03	.047
	All	0.90 (1.11)	1.15 (1.35)	+28%	2.71	.008
Friend	Younger	1.44 (1.16)	0.74 (1.15)	-49%	-4.86	<.001
	Older	1.51 (1.18)	1.32 (1.29)	-13%	1.11	.271
	All	1.47 (1.17)	1.05 (1.26)	-29%	-3.70	<.001
Negative interactions						
Mother	Younger	0.27 (0.35)	0.37 (0.41)	+37%	1.84	.071
	Older	0.38 (0.51)	0.26 (0.31)	-32%	-2.26	.028
	All	0.33 (0.44)	0.31 (0.36)	-6%	-0.46	.642
Father	Younger	0.20 (0.25)	0.33 (0.51)	+65%	1.94	.058
	Older	0.22 (0.38)	0.17 (0.33)	-23%	-1.02	.310
	All	0.21 (0.32)	0.25 (0.43)	+19%	0.90	.369
Sibling	Younger	0.71 (0.86)	0.76 (0.89)	+7%	0.44	.661
	Older	0.53 (0.94)	0.49 (0.84)	-7%	-0.40	.694
	All	0.62 (0.91)	0.62 (0.87)	0%	-0.01	.999
Friend	Younger	0.38 (0.60)	0.13 (0.22)	-66%	-3.42	.001
	Older	0.45 (0.57)	0.32 (0.44)	-29%	-1.83	.072
	All	0.42 (0.58)	0.23 (0.36)	-45%	-3.65	<.001

¹Younger=participants who were 9-12 during Wave 2; Older = participants who were 13-17 during Wave 2.

Table 2. *Day level positive spillover within and outside the family: Spillover moderated by data wave only*

						Moderation by wave		
		Estimate (SE)	t	df	p	t	df	p
(a) Outcome: positive	Intercept	-.02(.02)	-0.917	4790	.359	-0.521	4790	.603
	Previous day interactions mother	.14(.02)	5.709	4790	< .001	-1.198	4790	.231

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

		Moderation by wave						
		Estimate (SE)	t	df	p	t	df	p
interactions	Same day interactions father	.46(.04)	12.380	4790	< .001	2.369	4790	.018
mother	Mean interactions mother	.97(.01)	76.845	4790	< .001	2.009	4790	.045
	Mean interactions father	.03(.01)	2.309	4790	.021	-1.680	4790	.093
(b) Outcome: positive interactions friends	Intercept	0.01(.03)	0.471	4790	.638	-0.938	4790	.348
	Previous day interactions friends	0.25(.03)	9.313	4790	< .001	-2.816	4790	.005
	Same day interactions father	0.19(.03)	5.747	4790	< .001	-3.406	4790	.001
	Mean interactions friends	1.01(.01)	68.205	4790	< .001	-0.417	4790	.676
	Mean interactions father	0(.01)	-0.057	4790	.955	0.353	4790	.724
(c) Outcome: positive interactions friends	Intercept	0.02(.03)	0.546	4790	.585	-1.008	4790	.314
	Previous day interactions friends	0.26(.03)	9.592	4790	< .001	-3.214	4790	.001
	Same day interactions mother	0.15(.03)	5.667	4790	< .001	-2.244	4790	.025
	Mean interactions friends	1.01(.01)	71.937	4790	< .001	-0.504	4790	.614
	Mean interactions mother	0(.01)	-0.161	4790	.872	0.584	4790	.559
(d) Outcome: positive interactions friends	Intercept	.01(.03)	0.503	4790	.615	-1.067	4790	.286
	Previous day interactions friends	.24(.03)	8.863	4790	< .001	-2.284	4790	.022
	Same day interactions siblings	.23(.04)	5.817	4790	< .001	-3.091	4790	.002
	Mean interactions friends	> .99(.01)	71.705	4790	< .001	0.293	4790	.770
	Mean interactions siblings	.02(.02)	1.387	4790	.166	-0.396	4790	.692

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

Table 3. *Day level negative spillover within and outside the family: Spillover moderated by data wave only*

						Moderation by wave		
		Estimate (SE)	t	df	p	t	df	p
Outcome: negative interactions father	Intercept	0(.01)	-0.118	4790	.906	-0.316	4790	.752
	Previous day interactions							
	father	0.02(.03)	0.774	4790	.439	2.469	4790	.014
	Same day interactions							
	friends	-0.02(.02)	-0.842	4790	.400	2.395	4790	.017
	Mean interactions father	1.06(.03)	38.944	4790	< .001	-1.355	4790	.175
	Mean interactions friends	-0.01(.02)	-0.572	4790	.568	0.127	4790	.899

Table 4. *Person-level associations of positive social interactions within and outside the family, as well as differences during COVID-19*

		β	SE	df	t	95% CI
(a)	Intercept	0.08	0.11	141	0.71	-0.14, 0.31
Outcome:	Positive interactions mother	0.69	0.04	109	15.98***	0.60, 0.78
Positive	Wave ¹	-0.11	0.11	109	-1.02	-0.33, 0.11
interactions	Positive interactions mother * wave	0.03	0.04	109	0.76	-0.05, 0.12
father						
(b)	Intercept	0.62	0.10	141	6.10***	0.42, 0.83
Outcome:	Positive interactions siblings	0.93	0.07	109	13.09***	0.79, 1.07
Positive	Wave	-0.21	0.12	109	-1.79	-0.45, 0.02
interactions	Positive interactions siblings * Wave	-0.04	0.08	109	-0.46	-0.19, 0.12
father						
(c)	Intercept	1.20	0.13	141	9.36***	0.95, 1.46
Outcome:	Positive interactions siblings	0.86	0.09	109	9.75***	0.69, 1.04
Positive	Wave	-0.15	0.15	109	-1.05	-0.44, 0.14
interactions	Positive interactions siblings * Wave	-0.03	0.10	109	-0.37	-0.22, 0.15
mother						
(d)	Intercept	1.13	0.15	141	7.37***	0.83, 1.44
Outcome:	Positive interactions mother	0.15	0.06	109	2.40*	0.03, 0.27
Positive	Wave	-0.46	0.18	109	-2.57*	-0.81, -0.11
interactions	Positive interactions mother * wave	0.03	0.07	109	0.44	-0.11, 0.17
friends						
(e)	Intercept	1.09	0.13	141	8.17***	0.83, 1.35
Outcome:	Positive interactions father	0.23	0.06	109	3.55***	0.10, 0.36
Positive	Wave	-0.37	0.15	109	-2.42*	-0.68, -0.07
interactions	Positive interactions father * wave	-0.01	0.07	109	-0.18	-0.16, 0.13
friends						
(f)	Intercept	1.04	0.11	141	9.07***	0.82, 1.27
Outcome:	Positive interactions siblings	0.44	0.08	109	5.38***	0.28, 0.60
Positive	Wave	-0.43	0.13	109	-3.00**	-0.72, -0.15
interactions	Positive interactions siblings * Wave	-0.07	0.09	109	-0.73	-0.25, 0.12
friends						

¹Wave coded 1=during COVID-19, 0=before; * $p < .05$, ** $p < .01$, *** $p < .001$

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

Table 5. *Person-level associations of negative social interactions within and outside the family, as well as differences during COVID-19*

		β	SE	df	<i>t</i>	95% CI
(a)	Intercept	0.06	0.03	141	2.09*	0.003, 0.11
Outcome:	Negative interactions mother	0.47	0.05	109	8.95***	0.36, 0.57
Negative	Wave ¹	-0.07	0.04	109	-1.85	-0.14, 0.005
interactions	Negative interactions mother * wave	0.37	0.08	109	4.95***	0.22, 0.52
father						
(b)	Intercept	0.10	0.03	141	3.25**	0.04, 0.16
Outcome:	Negative interactions siblings	0.17	0.03	109	5.80***	0.11, 0.23
Negative	Wave	-0.03	0.04	109	-0.77	-0.11, 0.05
interactions	Negative interactions siblings * Wave	0.11	0.04	109	2.78**	0.03, 0.19
father						
(c)	Intercept	0.18	0.04	141	5.11***	0.11, 0.25
Outcome:	Negative interactions siblings	0.20	0.03	109	5.93***	0.13, 0.27
Negative	Wave	0.02	0.05	109	0.54	-0.06, 0.11
interactions	Negative interactions siblings * Wave	-0.06	0.04	109	-1.36	-0.14, 0.03
mother						
(d)	Intercept	0.25	0.05	141	4.92***	0.15, 0.34
Outcome:	Negative interactions mother	0.56	0.09	109	5.94***	0.37, 0.74
Negative	Wave	-0.01	0.06	109	-0.20	-0.14, 0.12
interactions	Negative interactions mother * wave	-0.52	0.14	109	-3.78***	-0.79, -0.25
friends						
(e)	Intercept	0.24	0.05	141	5.10***	0.15, 0.33
Outcome:	Negative interactions father	0.88	0.13	109	6.97***	0.63, 1.12
Negative	Wave	-0.05	0.06	109	-0.87	-0.17, 0.07
interactions	Negative interactions father * Wave	-0.66	0.15	109	-4.38***	-0.95, -0.36
friends						
(f)	Intercept	0.28	0.05	141	5.63***	0.18, 0.38
Outcome:	Negative interactions siblings	0.24	0.05	109	4.95***	0.14, 0.33
Negative	Wave	-0.10	0.06	109	-1.61	-0.22, 0.02
interactions	Negative interactions siblings * Wave	-0.14	0.06	109	-2.21*	-0.26, -0.01
friends						

¹Wave coded 1=during COVID-19, 0=before; * $p < .05$, ** $p < .01$, *** $p < .001$

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: 10.1037/dev0001211

Table 6. *Predicting change in depressive symptoms from before to after COVID-19 from positive and negative interactions with family members and friends.*

Predictor		B	SE	<i>t</i>	<i>p</i>
Step 1	Intercept	1.47	0.24	4.37	<.001
	Depressive symptoms before COVID-19	0.87	0.08	10.94	<.001
	$F[1, 110]=119.73, p<.001, R^2=.52$				
Step 2	Intercept	1.74	0.50	3.50	<.001
	Depressive symptoms before COVID-19	0.76	0.08	9.25	<.001
	Positive interactions with friends	0.05	0.22	0.24	.812
	Negative interactions with friends	0.59	0.76	0.78	.438
	Positive interactions with family members	-0.15	0.06	-2.34	.020
	Negative interactions with family members	0.47	0.18	2.64	.010
		$F_{change}[4, 106]=3.58, p=.009, R^2=.06$			

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: [10.1037/dev0001211](https://doi.org/10.1037/dev0001211)

Figure 1. *Day level spillover processes before and during COVID-19.*

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: [10.1037/dev0001211](https://doi.org/10.1037/dev0001211)

Figure 2. *Person level associations of negative interactions across social contexts before and during COVID-19.*

©American Psychological Association, 2021. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. Please do not copy or cite without author's permission. The final article is available, upon publication, at: [10.1037/dev0001211](https://doi.org/10.1037/dev0001211)

ⁱ We repeated this analysis with participants age and the interactions between social contexts and age as predictors; however interactions with age were not significant ($ps \geq .073$) and therefore the regression is presented in its simpler form.