Title Page

Full Article Title: Familiarity of visual perceptual styles: Local and global visual processing in autism and first-degree relatives

Authors: Kritika Nayar, M.S.,¹ Frederick Shic, Ph.D.,^{2,3} Molly Winston, M.S.,¹ Cassandra J. Stevens, B.S.,¹ Janna Guilfoyle, M.S.,¹ Molly Losh, Ph.D.¹

Affiliations:

¹Neurodevelopmental Disabilities Lab, Roxelyn and Richard Pepper Department of Communication Sciences and Disorders, Northwestern University, Evanston, IL, USA

²Center for Child Health, Behavior and Development, Seattle Children's Research Institute, Seattle, WA, USA

³Department of Pediatrics, University of Washington School of Medicine, Seattle, WA, USA

Corresponding Author: Molly Losh; m-losh@northwestern.edu; 847-491-2431; 2240 Campus Drive, Frances Searle Building, #2-366, Evanston, IL 60208

Abstract:

Background: Individuals with ASD often demonstrate a local visual processing bias, with some evidence showing concurrent impaired global processing. Atypicalities in global visual processing are associated with increased ASD symptom severity, shedding light into potential cognitive mechanisms contributing to the ASD phenotype. Visual perceptual and social attentional differences have also been documented more subtly among parents of individuals with ASD. However, prior studies of local/global visual processing in parents show mixed findings, and primarily use accuracy and reaction time measures alone, which may not confer enough specificity to capture sub-clinical differences. This study aimed to address this gap by implementing an objective measurement of local and global perception via eye tracking in ASD and their parents.

Methods: Participants included 161 individuals (n=32 individuals with ASD and n=30 controls; n=56 parents of individuals with ASD and n=43 control parents), who completed two interactive tasks that tap local/global visual processing, while their gaze was tracked on an eye tracker. To examine local/global visual processing, performance (i.e., accuracy and reaction time) and a suite of eye-tracking variables were calculated. A series of linear mixed effects models assessed ASD versus control and ASD parent versus control parent group differences. Within-family associations of local/global perception, and associations with ASD-related clinical and subclinical traits were also examined.

Results: Significant differences emerged for performance (i.e., accuracy and/or reaction time) indices during more complex tasks in both proband and parent groups. Eye-tracking results further revealed evidence of reduced global versus local visual processing in ASD and ASD parent groups, with the ASD group demonstrating heightened local processing, in addition to reduced global perception, and parents showing evidence of a diminished bias towards global stimuli, but without indications of heightened local processing. Within family similarities of processing styles were observed for ASD parent-child pairs, which were not found in control

families. Finally, heightened local processing was related to more severe restricted and repetitive behaviors in individuals with ASD; whereas heightened local processing was unexpectedly associated with better pragmatic language abilities was observed in both ASD proband and ASD parent groups.

Conclusions: Findings demonstrated robust local/global visual processing differences using eye tracking in individuals with ASD, alongside more subtle, and nuanced differences in parents that together could point to underlying neurobiological differences in visual perception. Within-family associations observed suggest that visual processing styles uniquely co-segregate in ASD families, and add to evidence that visual processing styles may be heritable, genetically meaningful features of the broader autism spectrum.

Key words: Local and global visual processing, weak central coherence, enhanced perceptual functioning, broad autism phenotype, endophenotype, eye-tracking

Introduction

Local and global processing in ASD

An essential component of visual perception is *global processing*, or seeing the gist, which involves the integration of local segments of a scene to create an integrated whole in a rapid and automatic fashion (Kimchi, 1992; Navon, 1983). This perceptual process is the typical and default perceptual strategy used by adults (Navon, 1977) and undergoes a protracted developmental trajectory from local to global processing, with adult-like global perceptual strategies emerging after age 7 years (Nayar, Franchak, Adolph, & Kiorpes, 2015). In contrast, *local perception* is the selective attention to parts of a scene (Kovacs, 1996; Navon, 1977), usually characteristic of young children's perceptual styles, and generally slower than global processing (Freeseaman, Colombo, & Coldren, 1993; Kimchi, 1992; Navon, 1977; Nayar et al., 2015). These processes in tandem support the fluid interpretation of our environment, including the complex nuances and dynamics defining social-communicative interactions. A reduced bias towards seeing the "big picture" (i.e., weak central coherence theory) have been related to weaker social skills, verbal and nonverbal communication, and cognition more generally (Behrmann, Thomas, & Humphreys, 2006; Burnette et al., 2005; Happe, 1999; Jarrold, Butler, Cottington, & Jimenez, 2000; Jolliffe & Baron-Cohen, 2000; Klin, Jones, Schultz, Volkmar, & Cohen, 2002; Van Eylen, Boets, Steyaert, Wagemans, & Noens, 2018). Differences in visual perceptual styles, including a local perceptual bias (i.e., enhanced perceptual functioning theory), are also thought to underlie some aspects of the social-communicative difficulties observed in autism spectrum disorder (ASD; a neurodevelopmental disorder characterized by deficits in social communication and the presence of repetitive behaviors or circumscribed interests) (APA, 2013). For example, local perception may be reflected in difficulties interpreting non-verbal

communicative cues, whereby an individual may fixate only on one part of the face, making it challenging to interpret facial expressions (Van der Hallen, Evers, Brewaeys, Van den Noortgate, & Wagemans, 2015; Van Eylen et al., 2018). Similarly, findings suggest that intact global processing (or, a reduced bias for details as defined by descriptions of paintings that were less detailed) relate to more optimal outcomes in those with ASD (Fitch, Fein, & Eigsti, 2015).

Conflicting findings of local/global perceptual abilities in ASD have been reported, with some findings suggesting no differences from controls and others revealing superior local processing and/or deficits in global processing relative to controls (Frith, 1989; Happe, 1996; Koldewyn, Jiang, Weigelt, & Kanwisher, 2013; Milne & Scope, 2008; Mottron, Burack, Iarocci, Belleville, & Enns, 2003; Mottron, Burack, Stauder, & Robaey, 1999; Plaisted, Swettenham, & Rees, 1999; Rinehart, Bradshaw, Moss, Brereton, & Tonge, 2000; Ropar & Mitchell, 1999, 2001). Similar reports of conflicting findings have also been documented among first-degree relatives of individuals with ASD (Bolte & Poustka, 2006; Briskman, Happe, & Frith, 2001; Losh et al., 2009; Neufeld et al., 2020; Van Eylen et al., 2017). Notable, however, is that these prior studies primarily relied upon performance-based outcome measures (i.e., accuracy and reaction time measures alone) that are highly susceptible to effects of methodological context, such as providing guided instructions (see meta-analysis Van der Hallen et al., 2015). A study combining performance and eye-tracking methodologies to assay global perception in ASD (Nayar, Voyles, Kiorpes, & Di Martino, 2017), documented reduced global processing abilities in the absence of heightened local perception in ASD compared to controls. These findings were captured primarily using eye tracking and revealed clear strategy differences that may have been missed from performance measures alone. Whereas performance-based measures reflect the final bias or ability of local/global processing, eye tracking provides not only a closer examination of

mechanism, but also intermediate levels of measurement capturing either mechanistic-relevant precursors to local/global processing or process-focused measurement that provides additional granularity in the understanding of variation within and across groups.

Eye movement differences in the broad autism phenotype (BAP)

Evidence also suggests that eye-movement patterns, including saccade trajectories, are heritable in the general population (Constantino et al., 2017) and across psychiatric disorders (Ettinger et al., 2004), and may be studied as endophenotypes, or intermediate traits that are along the pathway between behavior and biology that are also observed subtly among unaffected first-degree relatives. For instance, in ASD, differences in local/global visual processing identified via eye tracking have helped to indirectly inform an understanding of the neurobiological mechanisms involved in local/global processing, particularly within the visual cortex, which may interact with higher-order biological processes tapping social attention and social communication (Altmann, Bulthoff, & Kourtzi, 2003; Altschuler et al., 2014; Altschuler et al., 2012; Doniger, Foxe, Murray, Higgins, & Javitt, 2002; Doniger et al., 2000; Doniger et al., 2001; Fink et al., 1997; Foxe, Murray, & Javitt, 2005; Keehn et al., 2020; Stroganova, Orekhova, et al., 2007; Stroganova et al., 2012). Gaze and eye movement may therefore serve as particularly good candidate endophenotypes, given their known heritability and association with ASD (Constantino et al., 2017; Neufeld et al., 2020); however, few studies have explored local and global visual perception using eye tracking in unaffected family members who are at increased genetic liability to ASD.

Clinically unaffected relatives of individuals with ASD may demonstrate personality features (e.g., aloof or rigid personality traits, pragmatic language differences) that mirror the

social and repetitive behavior domains of impairment in ASD (known as the broad autism phenotype), as well as subtle differences in global processing and social cognition (Adolphs, Spezio, Parlier, & Piven, 2008; Bolte & Poustka, 2006; Briskman et al., 2001; Chouinard, Unwin, Landry, & Sperandio, 2016; Constantino et al., 2017; Cribb, Olaithe, Di Lorenzo, Dunlop, & Maybery, 2016; Hogan-Brown, Hoedemaker, Gordon, & Losh, 2014; M. Lee et al., 2019; Mosconi et al., 2010; Neufeld et al., 2020). Eye movements have also been found to differ among parents of individuals with ASD compared to controls across a range of tasks tapping social attention (Groen et al., 2012; M. Lee et al., 2019) and language processing (Nayar et al., 2018; Nayar et al., 2021). In particular, parents presenting with BAP features showed reduced language automaticity that coupled atypical, perseverative (i.e., getting “stuck”) and regressive (i.e., moving backwards) eye fixations, which related to patterns of restricted and repetitive behaviors (RRBs) in their children with ASD (Nayar et al., 2018). These studies provide intriguing evidence that certain features of a stimulus may “trap” the attention of individuals with ASD and the BAP, and which may reflect a local perceptual style.

Local and global processing in the BAP

Prior studies have highlighted potential differences in local and global processing among relatives of individuals with ASD mirroring the visual processing styles identified among those with ASD (Bolte & Poustka, 2006; Briskman et al., 2001; Neufeld et al., 2020; Van der Hallen et al., 2015; Van Eylen et al., 2017; Van Eylen et al., 2018). Specifically, heightened local perception has been documented among parents of individuals with ASD (Bolte & Poustka, 2006; Happe, Briskman, & Frith, 2001), although this finding was not replicated in a larger sample of parents (Losh et al., 2009). Van Eylen et al. (2017) found that parents of individuals

with ASD paid greater attention to detail in daily life according to a self-report questionnaire. The authors did not find differences in performance-based indices in experimental tasks of local and global processing, perhaps reflecting the limitations of accuracy and reaction time measures to reveal differences within a group exhibiting sub-clinical traits. A recent twin study of individuals with ASD revealed striking similarities among monozygotic twins in performance measures during tasks tapping local perception that also related to ASD severity (Neufeld et al., 2020), suggesting that visual perceptual styles may reflect underlying genetics contributing to ASD symptomatology. Such differences in local and global visual perception have even been found to vary as a function of BAP-related traits among the *typical* population (Chouinard et al., 2016; Cribb et al., 2016; DiCriscio, Hu, & Troiani, 2019; Hayward, Fenerci, & Ristic, 2018). Taken together, there is exists a lack of consistent methods examining at multilevel facets of local/global processing in parents, and that the bulk of evidence suggests that local/global processing is likely a heritable trait but that the strength of the signal likely varies by method and precision of the instruments involved.

Current study

The present study builds on existing performance-based (i.e., accuracy, reaction time) studies of local and global processing by providing detailed characterization of eye movement, in addition to traditional performance-based measures, in individuals with ASD and their parents. Both the Kanizsa Illusory Contours (Kanizsa, 1976) or KICs (*Fig. 1A*), and the Navon hierarchical figure (Navon, 1977) tasks were administered as established tests of processing style. The KIC, in which, an illusory shape is induced by strategically placed “pacman” elements, has shown particular utility in the neuropsychological literature to study the

development of global form processing in typical development (e.g., Altschuler et al., 2014; Bulf, Valenza, & Simion, 2009; Cox et al., 2013; Harris, Schwarzkopf, Song, Bahrami, & Rees, 2011), as well as in psychiatric populations (e.g., Keane, Joseph, & Silverstein, 2014) and ASD (Baruth, Casanova, Sears, & Sokhadze, 2010; Stroganova, Nygren, et al., 2007; Stroganova, Orekhova, et al., 2007; Stroganova et al., 2012), and also has direct neural correlates across the different levels of visual perception (Harris et al., 2011; T. S. Lee & Nguyen, 2001; Maertens & Pollmann, 2005; Murray et al., 2002; Ringach & Shapley, 1996; Scholte, Jolij, Fahrenfort, & Lamme, 2008; Stanley & Rubin, 2003; Wokke, Vandenbroucke, Scholte, & Lamme, 2013). The Navon hierarchical figure (Navon, 1977) depicts a large letter comprised of smaller letters, and has been one of the most extensively applied stimuli used to examine local and global processing and interference effects across neurotypical and neurodevelopmental populations.

First, we aim to replicate psychophysical performance measures of reaction time and accuracy (Nayar et al., 2015; Nayar et al., 2017), with the inclusion of a suite of eye-tracking variables applied to both tasks as the primary focus of this study, in order to clarify the mechanisms underlying performance metrics. It was predicted that individuals with ASD would demonstrate a style less reliant on global processing compared to controls, and that similar, but less pronounced, trends would be observed in parents, perhaps only evident through eye-tracking measures. Relationships between local/global processing and ASD symptom severity and BAP personality and language features were examined. Additionally, given prior work demonstrating ties between local/global processing and social cognition (Burnette et al., 2005; Happé, 2000; Martin & McDonald, 2003; Pellicano, Maybery, Durkin, & Maley, 2006) and, inconsistently, executive functioning (Bolte, Holtmann, Poustka, Scheurich, & Schmidt, 2007; Booth, Charlton, Hughes, & Happe, 2003; Happe & Frith, 2006; Rinehart et al., 2000; Van Eylen et al., 2017; Van

Eylen et al., 2018), these skills were additionally examined as correlates. Finally, potential familial effects of local/global processing style were examined through parent-child correlations.

Methods

Participants.

A total of 161 participants were included in this study, including 32 individuals with ASD and 30 controls, and 57 parents of individuals with ASD and 43 control parents. Parents were included in the ASD parent group if they had at least one child with idiopathic ASD. There were 16 dyadic pairs (i.e., parent-child) in ASD families and 21 control dyads. ASD and control families were recruited through prior and ongoing studies. Recruitment efforts additionally included advertisements to ASD clinics, advocacy organizations and participant registries and word of mouth. Inclusionary criteria for all participants included speaking English as their first language, a minimum full-scale IQ (FSIQ) of 70, age of 8 years or older, no uncorrected visual impairment (s) (e.g., strabismus, uncorrected double vision), no history of neurological conditions (e.g., brain injury, non-febrile seizures), and no known genetic syndrome associated with ASD (e.g., fragile X syndrome, Phelan-Mcdermid syndrome) or major psychiatric disorder (e.g., bipolar, schizophrenia, and related psychotic disorders). Control participants were included if they had no personal or family history of ASD or related genetic disorders, as well as no family member within their nuclear family with a history of depression or language-related delays. See *Table 2.1* for demographic characteristics of the participant sample. All study procedures were approved by the University's Institutional Review Board and written informed consent/assent was obtained for all participants.

Clinical-behavioral Characterization.

Assessment of Cognitive Ability. The Wechsler Abbreviated Scale of Intelligence (WASI) (Wechsler, 1999) or the Wechsler Intelligence Scale for Children—Third Edition (WISC-III) (Wechsler, 1991) was used to assess 2-subject or full-scale (FSIQ), verbal (VIQ), and performance (PIQ) IQ. The ASD group had a significantly lower FSIQ, VIQ, and PIQ than the control group ($ps < .0001$), and the ASD parent group had significantly lower FSIQ and PIQ than their respective controls ($ps < .01$).

Assessment of ASD Symptoms. The Autism Diagnostic Observation Schedule-General or 2nd Edition (ADOS) (Lord et al., 2012) and/or the Autism Diagnostic Interview-Revised (ADI-R) (Lord, Rutter, & Le Couteur, 1994) were used to confirm a diagnosis of ASD. ADOS calibrated severity scores (total, social affect, and restricted and repetitive behavior) (Gotham, Pickles, & Lord, 2009) were additionally calculated. Moreover, the Social Responsiveness Scale (SRS) (Constantino & Gruber, 2005), and the Repetitive Behavior Scales-Revised (RBS-R) (Lam & Aman, 2007) questionnaires were obtained to further characterize symptoms of social communication and restricted and repetitive behaviors.

Assessment of the Broad Autism Phenotype. Personality features associated with the BAP (i.e., sociability, rigidity, and pragmatic language) were assessed among parents of individuals with ASD. Assessments included a semi-structured interview, the Modified Personality Assessment Scale-Revised (MPAS-R) (Piven et al., 1997), which includes a series of questions specifically designed to tap the subclinical features related to the BAP including aloof, rigid, perfectionistic, and untactful personality traits. Coding of personality features followed methods outlined in prior work, such that raters were assigned scores ranging from 0 to 2 (trait absent, possibly present, definitely present) on a 5-point (0.5 increment) Likert scale. The broad

autism phenotype questionnaire (BAP-Q) (Hurley, Losh, Parlier, Reznick, & Piven, 2007) was used to further characterize BAP traits in a continuous manner including total scores, aloof and rigid traits, and pragmatic language violations.

Assessment of Pragmatic Language. Pragmatic language was assessed using the Pragmatic Rating Scale (PRS) (Landa et al., 1992) (for parents) and Pragmatic Rating Scale-School Age (PRS-SA) (Landa, 2011) (for ASD and controls). Raters blind to family diagnosis rated language and speech behaviors that were operationally defined based on a semi-structured conversation about the participant's Life History and a conversation that is incorporated into the ADOS, respectively. The PRS has been repeatedly shown to distinguish relatives of individuals with ASD from controls (Landa et al., 1992; Piven et al., 1990; Piven et al., 1994).

Assessment of Social Cognition. The Reading the Mind in the Eyes Test (Baron-Cohen, Wheelwright, Hill, Raste, & Plumb, 2001) assesses recognition of complex psychological states by viewing photographs depicting only the eye region of the face, presented with four complex mental state words, in which the participants must select which of the four words best represents the eyes in each photograph. Prior work has demonstrated deficits in ASD (Baron-Cohen, Wheelwright, & Jolliffe, 1997), and subtle differences have been reported in parents (Baron-Cohen & Hammer, 1997; Losh & Piven, 2007). Though inconsistent, prior work has additionally indicated relationships between local/global processing and theory of mind abilities in ASD (Burnette et al., 2005; Happé, 2000; Martin & McDonald, 2003; Pellicano et al., 2006).

Assessment of Executive Functioning. Given theories and literature documenting relationships with executive functioning and local/global processing (Booth et al., 2003; Happe & Frith, 2006; Van Eylen et al., 2017; Van Eylen et al., 2018), suggesting that biases for local versus global information processing reflect executive impairments rather than differences in

perception, two executive functioning measures were examined as correlates. The Flanker (Eriksen & Eriksen, 1974) test from the NIH Toolbox (Weintraub et al., 2013) was administered to a subset of ASD parents, individuals with ASD, and controls, as an experimental measure of selective attention and inhibitory control. The BRIEF-parent (Gioia, Isquith, Guy, & Kenworthy, 2000) and BRIEF-Adult (Roth, Lance, Isquith, Fischer, & Giancola, 2013) self-report questionnaires, specifically the Global Executive Composite (GEC) T-score, was used as a measure of overall executive functioning impairments in daily life among groups. The BRIEF-parent questionnaire was used for individuals younger than 18 years and is based on parent-informant reporting of child behaviors, and the BRIEF-Adult was used for any individual 18 years and older as a self-report measure of executive functions.

Design and Stimuli.

General Procedures

Participants completed two interactive similarity match-to-sample (i.e., selection of a target image that matches a sample) tasks involving KIC and Navon figures (see, “Kanizsa Illusory Contour Task” and “Navon Hierarchical Figure Task” below) to assess indices of global processing including accuracy, reaction time, and a suite of eye movements variables as outlined below. Tasks were presented on a 17-inch TFT LCD monitor (1,280 x 1,024 resolution) placed 50-60 cm away from the participant. Before each test run, all participants completed a practice run on which 80% accuracy was required to ensure mastery of task instructions. A Tobii T60 eye tracker (Tobii Technology AB, Danderyd, Sweden) was used to track gaze coordinates at a rate of 60 Hz. Prior to task administration, a standard 5-point calibration grid was used, which renders a typical visual angle accuracy of 0.5°. Participants were recalibrated following any large

movements during calibration, and tracking was additionally monitored during the task using the live view and track window options in Tobii Studio. Finally, calibration checks were embedded in the task (e.g., center cross-hair) to ensure tracking accuracy throughout the duration of the task.

Kanizsa Illusory Contour (KIC) task.

KIC Paradigm. KICs are easily quantifiable, affording correct responses (i.e., they elicit specific shapes like a square), and thus provide a fruitful model for investigating both local *and* global perceptual levels, as well as investigating contextual integration. The KIC task consisted of three interactive similarity match-to-sample conditions (*Fig. 1B*, descriptions below) presented in a fixed order across all participants: first, the *KIC practice* condition, then the two test conditions (*KIC basic* followed by *KIC noise*), based on Nayar and colleagues' previous work (Nayar et al., 2015; Nayar et al., 2017). On each trial across the three conditions, one of five non-illusory solid shapes (square, diamond, rectangle, triangle, or trapezoid) appeared for 1 second at the center of the screen, followed by a blank (black) screen for 1 second, after which two simultaneously presented figures appeared—the two figures were non-illusory solid shapes for the practice condition, and KIC figures for the test conditions. All participants were instructed to respond as quickly as possible. Verbal feedback indicating accuracy was only provided during the *KIC practice* condition; in contrast, neutral phrases of encouragement were periodically provided (approximately every 10 trials depending on the participant's level of attention and engagement), such as saying, “keep it going!”, “You’re moving along nicely!”, or “almost there!”. Across conditions, every participant was informed when the condition was at the mid-way point. Finally, at the beginning of each condition, participants were briefly reminded of the match-to-sample instructions, without mentioning illusory forms during instructions.

KIC Figure Parameters. Support ratio (the relative length of the inducing versus induced contour) of all the KIC figures was set at 60% and based on prior work (Nayar et al., 2015; Nayar et al., 2017). The size of the KIC figures was fixed and consistent with prior work (Nayar et al., 2017)—at a distance of 60cm, the pacman elements' radius was 1.5° and the illusory edge was 5° . As such, the total edge length was 8° . All figures were white on a black background with average luminance for each solid shape set at 180 cd/m^2 and the background at 35 cd/m^2 .

KIC Practice Condition. All participants completed a *KIC practice* condition, in which 24 trials of real (non-illusory) shapes were presented in order to prevent potential training effects on the illusory KIC figures. The *KIC practice* condition was repeated until mastery was achieved (80% accuracy, i.e. 20 out of 24 trials), up to three times. During this condition, a smiley face was displayed only after correct responses, otherwise a blank screen was displayed for the same amount of time. Participants were also prompted with standard instructions when incorrect to pay close attention during only the *KIC practice* condition. All participants passed the practice.

KIC Basic and KIC Noise Test Conditions. Following mastery of the match-to-sample concept via the *KIC practice* condition, two test conditions were administered in a fixed order: 1) a *KIC basic* test condition assessing basic KIC recognition, and 2) a *KIC noise* test condition that assessed KIC recognition in the presence of background “noise” consisting of randomly placed pacman elements. This condition therefore generated local interference as a means to assess how task-irrelevant information (i.e., local pacman distractors) may interfere with global processing of the KIC figure (Nayar et al., 2017; Van der Hallen et al., 2015). During both test conditions, participants were required to select one of two KIC figures that matched a previously presented non-illusory sample shape—one KIC figure induced the appearance of the non-illusory sample shape (i.e., “target KIC” or the match), and one that did *not* match the sample (i.e., “distractor

KIC”). The target and distractor KIC figures appeared on either the left or the right side of the screen’s vertical midline. The KIC figures were further randomized across these two halves, separating the screen into four quadrants—upper and lower left, and upper and lower right quadrants. To further avoid position biases, the target KIC figure was limited to presentation on the same side (i.e., left versus right) for four consecutive trials only. While the *KIC basic* condition included only two KIC figures, the *KIC noise* condition presented two KIC figures embedded in a background of “noise”. Unlike the *KIC practice* condition, a smiley face was presented following each click response regardless of accuracy to prevent learning effects. If participants inquired about accuracy, experimenters provided a standard neutral response. Further, to ensure that participants responded and to encourage engagement, the examiner presented the next trial only after the participant appeared ready (i.e., facing the screen, hand placed on the mouse, not talking). Both test conditions consisted of 40 trials.

[INSERT FIGURE 1 HERE]

Navon Hierarchical Figure (Navon) task.

Navon Paradigm. Similar to the KIC task, the Navon task consisted of two interactive match-to-sample conditions (*Fig. 1C*) presented in a fixed order across all participants: 1) a *Navon practice* condition and 2) the *Navon test* condition. The format of the match-to-sample paradigm was consistent to the KIC paradigm such that an image appeared at the center of the screen for 1 second, followed by a blank (white) screen for 1 second, followed by two simultaneously presented images. For both conditions, the two figures appeared in fixed positions, appearing on either side of the screen’s midline. The target (or match) figure did not repeat on the same side for more than four consecutive trials. Feedback for practice versus test conditions was consistent with that of the KIC Paradigm.

Navon Hierarchical Figure Parameters. The Navon figure is made up of a large letter, composed of smaller different letters. Navon figures were separated into two categories: those that were morphologically *similar* (e.g., F and E) and those that were morphologically *dissimilar* (e.g., O and N) (Lamb & Robertson, 1989; Mottron et al., 1999) (*Table 2*). Consistent with prior work (Koldewyn et al., 2013; Mottron et al., 1999), at a distance of 60 cm, the large Navon figure was 5° tall and 3.5° wide with the size of the smaller letters fixed at .5° x .5°. Letters were black against a white background.

Navon Practice Condition. All participants completed a *Navon practice* condition consisting of 10 trials prior to the test condition. Modeling Koldewyn and colleagues' (2013) work, all images included objects or pictures. Importantly, while Koldewyn's practice trials involved training their participants' to match categories with a clear correct versus incorrect match (e.g., dog sample, different dog target and flower distractor), the images presented during this practice condition involved unique images within the *same* category with a more ambiguous match (e.g., lion sample, lioness target and dog distractor). Given that neither of the two options are identical, participants are able to use whatever clues they would like to make their selection (global shapes, colors, detail differences), but the global category (i.e., animals) is not relevant. This ensured that the practice trial did not facilitate in training categorization of the global percept, but ensured mastery of the concept of a match-to-sample paradigm instead. All participants passed the practice.

Navon Test Condition. One test condition of 40 trials was administered, which included hierarchical Navon letters as described above. Participants were required to select one of two Navon hierarchical figures that matched the previously presented non-Navon letter (i.e., a real letter)—one Navon figure was the “global match” (i.e., the large letter matched the sample real

letter, and was comprised of smaller incongruent letters), and the other was the “local match” (i.e., the small letters matched the sample real letter, which were aligned to form an incongruent non-match larger letter). The Navon hierarchical figures always appeared on each side of the midline. The position of the Navon figures did not change across trials. Morphologically *similar* and *dissimilar* letter pairs were pseudo-randomized such that letter pairs were randomly presented in the same order for each participant. Letter pairs repeated only once across the 40 trials, with the target letter switching to represent the letter not previously presented (e.g., K for KM, and then M for KM letter pair). Category of letter pairs (i.e., morphologically *similar* or *dissimilar*) appeared no more than three consecutive trials. Smiley face reinforcement and verbal encouragement was administered similarly to the KIC test conditions.

[INSERT TABLE 2 HERE]

Eye-tracking Procedures.

Areas of Interest (AOI)

KIC Paradigm. Based on prior work (Guttman & Kellman, 2004; Nayar et al., 2017; Ringach & Shapley, 1996), attention allocated to the individual pacman elements comprising the KIC figure implicated local perceptual processes, while attention to the centers of the KIC forms indicated global perceptual processes. As such, there were two primary operationally defined AOIs (*Fig. 2*): the centers of the illusory contours and their pacman elements. Pacman AOI was defined consistent with our prior work (Nayar et al., 2017), such that attention allocated to the physical pacman element plus the empty pie was considered local processing. KIC center AOI was defined as the space induced by the pacman elements within the illusory boundaries and not including the empty pie. Gaze directed towards different components of the KIC stimuli reflect unique perceptual processes. For instance, there is evidence of a “pop out” effect in appreciating

the global form (i.e., the illusory shape), which prior work has demonstrated correspond to visual attention directed towards the “center” of the KIC indicating greater global perception, in contrast to looking towards the individual pacman elements, which reflects a greater local processing strategy (Guttman & Kellman, 2004; Ringach & Shapley, 1996). For the *KIC noise* test condition only, any attention to the background pacman elements was additionally considered to be local processing. Although the *KIC noise* test condition includes demands similar to those of visual search paradigms, it is not a classic visual search paradigm as it requires the integration of local pacman elements to create the global illusory form for both the target (match) and non-target (non-match) KICs, which are embedded within an array of jittered pacman elements.

Navon Paradigm. AOIs defined in the *Navon test* condition were relatively straightforward and only included 2 AOIs: the “global” match and the “local” match. A rectangular figure within the bounds of the stimulus surrounded the Navon figure and defined the global or local match AOIs.

[INSERT FIGURE 2 HERE]

Quality control procedures

Fixations were defined based on the I-VT fixation filter available in Tobii Studio, which included the following fixation settings: 1) minimum fixation duration of 100 ms; 2) fixation data were averaged across both eyes, 3) velocity threshold was set to 20°/s, 4) maximum duration and angle between each new fixation were set at 100 ms and 0.5°, respectively, 5) moving average window of 3 samples was included to reduce noise, and 6) missing data with gaps of maximum 150 ms were linearly interpolated. Additionally, a valid trial was defined as having a minimum of 1 fixation count and a valid participant was defined as having a minimum

of 50% valid trials of total trials. In order to maximize sample size to examine group differences, a participant was included in final analyses if they presented with at least one valid *KIC basic* and/or *KIC noise* trial for the *KIC paradigm*. Because the *Navon paradigm* only had one test condition, all participants with a valid *Navon test* condition were included in analyses.

No significant group differences emerged between groups in the average number of valid trials across conditions (*KIC Paradigm across test conditions*: ASD group—M (SD) = 36.6 (4.9), control group—M (SD) = 36.9 (4.3), $t(108) = .32, p = .75$; ASD parent group—M (SD) = 38.7 (2.7), Parent control group—M (SD) = 39.0 (2.0), $t(193) = 1.15, p = .25$. *Navon Paradigm*: ASD group—M (SD) = 35.1 (5.8), control group—M (SD) = 37.7 (3.5), $t(35.30) = 1.78, p = .08$; ASD parent group—M (SD) = 36.4 (5.4); Parent control group—M (SD) = 36.4 (4.5), $t(84) = -.02, p = .99$).

In addition to the above quality control settings, consistent with prior work (e.g., Anderson et al., 2006), each AOI had a “buffer” area based on proportionally expanding the AOI’s original size by 20%. For the *KIC basic* and *KIC noise* conditions, pacman elements were expanded by increasing the radius by 20% while maintaining the original (x, y) coordinates. For the area defined by the KIC center, a dilation ratio of 1.2 (i.e., 20%) was utilized to expand this area by 20% around the center’s (x, y) coordinates. Pacman AOIs were prioritized for overlapping AOIs after expansion in order to ensure that each fixation was assigned to only one AOI. For the *Navon test* condition, each rectangular AOI encompassing the Navon figure was expanded by a dilation ratio of 1.2 based on its center (x, y) coordinates.

Variables of Interest

Refer to Table 3 for condensed descriptions and overview of variables included in analyses across tasks.

Performance-based variables

1. Percent Accuracy or Global Match. For the KIC paradigm, accuracy was defined as the total number of correct responses (i.e., selecting the target KIC that matched the non-illusory sample shape), divided by the total number of trials included in the condition multiplied by 100. For the Navon paradigm, the percentage of trials in which the participant selected the “global match” was explored. Due to online data collection methods, accuracy was only determined for the entire Navon task (and not by similar/dissimilar Navon stimulus category).

2. Reaction Time. Reaction time (in seconds) was determined based on the time between the presentation of the two KIC target/non-target options or the two Navon figures and the instant the participant clicked the computer mouse to make their selection. Utilizing a touch screen (Nayar et al., 2015; Nayar et al., 2017) was not possible due to the interference hand movements in front of the screen would have had with simultaneous eye tracking.

Strategy-based gaze variables

1. Fixation Rate. The total number of fixations that occurred throughout the duration of the trial, divided by the total amount of time (in seconds) for which eye tracking data was collected during the trial (i.e., “track time”). Fixation rate generally indicates processing speed, attentional capacity, and cognitive load (Zagermann, Pfeil, & Reiterer, 2018), with elevated values reflecting serial/local processing strategies.

2. Percent Vacillation. The total number of times per trial that gaze moved from target to non-target KIC or from one Navon option to the other, was summed across all trials and divided

by the total number of fixations and multiplied by 100. In other words, percent vacillation is the proportion of fixations that were between the two KIC or Navon stimuli options (rather than within an individual KIC or Navon). The percentage of vacillations between two stimuli can point to levels of uncertainty and readiness to appreciate the global form, with greater vacillations indicating a higher level of uncertainty and reduced appreciation for the global percept, which are characteristic of younger children (Nayar et al., 2015).

Center-Pacman Difference Scores - for *KIC Test Conditions only*.

For the following variables, pacman elements for the *KIC basic* test condition included the pacman inducers to the KIC only, while for the *KIC noise* test condition, pacman elements also included the background individual pacman elements. The below were computed for each test condition separately. Additionally, higher values indicate greater center gaze allocation.

1. First Fixation. The average difference between the percentage of first fixations for each trial that was towards the KIC center minus pacman element (higher value indicates a greater percentage of center first fixations). The location of the first fixation is thought to index the utilization of peripheral visual information, which is associated with global or rapid and automatic visual information processing and generally reflects visual information preference (Kimchi, 1992).

2. Fixation. The average difference between the percentage of fixations for each trial made towards the KIC centers minus the percentage of fixations made towards the pacman elements (higher value indicates a greater percentage of center fixations). Increased levels of fixations or time spent fixating on the target (i.e., the stimulus that matches the sample previously presented) has been shown to index greater global processing that has also been

reflected in unique patterns of neural processing during local/global tasks with target and non-target stimuli (Baruth et al., 2010; Fink et al., 1997; Heinze, Hinrichs, Scholz, Burchert, & Mangun, 1998).

3. *Duration*. The average difference between the percentage of time spent for each trial looking towards the KIC centers minus the percentage of time spent looking at the pacman elements (higher value indicates a greater percentage of time spent looking at the center).

Target-NonTarget Difference Scores.

Higher values in the below variables reflect greater target gaze.

1. *First Fixation*. The average difference between the percentage of first fixations for each trial that was towards the target match minus the non-target option (higher value indicates a greater percentage of target first fixations).

2. *Fixation*. The average difference between the percentage of fixations for each trial made towards the target match minus the percentage of fixations made towards the non-target option (higher value indicates a greater percentage of target fixations).

3. *Duration*. The average difference between the percentage of time spent for each trial looking towards the target minus the percentage of time spent looking at the non-target stimulus (higher value indicates a greater percentage of time spent looking at the target).

Composite Gaze Variables.

Higher values in the below variables indicate greater center or target gaze (i.e., greater propensity for global processing).

1. *Center-Pacman Difference Score*: For the KIC paradigm only, the center-pacman difference composite score is the average of the sum of center percent first fixations, fixations, and duration minus the sum of the pacman first fixations, fixations, and duration.

2. *Target-NonTarget Difference Score*: The average of the sum of target percent first fixations, fixations, and duration minus the sum of the non-target first fixations, fixations, and duration.

Overall Composite Global-Local Difference Score

Overall Global-Local Difference Score (z-score difference): For both paradigms, this composite score was generated to reflect an overall difference between global and local processing styles. The score included all variables outlined above as follows. All variables were first converted to z-scores to ensure that all variables were of comparable scale. For the *KIC Paradigm*, this composite variable is the average sum of the z-scored accuracy, reaction time (reverse coded to ensure comparable theoretical directionality of z-scores across variables), target and center percent first fixations, fixations, and duration minus the sum of the z-scored non-target and pacman element first fixations, fixations, and duration as well as percent vacillations and fixation rate. For the *Navon Paradigm*, this composite variable is the average sum of the z-scored accuracy, reaction time (reverse coded), target percent first fixations, fixations, and duration minus the sum of the z-scored non-target first fixations, fixations, and duration as well as percent vacillations and fixation rate.

Statistical Analysis

Assumptions testing. Data were examined to assess model assumptions. Outlier examination revealed one ASD parent (father) as a relatively consistent extreme outlier across most KIC variables and this participant was therefore removed from subsequent analyses. There were no consistent extreme outliers in the Navon task. Assumptions for linear mixed effect models (see below) were appropriately met across most variables. Given failed homogeneity of variance on accuracy and reaction time measures overall, performance variables were re-analyzed using non-parametric tests (Mann-Whitney U).

Covariates. Covariates were examined using theoretical and data-driven approaches. For probands, given that IQ was significantly different between diagnostic groups (see *Table 1*), and there are known relationships between cognitive abilities and performance across neuropsychological tasks in ASD (Rommelse et al., 2015), IQ was added as a covariate for accuracy and reaction time analyses across tasks. Although IQ correlated with eye-tracking variables in proband groups, final models did not include IQ as a covariate for gaze indices, as 1) results did not differ when analyses were conducted with or without controlling IQ, and 2) given that differences in IQ are associated with the inherent heterogeneity in ASD, it is important to allow for variability within the group when assessing discriminability of potential diagnostic classifiers (Dennis et al., 2009). Age was not included as a covariate for proband analyses given lack of significant differences between groups or associations with outcome variables.

For parents only, age significantly correlated with outcome variables across both KIC and Navon paradigms. As such, analyses were conducted with and without age included as a covariate in all parent analyses. Given that significance did not differ between models (results not shown), and that all parents were over age 18 years when global perception is preferred (Motttron et al., 2003; Nayar et al., 2015) all reported parent analyses include models *without* age

as a control variable. Although IQ was significantly different between parent groups, it did not correlate with outcome variables and was therefore not controlled for in parent analyses.

1. Assessment of Group Differences in Local/Global Perception Tasks

KIC and Navon. For both paradigms, parent groups and proband groups were analyzed separately. Group differences in performance and gaze variables were assessed using a series of mixed effects linear regression models using the lmer package (Bates, Mächler, Bolker, & Walker, 2014) *R Studio* (Team, 2019) to assess the effect of condition, group, and the interaction. Models for the KIC paradigm included diagnostic group (control group as the reference group) and condition (*Basic KIC* and *KIC Noise*, with the *Basic KIC* condition as the reference group) as fixed effects, as well as their interaction term. For Navon, primary analyses included examining group differences only across all trials using ANCOVA or ANOVA, with the above-mentioned covariates. Secondary more exploratory analyses further examined effects of morphologically *similar* or *dissimilar* letters as follows: A series of linear mixed effect models with diagnostic group (control as the reference group) and letter similarity (*dissimilar* and *similar*, with dissimilar as the reference group) were included as the fixed effects, as well as their interaction. Across all models, the individual participant was included as the random effect. Interaction terms were only interpreted if there was a significant improvement in model fit, with the models being run with and without the interaction term. If there was no improvement in model fit with the addition of the interaction term, only main effects were interpreted. Finally, if the interaction term was significant, only this 2-way interaction finding was reported and interpreted, without main effects of group or condition separately.

Across Tasks. To explore cross-task differences, for each participant, overlapping variables were averaged first between the KIC conditions (*KIC basic* and *KIC noise* conditions)

and then between paradigms (KIC and Navon), to generate an average value for composite gaze and overall composite variables only (i.e., *Target-NonTarget Difference Composite score* and *Global-Local Difference Composite z-score*). A series of ANOVAs were examined to explore group differences in local or global processing across tasks.

Group differences including performance (i.e., accuracy and reaction time) and gaze composite variables (5 variables total for the KIC paradigm, and 4 for the Navon paradigm) were examined as a first level of analyses, followed by more exploratory analyses involving individual gaze variables comprising composite variables (i.e., fixation rate, vacillation, and difference scores). Group differences were adjusted using the Benjamini-Hochberg method (Benjamini & Hochberg, 1995) using a FDR of .10 to reduce false negatives and thereby potentially missing important effects (APA, 1994). Benjamini-Hochberg adjusted *p* values are reported in Tables 4 - 9; all other *p* values meeting an alpha criterion of $< .05$ are reported below.

Association between Performance and Gaze Variables. Finally, in order to examine how performance and gaze variables related to each other, within-task associations for KIC and Navon Paradigms between gaze composite variables (i.e., *Vacillation*, *Fixation Rate*, and composite *Center-Pacman Difference* and *Target-NonTarget Difference scores*) and performance variables (accuracy and reaction time) were examined across groups separately for parents and probands. For probands, to remain consistent with overall group models, IQ was included as a covariate, and a series of partial Pearson correlations were conducted. For the KIC paradigm, both conditions were averaged for the purpose of being parsimonious. Finally, accuracy and reaction time associations were explored to assess whether speed of response, impacted accuracy (particularly given that global processing is rapid). Due to most performance

variables failing Levene's test of homogeneity of variance, Pearson correlations were followed-up with bivariate Spearman's rank-order partial correlations.

2. Analyses of Parent-child Associations in Local/Global Perception Tasks

To explore whether profiles of local and global processing may show a pattern of familiarity, linear regression models were employed separately for each family diagnosis on composite variables across tasks only (i.e., using the across-task averaged composite variable—*Overall Global-Local Difference Score*) between parent and child pairs. All models included within-family relation nested within the model as a random effect to control for spousal relationships with the same child. These analyses were exploratory given small sample sizes (ASD dyads $n = 11$; Control dyads $n = 20$). Additionally, in order to examine whether local/global processing may confer familiarity or simply reflect ASD-risk more broadly, parent-child associations were re-run with unrelated, mismatched parent-child dyads. A randomization/Permutation (Katz, Lautenschlager, Blackburn, & Harris, 1990) test using a series of *Pearson's* correlations between unrelated mother-child dyads was applied to evaluate the specificity of familial relationships. Each parent value was randomly paired with a child's value for all possible permutations within each diagnostic group, which resulted in a sampling distribution of the correlation coefficients. Based on this test, the expected correlation coefficient is zero. The permutations test resulted in a distribution of all possible permutations of unrelated pairs within diagnostic groups. Subsequently, the strength of the true parent-child dyad correlation coefficient was compared against the distribution for the unrelated pairs (i.e., true ASD parent-child dyads versus all unrelated ASD pairings, and true control parent-child dyads versus all unrelated control pairings). Final metrics included a probability statistic that indicated

the likelihood of any random pairing producing a stronger correlation coefficient relative to the true parent-child dyad correlation coefficient.

3. Clinical-Behavioral and Cognitive Correlates in Local/Global Perception Tasks

Using the across-task averaged variables, Pearson correlations were examined between performance, gaze, and composite variables and clinical-behavioral and cognitive correlates. For probands, partial Pearson correlations were conducted with IQ added as a covariate for correlations between performance and clinical-behavioral characteristics. Associations with clinical-behavioral traits involving ASD or BAP measures were only examined in clinical groups (i.e., ASD and ASD parent groups).

Clinical-Behavioral Correlates

For individuals with ASD, the ADOS total and subscale (social affect and restricted and repetitive behaviors) calibrated severity scores, ADI current domain and total scores, SRS total T-score, and RBS-R Overall Sum were all utilized in correlations with cross-task variables. Rates of pragmatic language violations (PRS-SA Total and domain scores—e.g., *theory of mind domain*, which includes items that tap theory of mind skills during conversation such as being unable to clarify, failing to provide background, being redundant, being detail oriented) were examined in both ASD and control groups separately.

For parents of individuals with ASD, BAPQ total and sub-scores (i.e., pragmatic, aloof, and rigid) were examined in relation to local and global processing. Similar to individuals with ASD, the PRS total and factor scores (i.e., *dominating conversation* factor score—which is a sum of items including being overly detailed, being tangential, being frank, exhibiting odd humor, and demonstrating optic preoccupation—and *socially withdrawn* factor score, which includes

items such as being vague, being unable to clarify responses, failing to reciprocate, and failing to elaborate on their responses) were examined for both the ASD parent and parent control groups.

Cognitive Correlates

Social cognition (as measured by The Reading the Mind in the Eyes Test percent correct) and measures of executive functioning (Flanker age-corrected Standard Score and BRIEF GEC T-score) were also examined in both ASD and control families separately.

Results

Group differences are presented as follows: 1) overall composite gaze score (i.e., the *Overall Global-Local Composite Score*), 2) performance-based variables (accuracy and reaction time), and 3) strategy-based gaze variables. Within the strategy-based gaze variables results section, findings from individual gaze variables (i.e., center-pacman first fixation, fixation, and fixation duration difference scores and/or target-nontarget first fixation, fixation, and fixation duration difference scores) are described. All statistics are reported in relevant tables. Finally, given known differences between KIC test conditions (basic versus noise) (Nayar et al., 2017) and not central to primary aims of the study, condition effects are reported in Tables 4 and 5 only, and not in the text below. Additionally, although condition and group X condition effects during the Navon paradigm were exploratory in nature, results from condition effects are also only reported in tables 8 and 9.

1. Group Differences

KIC Paradigm (Tables 4 - 5; Fig. 3)

Overall Composite Global-Local Difference Score

ASD vs. Controls: A main effect of group emerged, showing that the ASD group demonstrated reduced global perception overall across conditions than did the control group.

ASD parent vs. control parents: No main effect of group was found between parent groups.

Performance-based variables

Accuracy.

ASD vs. Controls: There were no main effects of group or a group X condition interaction. Non-parametric analyses demonstrated significant group differences in the *KIC basic* condition, showing lower medians in the ASD ($M_{\text{rank}} = 21.25$) versus control ($M_{\text{rank}} = 31$) group ($U = 210, p < .05$). Similar trends were observed in the *KIC noise* condition showing relatively lower means among individuals with ASD ($M_{\text{rank}} = 25.38$) compared to the control group ($M_{\text{rank}} = 33.35$) ($U = 304.5, p = .05$).

ASD parent vs. control parents: Likewise, no group or interaction effects emerged for accuracy in parents. Non-parametric results revealed lower medians on the *KIC basic* condition in the ASD parent group ($M_{\text{rank}} = 43.72$) compared to the parent control group ($M_{\text{rank}} = 54.40$) ($U = 886, p < .05$). In contrast, no group differences emerged in the *KIC noise* condition (parent control group $M_{\text{rank}} = 51.31$, ASD parent group $M_{\text{rank}} = 43.72$, $U = 1058, p = .43$).

Reaction Time.

ASD vs. Controls: No group main effect or group X condition interaction effect emerged for reaction time in proband groups. Non-parametric Mann-Whitney U tests revealed significant group differences across both conditions, showing slower means in the ASD (*KIC basic* $M_{\text{rank}} =$

22.11; *KIC noise* $M_{\text{rank}} = 22.47$) compared to the control (*KIC basic* $M_{\text{rank}} = 31.63$; *KIC noise* $M_{\text{rank}} = 37.04$) group (*KIC basic* $U = 213, p < .05$; *KIC noise* $U = 209, p < .01$).

ASD parent vs. control parents: A main effect of group showed that the ASD parent group was slower to respond than the control parent group. There was no group X condition interaction. Non-parametric test similarly revealed significant group differences in mean ranks between groups, particularly during the *KIC basic* condition (parent control group $M_{\text{rank}} = 38.40$, ASD parent group $M_{\text{rank}} = 56.70, U = 1651, p < .01$). Similar trends were observed in the *KIC noise* condition showing that the ASD parent group ($M_{\text{rank}} = 53.56$) responded relatively slower than the parent control group ($M_{\text{rank}} = 43.02$) ($U = 904, p = .07$).

Strategy-based gaze variables

Exploration.

ASD vs. Controls: There was a main effect of group, showing that the ASD group had elevated exploration rates (i.e., made a significantly greater number of fixations per second) than the control group.

ASD parent vs. control parents: No main effects of group or group X condition interaction emerged for parents.

Percent Vacillation.

ASD vs. Controls: There was a main effect of group and group X condition interaction, showing that the ASD group vacillated more than the control group, particularly during the *KIC basic* condition.

ASD parent vs. control parents: No group or interaction effects emerged between parent groups.

Center-Pacman Difference Score.

ASD vs. Controls: While no overall main effects of group or interaction emerged for the overall composite score, a group X condition effect was found for the percentage of *fixation* and *fixation duration* center-pacman difference variables—both gaze variables showed a greater percentage of fixations and time spent on KIC centers versus pacman elements during the KIC basic versus KIC noise condition across groups, which was attenuated for the ASD group compared to the control group, and with more pronounced group differences emerging in the KIC noise condition.

ASD parent vs. control parents: A group X condition effect emerged for parents, showing a higher percentage of KIC center versus pacman perception during the KIC basic versus KIC noise condition across groups. Importantly, between-group patterns differed based on condition showing that the ASD parent group attended more towards the center versus pacman during the KIC basic condition compared to controls, which was flipped for the KIC noise condition (where both groups attended more towards pacman elements, with the ASD parent group showing even greater pacman versus center attention than the parent control group). This group X condition interaction was driven by the percentage of *fixation* and *fixation duration* variables.

Target-NonTarget Difference Score.

ASD vs. Controls: Across conditions, the ASD group attended less towards the target versus the non-target KIC compared to the control group, a pattern that was evident across all

variables comprising this composite variable. Additionally, a group X condition interaction revealed a greater percentage of target fixations and increased time spent looking at the target versus the non-target KIC during the KIC basic versus the KIC noise condition across groups, and that the ASD group overall demonstrated reduced target versus non-target fixations and fixation duration compared to the control group.

ASD parent vs. control parents: Parent findings were similar to those in probands, showing a main effect of group, such that the ASD parent group showed less attention towards target versus non-target KICs (particularly showing reduced *percent* fixation and *percent fixation duration*) compared to the parent control group across conditions.

[INSERT TABLES 4 AND 5 HERE]

[INSERT FIGURE 3 HERE]

Navon Paradigm (Tables 6-9; Fig 4)

Overall Global-Local Difference Score.

ASD vs. Controls: The ASD group showed a looking profile that was less global than local compared to the control group overall. No groups X condition interaction emerged.

ASD parent vs. control parents: No main effects of group or group X condition interactions were found.

Performance-based variables

Accuracy.

ASD vs. Controls: Overall, no group differences in accuracy emerged between groups. Non-parametric Mann-Whitney U also revealed no significant group differences in means (control group $M_{\text{rank}} = 24.20$, ASD group $M_{\text{rank}} = 19.05$, $U = 166$, $p = .13$).

ASD parent vs. control parents: Similarly, ASD parent and parent control groups performed similarly on the Navon test, with non-parametric findings also remaining non-significant (parent control group $M_{\text{rank}} = 45.29$, ASD parent group $M_{\text{rank}} = 42.01$, $U = 846.5$, $p = .48$).

Reaction Time.

ASD vs. Controls: The ASD and control groups demonstrated comparable reaction times overall as well as when examining letter similarity condition effects. Mann-Whitney U results were also not significant between groups (control group $M_{\text{rank}} = 20.90$, ASD group $M_{\text{rank}} = 22.05$, $U = 418$, $p = .77$).

ASD parent vs. control parents: Similarly, parent groups did not differ overall or when examining differences in performance during *similar* versus *dissimilar* letters. Mann-Whitney U non-parametric tests also revealed no group differences in reaction time overall (parent control group $M_{\text{rank}} = 43.83$, ASD parent group $M_{\text{rank}} = 43.22$, $U = 903.5$, $p = .91$).

Strategy-based gaze variables

Exploration.

ASD vs. Controls: No group differences emerged in the number of fixations per second of track time overall, or by letter similarity, with no group X condition interactions emerging.

ASD parent vs. control parents: Similarly, no overall group or interaction effects emerged.

Percent of Vacillations.

ASD vs. Controls: Overall and across conditions, there were no group differences in the percentage of vacillation between the two Navon options. No interaction effects emerged either.

ASD parent vs. control parents: Parent groups did not reveal any group or group X condition interaction effects.

Target-NonTarget Difference Score.

ASD vs. Controls: Overall, individuals with ASD demonstrated reduced attention towards the target versus non-target Navon option compared to controls, which was particularly driven by percentage of *fixations* and *fixation duration*. Additionally, a group X condition interaction ($Estimate = 11.30, \beta = .16, SE = 5.28, t(40) = 2.14, p < .05$) showed that the ASD group's target versus non-target looking pattern remained consistent between *similar* versus *dissimilar* letters, while the control group showed greater attention towards the target versus non-target Navon option during *dissimilar* letter trials versus trials with *similar* lettering. This interaction was primarily driven by the percentage of *fixations* towards the target versus non-target option ($Estimate = 10.54, \beta = .15, SE = 4.88, t(40) = 2.16, p < .05$).

ASD parent vs. control parents: Overall, while no differences emerged for target versus non-target looking, there were group differences in the percentage of first fixations that were made towards the target versus non-target option across conditions. In particular, the ASD parent group showed fewer first fixations towards the target stimulus than the parent control group.

[INSERT TABLES 6 AND 7 HERE]

[INSERT TABLES 8 AND 9 HERE]

[INSERT FIGURE 4 HERE]

Across Local/Global Perception Tasks (Tables 10 - 11; Figs. 5)

Target-NonTarget Difference Score.

ASD vs. Controls: The ASD group demonstrated significantly reduced attention towards the target versus non-target option compared to the control group.

ASD parent vs. control parents: Parents showed no overall group differences in attention towards the target versus non-target option.

Overall Global-Local Difference z-score.

ASD vs. Controls: Group differences emerged showing that control groups demonstrated greater global versus local processing compared to the ASD group across tasks.

ASD parent vs. control parents: Parent groups demonstrated comparable global versus local processing styles, though patterns trended in the same direction as proband group differences.

[INSERT TABLES 10 AND 11 HERE]

[INSERT FIGURE 5 HERE]

Association between Performance and Gaze Variables (Fig. 6)

ASD and Controls. With groups combined, a significant correlation emerged between the *Target-NonTarget Difference Score* and accuracy for KIC and Navon tasks ($r = .44, p < .01$ and $r = .67, p < .0001$) and reaction time for KIC only ($r = -.64, p < .0001$), demonstrating that the more accurate and quicker participants were in responding, the more they attended towards the target (versus non-target) KIC or Navon option. Similarly, the faster and more accurate participants performed, the less often they vacillated between the two KIC options ($r = .59, p < .0001$ and $r = -.47, p = .001$, respectively). Only accuracy significantly correlated with percent vacillations ($r = -.33, p < .05$) for the Navon Paradigm, showing fewer vacillations and greater

target (versus non-target) looking relating to higher accuracy. For the KIC paradigm, reaction time was significantly negatively associated with *Center-Pacman Difference Score* ($r = -.33, p < .05$) representing faster response times when there was greater attention directed to the KIC centers (versus Pacman). No associations with accuracy or reaction time with exploration were found in either task. Finally, higher accuracy and quicker response times were significantly correlated with one another for both KIC and Navon tasks ($r = -.54, p < .0001$ and $r = -.38, p < .05$, respectively). All significant and non-significant associations remained consistent upon re-analysis using non-parametric Spearman tests.

ASD parent and control parents. For parents (both ASD parent and parent control groups combined), a significant correlation between accuracy, reaction time and *Target-NonTarget Difference Score* during KIC revealed that the more parents attended to the target, the higher their accuracy ($r = .34, p = .001$) and the quicker their response times ($r = -.56, p < .0001$). Only accuracy was significantly correlated with *Target-NonTarget Difference Score* in the Navon paradigm ($r = -.22, p < .05$) (though, Spearman correlations revealed significant positive and negative relationships for accuracy and reaction time, respectively; $r_s > |.35|, p_s < .001$). A positive correlation also emerged between reaction time and vacillations ($r = .25, p < .05$) for the KIC paradigm (and for the Navon paradigm with Spearman tests; $r = .23, p < .05$), indicating slower response times when participants vacillated more between the two target/no-target options. Similarly, no associations with reaction time and exploration were found in either task, though, a positive correlation in the Navon task demonstrated elevated exploration rates relating to reduced accuracy ($r = -.22, p < .05$). Finally, a negative correlation emerged between reaction time and accuracy for the KIC ($r = -.24, p < .05$), but not for the Navon paradigm. Similar to the

proband groups, Spearman correlation results remained largely consistent with Pearson results reported here.

[INSERT FIGURE 6 HERE]

3. Parent-Child Associations Across Tasks (*Fig. 7*)

ASD families.

Significant parent-child associations emerged such that increased global looking patterns in parents predicted 34% of the variance of increased global looking patterns in their children ($estimate = .62, SE = .27, t = 2.26, p < .05$). In terms of the permutation test, *Pearson's* correlations for global versus local looking was significant for ASD families ($r = .60, p < .05$), with a 97.3% likelihood that the correlation coefficient derived from the true parent-child dyad correlation (r_{true}) is greater than the correlation coefficients derived from all permutations of the unrelated parent-child pairs (r_{random}) (i.e., probability $r_{true} > r_{random} = 97.3\%$).

Control families

Parent-child pairs. No significant parent-child associations emerged in the *Overall Global-Local Difference Score* composite for control families, with only 1.4% of variance in child global looking being explained by that of their parents ($estimate = .15, SE = .30, t = .49, p = .63$). The permutation test revealed non-significant *Pearson's* correlations for global versus local looking for control families ($r = .18, p = .45$), with a 76.2% likelihood that the correlation coefficient derived from the true parent-child dyad correlation is greater than the correlation coefficients derived from all permutations of the unrelated parent-child pairs (i.e., probability $r_{true} > r_{random} = 76.2\%$).

[INSERT FIGURE 7 HERE]

4. Relationship with Clinical-Behavioral and Cognitive Correlates (Fig. 8)

Probands.

Clinical-Behavioral Correlates. Negative associations between reaction time and RBS-R total score emerged, indicating slower response speeds correlating with greater rates of RRBs as measured by the RBS-R ($r = .79, p < .05$). Similarly, increased fixation rate and vacillation significantly positively correlated with greater rates of RRBs based on the ADI current scores ($r = .71, p < .05$ and $r = .85, p < .01$, respectively) (Figs. 9A and B). No associations between ADOS severity scores or SRS emerged among individuals with ASD. In the ASD group, significant positive associations between the *global-local difference composite z-score* and greater pragmatic language violations, particularly in speech and language behaviors (e.g., being formal, scripting) and theory of mind-related behaviors emerged (e.g., failing to clarify or to provide background information) ($r = .70, p < .01$ and $r = .61, p < .05$, respectively) (Fig. 9C). Significant correlations emerged for the control group with increased pragmatic language abilities and higher accuracy ($r = -.55, p < .05$).

Cognitive Correlates. No associations were found between local/global variables and social cognition or measures of executive function (age-corrected standard score on the Flanker task; GEC T-score from the BRIEF questionnaire) in either group.

Parents.

Clinical-Behavioral Correlates. No significant correlations emerged with measures of the broad autism phenotype, but a positive correlation of medium effect emerged between the *global/local composite z-score* with the PRS total score in the ASD parent group. Additionally, in parents of individuals with ASD, a negative correlation between number of vacillations

emerged with the PRS total score, indicating that greater pragmatic language violations occurred with greater global versus local processing ($r = .34, p < .05$ and $r = -.32, p < .05$). Specifically, among parents of individuals with ASD, the socially withdrawn factor score (which includes items such as being unable to clarify, failing to elaborate, failing to reciprocate) was related to greater global versus local processing ($r = .34, p < .05$) (Fig. 9D), greater rates of vacillations ($r = -.36, p < .05$), and greater target versus non-target attention ($r = .34, p < .05$). No significant associations with pragmatic language and local/global variables emerged in the parent control group.

Cognitive Correlates. No significant correlations emerged with The Eyes Test or either executive functioning measure, for ASD parent or control groups.

[INSERT FIGURE 8 HERE]

Discussion

This study applied a suite of eye-tracking metrics to two well-established tasks of local and global perception, in an attempt to study visual processing more deeply than standard performance-based measures of time and accuracy allow, to reveal potentially more nuanced patterns of visual processing styles among individuals with ASD and their parents. Robust differences were observed between individuals with ASD and controls, and more subtle, and nuanced differences emerged in parents, along with within-family similarities of processing styles that could point towards heritability of visual processing styles in ASD families. Finally, links between visual processing styles and ASD-related symptomatology were observed, as evidenced by relationships with elevated rates of restricted and repetitive behaviors in individuals with ASD and better pragmatic (i.e., social) language in both ASD proband and

parent groups. Associations suggest that local/global visual processing profiles may importantly relate to broader clinical and subclinical traits related to ASD. Together, these findings may point towards eye-tracking as a particularly revealing method for capturing often subtle, and potentially heritable and biologically meaningful styles of visual processing related to ASD that can help to further our understanding of potential underlying mechanisms of ASD and the broad autism phenotype (BAP). Findings are discussed in further detail, below.

Accuracy and Reaction Time

The ASD group tended to be less accurate and slower to respond than controls, across conditions in the KIC task in particular. Findings are consistent with prior work in school-age children of individuals with ASD, who demonstrated significant (but minimal) accuracy and reaction time differences during the more complex KIC noise condition, which included local interference (Nayar et al., 2017). Similarly, parents of individuals with ASD responded more slowly relative to control parents in the KIC task (although, all groups tended to respond within 1 second). Across all groups, quicker reaction times related to increased accuracy and greater global perception based on gaze variables; conversely, slower response speeds related to reduced global perception (i.e., greater vacillations between options, and fewer target/global looks in the KIC task). Such findings are consistent with original landmark findings from studies involving Navon Hierarchical Figures suggesting that global processing strategies are quicker and more efficient than effortful local perceptual processes (Navon, 1977) based on performance measures alone.

Gaze in ASD reflective of heightened local and weaker global processing

There is some debate on whether individuals with ASD exhibit heightened local processing in the absence or presence of impaired global processing (Simmons & Todorova, 2018; Van der Hallen et al., 2015). Findings from the present study showed increased vacillation and visual exploration (fixation rate) in ASD relative to controls in the KIC task. Greater vacillation between stimuli generally represents a less mature visual processing style, and is indicative of heightened local perception (alongside reduced global perception). These findings also point to greater levels of uncertainty, which has been shown to be particularly characteristic of younger children engaging in the processes of comparison or elimination before adult-like global processing strategies develop (Nayar et al., 2015). It may be that individuals with ASD are less readily able to appreciate the global form, and therefore more likely to rely on a developmentally less mature comparative strategy (between the target and non-target). This processing style is also consistent with the presence of slower or delayed global processing, whereby individuals with ASD have been shown to initially attend to information at the local level, shifting towards global processing strategies with prolonged looking, particularly during tasks with greater local interference such as the KIC noise condition (also see Nayar et al., 2017; Van der Hallen et al., 2015). This reduction in global processing among individuals with ASD was coupled with increased levels of detailed processing, as reflected in their exploration rates. Such findings mirror prior reports detailing elevated rates of exploration and perseverative fixations or refixations (i.e., repeat fixations) among individuals with ASD across contexts. These include non-social tasks (as in the present study), social information and non-social images (Sasson, Turner-Brown, Holtzclaw, Lam, & Bodfish, 2008; Swanson, Serlin, & Siller, 2013), social and non-social scenes (Au-Yeung, Benson, Castelhana, & Rayner, 2011), and a language processing task (Nayar et al., 2018), together highlighting the utility of studying

perseverative or detailed fixation patterns as a stable, cross-context biomarker for ASD. That increased vacillations and exploration related to increased rates of restricted and repetitive behaviors (RRBs) in ASD further suggests their specificity to traits inherent to ASD and may point to relevant biological connections between visual attention and RRBs.

In contrast to the KIC task, no group differences in vacillations and exploration were observed in the Navon paradigm. Given that this task was specifically designed to assess preference for local or global options, with a global cue (i.e., the real sample letter), fewer group differences for Navon suggest that individuals with ASD are able to engage in global processing adequately when cued, consistent with prior work showing that when individuals with ASD are primed or instructed to utilize global perception, they are able to efficiently do so (Koldewyn et al., 2013; Neufeld et al., 2020; Plaisted et al., 1999). Our findings in Navon additionally imply that the ASD group may be utilizing a different strategy to engage in global processing. In particular, individuals with ASD exhibited greater target versus non-target fixations, suggesting that perceptual strategies may differ between groups despite similarities in performance-based measures. A group by condition interaction for target fixations further highlights interference on efficient global processing from stimuli parameters consisting of similar versus dissimilar letters. Prior work has indicated that even typically developing individuals and high functioning individuals with ASD demonstrate slower response times when letters are morphologically similar (e.g., B and P) versus dissimilar (e.g., H and X) (Lamb & Robertson, 1989; Mottron et al., 1999). In the present study, those with ASD exhibited greater interference than controls during the morphologically similar versus dissimilar letter trials, evidenced by fewer target versus non-target fixations when letters were similar. Differences between KIC and Navon tasks

presented in this study, as well as key effects emerging from letter similarity can help to inform future local and global perception studies using Navon hierarchical figures.

Findings from more detailed eye-tracking variables in the KIC task have strong implications for our understanding of clinical and sub-clinical traits related to the weak central theory account in ASD. In KICs, the global form can be appreciated by integrating the local pacman elements to generate the coherent whole shape — this process is particularly automatized and occurs within the first 300 ms of perception according to studies examining neural responses to KICs (Foxy et al., 2005). Prior work has also suggested that attending to the center of the KIC indicates greater global perception, whereas looking to individual pacman elements reflects more detailed processing styles (Guttman & Kellman, 2004; Ringach & Shapley, 1996). The current study yielded robust group by condition effects in pacman versus center looking among proband groups. During the KIC basic condition, both groups exhibited comparable patterns of greater fixations to center versus pacman elements, but during the condition with local interference, the ASD group showed a pattern of looking that reflected heightened local processing relative to controls. The KIC basic and KIC noise conditions differ in complexity in that participants are required not only to integrate (i.e., fill in the illusion) but also to segregate (i.e., extract the relevant pacman elements for integration for both the target and non-target KICs), which confers a higher level of perceptual and attentional demand posed by the KIC noise condition. Findings of decreased target relative to non-target fixations during this KIC noise condition across groups, provides further evidence for the necessity to dual process (i.e., integrate and segregate) during the complex KIC noise condition - an effect that appears amplified for those with ASD, reflecting a more developmentally immature strategy (Nayar et al., 2015).

Together, findings demonstrate that individuals with ASD likely employ heightened local processing strategies (i.e., increased vacillations and exploration) coupled with reduced global processing strategies (i.e., decreased center and target fixations), in the relative absence of global impairments (i.e., no significant differences in accuracy or reaction time). These findings are consistent with studies documenting the ability of individuals with ASD to shift to global processing when required to do so (particularly in studies employing the Navon Hierarchical Figures) (Koldewyn et al., 2013; Mottron et al., 2003; Mottron et al., 1999; Neufeld et al., 2020; Van der Hallen et al., 2015), in line with the weak central coherence, local processing bias, or the enhanced local-processing efficiency theories (Bertone, Mottron, Jelenic, & Faubert, 2005; Frith & Happe, 1994; Happe, 1996; Happe & Frith, 2006; Jolliffe & Baron-Cohen, 1997; Joseph, Keehn, Connolly, Wolfe, & Horowitz, 2009; Kanner, 1943; Kemner, van Ewijk, van Engeland, & Hooge, 2008; Minshew, Goldstein, & Siegel, 1997; O'Riordan & Plaisted, 2001; Pellicano & Burr, 2012; Shah & Frith, 1983, 1993). As such, findings contribute to the historical debate surrounding local and global processing in ASD, as this study not only utilizes two well-established paradigms of local/global processing with defined neural substrates for the KIC stimuli in particular, but also help address the *strategies* being employed to better extrapolate local/global processing styles.

Gaze in ASD parents reflective of strategy-based differences in global perception

Parent groups did not demonstrate increased rates of vacillations or exploration across and within tasks, although greater exploration was associated with reduced accuracy in both groups in the Navon task. Given that vacillation and exploration are indicative of heightened local perceptual strategies, comparable rates of exploring or vacillating in parents point to the

absence of heightened local processing. These findings contradict literature demonstrating elevated levels of detail orientation among parents of individuals with ASD probed via questionnaires about daily life (Hurley et al., 2007; Van Eylen et al., 2017), described during semi-structured conversations (Losh, 2011; Piven et al., 1997), and evidenced by eye movements during a language processing task (Nayar et al., 2018). Important to note, however, is this study's focus on low-level visual perceptual differences, versus the higher level processes such as language and personality traits, as well as self-perceptions of general detail orientation in everyday life reported in this prior work. Given that parents generally do not present with any functional clinical impairment, lack of heightened local processing in the context of a local/global visual perception-based task is not unsurprising, and consistent with prior reports that parents did not differ in a block design or sentence completion task tapping local processing in an independent sample (Losh et al., 2009).

In contrast, clear differences in global perceptual strategies were observed among parents of individuals with ASD. As in the ASD group, parents too showed diminished target versus non-target looks compared to control parents, and reduced center versus pacman fixations during the KIC noise condition with local interference. Although parents' global processing was not impaired (given highly accurate responding), it was reduced in the presence of local distractors, suggesting a reliance on different strategies to complete the task, compared to control parents. Interestingly, differences observed in the Navon paradigm were subtle and nuanced — parents of individuals with ASD made fewer first fixations towards the target (i.e., global match) versus the non-target (i.e., local match), perhaps suggesting that ASD parents were less able to tap peripheral visual processes to engage in greater global processing, which reflects slower and less efficient processing more broadly (Kimchi, 1992; Nayar et al., 2015). Taken together, parallel

findings of reduced global perception across ASD parent and proband groups, and heightened local processing in ASD probands, suggest the potential for shared underlying genetic risk for ASD that contributes to visual perceptual styles documented here.

Clinical-behavioral and cognitive correlates of visual perceptual styles

The weak central coherence or enhanced perceptual functioning theories posit that differences in perceptual styles relate to downstream ASD symptomatology (Behrmann et al., 2006; Bertone et al., 2005; Burnette et al., 2005; Fitch et al., 2015; Happe, 1996, 1999; Happe & Frith, 2006; Jarrold et al., 2000; Jolliffe & Baron-Cohen, 1997, 2000; Joseph et al., 2009; Kemner et al., 2008; Klin et al., 2002; Minshew et al., 1997; Neufeld et al., 2020; O'Riordan & Plaisted, 2001; Shah & Frith, 1983, 1993; Van der Hallen et al., 2015; Van Eylen et al., 2018). Robust positive relationships emerged between vacillations and exploration and the severity of RRBs in ASD, suggesting that local processing and RRBs in ASD are intricately related, and perhaps stem from a common neurobiological origin. Individuals with ASD also exhibited reduced global visual processing overall, which was surprisingly related to *better* pragmatic language skills during conversational interactions. Parents of individuals with ASD showed parallel findings of reduced global perception relating to better pragmatic language abilities in conversation. Conversational skills require the utilization of global strategies such as integrating local, detailed information into global conversational themes. However, local strategies may also be helpful as an alternative strategy for tracking the details of a communication exchange. This underscores the notion that individuals with ASD, and their family members, may be using alternative strategies to achieve the same outcome as their respective controls. Certainly, the items comprising the domains of pragmatic language in which these links were found included

items such as being vague, being unable to clarify, being informal, failing to reciprocate, and failing to elaborate. Perhaps applying local strategies in a conversation that taps these global skills is compensatory for the ASD groups (e.g., being vague might be being "too" global).

There were no significant relationships with measures of social cognition or executive functioning and local/global processing across groups. Lack of relationships with executive functions help to address conflicting findings from studies that assessed the independent or dependent role of executive functioning in local/global processing, and that they are inherently separate entities (Bolte et al., 2007; Rinehart et al., 2000; Van Eylen et al., 2017) rather than local/global perception being an executive (versus a perceptual) impairment (Booth et al., 2003; Happe & Frith, 2006; Van Eylen et al., 2017; Van Eylen et al., 2018).

Finally, potentially important relationships in visual-perceptual styles emerged between parents and their children diagnosed with ASD, that were not evident in control parent-child pairs or in unrelated parent-child pairs across diagnostic groups. Though prior work has demonstrated shared gaze trajectories in families with typically developing individuals (Constantino et al., 2017), lack of relationships within control families in the present study is not surprising given lack of impairments in low level local/global processing in this group. Relationships of gaze in families with an affected individual are not only consistent with prior work documenting the heritability of gaze or local/global processing in ASD (Constantino et al., 2017; Neufeld et al., 2020), but also imply shared underlying biological mechanisms of local/global perception specific to families with an affected member.

Conclusions and Future Directions

This study is among the first of its kind to examine local/global perception in parents of individuals with ASD using eye tracking as a means to identify potential objective ASD-related endophenotypes and biological markers. Results highlight the objective and quantifiable nature of eye tracking to capture *strategies* related to local/global processing, and they also raise important questions regarding the types of stimuli utilized in prior and ongoing work of local/global processing in ASD. The use of KICs has been extensively applied to studies of schizophrenia and typical development within the realms of vision and cognitive science, particularly given their distinct neural correlates (Altschuler et al., 2014; Altschuler et al., 2012; Foxe, Doniger, & Javitt, 2001; Foxe et al., 2005; Foxe & Simpson, 2002). Yet, studies are lacking in their application to ASD. With the addition of eye tracking, the present study was able to not only document reduced global processing in ASD, but also was able to demonstrate heightened local processing in the same study. As such, the use of KICs offers a unique stance in the examination of both levels of processing simultaneously without one interfering with the other, which is often the case in much of the work already conducted in ASD and confounds the examination of baseline perceptual strategies (Van der Hallen et al., 2015). Interestingly, underlying neural correlates of KICs documented in human subjects have also been extended to non-human primates (Feltner & Kiorpes, 2010; Sary et al., 2008), thus shedding light into its utility as a marker of underlying neurobiology that is devoid of culture, language, and higher cognitive abilities (unlike, for example, social stimuli). As such, KIC perception may be further examined in the context of an ASD-family study in cross-cultural work, which has the potential to span cultural boundaries to reveal a cross-cultural ASD-related endophenotype, and which may be used in future studies examining underlying neurobiological and molecular mechanisms of ASD.

Future studies may also consider the perceptual maturation processes among those with ASD more directly, as well as studying these phenomena in lower functioning individuals. Due to the demands of the match-to-sample task in the present study, participants included high functioning individuals with ASD with intact cognitive ability, reducing the generalizability of the findings to the full spectrum of ASD severity. Finally, given that BAP traits are subtly expressed in a subset of parents of individuals with ASD (Losh, Adolphs, & Piven, 2011; Losh et al., 2009; Losh, Childress, Lam, & Piven, 2008), examination of underlying neural correlates may be more fruitful than eye tracking or performance measures alone. Indeed, recent work has documented atypicalities of local/global visual processing using KICs in individuals with ASD compared to controls (Baruth et al., 2010; Stroganova, Orekhova, et al., 2007; Stroganova et al., 2012), and more robust differences in underlying neural correlates during facial processing tasks observed among parents of individuals with ASD (Billeci et al., 2016; Yucel et al., 2015). As such, given well-studied neural circuitry of local and global visual processing (Altmann et al., 2003; Altschuler et al., 2014; Altschuler et al., 2012; Doniger et al., 2002; Doniger et al., 2000; Doniger et al., 2001; Fink et al., 1997; Foxe et al., 2005; Stroganova, Orekhova, et al., 2007; Stroganova et al., 2012), future investigation to better understand the neural underpinnings of visual processing styles in first-degree relatives of individuals with ASD is warranted and may provide important insights into the gene-brain-behavior relationships of visual processing and ASD.

Declarations

Ethics approval and consent to participate. *Ethical approval.* All procedures performed in this study were in accordance with the ethical standards of the institutional research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. *Informed consent.* Informed consent was obtained from all individual participants included in the study.

Competing interests. FS is a paid consultant for Roche and Janssen pharmaceutical companies. All other authors declare that they have no disclosures or conflict of interest.

Funding. This research was supported by grants from the National Institutes of Health (R01DC010191), the National Institutes of Mental Health (R03MH107834); and the National Science Foundation (BCS-0820394).

Authors' contributions. ML conceived the study and oversaw all data collection, analyses, and writing of the manuscript. KN conceived the study, assisted in study and stimulus design, and acquisition and processing of data, and led the development of eye-tracking variables, data analyses, and manuscript preparation. MW assisted with acquisition of data, and contributed to manuscript preparation. CS assisted with acquisition and management of data and manuscript preparation. JG assisted with manuscript preparation. All authors read and approved the final manuscript.

Acknowledgments. We are grateful to the individuals and families who participated in this study. We are additionally grateful to Nell Maltman and Jamie Barstein for their substantial contributions to data acquisition and participant recruitment. We thank Adriana Di Martino, M.D. and Lynne Kiorpes, Ph.D. for their consultation on stimulus design and paradigm. We thank Chao Tang and Chun Liang Chan and for their substantive contributions and expertise in computer programming for the stimuli design and data processing.

Table 1: Sample Characteristics

KIC	Control Group			ASD Group			Group Difference		
	M	Range	SD	M	Range	SD	t	df	p
Probands									
n (M/F)	30 (18/12)	-	-	31 (25/6)	-		-	-	-
Age (years)	14.72	8.25 - 35.64	6.04	16.03	9.53 - 32.42	4.73	-0.95	59	0.347
FSIQ	116.6	88 - 145	13.8	98.1	73 - 135	16.1	4.82	59	<0.0001
VIQ	116.8	93 - 142	13.3	99.3	69 - 142	18.9	4.17	59	<0.0001
PIQ	115.8	79 - 149	16.7	97.7	68 - 136	17.8	4.07	59	<0.0001
ADOS Total Severity Score^	-	-	-	8.4	5 - 10	1.4	-	-	-
SA Severity Score	-	-	-	8.3	4 - 10	1.5	-	-	-
RRB Severity Score	-	-	-	7.9	1 - 10	1.9	-	-	-
Parents									
n (M/F)	43 (11/32)			55 (17/38)			-	-	-
Age (years)	45.36	31.85 - 67.25	9.29	48.45	35.53 - 61.8	6.84	-1.83	74.73	0.072
FSIQ	118.3	83 - 134	11.4	111.7	85 - 136	12.2	2.75	96	0.007
VIQ	115.0	84 - 132	10.8	110.3	80 - 131	12.1	2.01	96	0.047
PIQ	116.4	86 - 136	11.1	110.1	83 - 137	11.9	2.70	96	0.008

Bold indicates significance $p < .05$; Italics indicates unequal variance assumed; ^Comparison severity score labels are as follows: 0-2 = “minimal-to-no evidence”, 3-4 = “low”, 5-7 = “moderate”, 8-10 = “high”. ADOS, Autism Diagnostic Observation Scale; FSIQ, Full-Scale IQ; PIQ, Performance IQ; RRB, Restricted and Repetitive Behaviors and Interests; VIQ, Verbal IQ.

Table 1: Sample Characteristics

Navon	Control Group			ASD Group			Group Difference		
	M	Range	SD	M	Range	SD	t	df	p
Probands									
n (M/F)	20 (11/9)	-	-	22 (18/4)	-		-	-	-
Age (years)	13.88	8.25 - 35.64	6.01	17.09	10.88 - 32.42	5.03	-1.88	40	0.067
FSIQ	116.8	88 - 145	13.5	97.2	73 - 135	14.8	4.47	40	<0.0001
VIQ	117.5	93 - 142	13.2	100.0	71 - 136	17.4	3.64	40	0.001
PIQ	116.2	79 - 149	17.4	95.4	68 - 135	17.8	3.82	40	<0.0001
ADOS Total Severity Score [^]	-	-	-	8.4	5 - 10	1.4	-	-	-
SA Severity Score	-	-	-	8.1	4 - 10	1.5	-	-	-
RRB Severity Score	-	-	-	8.1	1 - 10	2.0	-	-	-
Parents									
n (M/F)	39 (9/30)			47 (15/32)			-	-	-
Age (years)	45.52	31.85 - 68.4	9.72	48.54	35.71 - 62.06	6.94	<i>-1.63</i>	67	<i>0.108</i>
FSIQ	117.6	83 - 134	11.6	111.3	85 - 136	12.2	2.42	84	0.018
VIQ	114.6	84 - 132	11.1	109.6	80 - 130	12.2	1.95	84	0.055
PIQ	115.5	86 - 136	11.0	110.2	83 - 137	12.0	2.12	84	0.037

Bold indicates significance $p < .05$; Italics indicates unequal variance assumed; [^]Comparison severity score labels are as follows: 0-2 = “minimal-to-no evidence”, 3-4 = “low”, 5-7 = “moderate”, 8-10 = “high”. ADOS, Autism Diagnostic Observation Scale; FSIQ, Full-Scale IQ; PIQ, Performance IQ; RRB, Restricted and Repetitive Behaviors and Interests; VIQ, Verbal IQ.

Table 2. Morphologically similar and dissimilar letters presented in a pseudo-random order (randomized within a participant in the same order across participants)

Navon Hierarchical Figure Letter Pairs	
Morphologically Similar	Morphologically Dissimilar
FE	ON
JU	LC
VW	KM
MN	ET
AH	XS
NM	AJ
BP	EO
IT	RN
TI	HZ
UJ	NR
EF	OE
LE	ZH
OC	NO
XK	MK
WV	HX
HA	CL
CO	TE
KX	SX
PB	XH
EL	JA

Table 3: Definitions of KIC and Navon performance and gaze variables and composites

	Variable Definition (KIC)	Variable Definition (Navon)
Performance-based variables		
Accuracy	Percent correct based on target click	Percent correct based on target (i.e., global) click (only present for overall Navon analyses)
Reaction Time	Average reaction time in seconds per trial	Average reaction time in seconds per trial
Strategy-based gaze variables		
Fixation Rate (exploration)	Average number of fixations per second per trial	Average number of fixations per second per trial
Vacillations	Average percent of transition fixations between target versus non-target stimuli	Average percent of transition fixations between target versus non-target stimuli
Center-Pacman Difference (i.e., Diff) Composite	Average percent difference between center and pacman looks (higher value indicates a greater percentage of center looks)	-
Center-Pacman Diff First Fixation	Average percent difference of first fixations made towards KIC centers versus pacman first fixations (higher value indicates a greater percentage of center first fixations)	-
Center-Pacman Diff Fixation	Average percent difference of fixations made towards KIC centers versus pacman fixations (higher value indicates a greater percentage of center fixations)	-
Center-Pacman Diff Fixation Duration	Average percent difference of fixation duration spent attending towards KIC centers versus pacman elements (higher value indicates a greater percentage of time allocated towards KIC centers)	-
Target-Non-Target Diff Composite	Average percent difference between target and non-target looks (higher value indicates a greater percentage of target looks)	Average percent difference between target and non-target looks (higher value indicates a greater percentage of target looks)
Target-Non-Target Diff First Fixation	Average percent difference of first fixations made towards the target KIC versus the non-target KIC (higher value indicates a greater percentage of target first fixations)	Average percent difference of first fixations made towards the target/global Navon versus the non-target/local Navon (higher value indicates a greater percentage of target first fixations)

Target-Non-Target Diff Fixation	Average percent difference of fixations made towards the target KIC versus the non-target KIC (higher value indicates a greater percentage of target match fixations)	Average percent difference of fixations made towards the target/global Navon versus the non-target/local Navon (higher value indicates a greater percentage of target fixations)
Target-Non-Target Diff Fixation Duration	Average percent difference of fixation duration spent attending towards the target KIC versus the non-target KIC (higher value indicates a greater percentage of time allocated toward the target match).	Average percent difference of fixation duration spent attending towards the target/global Navon versus the non-target/local Navon (higher value indicates a greater percentage of time allocated toward the target Navon stimulus)
<hr/>		
Composite score		
Global-Local Diff Composite	Average z-scored sum of accuracy, reaction time, target and center percent first fixations, fixations, and duration minus the z-scored sum of the non-target and pacman element first fixations, fixations, and duration as well as percent vacillations and fixation rate (higher value indicates greater "global" than "local" processing)	Average z-scored sum of accuracy, reaction time, target percent first fixations, fixations, and duration minus the z-scored sum of the non-target first fixations, fixations, and duration as well as percent vacillations and fixation rate (higher value indicates greater "global" than "local" processing)
<hr/>		

Table 4: Summary of KIC Results- ASD versus Control groups

	Control Group		ASD Group	
	KIC Basic M (SE)	KIC Noise M (SE)	KIC Basic M (SE)	KIC Noise M (SE)
Performance-based variables				
Accuracy^	97.80 (0.89)	97.30 (0.86)	96.00 (0.96)	96.00 (0.91)
Reaction Time^ (s)	.84 (0.08)	1.24 (0.08)	1.06 (0.09)	1.63 (0.08)
Strategy-based gaze variables				
Fixation Rate	3.38 (0.10)	3.63 (0.10)	3.61 (0.10)	4.00 (0.11)
Vacillations	20.12 (1.41)	14.29 (0.89)	25.91 (2.13)	15.05 (0.60)
Center-Pacman Diff Composite	13.73 (4.85)	-27.38 (3.6)	8.27 (3.91)	-44.72 (3.97)
Center-Pacman Diff First Fixation	19.14 (4.44)	-22.58 (3.31)	15.98 (4.27)	-39.84 (3.78)
Center-Pacman Diff Fixation	23.81 (4.79)	-13.16 (3.69)	18.85 (4.88)	-33.35 (4.68)
Center-Pacman Diff Fixation Duration	9.02 (4.33)	-49.29 (4.45)	-0.7 (3.05)	-61.31 (3.31)
Target-Non-Target Diff Composite	50.25 (2.76)	41.41 (2.14)	34.83 (2.51)	28.29 (2.30)
Target-Non-Target Diff First Fixation	49.68 (2.36)	42.42 (2.07)	36.73 (2.59)	30.59 (2.25)
Target-Non-Target Diff Fixation	63.14 (2.51)	56.57 (1.99)	47.79 (3.05)	40.33 (2.70)
Target-Non-Target Diff Fixation Duration	37.92 (4.02)	25.24 (2.90)	19.97 (2.54)	13.94 (2.62)
Composite z-score				
Global-Local Diff Composite	0.98 (0.16)	0.05 (0.15)	-0.02 (0.20)	-1.02 (0.18)

Table 5: Summary of KIC Results - ASD parent versus parent control groups

	Parent Control Group		ASD Parent Group	
	KIC Basic M (SE)	KIC Noise M (SE)	KIC Basic M (SE)	KIC Noise M (SE)
Performance-based variables				
Accuracy	99.00 (0.3)	99.00 (0.23)	98.32 (0.32)	98.38 (0.32)
Reaction Time	0.83 (0.03)	1.35 (0.05)	0.96 (0.02)	1.55 (0.07)
Strategy-based gaze variables				
Fixation Rate	3.43 (0.07)	3.73 (0.07)	3.50 (0.07)	3.75 (0.06)
Vacillations	24.74 (1.40)	14.85 (0.47)	22.37 (1.44)	16.11 (0.48)
Center-Pacman Diff Composite	6.66 (2.68)	-40.5 (2.39)	9.58 (2.42)	-44.82 (2.01)
Center-Pacman Diff First Fixation	12.69 (2.85)	-33.02 (2.45)	15.14 (2.58)	-37.94 (2.12)
Center-Pacman Diff Fixation	17.35 (3.10)	-24.43 (2.74)	20.29 (2.87)	-29.48 (2.42)
Center-Pacman Diff Fixation Duration	-2.07 (2.52)	-67.47 (2.49)	-1.23 (2.43)	-70.99 (2.12)
Target-Non-Target Diff Composite	39.3 (1.97)	32.37 (1.01)	34.38 (1.63)	29.23 (1.10)
Target-Non-Target Diff First Fixation	40.98 (1.94)	35.36 (1.14)	36.22 (1.53)	32.2 (1.18)
Target-Non-Target Diff Fixation	56.74 (1.86)	48.45 (1.23)	50.66 (1.77)	44.45 (1.39)
Target-Non-Target Diff Fixation Duration	20.18 (2.61)	13.31 (1.42)	16.25 (2.17)	11.03 (1.40)
Composite z-score				
Global-Local Diff Composite	0.60 (0.14)	-0.32 (0.09)	0.42 (0.11)	-0.64 (0.10)

ASD versus control groups	<i>Est.</i>	β	<i>SE</i>	<i>df</i>	<i>t</i>	<i>p</i>	<i>padj</i>
Main Effect of Group							
Performance-based variables							
Accuracy^	-1.82	-0.19	1.39	82.14	-1.32	0.192	0.250
Reaction Time^ (s)	0.22	0.19	0.13	76.63	1.73	0.088	0.132
Strategy-based gaze variables							
Fixation Rate	0.34	0.29	0.15	80.67	2.25	0.027	0.048
Vacillations	5.64	0.35	1.90	103.81	2.96	0.004	0.008
Center-Pacman Diff Composite	-5.32	-0.08	5.88	95.31	-0.91	0.368	0.463
Center-Pacman Diff First Fixation	-9.31	-0.13	5.66	97.72	-1.64	0.104	0.144
Center-Pacman Diff Fixation	-2.93	-0.05	5.64	96.27	-0.52	0.604	0.693
Center-Pacman Diff Fixation Duration	-5.25	-0.08	6.37	91.06	-0.82	0.412	0.503
Target-Non-Target Diff Composite	-14.28	-0.48	3.45	88.76	-4.14	0.000	0.000
Target-Non-Target Diff First Fixation	-17.00	-0.46	4.47	95.89	-3.80	0.000	0.001
Target-Non-Target Diff Fixation	-11.60	-0.42	3.27	89.17	-3.54	0.001	0.002
Target-Non-Target Diff Fixation Duration	-14.44	-0.46	3.57	85.25	-4.04	0.000	0.001
Composite z-score							
Global-Local Diff Composite	-0.99	-0.43	0.24	86.24	-4.07	0.000	0.000
Main Effect of Condition							
Performance-based variables							
Accuracy^	-0.51	-0.05	0.80	46.33	-0.64	0.526	0.622
Reaction Time^ (s)	0.41	0.46	0.07	44.53	6.13	0.000	0.000
Strategy-based gaze variables							
Fixation Rate	0.28	0.24	0.08	45.57	3.44	0.001	0.003
Vacillations	-5.92	-0.36	1.59	49.99	-3.72	0.001	0.002
Center-Pacman Diff Composite	-41.09	-0.64	4.05	49.80	-10.16	0.000	0.000
Center-Pacman Diff First Fixation	-58.60	-0.80	4.16	47.06	-14.07	0.000	0.000
Center-Pacman Diff Fixation	-41.49	-0.64	3.95	49.97	-10.50	0.000	0.000
Center-Pacman Diff Fixation Duration	-36.96	-0.56	4.04	49.13	-9.15	0.000	0.000
Target-Non-Target Diff Composite	-8.77	-0.29	2.11	48.08	-4.16	0.000	0.001
Target-Non-Target Diff First Fixation	-12.64	-0.34	3.14	48.76	-4.03	0.000	0.001
Target-Non-Target Diff Fixation	-6.96	-0.25	2.02	48.17	-3.45	0.001	0.003
Target-Non-Target Diff Fixation Duration	-6.74	-0.21	2.01	48.58	-3.35	0.002	0.004
Composite z-score							
Global-Local Diff Composite	-0.93	-0.41	0.14	48.40	-6.61	0.000	0.000
Group X Condition Interaction							
Performance-based variables							
Accuracy^	0.56	0.05	1.20	49.84	0.47	0.643	0.697
Reaction Time^ (s)	0.17	0.09	0.10	47.53	1.67	0.101	0.144

Strategy-based gaze variables

Fixation Rate	0.03	0.02	0.12	48.06	0.23	0.818	0.818
Vacillations	-5.03	-0.27	2.34	54.66	-2.15	0.036	0.061
Center-Pacman Diff Composite	-11.86	-0.16	6.02	53.60	-1.97	0.054	0.085
Center-Pacman Diff First Fixation	-3.07	-0.04	6.18	51.18	-0.50	0.622	0.693
Center-Pacman Diff Fixation	-14.25	-0.19	5.88	53.87	-2.43	0.019	0.037
Center-Pacman Diff Fixation Duration	-14.58	-0.19	6.04	52.51	-2.42	0.019	0.037
Target-Non-Target Diff Composite	1.15	0.03	3.16	51.25	0.36	0.717	0.736
Target-Non-Target Diff First Fixation	5.71	0.14	2.19	56.40	-1.49	0.141	0.190
Target-Non-Target Diff Fixation	-0.20	-0.01	1.61	53.46	-2.12	0.038	0.062
Target-Non-Target Diff Fixation Duration	-1.84	-0.05	1.46	53.66	-2.28	0.026	0.048
Composite z-score							
Global-Local Diff Composite	-0.08	-0.03	0.21	51.30	-0.39	0.699	0.736

Bold $p < .05$; padj reflects the Benjamini-Hochberg correction at a false discovery rate of .10; ^IQ added as a covariate

ASD parent versus parent control groups	<i>Est.</i>	β	<i>SE</i>	<i>df</i>	<i>t</i>	<i>p</i>	<i>padj</i>
Main Effect of Group							
Performance-based variables							
Accuracy	-0.72	-0.17	0.43	171.03	-1.65	0.101	0.187
Reaction Time (s)	0.17	0.19	0.08	121.51	2.25	0.027	0.065
Strategy-based gaze variables							
Fixation Rate	0.09	0.09	0.09	131.70	0.92	0.361	0.522
Vacillations	-2.37	-0.14	1.52	188.99	-1.56	0.121	0.215
Center-Pacman Diff Composite	2.79	0.05	3.36	154.75	0.83	0.408	0.548
Center-Pacman Diff First Fixation	0.97	0.01	3.40	174.64	0.29	0.775	0.817
Center-Pacman Diff Fixation	2.12	0.03	3.55	147.30	0.60	0.551	0.632
Center-Pacman Diff Fixation Duration	2.59	0.04	3.95	146.37	0.66	0.513	0.606
Target-Non-Target Diff Composite	-5.06	-0.23	2.09	157.10	-2.42	0.017	0.047
Target-Non-Target Diff First Fixation	-3.82	-0.14	2.78	172.00	-1.38	0.171	0.290
Target-Non-Target Diff Fixation	-4.99	-0.24	2.09	159.64	-2.39	0.018	0.047
Target-Non-Target Diff Fixation Duration	-6.46	-0.27	2.28	148.01	-2.83	0.005	0.017
Composite z-score							
Global-Local Diff Composite	-0.20	-0.11	0.16	152.51	-1.24	0.216	0.352
Main Effect of Condition							
Performance-based variables							
Accuracy	0.01	0.00	0.38	92.53	0.02	0.983	0.983
Reaction Time (s)	0.52	0.60	0.05	78.12	10.78	<0.0001	<0.0001

Strategy-based gaze variables

Fixation Rate	0.29	0.30	0.06	92.95	5.01	<0.0001	<0.0001
Vacillations	-9.89	-0.59	1.60	96.06	-6.18	<0.0001	<0.0001
Center-Pacman Diff Composite	-46.86	-0.77	2.56	94.04	-18.33	<0.0001	<0.0001
Center-Pacman Diff First Fixation	-65.19	-0.86	3.02	93.48	-21.58	<0.0001	<0.0001
Center-Pacman Diff Fixation	-45.51	-0.75	2.53	93.87	-18.00	<0.0001	<0.0001
Center-Pacman Diff Fixation Duration	-41.53	-0.69	2.79	93.92	-14.88	<0.0001	<0.0001
Target-Non-Target Diff Composite	-6.64	-0.31	1.64	92.06	-4.05	<0.0001	<0.0001
Target-Non-Target Diff First Fixation	-6.59	-0.24	2.43	91.60	-2.71	0.008	0.024
Target-Non-Target Diff Fixation	-5.43	-0.26	1.67	92.27	-3.26	0.002	0.005
Target-Non-Target Diff Fixation Duration	-8.00	-0.34	1.67	91.06	-4.80	<0.0001	<0.0001
Composite z-score							
Global-Local Diff Composite	-0.89	-0.48	0.12	90.37	-7.33	<0.0001	<0.0001

Group X Condition Interaction**Performance-based variables**

Accuracy	0.09	0.03	0.50	92.78	0.18	0.860	0.882
Reaction Time (s)	0.03	0.02	0.06	78.27	0.47	0.643	0.716
Strategy-based gaze variables							
Fixation Rate	-0.05	-0.05	0.08	93.10	-0.67	0.502	0.606
Vacillations	3.63	0.20	2.14	96.34	1.70	0.093	0.182
Center-Pacman Diff Composite	-7.41	-0.11	3.42	94.25	-2.17	0.033	0.075
Center-Pacman Diff First Fixation	-4.70	-0.06	4.04	93.74	-1.16	0.247	0.386
Center-Pacman Diff Fixation	-7.23	-0.11	3.38	94.07	-2.14	0.035	0.076
Center-Pacman Diff Fixation Duration	-7.89	-0.12	3.73	94.11	-2.11	0.037	0.077
Target-Non-Target Diff Composite	1.63	0.07	2.19	92.28	0.74	0.459	0.579
Target-Non-Target Diff First Fixation	1.26	0.04	3.26	91.85	0.39	0.700	0.758
Target-Non-Target Diff Fixation	1.65	0.07	2.23	92.49	0.74	0.460	0.579
Target-Non-Target Diff Fixation Duration	2.17	0.08	2.23	91.26	0.97	0.334	0.501
Composite z-score							
Global-Local Diff Composite	-0.14	-0.07	0.16	90.58	-0.87	0.385	0.536

Bold $p < .05$; padj reflects the Benjamini-Hochberg correction at a false discovery rate of .10.

Table 6: Summary of Navon Results - ASD versus control groups.

	Control Group	ASD Group	Main Effect of Group				
	M (SD)	M (SD)	<i>F</i>	<i>df</i>	<i>p</i>	<i>padj</i>	partial η^2
Performance-based variables							
Accuracy [^]	97.02 (6.85)	75.66 (38.90)	3.81	39	0.058	0.104	0.089
Reaction Time [^]	2.09 (1.54)	1.95 (1.31)	0.08	39	0.786	0.786	0.002
Strategy-based gaze variables							
Fixation Rate	2.89 (0.61)	3.17 (0.58)	2.37	40	0.131	0.168	0.056
Vacillations	25.23 (12.09)	26.84 (9.76)	0.23	40	0.637	0.717	0.006
Target-Non-Target Diff Composite	45.50 (25.32)	24.74 (30.19)	5.76	40	0.021	0.048	0.126
Target-Non-Target Diff First Fixation	34.65 (27.96)	22.50 (21.07)	2.56	40	0.117	0.168	0.060
Target-Non-Target Diff Fixation	45.01 (23.89)	23.20 (31.59)	6.27	40	0.016	0.048	0.135
Target-Non-Target Diff Fixation Duration	56.82 (28.04)	28.53 (43.34)	6.17	40	0.017	0.048	0.134
Composite z-score							
Global-Local Diff Composite	0.25 (1.15)	-0.75 (1.48)	5.90	40	0.020	0.048	0.129

Bold $p < .05$; *padj* reflects the Benjamini-Hochberg correction at a false discovery rate of .10; [^]IQ added as a covariate

Table 7: Summary of Navon Results - ASD parent versus parent control groups.

	Parent Control Group	ASD Parent Group	Main Effect of Group				
	M (SD)	M (SD)	<i>F</i>	<i>df</i>	<i>p</i>	<i>padj</i>	partial η^2
Performance-based variables							
Accuracy	94.55 (20.30)	89.84 (26.53)	0.83	84	0.366	0.412	0.010
Reaction Time	2.05 (2.35)	1.61 (1.60)	1.08	84	0.301	0.412	0.013
Strategy-based gaze variables							
Fixation Rate	3.23 (0.67)	3.48 (0.73)	2.56	84	0.113	0.367	0.030
Vacillations	21.75 (11.68)	22.69 (11.74)	0.14	84	0.710	0.710	0.002
Target-Non-Target Diff Composite	47.90 (23.55)	39.46 (26.45)	2.39	84	0.126	0.367	0.028
Target-Non-Target Diff First Fixation	43.18 (21.36)	31.51 (22.035)	6.15	84	0.015	0.136	0.068
Target-Non-Target Diff Fixation	45.10 (24.27)	38.37 (26.77)	1.46	84	0.230	0.412	0.017

Target-Non-Target Diff Fixation Duration	55.41 (29.02)	48.49 (34.65)	0.98	84	0.325	0.412	0.012
Composite z-score							
Global-Local Diff Composite	0.35 (1.17)	-0.05 (1.37)	1.98	84	0.163	0.367	0.023

Bold $p < .05$; padj reflects the Benjamini-Hochberg correction at a false discovery rate of .10

Table 8: Summary of Navon Condition Results- ASD versus Control Groups

	Control Group		ASD Group	
	NAV Dissimilar	NAV Similar	NAV Dissimilar	NAV Similar
	M (SE)	M (SE)	M (SE)	M (SE)
Performance-based variables				
Reaction Time^	1.33 (0.18)	1.36 (0.18)	1.49 (0.17)	1.52 (0.17)
Strategy-based gaze variables				
Fixation Rate	2.88 (0.13)	2.90 (0.14)	3.14 (0.12)	3.19 (0.14)
Vacillations	24.19 (2.98)	26.37 (2.79)	26.67 (2.19)	26.88 (2.39)
Target-Non-Target Diff Composite	51.48 (5.86)	39.94 (6.14)	25.03 (7.25)	24.80 (6.00)
Target-Non-Target Diff First Fixation	50.73 (5.76)	40.03 (5.66)	23.58 (7.43)	23.42 (6.40)
Target-Non-Target Diff Fixation	60.76 (6.58)	53.55 (6.41)	28.5 (10.13)	29.15 (8.61)
Target-Non-Target Diff Fixation Duration	42.94 (6.22)	26.25 (7.45)	23.02 (5.46)	21.83 (5.16)
Composite score				
Global-Local Diff Composite	0.47 (0.30)	-0.06 (1.39)	-0.82 (0.35)	-0.86 (0.30)

Table 9: Summary of Navon Condition Results - ASD Parent versus Parent Control Groups

	Parent Control Group		ASD Parent Group	
	NAV Dissimilar	NAV Similar	NAV Dissimilar	NAV Similar
	M (SE)	M (SE)	M (SE)	M (SE)
Performance-based variables				
Reaction Time	1.01 (0.04)	1.01 (0.03)	1.08 (0.03)	1.13 (0.04)
Strategy-based gaze variables				
Fixation Rate	3.20 (0.11)	3.24 (0.12)	3.45 (0.11)	3.49 (0.11)
Vacillations	19.61 (2.00)	23.84 (2.00)	22.26 (1.88)	23.07 (1.74)
Target-Non-Target Diff Composite	52.08 (3.97)	43.96 (3.92)	42.5 (4.00)	36.61 (4.03)
Target-Non-Target Diff First Fixation	49.02 (4.07)	41.71 (4.05)	41.06 (4.00)	36.10 (4.13)
Target-Non-Target Diff Fixation	58.68 (4.84)	52.39 (4.71)	51.29 (5.16)	46.00 (5.15)
Target-Non-Target Diff Fixation Duration	48.55 (3.76)	37.78 (3.93)	35.15 (3.5)	27.72 (3.73)
Composite score				
Global-Local Diff Composite	0.63 (0.19)	0.21 (0.19)	.10 (0.20)	-0.19 (0.19)

ASD versus control groups	<i>Estimate</i>	β	<i>SE</i>	<i>df</i>	<i>t value</i>	<i>p</i>	<i>padj</i>
Main Effect of Condition							
Performance-based variables							
Reaction Time [^]	0.04	0.00	0.03	40.00	1.19	0.240	0.350
Strategy-based gaze variables							
Fixation Rate	0.02	0.02	0.07	40.00	0.31	0.761	0.802
Vacillations	2.18	0.09	1.94	40.00	1.12	0.268	0.358
Target-Non-Target Diff Composite	-11.53	-0.19	3.82	40.00	-3.02	0.004	0.024
Target-Non-Target Diff First Fixation	-16.69	-0.29	5.92	40.00	-2.82	0.007	0.030
Target-Non-Target Diff Fixation	-10.70	-0.17	3.54	40.00	-3.03	0.004	0.024
Target-Non-Target Diff Fixation Duration	-7.21	-0.09	3.61	40.00	-2.00	0.053	0.106
Composite score							
Global-Local Diff Composite	-0.53	-0.17	0.18	40.00	-3.02	0.004	0.024
Group X Condition Interaction							
Performance-based variables							
Reaction Time [^]	-0.01	0.00	0.04	40.00	-0.25	0.802	0.802
Strategy-based gaze variables							
Fixation Rate	0.03	0.02	0.09	40.00	0.35	0.728	0.802
Vacillations	-1.98	-0.08	2.69	40.00	-0.74	0.466	0.573
Target-Non-Target Diff Composite	11.30	0.16	5.28	40.00	2.14	0.039	0.103
Target-Non-Target Diff First Fixation	15.49	0.24	8.18	40.00	1.89	0.066	0.117
Target-Non-Target Diff Fixation	10.54	0.15	4.88	40.00	2.16	0.037	0.103
Target-Non-Target Diff Fixation Duration	7.86	0.09	4.99	40.00	1.57	0.123	0.197
Composite score							
Global-Local Diff Composite	0.48	0.14	0.24	40.00	1.99	0.053	0.106

Bold $p < .05$; padj reflects the Benjamini-Hochberg correction at a false discovery rate of .10; [^]IQ added as a covariate

ASD parent versus parent control groups	<i>Estimate</i>	β	<i>SE</i>	<i>df</i>	<i>t value</i>	<i>p</i>	<i>padj</i>
Main Effect of Condition							
Performance-based variables							
Reaction Time	0.01	-0.01	0.03	84.00	0.21	0.837	0.893
Strategy-based gaze variables							
Fixation Rate	0.04	0.02	0.06	84.00	0.54	0.590	0.726
Vacillations	4.24	0.17	1.36	84.00	3.12	0.002	0.010
Target-Non-Target Diff Composite	-8.12	-0.15	2.34	84.00	-3.47	0.001	0.007
Target-Non-Target Diff First Fixation	-10.77	-0.21	3.54	84.00	-3.04	0.003	0.010
Target-Non-Target Diff Fixation	-7.30	-0.14	2.37	84.00	-3.08	0.003	0.010
Target-Non-Target Diff Fixation Duration	-6.29	-0.10	2.18	84.00	-2.88	0.005	0.013
Composite score							

Global-Local Diff Composite	-0.41	-0.16	0.11	84.00	-3.73	< 0.0001	0.006
Group X Condition Interaction							
Performance-based variables							
Reaction Time	0.05	0.08	0.04	84.00	1.33	0.186	0.372
Strategy-based gaze variables							
Fixation Rate	0.01	0.01	0.09	84.00	0.14	0.893	0.893
Vacillations	-3.43	-0.12	1.84	84.00	-1.87	0.065	0.149
Target-Non-Target Diff Composite	2.23	0.04	3.17	84.00	0.70	0.483	0.649
Target-Non-Target Diff First Fixation	3.35	0.06	4.79	84.00	0.70	0.487	0.649
Target-Non-Target Diff Fixation	2.34	0.04	3.21	84.00	0.73	0.467	0.649
Target-Non-Target Diff Fixation Duration	1.00	0.01	2.95	84.00	0.34	0.736	0.842
Composite score							
Global-Local Diff Composite	0.12	0.04	0.15	84.00	0.80	0.427	0.649

Bold $p < .05$; p_{adj} reflects the Benjamini-Hochberg correction at a false discovery rate of .10

Table 10: Summary of cross-task Results - ASD versus control groups.

	Control Group	ASD Group	Main Effect of Group				
	M (SD)	M (SD)	<i>F</i>	<i>df</i>	<i>p</i>	<i>p</i> _{adj}	partial η^2
Strategy-based gaze variables							
Target-Non-Target Diff Composite	44.27(15.08)	33.11 (15.87)	4.38	32	0.044	0.044	0.120
Composite z-score							
Global-Local Diff Composite	0.33 (0.84)	-0.42 (1.03)	5.50	32	0.025	0.044	0.147

Bold $p < .05$; p_{adj} reflects the Benjamini-Hochberg correction at a false discovery rate of .10

Table 11: Summary of cross-task Results - ASD parent versus parent control groups.

	Parent Control Group	ASD Parent Group	<i>F</i>	<i>df</i>	Main Effect of Group		
	M (SD)	M (SD)			<i>p</i>	<i>padj</i>	partial η^2
Strategy-based gaze variables							
Target-Non-Target Diff Composite	41.51 (13.69)	34.79 (13.87)	4.84	80	0.031	0.061	0.057
Composite z-score							
Global-Local Diff Composite	0.23 (0.82)	-0.11 (0.85)	3.27	80	0.074	0.074	0.039

Bold $p < .05$; p_{adj} reflects the Benjamini-Hochberg correction at a false discovery rate of .10

Figure 1. Stimuli presented. a) an example of a Kanizsa Illusory Contour (KIC) with an illusory square percept; b) match-to-sample sequence of test stimuli for the KIC Paradigm (KIC basic and KIC noise, respectively) using KICs; c) match-to-sample sequence of test stimuli for the Navon Paradigm, using hierarchical Navon figures.

Figure 2. A series of images depicting eye tracking trajectories during the KIC Paradigm between an individual with ASD compared to a control individual. Red KIC is the non-target image and the green KIC is the target match. Star depicts location of fixation. A) A panel depicting a control individual making 3 consecutive fixations to the target before selecting the match. B) panel depicting an individual with ASD making 3 consecutive fixations to the non-target before fixating on the target.

Figure 3. Performance and gaze for the KIC basic and KIC noise condition for A) Accuracy, B) Reaction time, C) Visual exploration, D) Vacillation between target and non-target, E) Center-Pacman composite gaze score, F) Target-NonTarget composite gaze score, G) Global-Local Composite z-score.

Figure 4. Group differences for the Navon Paradigm overall for A) Accuracy, B) Reaction time, C) Visual exploration, D) Vacillation between target and non-target, E) Target-NonTarget composite gaze score, F) Global-Local Composite z-score

Figure 5. Group differences across tasks indicate A) ASD and ASD parents looked more comparably to target and non target stimuli than control counterparts, who looked more to target stimuli and B) Overall, the ASD group processed more locally than their TD counterparts (more global) with a similar but nonsignificant pattern emerging for parent groups.

Figure 6. Grids depict inter-variable correlations for A) the KIC Paradigm (KIC basic and KIC noise combined), and B) the Navon Paradigm between performance indices (accuracy and reaction time) and gaze variables.

Figure 7. Parent child correlations in the Overall Global-Local Difference z-score. A) Correlations for matched pairs of parents and their children indicate a moderate-strong relationship of gaze patterns in ASD families only. B) This effect is not seen in ASD or Control families when a child is paired with a different parent in the respective parent group. Distributions produced from randomization test to examine familiarity of local/global processing in parent and child dyads. Frequency distribution reflects the frequency of obtaining a correlation coefficient value for random parent-child pairings; arrow signifies true correlation coefficient between parent and child dyad. For ASD families, probability $r_{\text{true}} > r_{\text{random}} = 97.3\%$; for Control families, probability $r_{\text{true}} > r_{\text{random}} = 76.2\%$.

Figure 8. Clinical-behavioral correlates of local/global processing. A) and B) depict significant associations between elevated rates of restricted and repetitive behaviors (RRB) based on the Autism Diagnostic Interview-Revised (ADI-R) with increased vacillation and exploration across tasks; C) individuals with ASD and D) parents of individuals with ASD show greater global processing abilities relating to poorer pragmatic language abilities (higher score = greater impairment).

Citations

- Adolphs, R., Spezio, M. L., Parlier, M., & Piven, J. (2008). Distinct face-processing strategies in parents of autistic children. *Curr Biol*, 18(14), 1090-1093. doi:10.1016/j.cub.2008.06.073
- Altmann, C. F., Bulthoff, H. H., & Kourtzi, Z. (2003). Perceptual organization of local elements into global shapes in the human visual cortex. *Curr Biol*, 13(4), 342-349. doi:10.1016/s0960-9822(03)00052-6
- Altschuler, T. S., Molholm, S., Butler, J. S., Mercier, M. R., Brandwein, A. B., & Foxe, J. J. (2014). The effort to close the gap: tracking the development of illusory contour processing from childhood to adulthood with high-density electrical mapping. *Neuroimage*, 90, 360-373. doi:10.1016/j.neuroimage.2013.12.029
- Altschuler, T. S., Molholm, S., Russo, N. N., Snyder, A. C., Brandwein, A. B., Blanco, D., & Foxe, J. J. (2012). Early electrophysiological indices of illusory contour processing within the lateral occipital complex are virtually impervious to manipulations of illusion strength. *Neuroimage*, 59(4), 4074-4085. doi:10.1016/j.neuroimage.2011.10.051
- APA. (1994). *Publication manual of the American Psychological Association* (4th Edition ed.). Washington, DC: American Psychiatric Association.
- APA. (2013). *Diagnostic and statistical manual of mental disorders (DSM-5)* (5th ed.). Arlington, VA: American Psychiatric Association.
- Au-Yeung, S. K., Benson, V., Castelhamo, M., & Rayner, K. (2011). Eye movement sequences during simple versus complex information processing of scenes in autism spectrum disorder. *Autism Res Treat*, 2011, 657383. doi:10.1155/2011/657383
- Baron-Cohen, S., & Hammer, J. (1997). Parents of Children with Asperger Syndrome: What is the Cognitive Phenotype? *J Cogn Neurosci*, 9(4), 548-554. doi:10.1162/jocn.1997.9.4.548
- Baron-Cohen, S., Wheelwright, S., Hill, J., Raste, Y., & Plumb, I. (2001). The "Reading the Mind in the Eyes" Test revised version: a study with normal adults, and adults with Asperger syndrome or high-functioning autism. *J Child Psychol Psychiatry*, 42(2), 241-251. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/11280420>
- Baron-Cohen, S., Wheelwright, S., & Jolliffe, T. (1997). Is there a "language of the eyes"? Evidence from normal adults, and adults with autism or Asperger syndrome. *Visual cognition*, 4(3), 311-331.
- Baruth, J. M., Casanova, M. F., Sears, L., & Sokhadze, E. (2010). Early-stage visual processing abnormalities in high-functioning autism spectrum disorder (ASD). *Transl Neurosci*, 1(2), 177-187. doi:10.2478/v10134-010-0024-9
- Bates, D., Mächler, M., Bolker, B., & Walker, S. (2014). Fitting linear mixed-effects models using lme4. *arXiv preprint arXiv:1406.5823*.

- Behrmann, M., Thomas, C., & Humphreys, K. (2006). Seeing it differently: visual processing in autism. *Trends Cogn Sci*, 10(6), 258-264. doi:10.1016/j.tics.2006.05.001
- Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate - a Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society Series B-Statistical Methodology*, 57(1), 289-300. doi:DOI 10.1111/j.2517-6161.1995.tb02031.x
- Bertone, A., Mottron, L., Jelenic, P., & Faubert, J. (2005). Enhanced and diminished visuo-spatial information processing in autism depends on stimulus complexity. *Brain*, 128(Pt 10), 2430-2441. doi:10.1093/brain/awh561
- Billeci, L., Calderoni, S., Conti, E., Gesi, C., Carmassi, C., Dell'Osso, L., . . . Guzzetta, A. (2016). The broad autism (endo)phenotype: neurostructural and neurofunctional correlates in parents of individuals with autism spectrum disorders. *Front Neurosci*, 10, 346. doi:10.3389/fnins.2016.00346
- Bolte, S., Holtmann, M., Poustka, F., Scheurich, A., & Schmidt, L. (2007). Gestalt perception and local-global processing in high-functioning autism. *J Autism Dev Disord*, 37(8), 1493-1504. doi:10.1007/s10803-006-0231-x
- Bolte, S., & Poustka, F. (2006). The broader cognitive phenotype of autism in parents: how specific is the tendency for local processing and executive dysfunction? *J Child Psychol Psychiatry*, 47(6), 639-645. doi:10.1111/j.1469-7610.2006.01603.x
- Booth, R., Charlton, R., Hughes, C., & Happe, F. (2003). Disentangling weak coherence and executive dysfunction: planning drawing in autism and attention-deficit/hyperactivity disorder. *Philos Trans R Soc Lond B Biol Sci*, 358(1430), 387-392. doi:10.1098/rstb.2002.1204
- Briskman, J., Happe, F., & Frith, U. (2001). Exploring the cognitive phenotype of autism: weak "central coherence" in parents and siblings of children with autism: II. Real-life skills and preferences. *J Child Psychol Psychiatry*, 42(3), 309-316.
- Bulf, H., Valenza, E., & Simion, F. (2009). The visual search of an illusory figure: a comparison between 6-month-old infants and adults. *Perception*, 38(9), 1313-1327. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/19911629>
- Burnette, C. P., Mundy, P. C., Meyer, J. A., Sutton, S. K., Vaughan, A. E., & Charak, D. (2005). Weak central coherence and its relations to theory of mind and anxiety in autism. *J Autism Dev Disord*, 35(1), 63-73. doi:10.1007/s10803-004-1035-5
- Chouinard, P. A., Unwin, K. L., Landry, O., & Sperandio, I. (2016). Susceptibility to optical illusions varies as a function of the autism-spectrum quotient but not in ways predicted by local-global biases. *J Autism Dev Disord*, 46(6), 2224-2239. doi:10.1007/s10803-016-2753-1
- Constantino, J. N., & Gruber, C. (2005). *Social Responsiveness Scale*. Los Angeles: Western Psychological Services.

- Constantino, J. N., Kennon-McGill, S., Weichselbaum, C., Marrus, N., Haider, A., Glowinski, A. L., . . . Jones, W. (2017). Infant viewing of social scenes is under genetic control and is atypical in autism. *Nature*, 547(7663), 340-344. doi:10.1038/nature22999
- Cox, M. A., Schmid, M. C., Peters, A. J., Saunders, R. C., Leopold, D. A., & Maier, A. (2013). Receptive field focus of visual area V4 neurons determines responses to illusory surfaces. *Proc Natl Acad Sci U S A*, 110(42), 17095-17100. doi:10.1073/pnas.1310806110
- Cribb, S. J., Olaithe, M., Di Lorenzo, R., Dunlop, P. D., & Maybery, M. T. (2016). Embedded figures test performance in the broader autism phenotype: a meta-analysis. *J Autism Dev Disord*, 46(9), 2924-2939. doi:10.1007/s10803-016-2832-3
- Dennis, M., Francis, D. J., Cirino, P. T., Schachar, R., Barnes, M. A., & Fletcher, J. M. (2009). Why IQ is not a covariate in cognitive studies of neurodevelopmental disorders. *J Int Neuropsychol Soc*, 15(3), 331-343. doi:10.1017/S1355617709090481
- DiCriscio, A. S., Hu, Y., & Troiani, V. (2019). Brief Report: Pupillometry, Visual Perception, and ASD Features in a Task-Switching Paradigm. *J Autism Dev Disord*, 49(12), 5086-5099. doi:10.1007/s10803-019-04213-8
- Doniger, G. M., Foxe, J. J., Murray, M. M., Higgins, B. A., & Javitt, D. C. (2002). Impaired visual object recognition and dorsal/ventral stream interaction in schizophrenia. *Arch Gen Psychiatry*, 59(11), 1011-1020. doi:10.1001/archpsyc.59.11.1011
- Doniger, G. M., Foxe, J. J., Murray, M. M., Higgins, B. A., Snodgrass, J. G., Schroeder, C. E., & Javitt, D. C. (2000). Activation timecourse of ventral visual stream object-recognition areas: high density electrical mapping of perceptual closure processes. *J Cogn Neurosci*, 12(4), 615-621. doi:10.1162/089892900562372
- Doniger, G. M., Foxe, J. J., Schroeder, C. E., Murray, M. M., Higgins, B. A., & Javitt, D. C. (2001). Visual perceptual learning in human object recognition areas: a repetition priming study using high-density electrical mapping. *Neuroimage*, 13(2), 305-313. doi:10.1006/nimg.2000.0684
- Eriksen, B. A., & Eriksen, C. W. (1974). Effects of noise letters upon the identification of a target letter in a nonsearch task. *Perception & psychophysics*, 16(1), 143-149.
- Ettinger, U., Kumari, V., Crawford, T. J., Corr, P. J., Das, M., Zachariah, E., . . . Sharma, T. (2004). Smooth pursuit and antisaccade eye movements in siblings discordant for schizophrenia. *J Psychiatr Res*, 38(2), 177-184. doi:10.1016/s0022-3956(03)00105-5
- Feltner, K. A., & Kiorpes, L. (2010). Global visual processing in macaques studied using Kanizsa illusory shapes. *Visual Neuroscience*, 27(3-4), 131-138. doi:10.1017/S0952523810000088
- Fink, G. R., Halligan, P. W., Marshall, J. C., Frith, C. D., Frackowiak, R. S., & Dolan, R. J. (1997). Neural mechanisms involved in the processing of global and local aspects of hierarchically organized visual stimuli. *Brain*, 120 (Pt 10), 1779-1791. doi:10.1093/brain/120.10.1779

- Fitch, A., Fein, D. A., & Eigsti, I. M. (2015). Detail and gestalt focus in individuals with optimal outcomes from autism spectrum disorders. *J Autism Dev Disord*, 45(6), 1887-1896. doi:10.1007/s10803-014-2347-8
- Foxe, J. J., Doniger, G. M., & Javitt, D. C. (2001). Early visual processing deficits in schizophrenia: impaired P1 generation revealed by high-density electrical mapping. *Neuroreport*, 12(17), 3815-3820. doi:10.1097/00001756-200112040-00043
- Foxe, J. J., Murray, M. M., & Javitt, D. C. (2005). Filling-in in schizophrenia: a high-density electrical mapping and source-analysis investigation of illusory contour processing. *Cereb Cortex*, 15(12), 1914-1927. doi:10.1093/cercor/bhi069
- Foxe, J. J., & Simpson, G. V. (2002). Flow of activation from V1 to frontal cortex in humans. A framework for defining "early" visual processing. *Exp Brain Res*, 142(1), 139-150. doi:10.1007/s00221-001-0906-7
- Freeseaman, L. J., Colombo, J., & Coldren, J. T. (1993). Individual differences in infant visual attention: four-month-olds' discrimination and generalization of global and local stimulus properties. *Child Dev*, 64(4), 1191-1203. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/8404264>
- Frith, U. (1989). A new look at language and communication in autism. *Br J Disord Commun*, 24(2), 123-150. doi:10.3109/13682828909011952
- Frith, U., & Happe, F. (1994). Autism: beyond "theory of mind". *Cognition*, 50(1-3), 115-132. doi:10.1016/0010-0277(94)90024-8
- Gioia, G. A., Isquith, P. K., Guy, S. C., & Kenworthy, L. (2000). *Behavior rating inventory of executive function: BRIEF*: Psychological Assessment Resources Odessa, FL.
- Gotham, K., Pickles, A., & Lord, C. (2009). Standardizing ADOS scores for a measure of severity in autism spectrum disorders. *J Autism Dev Disord*, 39(5), 693-705. doi:10.1007/s10803-008-0674-3
- Groen, W. B., Rommelse, N., de Wit, T., Zwiers, M. P., van Meerendonck, D., van der Gaag, R. J., & Buitelaar, J. K. (2012). Visual scanning in very young children with autism and their unaffected parents. *Autism Res Treat*, 2012, 748467. doi:10.1155/2012/748467
- Guttman, S. E., & Kellman, P. J. (2004). Contour interpolation revealed by a dot localization paradigm. *Vision Res*, 44(15), 1799-1815. doi:10.1016/j.visres.2004.02.008
- Happe, F. (1996). Studying weak central coherence at low levels: children with autism do not succumb to visual illusions. A research note. *J Child Psychol Psychiatry*, 37(7), 873-877. doi:10.1111/j.1469-7610.1996.tb01483.x
- Happe, F. (1999). Autism: cognitive deficit or cognitive style? *Trends Cogn Sci*, 3(6), 216-222. doi:10.1016/s1364-6613(99)01318-2

- Happé, F. (2000). Parts and wholes, meaning and minds: Central coherence and its relation to theory of mind. In S. Baron-Cohen, H. Tager-Flusberg, & D. J. Cohen (Eds.), *Understanding other minds: Perspectives from developmental cognitive neuroscience* (pp. 203–221): Oxford University Press.
- Happe, F., Briskman, J., & Frith, U. (2001). Exploring the cognitive phenotype of autism: weak "central coherence" in parents and siblings of children with autism: I. Experimental tests. *J Child Psychol Psychiatry*, 42(3), 299-307. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/11321199>
- Happe, F., & Frith, U. (2006). The weak coherence account: detail-focused cognitive style in autism spectrum disorders. *J Autism Dev Disord*, 36(1), 5-25. doi:10.1007/s10803-005-0039-0
- Harris, J. J., Schwarzkopf, D. S., Song, C., Bahrami, B., & Rees, G. (2011). Contextual illusions reveal the limit of unconscious visual processing. *Psychol Sci*, 22(3), 399-405. doi:10.1177/0956797611399293
- Hayward, D. A., Fenerci, C., & Ristic, J. (2018). An investigation of global-local processing bias in a large sample of typical individuals varying in autism traits. *Conscious Cogn*, 65, 271-279. doi:10.1016/j.concog.2018.09.002
- Heinze, H. J., Hinrichs, H., Scholz, M., Burchert, W., & Mangun, G. R. (1998). Neural mechanisms of global and local processing. A combined PET and ERP study. *J Cogn Neurosci*, 10(4), 485-498. doi:10.1162/089892998562898
- Hogan-Brown, A. L., Hoedemaker, R. S., Gordon, P. C., & Losh, M. (2014). Eye-voice span during rapid automatized naming: evidence of reduced automaticity in individuals with autism spectrum disorder and their siblings. *J Neurodev Disord*, 6(1), 33. doi:10.1186/1866-1955-6-33
- Hurley, R. S., Losh, M., Parlier, M., Reznick, J. S., & Piven, J. (2007). The broad autism phenotype questionnaire. *J Autism Dev Disord*, 37(9), 1679-1690. doi:10.1007/s10803-006-0299-3
- Jarrold, C., Butler, D. W., Cottington, E. M., & Jimenez, F. (2000). Linking theory of mind and central coherence bias in autism and in the general population. *Dev Psychol*, 36(1), 126-138.
- Jolliffe, T., & Baron-Cohen, S. (1997). Are people with autism and Asperger syndrome faster than normal on the Embedded Figures Test? *J Child Psychol Psychiatry*, 38(5), 527-534. doi:10.1111/j.1469-7610.1997.tb01539.x
- Jolliffe, T., & Baron-Cohen, S. (2000). Linguistic processing in high-functioning adults with autism or Asperger's syndrome. Is global coherence impaired? *Psychol Med*, 30(5), 1169-1187. doi:10.1017/s003329179900241x

- Joseph, R. M., Keehn, B., Connolly, C., Wolfe, J. M., & Horowitz, T. S. (2009). Why is visual search superior in autism spectrum disorder? *Dev Sci*, 12(6), 1083-1096. doi:10.1111/j.1467-7687.2009.00855.x
- Kanizsa, G. (1976). Subjective contours. *Sci Am*, 234(4), 48-52. doi:10.1038/scientificamerican0476-48
- Kanner, L. (1943). Autistic disturbances of affective contact. *Nervous child*, 2(3), 217-250.
- Keane, B. P., Joseph, J., & Silverstein, S. M. (2014). Late, not early, stages of Kanizsa shape perception are compromised in schizophrenia. *Neuropsychologia*, 56, 302-311. doi:10.1016/j.neuropsychologia.2014.02.001
- Keehn, R. J. J., Pueschel, E. B., Gao, Y., Jahedi, A., Alemu, K., Carper, R., . . . Müller, R. A. (2020). Underconnectivity between visual and salience networks and links with sensory abnormalities in autism spectrum disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*.
- Kemner, C., van Ewijk, L., van Engeland, H., & Hooge, I. (2008). Brief report: eye movements during visual search tasks indicate enhanced stimulus discriminability in subjects with PDD. *J Autism Dev Disord*, 38(3), 553-557. doi:10.1007/s10803-007-0406-0
- Kimchi, R. (1992). Primacy of wholistic processing and global/local paradigm: a critical review. *Psychol Bull*, 112(1), 24-38. doi:10.1037/0033-2909.112.1.24
- Klin, A., Jones, W., Schultz, R., Volkmar, F., & Cohen, D. (2002). Visual fixation patterns during viewing of naturalistic social situations as predictors of social competence in individuals with autism. *Arch Gen Psychiatry*, 59(9), 809-816. doi:10.1001/archpsyc.59.9.809
- Koldewyn, K., Jiang, Y. V., Weigelt, S., & Kanwisher, N. (2013). Global/local processing in autism: not a disability, but a disinclination. *J Autism Dev Disord*, 43(10), 2329-2340. doi:10.1007/s10803-013-1777-z
- Kovacs, I. (1996). Gestalten of today: early processing of visual contours and surfaces. *Behav Brain Res*, 82(1), 1-11. doi:10.1016/s0166-4328(97)81103-5
- Lam, K. S., & Aman, M. G. (2007). The Repetitive Behavior Scale-Revised: independent validation in individuals with autism spectrum disorders. *J Autism Dev Disord*, 37(5), 855-866. doi:10.1007/s10803-006-0213-z
- Lamb, M. R., & Robertson, L. C. (1989). Do Response-Time Advantage and Interference Reflect the Order of Processing of Global-Level and Local-Level Information. *Perception & psychophysics*, 46(3), 254-258. doi:Doi 10.3758/Bf03208087
- Landa, R. (2011). *The Pragmatic Rating Scale for School-Age Children (PRS-SA)*. Unpublished manual.

- Landa, R., Piven, J., Wzorek, M. M., Gayle, J. O., Chase, G. A., & Folstein, S. E. (1992). Social language use in parents of autistic individuals. *Psychol Med*, 22(1), 245-254. doi:10.1017/s0033291700032918
- Lee, M., Nayar, K., Maltman, N., Hamburger, D., Martin, G. E., Gordon, P. C., & Losh, M. (2019). Understanding social communication differences in autism spectrum disorder and first-degree relatives: a study of looking and speaking. *J Autism Dev Disord*. doi:10.1007/s10803-019-03969-3
- Lee, T. S., & Nguyen, M. (2001). Dynamics of subjective contour formation in the early visual cortex. *Proc Natl Acad Sci U S A*, 98(4), 1907-1911. doi:10.1073/pnas.031579998
- Lord, C., Rutter, M., DiLavore, P., Risi, S., Gotham, K., & Bishop, S. (2012). Autism diagnostic observation schedule—2nd edition (ADOS-2). *Los Angeles, CA: Western Psychological Corporation*.
- Lord, C., Rutter, M., & Le Couteur, A. (1994). Autism Diagnostic Interview-Revised: a revised version of a diagnostic interview for caregivers of individuals with possible pervasive developmental disorders. *Journal of autism and developmental disorders*, 24(5), 659-685.
- Losh, M. (2011). The broad autism phenotype. In *Autism spectrum disorders*: Oxford University Press.
- Losh, M., Adolphs, R., & Piven, J. (2011). The broad autism phenotype. In *Autism spectrum disorders*: Oxford University Press.
- Losh, M., Adolphs, R., Poe, M. D., Couture, S., Penn, D., Baranek, G. T., & Piven, J. (2009). Neuropsychological profile of autism and the broad autism phenotype. *Arch Gen Psychiatry*, 66(5), 518-526. doi:10.1001/archgenpsychiatry.2009.34
- Losh, M., Childress, D., Lam, K., & Piven, J. (2008). Defining key features of the broad autism phenotype: a comparison across parents of multiple- and single-incidence autism families. *Am J Med Genet B Neuropsychiatr Genet*, 147B(4), 424-433. doi:10.1002/ajmg.b.30612
- Losh, M., & Piven, J. (2007). Social-cognition and the broad autism phenotype: identifying genetically meaningful phenotypes. *J Child Psychol Psychiatry*, 48(1), 105-112. doi:10.1111/j.1469-7610.2006.01594.x
- Maertens, M., & Pollmann, S. (2005). fMRI reveals a common neural substrate of illusory and real contours in V1 after perceptual learning. *J Cogn Neurosci*, 17(10), 1553-1564. doi:10.1162/089892905774597209
- Martin, I., & McDonald, S. (2003). Weak coherence, no theory of mind, or executive dysfunction? Solving the puzzle of pragmatic language disorders. *Brain and Language*, 85(3), 451-466. doi:10.1016/S0093-934x(03)00070-1
- Milne, E., & Scope, A. (2008). Are children with autistic spectrum disorders susceptible to contour illusions? *British Journal of Developmental Psychology*, 26(1), 91-102.

- Minshew, N. J., Goldstein, G., & Siegel, D. J. (1997). Neuropsychologic functioning in autism: profile of a complex information processing disorder. *J Int Neuropsychol Soc*, 3(4), 303-316. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/9260440>
- Mosconi, M. W., Kay, M., D'Cruz, A. M., Guter, S., Kapur, K., Macmillan, C., . . . Sweeney, J. A. (2010). Neurobehavioral abnormalities in first-degree relatives of individuals with autism. *Arch Gen Psychiatry*, 67(8), 830-840. doi:10.1001/archgenpsychiatry.2010.87
- Mottron, L., Burack, J. A., Iarocci, G., Belleville, S., & Enns, J. T. (2003). Locally oriented perception with intact global processing among adolescents with high-functioning autism: evidence from multiple paradigms. *J Child Psychol Psychiatry*, 44(6), 904-913. doi:10.1111/1469-7610.00174
- Mottron, L., Burack, J. A., Stauder, J. E., & Robaey, P. (1999). Perceptual processing among high-functioning persons with autism. *J Child Psychol Psychiatry*, 40(2), 203-211.
- Murray, M. M., Wylie, G. R., Higgins, B. A., Javitt, D. C., Schroeder, C. E., & Foxe, J. J. (2002). The spatiotemporal dynamics of illusory contour processing: combined high-density electrical mapping, source analysis, and functional magnetic resonance imaging. *J Neurosci*, 22(12), 5055-5073.
- Navon, D. (1977). Forest before trees: the precedence effect of global features in visual perception. *Cognitive Psychol.*, 7, 476-484.
- Navon, D. (1983). How many trees does it take to make a forest? *Perception*, 12(3), 239-254. doi:10.1068/p120239
- Nayar, K., Franchak, J., Adolph, K., & Kiorpes, L. (2015). From local to global processing: the development of illusory contour perception. *J Exp Child Psychol*, 131, 38-55. doi:10.1016/j.jecp.2014.11.001
- Nayar, K., Gordon, P. C., Martin, G. E., Hogan, A. L., La Valle, C., McKinney, W., . . . Losh, M. (2018). Links between looking and speaking in autism and first-degree relatives: insights into the expression of genetic liability to autism. *Mol Autism*, 9, 51. doi:10.1186/s13229-018-0233-5
- Nayar, K., Kang, X., Xing, J., Gordon, P. C., Wong, P. C. M., & Losh, M. (2021). A cross-cultural study showing deficits in gaze-language coordination during rapid automatized naming among individuals with ASD. *Sci Rep*, 11(1), 13401. doi:10.1038/s41598-021-91911-y
- Nayar, K., Voyles, A. C., Kiorpes, L., & Di Martino, A. (2017). Global and local visual processing in autism: An objective assessment approach. *Autism Res*, 10(8), 1392-1404. doi:10.1002/aur.1782
- Neufeld, J., Hagstrom, A., Van't Westeinde, A., Lundin, K., Cauvet, E., Willfors, C., . . . Bolte, S. (2020). Global and local visual processing in autism - a co-twin-control study. *J Child Psychol Psychiatry*, 61(4), 470-479. doi:10.1111/jcpp.13120

- O'Riordan, M., & Plaisted, K. (2001). Enhanced discrimination in autism. *Q J Exp Psychol A*, 54(4), 961-979. doi:10.1080/713756000
- Pellicano, E., & Burr, D. (2012). When the world becomes 'too real': a Bayesian explanation of autistic perception. *Trends Cogn Sci*, 16(10), 504-510. doi:10.1016/j.tics.2012.08.009
- Pellicano, E., Maybery, M., Durkin, K., & Maley, A. (2006). Multiple cognitive capabilities/deficits in children with an autism spectrum disorder: "weak" central coherence and its relationship to theory of mind and executive control. *Dev Psychopathol*, 18(1), 77-98. doi:10.1017/S0954579406060056
- Piven, J., Gayle, J., Chase, G. A., Fink, B., Landa, R., Wzorek, M. M., & Folstein, S. E. (1990). A family history study of neuropsychiatric disorders in the adult siblings of autistic individuals. *J Am Acad Child Adolesc Psychiatry*, 29(2), 177-183. doi:10.1097/00004583-199003000-00004
- Piven, J., Palmer, P., Landa, R., Santangelo, S., Jacobi, D., & Childress, D. (1997). Personality and language characteristics in parents from multiple-incidence autism families. *Am J Med Genet*, 74(4), 398-411.
- Piven, J., Wzorek, M., Landa, R., Lainhart, J., Bolton, P., Chase, G. A., & Folstein, S. (1994). Personality characteristics of the parents of autistic individuals. *Psychol Med*, 24(3), 783-795. doi:10.1017/s0033291700027938
- Plaisted, K., Swettenham, J., & Rees, L. (1999). Children with autism show local precedence in a divided attention task and global precedence in a selective attention task. *J Child Psychol Psychiatry*, 40(5), 733-742. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/10433407>
- Rinehart, N. J., Bradshaw, J. L., Moss, S. A., Brereton, A. V., & Tonge, B. J. (2000). Atypical interference of local detail on global processing in high-functioning autism and Asperger's disorder. *J Child Psychol Psychiatry*, 41(6), 769-778. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/11039689>
- Ringach, D. L., & Shapley, R. (1996). Spatial and temporal properties of illusory contours and amodal boundary completion. *Vision Res*, 36(19), 3037-3050. doi:10.1016/0042-6989(96)00062-4
- Rommelse, N., Langerak, I., van der Meer, J., de Bruijn, Y., Staal, W., Oerlemans, A., & Buitelaar, J. (2015). Intelligence may moderate the cognitive profile of patients with ASD. *PLoS One*, 10(10), e0138698. doi:10.1371/journal.pone.0138698
- Ropar, D., & Mitchell, P. (1999). Are individuals with autism and Asperger's syndrome susceptible to visual illusions? *J Child Psychol Psychiatry*, 40(8), 1283-1293. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/10604406>

- Ropar, D., & Mitchell, P. (2001). Susceptibility to illusions and performance on visuospatial tasks in individuals with autism. *J Child Psychol Psychiatry*, 42(4), 539-549. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/11383970>
- Roth, R. M., Lance, C. E., Isquith, P. K., Fischer, A. S., & Giancola, P. R. (2013). Confirmatory factor analysis of the Behavior Rating Inventory of Executive Function-Adult version in healthy adults and application to attention-deficit/hyperactivity disorder. *Arch Clin Neuropsychol*, 28(5), 425-434. doi:10.1093/arclin/act031
- Sary, G., Koteles, K., Kaposvari, P., Lenti, L., Csifcsak, G., Franko, E., . . . Tompa, T. (2008). The representation of Kanizsa illusory contours in the monkey inferior temporal cortex. *Eur J Neurosci*, 28(10), 2137-2146. doi:10.1111/j.1460-9568.2008.06499.x
- Sasson, N. J., Turner-Brown, L. M., Holtzclaw, T. N., Lam, K. S., & Bodfish, J. W. (2008). Children with autism demonstrate circumscribed attention during passive viewing of complex social and nonsocial picture arrays. *Autism Res*, 1(1), 31-42. doi:10.1002/aur.4
- Scholte, H. S., Jolij, J., Fahrenfort, J. J., & Lamme, V. A. (2008). Feedforward and recurrent processing in scene segmentation: electroencephalography and functional magnetic resonance imaging. *J Cogn Neurosci*, 20(11), 2097-2109. doi:10.1162/jocn.2008.20142
- Shah, A., & Frith, U. (1983). An islet of ability in autistic children: a research note. *J Child Psychol Psychiatry*, 24(4), 613-620. doi:10.1111/j.1469-7610.1983.tb00137.x
- Shah, A., & Frith, U. (1993). Why do autistic individuals show superior performance on the block design task? *J Child Psychol Psychiatry*, 34(8), 1351-1364. doi:10.1111/j.1469-7610.1993.tb02095.x
- Simmons, D. R., & Todorova, G. K. (2018). Local versus global processing in autism: special section editorial. *J Autism Dev Disord*, 48(4), 1338-1340. doi:10.1007/s10803-017-3452-2
- Stanley, D. A., & Rubin, N. (2003). fMRI activation in response to illusory contours and salient regions in the human lateral occipital complex. *Neuron*, 37(2), 323-331. doi:10.1016/s0896-6273(02)01148-0
- Stroganova, T. A., Nygren, G., Tsetlin, M. M., Posikera, I. N., Gillberg, C., Elam, M., & Orekhova, E. V. (2007). Abnormal EEG lateralization in boys with autism. *Clin Neurophysiol*, 118(8), 1842-1854. doi:10.1016/j.clinph.2007.05.005
- Stroganova, T. A., Orekhova, E. V., Prokofyev, A. O., Posikera, I. N., Morozov, A. A., Obukhov, Y. V., & Morozov, V. A. (2007). Inverted event-related potentials response to illusory contour in boys with autism. *Neuroreport*, 18(9), 931-935. doi:10.1097/WNR.0b013e32811e151b
- Stroganova, T. A., Orekhova, E. V., Prokofyev, A. O., Tsetlin, M. M., Gratchev, V. V., Morozov, A. A., & Obukhov, Y. V. (2012). High-frequency oscillatory response to illusory contour

- in typically developing boys and boys with autism spectrum disorders. *Cortex*, 48(6), 701-717. doi:10.1016/j.cortex.2011.02.016
- Swanson, M. R., Serlin, G. C., & Siller, M. (2013). Broad autism phenotype in typically developing children predicts performance on an eye-tracking measure of joint attention. *J Autism Dev Disord*, 43(3), 707-718. doi:10.1007/s10803-012-1616-7
- Team, R. C. (2019). R: A Language and Environment for Statistical Computing. In R. F. f. S. Computing (Ed.). Vienna, Austria.
- Van der Hallen, R., Evers, K., Brewaeys, K., Van den Noortgate, W., & Wagemans, J. (2015). Global processing takes time: a meta-analysis on local-global visual processing in ASD. *Psychol Bull*, 141(3), 549-573. doi:10.1037/bul0000004
- Van Eylen, L., Boets, B., Cosemans, N., Peeters, H., Steyaert, J., Wagemans, J., & Noens, I. (2017). Executive functioning and local-global visual processing: candidate endophenotypes for autism spectrum disorder? *J Child Psychol Psychiatry*, 58(3), 258-269. doi:10.1111/jcpp.12637
- Van Eylen, L., Boets, B., Steyaert, J., Wagemans, J., & Noens, I. (2018). Local and Global Visual Processing in Autism Spectrum Disorders: Influence of Task and Sample Characteristics and Relation to Symptom Severity. *J Autism Dev Disord*, 48(4), 1359-1381. doi:10.1007/s10803-015-2526-2
- Wechsler, D. (1991). Manual for the Wechsler Intelligence Scale for Children—third edition. *San Antonio, TX: The Psychological Corporation*.
- Wechsler, D. (1999). *Wechsler Abbreviated Scale of Intelligence (WASI)*. New York, NY: The Psychological Corporation: Harcourt Brace & Company.
- Weintraub, S., Dikmen, S. S., Heaton, R. K., Tulsky, D. S., Zelazo, P. D., Bauer, P. J., . . . Gershon, R. C. (2013). Cognition assessment using the NIH Toolbox. *Neurology*, 80(11 Suppl 3), S54-64. doi:10.1212/WNL.0b013e3182872ded
- Wokke, M. E., Vandenbroucke, A. R., Scholte, H. S., & Lamme, V. A. (2013). Confuse your illusion: feedback to early visual cortex contributes to perceptual completion. *Psychol Sci*, 24(1), 63-71. doi:10.1177/0956797612449175
- Yucel, G. H., Belger, A., Bizzell, J., Parlier, M., Adolphs, R., & Piven, J. (2015). Abnormal neural activation to faces in the parents of children with autism. *Cereb Cortex*, 25(12), 4653-4666. doi:10.1093/cercor/bhu147
- Zagermann, J., Pfeil, U., & Reiterer, H. (2018). *Studying eye movements as a basis for measuring cognitive load*. Paper presented at the Extended Abstracts of the 2018 CHI Conference on Human Factors in Computing Systems.