The Parents are Not Alright: A Call for Parental Mental Health Screening During the COVID-19 Pandemic

Tara S. Peris, Ph.D.^a & Jill Ehrenreich-May, Ph.D.^b

Affiliations: ^aUniversity of California, Los Angeles, Los Angeles, California, and
^bUniversity of Miami, Miami, Florida

Address correspondence to: Tara Peris, 760 Westwood Plaza, Room 67-463, Los Angeles, CA, 90024; tperis@mednet.ucla.edu

Short title: Screening Caregivers for Emotional Distress

Conflict of Interest Disclosures: Drs. Peris and Ehrenreich-May have no conflicts of interest relevant to this article to disclose.

Funding/Support: No funding was secured for this study.

Abbreviations: None.

Word Count: 1272

Contributors' Statement Page

Dr. Peris and Dr. Ehrenreich-May drafted the manuscript, critically reviewed the submission content, and reviewed and revised for publication.

Each author approved the final manuscript as submitted and agrees to be accountable for all aspects of the work.

Acknowledgements

The authors wish to thank Drs. Eraka Bath and Shawn CT Jones for their willingness to review and comment on this work.

Nearly a year into the pandemic, conversations about the impact of COVID-19 on children and families have shifted. Initial advice for parents stressed topics such as how to talk about the pandemic with children or cope with illness-related distress. They now focus on youth adjustment to a heavily disrupted school year and on strategies for building long-term resilience. Although these conversations often center on youth adjustment, they have—at last – started to consider the wellbeing of parents (and other caregivers) as well. This shift in focus is crucial given the enormous challenges that parents face right now and the direct links between their wellbeing and that of their children. What continue to lag, even well into the pandemic, however, are workable solutions for addressing parent mental health. While we applaud the renewed focus on parenting stress and wellbeing, we remain deeply concerned by the absence of a plan for intervening.

Early data suggest that in the wake of the pandemic, rates of psychiatric illness have surged, with nearly a quarter of adults meeting key criteria for major depressive disorder and nearly a third meeting such for generalized anxiety disorder.¹ Data from both past quarantines and the current pandemic suggest that primary caregivers are at even greater risk than those in the general population.^{2,3} There are also new data on COVID-19 adjustment suggesting that parents find themselves less emotionally supported than they were prior to its onset.⁴ The risks posed by parental psychopathology are not trivial during the best of times: they shape numerous aspects of the home environment and elevate risk of youth psychopathology considerably, even when genetic factors are considered. During times of high stress and uncertainty, it's potentially worse. For example, during the H1N1 outbreak caregiver fear and experience of threat impacted

child anxiety directly.⁵ For youth who need psychiatric treatment during the pandemic, there is evidence that parental psychopathology attenuates outcomes and that it is not adequately addressed by child treatment.⁶

Even among parents who do not suffer from diagnosable mental health problems, the toll of these extraordinary circumstances should not be overlooked. Many parents are caring not only for their children, but for aging parents as well. Some are grieving loved ones lost. Many are coping with job loss that has precipitated financial instability and caused food and housing insecurity. Juggling the demands of work, virtual schooling, and restless children amidst intense isolation and poor social support do not make for better parenting. They create stress, irritability, and guilt that will lead to greater mental health problems, chaotic home environments, and the potential for harsh parenting. For parents of youth with special needs, these challenges are particularly pronounced. All of this comes amidst reduced contact with schools and social services agencies for families in need.

These challenges exist for all parents, but they are the tip of the iceberg for those disproportionately affected by the pandemic. In Black and Latinx communities with markedly higher rates of COVID-19 morbidity and mortality, limited access to care, and systemic barriers to quality care that lead to disparate health outcomes, the pandemic is likely to bring far greater parenting stress. These challenges are amplified further by our national reckoning with generations of racism, a long overdue process that may bring additional stress and exhaustion for Black families as a spotlight is directed at deeply painful and personal issues.

If the stressors and their consequences are well known, what then do we do? The first step: seize every opportunity to evaluate and promote parent wellbeing, *even when it falls outside of traditional models of care*. Universal screening for youth mental health problems has gained traction over the past decade, and it has a demonstrated record of success for improving early identification and access to care among youth with common conditions such as ADHD and depression. Many practices now administer measures such as the Pediatric Symptom Checklist-17 as part of routine visits. Why not do the same for parents? Brief inventories exist to capture parenting stress and adjustment, and they provide not only information about parent mental health, but critical context for understanding the ecosystem in which pediatric patients live. They are a valuable treatment-planning tool and can be used to guide discussions on how to promote wellbeing. When these measures prove burdensome or unfeasible, routinely asking parents “How are you holding up?” or expressing empathy about the strain of their multiple caregiving roles can create valuable openings to addressing parent mental health. Such questions can be posed in the context of child psychiatry visits or during adult psychiatric visits where the focus may not typically be on parenting. These family-based approaches to screening and triage are not new. Strategies such as the Vermont Family Treatment Model have well-documented success⁷, but they bear further translation, particularly right now.

Amidst a pandemic that has already consumed the better part of a year and will be with us for some time, we already know enough about the impact on parents. The impact on children is not far behind. To offset the damage, we encourage a broader screening approach that captures parents and caregivers during any point of contact with psychiatric

care providers whether it be child- or parent-directed services, across settings (See Fig. 1). We recommend widely available, reliable and valid tools for assessment of caregiver distress, parenting behavior and/or symptomatology. Many are free and easily administered electronically or during in-person visits. For those who screen “positive”, there are now brief and “single-session interventions”^{8,9} that target key mechanisms of change in anxiety and depression, many in telehealth and compelling digital formats that are low burden and cost effective. They can be purchased by health systems or quickly deployed on a more individual or small group level by psychiatric or psychological providers and broader community service agencies to prevent further symptoms. These brief treatments can provide meaningful support, often using evidence-based treatment strategies found in lengthier psychotherapy protocols. For those whose distress persists, psychological services nationwide, including Medicaid-funded services, are now available via telehealth, phone or both. Given changes to HIPAA privacy and billing coverage laws, they may be accessed by providers outside an individual’s immediate local community. Finally, we strongly recommend that clinical practices develop resource lists that can be shared with parents as part of routine child visits and link them to bibliotherapy and other low cost, low burden intervention options for their own care. For those who need more, referral lists for practitioners who provide evidence-based psychotherapy can be shared. A list of primary care and psychiatric providers that are comfortable with prescribing effective pharmacotherapy is also vital. Finally, on-going screening of caregivers by pediatric and family or internal medicine providers throughout this pandemic may be needed to meet the mental health needs of caregivers. As with all universal screening efforts, there are bound to be implementation challenges. Time is

limited and virtual sessions complicate data collection. Nonetheless, advanced planning and even informal probes can pave the way for brief systematic assessment that is also done virtually. Extant data on pandemic mental health outcomes already tells us that this would be time well spent.

Although the full mental health impact of these tumultuous times remains to be seen, we are clear that rates of anxiety and depression were rising well before these challenges arrived. Without quick and responsive deployment of effective screening, prevention, and evidence-based intervention, we risk leaving rising numbers of children and parents in increasingly stressful environments ripe to leave a pandemic legacy of toxic stress, mental illness, and potentially, of abuse and neglect. We can act now with proven prevention measures to ensure the best possible outcomes for children and families.

References

1. U.S. Census Bureau, Data reported from May 7-12, 2020
2. Sprang G, Silman M. Posttraumatic Stress Disorder in Parents and Youth After Health-Related Disasters. *Disaster Med Public Health Prep.* 2013; 7(1):105-110.
3. Lei S, Jiang F, Su W, et al. *Clinical characteristics and outcomes of patients undergoing surgeries during the incubation period of COVID-19 infection.* April 2020.
4. Center for Translational Neuroscience. *RAPID-EC Project.* 2020. Retrieved from <https://medium.com/rapid-ec-project>
5. Remmerswaal D, Muris P. Children's Fear Reactions to the 2009 Swine Flu Pandemic: The Role of Threat Reinforcement as Provided by Parents. *Journal of Anxiety Disorders.* 17 Sept 2010; 25:444-449.
6. Maliken AC, Katz LF. Exploring the impact of parental psychopathology and emotion regulation on evidence-based parenting interventions: a transdiagnostic approach to improving treatment effectiveness. *Clin Child Fam Psychol Rev.* 2013;16(2):173-186.
7. Hudziak J, & Ivanova MY (2016). The Vermont family based approach: Family based health promotion, illness prevention, and intervention. *Child Adolesce Psychiatr Clin N Am*, 25, 167-78.
8. Schleider, JL, Weisz, JR (2017). Little treatments, promising effects? Meta-analysis of single-session interventions for youth psychiatric problems. *Journal of the American Academy of Child and Adolescent Psychiatry*, 56, 2, 107-115.
9. Schleider, JL, Dobias, ML, & Pati, S (2020). Promoting treatment access following pediatric primary care depression screening: Randomized trial of web-based, single-session interventions for parents and youths. *Journal of the American Academy of Child and Adolescent Psychiatry.*

FIGURE 1. SCREENING, PREVENTION AND INTERVENTION FOR CAREGIVERS OF YOUTH DURING COVID-19