Sleep and Next-Day Negative Affect and Suicidal Ideation in Borderline Personality Disorder

Aleksandra Kaurin^a, Garrett Hisler^b, Alexandre Y. Dombrovski^b, Michael N. Hallquist^c & Aidan G.C. Wright^a

^aDepartment of Psychology, University of Pittsburgh

^bDepartment of Psychiatry, University of Pittsburgh School of Medicine

^cDepartment of Psychology and Neuroscience, University of North Carolina, Chapel Hill

© 2021, American Psychological Association. This paper is not the copy of record and may not exactly replicate the final, authoritative version of the article. Please do not copy or cite without authors' permission. The final article will be available, upon publication, via its DOI: 10.1037/per0000496

This study was supported by the National Institute of Mental Health (R01 MH048463).

The authors certify that they have no affiliations with or involvement in any organization or entity with any financial interest, or non-financial interest in the subject matter or materials discussed in this manuscript. We are grateful to the patients who took part in this study and to the clinical and research teams who made this work possible.

Correspondence concerning this article should be addressed to Aleksandra Kaurin, Department of Psychology, University of Pittsburgh, 3213 Sennott Square, 210 S. Bouquet St., Pittsburgh, PA, 15260. E-Mail: kaurinaleksandra@pitt.edu

Abstract

Sleep disturbance is associated with elevated suicidal ideation and negative affect. To date, however, no study has investigated the temporal relationship between sleep and suicidality among those diagnosed with borderline personality disorder (BPD). This preregistered (<https://osf.io/4vugk>) study tested whether nightly sleep (self-reported sleep duration, sleep onset latency, and subjective sleep quality) represents a (within-person) short-term risk factor for affective dysregulation and increases in suicide risk from day-to-day, as well as whether between-person differences in sleep, negative affect, and suicidality were associated. We used a 21-day ecological momentary assessment protocol in a sample of 153 people diagnosed with BPD, 105 of which had a history of serious suicide attempts, and 52 healthy controls ($N=4086$ days). We found a within-person association between worse subjective sleep quality and greater next day negative affect. At the between-person level, we found positive relationships between sleep latency and suicidal ideation, and a negative association between subjective sleep quality and negative affect. BPD severity did not significantly moderate the strength of any within-person associations, although BPD was positively associated with average levels of suicidal ideation, sleep latency, and negative affect, and negatively related to subjective sleep quality. These findings suggest that the association of sleep with suicidal ideation and BPD exists largely at the between-persons rather than the within-person level. Disturbed sleep, therefore, seems to largely coincide, rather than specifically contribute to the exacerbation of suicidal crises in BPD.

Key words: suicidal ideation; ecological momentary assessment; borderline personality disorder; multilevel structural equation modelling; sleep

Sleep and Next-Day Negative Affect and Suicidal Ideation in Borderline Personality Disorder

Research suggests that the association between sleep disturbances and suicidal ideation may be potentiated by Borderline Personality Disorder (BPD; Winsper & Tang, 2014; Winsper et al., 2017). Previous work, however, has returned contradictory results (e.g., Oltmanns & Oltmanns, 2015), and the temporal, within-person link of sleep and suicidal ideation in BPD has received limited clinical and research attention (Winsper & Tang, 2014; *see* Liu et al. (2020) for a recent systematic review and meta-analysis of longitudinal studies). In this study, we address this gap in the literature by asking: Are sleep disturbances contributing factors to exacerbating short-term suicide risk, or an epiphenomenal aspect of BPD? Our analyses are based on a preregistered study protocol (<https://osf.io/4vugk>), and a 21-day ambulatory assessment procedure, in a carefully sampled population of suicide attempters and non-attempters diagnosed with BPD.

Meta-analytic (Winsper et al., 2017), and narrative reviews (Oltmanns & Oltmanns, 2015) present strong evidence that BPD is linked to sleep disturbance (Oltmanns, 2019; Winsper et al., 2017). Compared to healthy controls, people diagnosed with BPD experience shorter sleep duration, worse sleep efficiency, and reduced REM latency (e.g., Akiskal et al., 1985; McNamara et al., 1984; Reynolds et al., 1985; Simor & Horváth, 2013). Yet, inconsistent results within this set of studies make it difficult to articulate clear conclusions regarding the relationship of BPD and individual sleep parameters (Hafizi, 2013; Oltmanns & Oltmanns, 2015; Winsper et al., 2017). These discrepancies could potentially be attributed to the fact that previous related research has been methodologically diverse and conducted in small samples of phenotypically heterogeneous groups of individuals with BPD (Harty et al., 2010; Winsper et al., 2017).

One reason why it is important to understand sleep in BPD is that suicidal ideation, suicidal attempts, and even death by suicide have been linked to a variety of indicators of poor sleep, including insomnia, nightmares, poor subjective sleep quality, short self-reported sleep duration or low sleep efficiency (Bromundt et al., 2013; Grove et al., 2020; Hafizi, 2013; Oltmanns, Weinstein, & Oltmanns, 2014; Sansone, Edwards, & Forbis, 2010; Selby, 2013; Wall et al, 2020).

Conventionally, poor sleep has been viewed to be reflective of psychiatric burden, rather than representing a stable risk factor of psychopathology. This view, however, is amended by findings suggesting that sleep disturbance may *uniquely* contribute to the exacerbation of psychopathological symptoms (Tkachenko et al., 2014; Nutt et al., 2008), which is based on the notion that the successful treatment of primary psychopathology does not automatically seem to improve sleep problems (Thase et al. 2002; Mouchabac et al. 2003; Carney et al. 2007). Some cross-sectional evidence also suggests that elevated insomnia symptoms may account for the link between BPD traits and an increased risk for suicide (DeShong & Tucker, 2019).

Relatedly, a recent set of findings implies that sleep disruption potentially charts the progression of BPD symptom severity and suicidality by impairing emotional functioning (Balestrieri et al., 2006; Fleischer et al., 2012; Grove et al., 2017; Lereya et al., 2017; Morales-Muñoz et al., 2020; Sjostrom et al., 2009; Wojnar et al., 2009; Wall et al., 2020). On a conceptual level, the Emotional Cascade Model (Selby & Joiner, 2009) provides one plausible explanatory framework for this relationship. The model suggests that upsetting emotional experiences may be potentiated via poor affect regulatory skills that, in turn, may exacerbate in the context of poor sleep in those with a BPD diagnosis. This positive feedback loop contributes to the escalation of negative affect, which then results in exceptionally aversive emotional states

that are met by impulsive and self-harming behaviors in attempts to soothe those affective experiences. Therefore, understanding the clinical significance of sleep for affective dysregulation is particularly relevant to BPD, because cumulative evidence implies that the within-person association between negative affective states and the severity of suicidal ideation is particularly pronounced in BPD (Kaurin et al., 2020; Links et al., 2007; Mou et al., 2018).

What cannot be ascertained from previous, largely cross-sectional studies, however, is whether sleep problems precede, thus exacerbate affective dysregulation and suicidal ideation in BPD, or whether sleep problems largely coincide with BPD. Moreover, correlates obtained from cross-sectional designs may further be driven by individual differences or comorbidity rather than dynamical within-person processes, and cannot be ruled out as concomitants to or epiphenomena of suicidal ideation (Liu et al., 2020). To disambiguate both possibilities, longitudinal studies are needed that are designed to characterize temporal aspects of the relationship between sleep and suicidality (Bernert et al., 2015; Littlewood et al., 2017), both of which fluctuate over time and may influence each other (Ben-Zeev et al., 2012; Kleiman et al., 2017; Lemola et al. 2013; Vallieres et al. 2005).

A few longitudinal studies have sought to address this issue, but none of them in BPD samples in particular. For instance, Ribeiro et al. (2012) found that insomnia predicted greater suicidal ideation one month later in a sample of young military recruits, but not vice versa. These findings were replicated by Zuromski et al. (2017) in an online sample of individuals endorsing lifetime experience of suicidal behavior. Additionally, a one week-long ecological momentary assessment study reported that subjective and actigraphic sleep duration as well as subjective sleep quality predicted next day ideation (Littlewood et al., 2019). Again, suicidal ideation did not predict next night sleep. Finally, a unidirectional relationship was reported in which

nightmares significantly predicted post-sleep self-harm via increased daytime negative affect in a sample of university students (Hochard et al. 2015).

Although it appears from this brief review that disturbed sleep is prospectively linked to elevated levels of suicide risk, it should be noted that the majority of studies assessed sleep disturbance as a long-term risk factor (Liu et al., 2020). Thus, these studies generally provide limited insight into the temporal association between sleep, suicidal ideation, and negative affect, and do not permit disentangling individual difference level associations from the estimates of temporal associations (e.g., person-centering day-level predictors; Bolger & Laurenceau, 2013). In fact, a recent meta-analysis concluded that sleep disturbance prospectively predicts suicide outcomes, but more work is needed to examine sleep as a short term risk factor and to differentiate the associations of acute from chronic sleep disturbance (Liu et al., 2020). Therefore, it remains unclear to what degree disrupted sleep predicts short-term suicide risk. Narrowing this gap is of substantial clinical importance, because clinical risk assessments are focused on determining if individuals may engage in suicidal behavior in the near future (Liu et al., 2020).

To this end, the present study expands upon prior research by disentangling, and then separately estimating the day-level and person-level associations among sleep, negative affect, and suicidal ideation. Such methodological distinction is key to the question whether poor sleep largely coincides with BPD, or significantly contributes to the short-term exacerbation of BPD symptoms (including negative affectivity and suicidality) from day to day. As such, we aim to contribute to a better understanding of whether temporal changes in sleep predict short-term changes of suicide risk, and whether that relationship is accounted for by negative affect.

Preregistered Analyses

Although previous research has found sleep disturbance to contribute to elevated negative affect, and suicidal ideation, no study has investigated the temporal relationship between sleep and suicidality in BPD, whether negative affect accounts for this relationship, nor directly examined the separate day-level and person-level associations. The current preregistered (<https://osf.io/4vugk>) study responds to a recent call for more methodologically rigorous studies of the sleep-suicidal ideation relationship (Bernert et al., 2015; Littlewood et al., 2017; Liu et al., 2020) by investigating the temporal relationships among sleep, negative affect, and suicidal ideation using a 21-day ecological momentary assessment protocol in a sample of 142 people diagnosed with BPD, 98 of which had a history of serious suicide attempts, and 49 healthy controls. This sampling approach increases the likelihood of participants reporting suicidal ideation during the study period, because past suicide attempts represent the most reliable predictors of future suicide attempts (O'Connor & Nock, 2014). Moreover, the design of this study allows to improve limitations in the suicide literature by utilizing a brief follow-up period and focusing on sleep as a less studied short-term risk factor for suicidal ideation in BPD.

Based on previous work reviewed above, we set out to disambiguate whether sleep problems antecede and potentially aggravate short-term suicidal ideation in BPD, or whether disturbed sleep and suicidal ideation largely coincide with BPD. We tested a series of hypotheses related to whether nightly sleep predicted next-day suicidal ideation or negative affect, and whether BPD severity moderated the dynamic links among those variables as they fluctuate, within-person, from day to day. Specifically, our hypotheses were:

Tests of Within-Person Associations

H1a: Worse nightly sleep would predict greater next-day suicidal ideation.

H1b: Worse nightly sleep would predict greater next-day negative affect.

H1c: The within-person association between worse sleep and next-day suicidal ideation would be accounted for by the next-day negative affect.

Cross-Level Interaction Effects

H2a: BPD features would magnify the within-person association between sleep disturbance and suicidal ideation.

H2b: BPD features would magnify the within-person association between sleep disturbance and negative affect.

Our hypotheses are presented in simplified form for clarity. Estimated models included covariates (i.e., between-level: age, gender; within-level: day in study, and weekday vs. weekend). Sleep onset latency, shorter sleep duration, and worse subjective quality were examined each night and models were conducted separately for each sleep parameter to avoid multicollinearity. Of note, we have tested the association between negative affect and suicidality at both, the within- and between-person level in previous analyses. Results from these analyses revealed that negative affect significantly predicts daily levels of suicidality as operationalized in the models below (Kaurin et al., 2020).

Recent meta-analytic evidence suggests that the link between disturbed sleep and BPD features is likely independent of comorbid depression (Winsper et al., 2017). However, given the importance of persistent negative affectivity for the prediction of suicidal behavior (Zanarini et al., 2016), we exploratorily probed the robustness of our analyses by controlling for current depressive symptoms at the between-person level (osf.io/cxy9b).

Methods

All study procedures were approved by the Institutional Review Boards of the University of Pittsburgh (STUDY19050210), and all data, and analyses have been made publicly available at the OSF.

Sample. Participants were drawn from a longitudinal study, and were recruited from in- and outpatient clinics as well as from the nearby community by advertisement. At enrollment, participants had to be between 18 and 45 years and discharged from any inpatient treatment setting. Exclusion criteria included a lifetime diagnosis of any psychotic or bipolar disorder, clinical evidence of organic brain disease, physical disorders or treatments with known psychiatric consequence (e.g., lupus, steroids), and $IQ < 70$ measured by WAIS. A more detailed description of the parent study protocol has been reported elsewhere (Soloff et al., 2017). The sample comprised 153 individuals diagnosed with BPD and 52 healthy control participants (mean age: 33.64, $SD = 9.50$; range 20-60 years; 80% female; 74% Caucasian, 12% African American, 5% Asian, 5% Multi-Race, 4% Pacific Islander or non-specified). Of the 153 BPD patients, $n=35$ had a history of high-lethality, $n=70$ had past low-lethality past suicide attempts, and $n=48$ reported no past suicide attempts. As preregistered, to increase the consistency of our EMA measurements, participants with less than 5 days of sleep were excluded from data analyses. Moreover, data on bed and waketime variables were excluded from analyses if identified as an outlier at 4 standard deviations from the mean. This resulted in a final sample size of $N=201$ participants, which did not significantly differ from the original sample in key outcomes.

The diagnosis of BPD was based on the International Personality Disorder Examination (IPDE; Loranger et al., 1996), and participants had to meet probable or definite criteria for a lifetime diagnosis. Participants diagnosed with BPD were further divided into two groups based

on history of reported suicide attempts. Suicide attempters were required to have clinically significant attempts, defined by a score of >1 on the Beck Lethality Scale (Beck, Beck, & Kovacs, 1975). For participants with multiple attempts, data for the highest-lethality attempt were used.

Participants of the healthy control group and were required to have no lifetime Axis I or II diagnoses determined by the Structured Clinical Interview for DSM-IV or IPDE. As preregistered, participants from the three groups were merged to represent a range of symptom severity, which was further specified with the Personality Assessment Inventory-Borderline Scale (PAI-BOR; Morey, 1991), which demonstrated excellent ($\alpha=.81$) reliability in our sample. The PAI-BOR contains four subscales that tap core features of BPD: affective instability, identity problems, negative relationships, and self-harm. The self-harm subscale taps impulsivity and impulsive behaviors more broadly and is not limited to self-harm behaviors.

Momentary Assessments. Participants completed a 21-day EMA protocol within pre-defined time windows, using the MetricWire smartphone application, which randomly reminded them to complete surveys via push notifications approximately six times per day. This resulted in $N=4086$ days (based on $N=24916$ momentary assessments), with an average number of 19.9 days per participant, ranging from 1-22 days overall. Based on data preregistered exclusion procedures detailed above, the final sample comprised $N=4076$ days and an average number of 20 days per participant.

Each morning, participants reported on last night's sleep characteristics. Specifically, they reported what time they went to bed the prior night, how long it took them to fall asleep (i.e., sleep onset latency), what time they woke up, and the quality of the prior night sleep (i.e., subjective sleep quality).

Sleep duration was calculated from bedtime, waketime, and sleep onset latency reports. First, bedtime and waketime reports were checked for reporting errors (e.g., participants reporting a bedtime of 10:00 instead of 22:00) and either transformed to their appropriate time or were removed from the data. After addressing potential bed and waketime reporting errors, sleep duration was calculated by the difference, in hours, between self-reported waketime and bedtime.

In order to calculate sleep duration, we added 24 hours to the waketime and subtracted bedtime from the transformed waketime variable. To illustrate, if a participant reported a bedtime of 23:00 and a waketime of 8:00, the waketime of 8:00 would be transformed to 32:00 by adding 24 hours. Then, the bedtime of 23:00 was subtracted from 32:00 to calculate a sleep duration of 9 hours.

Sleep onset latency was reported on a scale in which each response was a 15 chunk of time (e.g., 1 = “under 15 minutes”, 2 = “15-30 minutes”). This sleep onset latency scale ranged from “under 15 minutes” to “1 hour 45 minutes to 2 hours.” Sleep onset latency was recoded so that participant responses reflected the mean of the selected response option (e.g., responses of between 15-30 minutes were recoded to equal 22.5 minutes). Note that the 15-minute interval of 1 hour 30 minutes to 1 hour 45 minutes was erroneously left out of the scale during data collection. Subjective sleep quality was measured with the question “Last night my sleep quality was...”, based on a five-point rating scale, ranging from 1 (“very poor”) 5 (“very good”).

Participants reported how strongly they experienced negative (i.e., nervous, sad, irritated, angry; $\omega_{\text{within-person}} = .81$; $\omega_{\text{between-person}} = .91$) derived from the Positive and Negative Affect Schedule (Watson et al., 1988), on a scale from 0 (“Not at All”) to 100 (“Extremely”) for each adjective.

Suicidal ideation was assessed with six dichotomous items (1=yes, 0=no; range: 0-5.4) derived from the suicidal ideation subscale of the Columbia-Suicide Severity Rating Scale (C-

SSRS; Posner et al., 2008). All items were used to form a daily mean score of suicidal intent severity, and read as follows: “Have you wished you were dead or wished you could go to sleep and not wake up?”, “Have you actually had any thoughts of killing yourself?”, “Have you been thinking about how you might do this?”, “Have you had these thoughts and had some intention of acting on them?”, “Have you started to work out or worked out the details of how to kill yourself?”, “Do you intend to carry out this plan?” ($\omega_{\text{within}} = .67$; $\omega_{\text{between}} = .83$). The examination of intraclass correlation returned that approximately 60% of the variability in ratings of suicidal ideation was explained by between-person differences ($\text{ICC} = .601$).

On 11.4% of days participants reported suicidal ideation at least once during the day. More specifically, participants in the BPD group endorsed suicidal ideation on 6.2%, and suicide attempters, on 19.5% of all study days. Among the healthy controls, one participant endorsed a single episode of suicidal ideation.

Data Analysis. As preregistered, multi-level structural equation models (MSEMs) were used to test our hypotheses of the links between daily sleep, affect and suicidal ideation. All models were estimated in Mplus (version 8.4; Muthén & Muthén, 1998-2019). We used Bayesian estimation because it provides a true latent decomposition into within- and between-person variance for both the predictors and outcome variables when random effects are specified. Significance for all model parameters was based on 95% Credibility Intervals (CIs), with CIs that excluded zero indicative of a parameter that differed significantly from zero. Figure 1 provides a hypothesis-by-hypothesis schematic overview of the estimated models, along with path annotations as referenced in the results section.

Total variance in the observed variables of sleep and suicidal ideation was decomposed into their constituent pieces. Because sleep is a daily level variable nested within people, it was

decomposed into its day-level and person-level variables using latent decomposition. Accordingly, both assessments of suicidal ideation and negative affect were averaged over each day, and decomposed into their day-level, and person-level latent variables. This decomposition is akin to the partitioning of variance that occurs with the outcome variable in standard multilevel modeling, however in MSEM it is conducted for both outcomes and predictors that have repeated observations within a person. The between person-level latent variables are an estimate of an individual's average (i.e., random intercepts), and resemble the portion of variance that is usually captured in cross-sectional studies. The day-level variables reflect the deviations from their average on a given day, for that person, respectively.

Based on our preregistered hypotheses, two sets of models were estimated with nightly sleep (i.e., latency, duration, quality) predicting daily suicidal ideation or negative affect. Model Set 1 (*H1a*) regressed daily suicidal ideation on nightly sleep at Level 1 (i.e., sleep and daily suicidal ideation), and Level 2 (i.e., associations among individual differences in average sleep parameters and suicidal ideation). Model Set 2 (*H1b*), regressed daily negative affect on nightly sleep at both levels. To test *H1c*, Model Set 3 combined Models 1 and 2 by introducing negative affect as intermediate variable for the relationship between sleep and suicidal ideation at Level 1 and Level 2.

To test the second set of hypotheses, Models 4 (*H2a*) and 5 (*H2b*) introduced BPD symptom severity as moderator of the link between sleep and suicidal ideation, or sleep and negative affect (i.e., cross-level interactions), while also predicting individual differences in ratings of one's nightly sleep, negative affect and suicidal ideation.

Age and gender (0=female; 1=male) were modeled as covariates at the between-person level, and random slopes and intercepts were regressed on these covariates. Study day and

weekend (vs. weekday) were modeled as covariates at the within-person level to control for any confounding influence of time and weekend effects. No prior information about parameter values were specified (i.e., we used non-informative/diffuse priors that are the default in MPlus).

Missing data was assumed to be missing at random. A Bayesian approach to SEM uses all available data in estimation: with increasingly large samples, it provides similar results to Full Information Maximum Likelihood to address missing data (Asparouhov & Muthén, 2010).

Along with other parameters not reported in the tables (e.g., residual variances), covariances among between-persons variables are not depicted in the diagrams, but full specifications and detailed output from all models can be found online at osf.io/cxy9b. Table 1 summarizes pooled within-person correlations among the variables along with correlations among the random intercepts (i.e., person averages) at the between-person level.

Results

Model Set 1: Does nightly sleep predict next-day suicidal ideation?

Results for models examining the effects of sleep (i.e., sleep latency, duration, and quality) on next-day suicidal ideation can be found in the first column of Table 2. None of the three model specifications returned a significant link between sleep and next day suicidal ideation at the within-person level. That is, on days when participants reported worse than average nightly sleep, they did not also report greater suicidal ideation. At the between-person level, however, we found a relationship between sleep latency and suicidal ideation suggesting that participants who reported taking a longer time to fall asleep also reported higher average levels of suicidal ideation.

Model Set 2: Does nightly sleep predict next-day negative affect?

Key estimates for models testing the effects of sleep (i.e., sleep latency, duration, and quality) on next-day negative affect are detailed in the central column of Table 2. Overall, we only found a significant positive within-person link between subjective sleep quality and next day negative affect. Thus, on average, when individuals reported worse nightly subjective sleep quality, they also tended to report elevated levels of negative affect on the subsequent day.

At the between-persons level, a positive relationship emerged for negative affect and sleep latency, and a negative association for subjective sleep quality and negative affect. That is, those who generally tended to report longer sleep onset latency and worse subjective sleep quality, also reported higher levels of negative affect on average.

Model 3: Does negative affect explain the relation between nightly sleep and next-day suicidal ideation?

As noted in our preregistration, our aim was to combine both Model 1 and Model 2 to test whether negative affect would account for the relationship between poor nightly sleep and suicidal ideation. Unsurprisingly given the nonsignificant direct associations between sleep variables and suicidal ideation at the within-person level, negative affect did not statistically mediate these associations (see right column of Table 2, Model 3).

Model 4: Does BPD severity moderate the associations among sleep, negative affect, and suicidal ideation?

Finally, we tested whether BPD severity features moderated the paths of the within-person Models 1 and 2. All within-person paths exhibited significant variability across individuals. Individual differences in the strength of these within-person paths were regressed on BPD severity, sex, and age. Coefficients from these models are displayed in Table 3, organized

by sleep variables. Results indicated that BPD severity had no significant effect on any within-person pathway, although it did have associations with between-person differences.

Specifically, BPD severity was positively associated with individual differences in suicidal ideation, except for the model where suicidal ideation was regressed on sleep latency. BPD severity was also positively related to sleep latency and negatively to subjective sleep quality. Additionally, in models where negative affect was regressed on sleep, we found that BPD severity was associated with greater negative affect (see Table 3 for an overview).

Discussion

Whereas previous studies reported long-term associations between disturbed sleep and suicide risk, it remained unclear whether these associations were driven by individual differences or dynamic within-person processes. Therefore, sleep disturbances could not be ruled out as concomitant to or an epiphenomenon of BPD, and to what degree disrupted sleep predicts short-term suicide risk in contrast (*see Liu et al., 2020 for a systematic review and meta-analysis of longitudinal studies*). To narrow this gap in the literature, we obtained data from a large, well-characterized BPD sample oversampled for past suicide attempts, and preregistered our analyses.

In contrast to our first set of preregistered hypotheses, we did not find a significant within-person association between daytime suicidal ideation and self-reports of how a participant slept the night before. Overall, this pattern contrasts with sparse previous evidence that prior sleep disruption predicts suicidal ideation (Hochard et al., 2015; Ribeiro et al., 2012; Zuromski et al., 2017). It particularly contrasts with Littlewood and colleagues (2019) who found that both self-reported sleep duration and subjective sleep quality predicted greater next-day suicidal ideation. One explanation may be that these prior studies did not appear to explicitly disentangle and separately model within-person and between-person associations between sleep and suicidal

ideation. Moreover, the use of community or non-BPD samples in these studies limits the ability to generalize results to BPD samples, like the sample reported in this study. It is also worth noting that the current study collected substantially more repeated assessments per person than most prior studies, and obtained a significantly larger sample size (Hochard et al., 2015; Littlewood et al., 2019; Ribeiro et al., 2012), affording more accuracy when approximating links between BPD, suicidal ideation and sleep (e.g., Oltmanns, 2019).

At the between-person level, however, we found that people who generally take longer to fall asleep also tended to report greater suicidal ideation. The relationship between sleep latency and suicidal ideation could be reflective of more general, aberrant self-regulatory processes in BPD that are shared, potentially etiological and maintaining factors of poor subjective sleep quality in BPD. However, because these links emerged at the between- rather than the within-person level, sleep disturbances likely reflect concomitants or epiphenomena of BPD rather than a stable risk factor (Liu et al., 2020). In that regard, it is reasonable to assume that effects of sleep on BPD symptom exacerbation unfold over longer periods of time, and thus more likely resemble stable predispositions to affective dysregulation.

Accordingly, tests of the second set of hypotheses revealed that at the between-person level, people with greater average negative affect had longer sleep onset latency and worse subjective sleep quality on average. Moreover, a significant within-person link between subjective sleep quality and negative affect emerged, wherein a night of better quality of sleep than typical predicted less than usual next day negative affect. Importantly, this effect may be interpreted as a check of our study's ability to detect within-person relationships in general. Overall, these findings generally replicate findings that sleep and negative affect are linked at both the within- and between-person level (Kahn et al., 2013; Palmer & Alfano, 2017). They also

converge with evidence that a number of symptoms and behaviors central to BPD are known to disrupt sleep patterns (Fleischer et al., 2012; Winsper & Tang, 2014). Relatedly, Harvey et al. (2011) proposed that it is important to address both sleep disturbance and emotion dysregulation, in treatment to produce durable treatment gains. For instance, helping to stabilize sleep prior to addressing emotion regulation skills might facilitate skill acquisition and improve overall treatment outcome (*see also* Nutt et al., 2008).

Regarding our third hypothesis, we did not find that negative affect accounted for the within-person link between sleep and suicidal ideation. Wall et al (2020) reported a similar pattern in cross-sectional data, where adolescent BPD and affective dysregulation were significantly related to sleep disturbance, but affective dysregulation did not account for the link between sleep disturbance and BPD.

Finally, tests of our second set of hypotheses revealed, that none of the significant within-person links was moderated by BPD severity. In line with previous work (Winsper et al., 2017), however, we found that BPD severity was generally positively associated with suicidal ideation, and sleep latency, while its was negatively associated with subjective sleep quality. Additionally, in models where negative affect was regressed on sleep, we found that BPD severity was positively related to negative affect. Thus, people with greater severity of BPD had worse sleep, greater negative affect, and greater suicidal ideation on average. However, these participants did not have stronger within-person links among these variables. Moreover, these effects were unique to BPD, even after controlling for depression, which we demonstrated in an exploratory, not preregistered set of analyses (osf.io/cxy9b).

Overall, these findings suggest that the associations of sleep with suicidal ideation and BPD largely reside at the individual difference level rather than the within-person level. Those

who experience more negative affect in general also tend to report lower subjective sleep quality or greater sleep latency. Additionally, those who reported longer average sleep latency also endorsed higher levels of suicidal ideation. In addition, poor sleep and greater suicidal ideation tended to characterize individuals higher in BPD severity rather than poor sleep exacerbating suicidal ideation and negative emotions in BPD patient populations. It is possible that the pattern of results reported in this study reflects more general self-regulatory impairments in BPD, that have the potential to spill over into disturbed nightly sleep (e.g., Harty et al., 2010; Selby et al., 2013) that may largely coincide, though not specifically contribute to the exacerbation of suicidal crises in BPD.

Study Limitations

The current research has limitations that are important to consider. First, findings are based upon self-report data which may upwardly bias associations due to common method variance (Podsakoff et al., 2003). Additionally, self-reports of sleep demonstrate moderate convergence with actigraphy and polysomnographic measures of sleep and future research should strive to use other methods to measure sleep. Along these lines, participant raw reports of bedtime and waketime, which were used to calculate sleep duration, had reporting errors (a frequent, yet infrequently discussed problem with self-reports of bed and waketimes) that either had to be corrected or resulted in data loss. Using an actigraphic measure of sleep would avoid such reporting errors, (though it is still prone to other data cleaning problems, see Fekedulegn et al., 2020). Despite the limitation of using self-reported sleep data, it should be noted that subjective sleep quality is a unique sleep characteristic that can only be measured by self-report and which is an important, distinct element of sleep that is tied to health and functioning (Buysse, 2014).

Even in real-world, every day contexts and representative, high-risk populations, the base rate of momentary endorsements of suicidal ideation is low, and the levels reported in this sample were slightly higher than averages reported in previous studies (7.8% of reports in Husky et al., 2014). This low variance in suicidal ideation likely reduced the ability to detect and downwardly biased associations between sleep and suicidal ideation. Thus, to increase the reliability of our statistical models, we averaged momentarily assessed suicidal ideation over each day. This allowed us to create a detailed measure of suicidal ideation severity, encompassing a wide range of suicidal thoughts and potential behaviors that no previous momentary study has assessed. Future studies could, therefore, explore whether different item and scaling choices would affect response rates, and also, which temporal resolution would yield most reliable estimates.

Further, our findings at the between-person level are cross-sectional in nature and therefore preclude causal conclusions. Thus, while an individual's BPD severity and average levels of sleep, negative affect, and suicidal ideation all tended to be associated with each other, it is unclear whether any of these characteristics precede or cause one another. There is limited evidence supporting the notion that sleep problems may be evident before the development of BPD and contribute to its etiology (Lereya et al., 2017; Selby et al., 2013; Wall et al., 2020). There is also tentative evidence to suggest that ongoing sleep problems can contribute to the chronicity of the disorder (Plante et al., 2013). Intensive longitudinal studies focused on more chronic levels of these characteristics, across different developmental stages are needed to determine how poor sleep and short-term fluctuations in suicidal ideation relate to each other in daily life.

Although we have probed and illustrated the robustness of our study results via sensitivity analyses controlling for concurrent depression (osf.io/cxy9b), the present study is – to some extent – limited by a comprehensive control strategy of important confounders of sleep disturbances, such as other relevant comorbid disorders or socio-economic features (e.g., working status). Data derived from ambulatory assessment protocols allow to characterize temporal aspects of the relationship between sleep, affect and suicidality as they fluctuate over time as well as their reciprocal influence. Cross-sectional analyses, in contrast, focus on epiphenomena of suicidal ideation. Thus, although it is plausible to assume that it is predominantly the latter relationship that is more susceptible to biases by unobserved confounds, future studies should investigate whether the dynamic coupling between variables of interest may be weakened or enhanced in the context of theoretically justifiable control variables.

Finally, it is important to note that our set of sleep measures did not include an indicator of nightmares, which are known to occur frequently in BPD (Winsper et al., 2017). This is a critical limitation, because preliminary evidence suggests a prospective association between nightmares and BPD symptoms based on intensive longitudinal data in a small sample (Selby et al., 2013) as well as over longer developmental periods (Lereya et al., 2016). There is also growing evidence showing that nightmares are associated with increased risk of suicidal behavior or self-harm in clinical and community samples (e.g., Hochard et al., 2015; Russel, Rasmussen, & Hunter, 2017), independently of a history of traumatic experience (Sandmann et al., 2017). Possibly, nightmares aggravate daytime affective dysregulation which may manifest in symptom exacerbation; alternatively, nightmares may also be reflective of daily affect dysregulatory experiences (Selby et al., 2013). Thus, more research is needed to clarify the direction and magnitude of the relationship between nightmares and affect-regulatory processes

in daily life, and how each of these processes as well as their interactive effects may contribute to states of heightened risk for death by suicide.

Conclusion

This is the first preregistered study examining sleep as a short-term risk factor of suicidal ideation in a well-characterized BPD sample. We believe that it represents an important first step toward rigorous research on how sleep and suicide are related on a narrow timescale in BPD (*see also* Oltmanns, 2019), and emphasize the importance of disentangling within- and between-person associations in this context. Because our results suggest that the link between sleep disturbance and suicidal ideation in BPD largely exists at the individual difference level rather than the within-person level, it is reasonable to assume that sleep disturbance generally emerges as part of the disorder. Those who experience generally elevated levels of negative affect, also tend to report lower subjective sleep quality or greater sleep latency, while those who experience generally more pronounced levels of sleep latency, also endorse higher levels of suicidal ideation. In other words, poor sleep and greater suicidal ideation tend to characterize individuals higher in BPD severity rather than poor sleep exacerbating suicidal ideation and negative emotions in BPD. Possibly, effects of sleep on BPD symptom exacerbation unfold over longer periods of time, which is why they phenomenologically resemble stable predispositions to affective dysregulation, and more general self-regulatory impairments likely manifest as disturbed nightly sleep in BPD.

Disturbed sleep, therefore, seems to largely coincide, rather than specifically contribute to the exacerbation of suicidal crises in BPD.

References

- Akiskal, H.S., Yerevanian, B.I., Davis, G.C., King, D., & Lemmi, H. (1985). The nosologic status of borderline personality: Clinical and polysomnographic study. *The American Journal of Psychiatry*, 142, 192–198.
- Asparouhov, T., & Muthén, B. (2010). Bayesian analysis of latent variable models using Mplus.
- Balestrieri, M., Rucci, P., Sbrana, A., Ravani, L., Benvenuti, A., Gonnelli, C., ... & Cassano, G. B. (2006). Lifetime rhythmicity and mania as correlates of suicidal ideation and attempts in mood disorders. *Comprehensive psychiatry*, 47(5), 334-341.
- Beck, A.T., Beck, R., & Kovacs, M. (1975). Classification of suicidal behaviors: I. Quantifying intent and medical lethality. *The American Journal of Psychiatry*, 132(3), 285–287.
- Ben-Zeev, D., Young, M.A., & Depp, C.A. (2012). Real-time predictors of suicidal ideation: mobile assessment of hospitalized depressed patients. *Psychiatry Research*, 197(1-2), 55-59.
- Bernert, R.A., Kim, J.S., Iwata, N.G., & Perlis, M.L. (2015). Sleep disturbances as an evidence-based suicide risk factor. *Current Psychiatry Reports*, 17(3), 15.
- Bolger, N., & Laurenceau, J.P. (2013). *Intensive longitudinal methods: An introduction to diary and experience sampling research*. Guilford Press.
- Bromundt, V., Wirz-Justice, A., Kyburz, S., Opwis, K., Dammann, G., & Cajochen, C. (2013). Circadian sleep-wake cycles, well-being, and light therapy in borderline personality disorder. *Journal of Personality Disorders*, 27(5), 680-696.
- Buyse, D.J. (2014). Sleep health: can we define it? Does it matter?. *Sleep*, 37(1), 9-17.
- DeShong, H.L., & Tucker, R.P. (2019). Borderline personality disorder traits and suicide risk:

- 523 The mediating role of insomnia and nightmares. *Journal of Affective Disorders*, 244, 85-
524 91.
- 525 Fekedulegn, D., Andrew, M.E., Shi, M., Violanti, J.M., Knox, S., & Innes, K.E. (2020).
526 Actigraphy-Based Assessment of Sleep Parameters. *Annals of Work Exposures and*
527 *Health*, 64(4), 350-367.
- 528 Fleischer, M., Schäfer, M., Coogan, A., Häbeler, F., & Thome, J. (2012). Sleep disturbances and
529 circadian CLOCK genes in borderline personality disorder. *Journal of Neural*
530 *Transmission*, 119(10), 1105-1110.
- 531 Grove, J.L., Smith, T.W., Carlson, S.E., Bryan, C.J., Crowell, S.E., Czajkowski, L., ... &
532 Parkhurst, K. (2020). Prospective association between suicide cognitions and emotional
533 responses to a laboratory stressor: The mediating role of nightly subjective sleep
534 quality. *Journal of Affective Disorders*, 265, 77-84.
- 535 Grove, J.L., Smith, T.W., Crowell, S.E., & Ellis, J.H. (2017). Preliminary evidence for
536 emotion dysregulation as a mechanism underlying poor sleep quality in borderline
537 personality disorder. *Journal of Personality Disorders*, 31(4), 542-552.
- 538 Hafizi, S. (2013). Sleep and borderline personality disorder: A review. *Asian Journal of*
539 *Psychiatry*, 6(6), 452-459.
- 540 Harty, L., Forkner, R.D., Thompson, A., Stuewig, J., & Tangney, J.P. (2010). Are inmates'
541 subjective sleep problems associated with borderline personality, psychopathy, and
542 antisocial personality independent of depression and substance dependence?. *Journal of*
543 *Forensic Psychiatry & Psychology*, 21(1), 23-38.
- 544 Hochard, K.D., Heym, N., & Townsend, E. (2015). The unidirectional relationship of
545 nightmares on self-harmful thoughts and behaviors. *Dreaming*, 25(1), 44-58.

- 546 Husky, M., Olié, E., Guillaume, S., Genty, C., Swendsen, J., & Courtet, P. (2014). Feasibility
547 and validity of ecological momentary assessment in the investigation of suicide risk.
548 *Psychiatry Research*, 220(1-2), 564-570.
- 549 Kahn, M., Sheppes, G., & Sadeh, A. (2013). Sleep and emotions: bidirectional links and
550 underlying mechanisms. *International Journal of Psychophysiology*, 89(2), 218-228.
- 551 Kaurin, A., Dombrovski, A., Hallquist, M., & Wright, A.G. (2020, July 22). Daily Suicidal
552 Surge and Attempted Suicide in Borderline Personality Disorder.
553 <https://doi.org/10.31234/osf.io/hswea>
- 554 Kleiman, E.M., Turner, B.J., Fedor, S., Beale, E.E., Huffman, J.C., & Nock, M.K. (2017).
555 Examination of real-time fluctuations in suicidal ideation and its risk factors: Results
556 from two ecological momentary assessment studies. *Journal of Abnormal Psychology*,
557 126(6), 726–738.
- 558 Lemola, S., Ledermann, T., & Friedman, E.M. (2013). Variability of sleep duration is related to
559 subjective sleep quality and subjective well-being: an actigraphy study. *PloS one*, 8(8),
560 e71292.
- 561 Lereya, S.T., Winsper, C., Tang, N.K., & Wolke, D. (2017). Sleep problems in childhood and
562 borderline personality disorder symptoms in early adolescence. *Journal of Abnormal*
563 *Child Psychology*, 45(1), 193-206.
- 564 Links, P.S., Eynan, R., Heisel, M.J., Barr, A., Korzekwa, M., McMain, S., & Ball, J.S. (2007).
565 Affective instability and suicidal ideation and behavior in patients with borderline
566 personality disorder. *Journal of Personality Disorders*, 21(1), 72-86.
- 567 Littlewood, D.L., Kyle, S.D., Carter, L.A., Peters, S., Pratt, D., & Gooding, P. (2019). Short

- 568 sleep duration and poor sleep quality predict next-day suicidal ideation: an ecological
569 momentary assessment study. *Psychological Medicine*, 49(3), 403-411.
- 570 Liu, R. T., Steele, S. J., Hamilton, J. L., Do, Q. B., Furbish, K., Burke, T. A., ... & Gerlus, N.
571 (2020). Sleep and suicide: A systematic review and meta-analysis of longitudinal
572 studies. *Clinical Psychology Review*, 101895.
- 573 Loranger, A., Sartorius, N., & Janca, A. (1996). Assessment and diagnosis of personality
574 disorders: The International Personality Disorder Examination (IPDE). New York, NY:
575 Cambridge University Press.
- 576 McNamara, E., Reynolds, C.F., III, Soloff, P. H., Mathias, R., Rossi, A., Spiker, D., ... Kupfer,
577 D. J. (1984). EEG sleep evaluation of depression in borderline patients. *The American*
578 *Journal of Psychiatry*, 141, 182–186.
- 579 Morales-Muñoz, I., Broome, M. R., & Marwaha, S. (2020). Association of Parent-Reported
580 Sleep Problems in Early Childhood With Psychotic and Borderline Personality Disorder
581 Symptoms in Adolescence. *JAMA Psychiatry*. Published online July 01, 2020.
- 582 Morey, L. C. (1991). Personality assessment screener. *Odessa, FL: Psychological Assessment*
583 *Resources*.
- 584 Mou, D., Kleiman, E.M., Fedor, S., Beck, S., Huffman, J.C., & Nock, M.K. (2018). Negative
585 affect is more strongly associated with suicidal thinking among suicidal patients with
586 borderline personality disorder than those without. *Journal of Psychiatric Research*, 104,
587 198-201.
- 588 Muthén, L., & Muthén, B. (2019). Mplus User's Guide (2019). *Muthén & Muthén: Los Angeles,*
589 *CA*.
- 590 Nutt, D., Wilson, S., & Paterson, L. (2008). Sleep disorders as core symptoms of

- 591 depression. *Dialogues in Clinical Neuroscience*, 10(3), 329-336.
- 592 O'Connor, R.C., & Nock, M.K. (2014). The psychology of suicidal behaviour. *The Lancet*
593 *Psychiatry*, 1(1), 73-85.
- 594 Oltmanns, J.R. (2019). Personality Disorders, Maladaptive Personality Traits, and Sleep
595 Problems: Findings and Future Directions. In *Sleep, Personality, and Social*
596 *Behavior* (pp. 239-254). Springer, Cham.
- 597 Oltmanns, J.R., & Oltmanns, T.F. (2015). Borderline personality pathology, polysomnography,
598 and self-reported sleep problems: a review. *Current Sleep Medicine Reports*, 1(2), 141-
599 149.
- 600 Oltmanns, J.R., Weinstein, Y., & Oltmanns, T.F. (2014). Borderline personality pathology and
601 insomnia symptoms in community-dwelling older adults. *Personality and Mental*
602 *Health*, 8(3), 178-187.
- 603 Palmer, C.A., & Alfano, C.A. (2017). Sleep and emotion regulation: an organizing, integrative
604 review. *Sleep Medicine Reviews*, 31, 6-16.
- 605 Pandi-Perumal, S.R., Monti, J.M., Burman, D., Ramanujam, K., BaHammam, A.S., Spence,
606 D.W., Brown, G.M & Narashimhan, M. (2020). Clarifying the role of sleep in
607 depression: A narrative review. *Psychiatry Research*, 113239.
- 608 Plante, D.T., Frankenburg, F.R., Fitzmaurice, G.M., & Zanarini, M.C. (2013). Relationship
609 between maladaptive cognitions about sleep and recovery in patients with borderline
610 personality disorder. *Psychiatry Research*, 210(3), 975-979.
- 611 Podsakoff, P.M., MacKenzie, S.B., Lee, J.Y., & Podsakoff, N.P. (2003). Common method
612 biases in behavioral research: a critical review of the literature and recommended
613 remedies. *Journal of Applied Psychology*, 88(5), 879-903.

- 614 Posner, K., Brent, D., Lucas, C., Gould, M., Stanley, B., Brown, G., ... & Mann, J. (2008).
615 Columbia-Suicide Severity Rating Scale (C-SSRS). *New York, NY: Columbia University*
616 *Medical Center, 2008.*
- 617 Reynolds, C.F., III, Soloff, P.H., Kupfer, D.J., Taska, L.S., Restifo, K., Coble, P.A., &
618 McNamara, M.E. (1985). Depression in borderline patients: A prospective EEG sleep
619 study. *Psychiatry Research, 14*, 1–15.
- 620 Ribeiro, J.D., Pease, J.L., Gutierrez, P.M., Silva, C., Bernert, R.A., Rudd, M.D., & Joiner Jr,
621 T.E. (2012). Sleep problems outperform depression and hopelessness as cross-sectional
622 and longitudinal predictors of suicidal ideation and behavior in young adults in the
623 military. *Journal of Affective Disorders, 136*(3), 743-750.
- 624 Russell, K., Rasmussen, S., & Hunter, S. C. (2018). Insomnia and nightmares as markers of risk
625 for suicidal ideation in young people: investigating the role of defeat and
626 entrapment. *Journal of Clinical Sleep Medicine, 14*(5), 775-784.
- 627 Sandman, N., Valli, K., Kronholm, E., Vartiainen, E., Laatikainen, T., & Paunio, T. (2017).
628 Nightmares as predictors of suicide: an extension study including war veterans. *Scientific*
629 *Reports, 7*(1), 1-7.
- 630 Sansone, R.A., Edwards, H.C., & Forbis, J.S. (2010). Sleep quality in borderline personality
631 disorder: a cross-sectional study. *Primary Care Companion to the Journal of Clinical*
632 *Psychiatry, 12*(5): PCC.09m00919.
- 633 Selby, E.A. (2013). Chronic sleep disturbances and borderline personality disorder
634 symptoms. *Journal of Consulting and Clinical Psychology, 81*(5), 941–947.
- 635 Selby, E. A., & Joiner Jr, T. E. (2009). Cascades of emotion: The emergence of borderline

- 636 personality disorder from emotional and behavioral dysregulation. *Review of General*
637 *Psychology*, 13(3), 219-229.
- 638 Simor, P., & Horváth, K. (2013). Altered sleep in borderline personality disorder in relation to
639 the core dimensions of psychopathology. *Scandinavian Journal of Psychology*, 54(4),
640 300-312.
- 641 Sjöström, N., Hetta, J., & Waern, M. (2009). Persistent nightmares are associated with repeat
642 suicide attempt: a prospective study. *Psychiatry Research*, 170(2-3), 208-211.
- 643 Soloff, P. H., & Chiappetta, L. (2017). Suicidal behavior and psychosocial outcome in
644 borderline personality disorder at 8-year follow-up. *Journal of Personality Disorders*, 31
645 (6), 774-789.
- 646 Tkachenko, O., Olson, E. A., Weber, M., Preer, L. A., Gogel, H., & Killgore, W. D. (2014).
647 Sleep difficulties are associated with increased symptoms of psychopathology.
648 *Experimental Brain Research*, 232(5), 1567-1574.
- 649 Vallières, A., Ivers, H., Bastien, C.H., Beaulieu-Bonneau, S., & Morin, C.M.
650 (2005). Variability and predictability in sleep patterns of chronic insomniacs. *Journal of*
651 *Sleep Research*, 14(4), 447-453.
- 652 Wall, K., Vanwoerden, S., Penner, F., Patriquin, M., Alfano, C.A., & Sharp, C. (2020).
653 Adolescent Sleep Disturbance, Emotion Regulation and Borderline Features in an
654 Inpatient Setting. *Journal of Clinical Child & Adolescent Psychology*, 1-15.
- 655 Winsper, C., & Tang, N.K. (2014). Linkages between insomnia and suicidality: prospective
656 associations, high-risk subgroups and possible psychological mechanisms. *International*
657 *Review of Psychiatry*, 26(2), 189-204.
- 658 Winsper, C., Tang, N.K., Marwaha, S., Lereya, S.T., Gibbs, M., Thompson, A., & Singh, S.P.

- 659 (2017). The sleep phenotype of Borderline Personality Disorder: A systematic review and
660 meta-analysis. *Neuroscience & Biobehavioral Reviews*, 73, 48-67.
- 661 Wojnar, M., Ilgen, M.A., Wojnar, J., McCammon, R.J., Valenstein, M., & Brower, K.J. (2009).
662 Sleep problems and suicidality in the National Comorbidity Survey Replication. *Journal*
663 *of Psychiatric Research*, 43(5), 526-531.
- 664 Zanarini, M.C., Frankenburg, F.R., Reich, D.B., & Fitzmaurice, G.M. (2016). Fluidity of the
665 subsyndromal phenomenology of borderline personality disorder over 16 years of
666 prospective follow-up. *American Journal of Psychiatry*, 173(7), 688-694.
- 667 Zuromski, K.L., Cero, I., & Witte, T.K. (2017). Insomnia symptoms drive changes in suicide
668 ideation: A latent difference score model of community adults over a brief
669 interval. *Journal of Abnormal Psychology*, 126(6), 739.

Figure 1. Overview of quantitative model specifications used to test preregistered hypotheses, including decomposition of observed daily variables into between- (subscript *i*) and within-person (subscript *t*) variance. Note, not all parameters are diagrammed. All models controlled for age and gender at the between-person level, and for time and weekday at the within-person level. Coefficients for covariates are not presented for parsimony. Single headed arrows indicate regression paths. Solid lines represent paths for hypotheses 1a-c, and dashed for hypotheses 2a and b. Filled dots represent random effects. NA = negative affect; SI = suicidal ideation.

Table 1. Correlations among study variables at within and between-person levels.

	quality	latency	duration	negative affect	suicidal ideation	BPD
within-person						
quality	-					
latency	-.34	-				
duration	.30	-.31	-			
negative affect	-.06	.02	-.06	-		
suicidal ideation	-.03	.07	-.04	.15	-	-
between-person						
quality						
latency	-.46					
duration	.21	-.14				
negative affect	-.04	.32	.10			
suicidal ideation	.07	.30	.06	.23		
BPD	-.52	.46	-.12	.37	.21	
<i>M</i>	3.18	22.66	7.35	1.62	.11	39.19
<i>SD</i>	1.01	22.67	1.85	.07	.04	19.50

Note. $N=201$ (between), $N=4076$ (within); Bolded values indicate the credibility interval does not contain zero.

679 Table 2.

680 Key estimates from three individual multi-level structural equation models showing associations among sleep, negative affect and
 681 suicidal ideation.

	Model Set 1 Hypothesis 1a		Model Set 2 Hypothesis 1b		Model Set 3 Hypothesis 1c	
	<i>Est</i> (CI)	β	<i>Est</i> (CI)	β	<i>Est</i> (CI)	β
sleep latency						
within-person						
sleep → suicidal ideation (c)	.00 [-.00, .01]	.14	—	—	.00 [00, .00]	.13
sleep → NA (a)	—	—	.00 [-.00, .01]	.06	.00 [-.00, .00]	.02
NA → suicidal ideation (b)	—	—	—	—	.14 [.08, .20]	.45
indirect effect	—	—	—	—	.00 [-.00, -.01]	—
between-person						
sleep → suicidal ideation	.01 [.00, .01]	.24	—	—	.99 [.38, 1.70]	.24
sleep → NA	—	—	.02 [.01, .02]	.44	3.92 [2.50, 5.60]	.44
NA → suicidal ideation	—	—	—	—	.06 [.04, .10]	.36
sleep duration						
within-person						
sleep → suicidal ideation (c)	-.00 [-.01, .01]	-.04	—	—	-.01 [-.02, .01]	-.07
sleep → NA (a)	—	—	-.01 [-.02, .01]	-.10	-.02 [-.03, .00]	-.15
NA → suicidal ideation (b)	—	—	—	—	.15 [.09, .20]	.48
indirect effect	—	—	—	—	.00 [-.01, .01]	—
between-person						
sleep → suicidal ideation	-.02 [-.03, .07]	.06	—	—	.01 [-.04, .07]	.04
sleep → NA	—	—	-.03 [-.12, .06]	-.07	-.04 [-.16, .08]	-.05
NA → suicidal ideation	—	—	—	—	.07 [.04, .10]	.37

(continued)

682

subjective sleep quality						
within-person						
sleep → suicidal ideation (c)	-.01 [-.03, .00]	-.16	—	—	-.01 [-.03, .00]	-.21
sleep → NA (a)	—	—	-.05 [-.10, -.02]	-.33	-.08 [-.11, -.05]	-.49
NA → suicidal ideation (b)	—	—	—	—	.13 [.08, .19]	.45
indirect effect	—	—	—	—	-.01 [-.02, .01]	—
between-person						
sleep → suicidal ideation	-.03 [-.10, .03]	-.06	—	—	.00 [-.03, .03]	.00
sleep → NA	—	—	-.24 [-.37, -.12]	-.29	-.13 [-.20, -.07]	-.32
NA → suicidal ideation	—	—	—	—	.07 [.04, .10]	.36

Note. $N=201$ (between), $N=4076$ (within); → indicates regression; β parameter estimates are standardized. 95% credibility intervals of unstandardized parameter estimates are in parentheses. Bolded values indicate the credibility interval does not contain zero; NA = negative affect.

Table 3.

Key estimates from multilevel models with sleep predicting suicidal ideation or negative affect and moderation of these within-person effects by BPD severity.

Model Set 4				
	Hypothesis 2a		Hypothesis 2b	
	<i>Est</i> (CI)	β	<i>Est</i> (CI)	β
sleep latency				
BPD → sleep	.31 [.21,.42]	.42	.32 [.22, .43]	.42
BPD → NA	—	—	.02 [.01, .02]	.48
BPD → suicidal ideation	.00 [.00, .00]	.14	—	—
BPD x (sleep → suicidal ideation)	.00 [.00,.00]	.06	—	—
BPD x (sleep → NA)	—	—	.00 [.00, .00]	-.01
sleep duration				
BPD → sleep	-.01 [-.02, .01]	-.09	-.01 [-.02 .00]	-.12
BPD → NA	—	—	.02 [.01, .02]	.58
BPD → suicidal ideation	.00 [.00, .01]	.22	—	—
BPD x (sleep → suicidal ideation)	.00 [-.00, .00]	-.05	—	—
BPD x (sleep → NA)	—	—	-.00 [.00 .00]	-.10
BPD x (NA → suicidal ideation)	—	—	—	—
subjective sleep quality				
BPD → sleep	-.02 [-.02, -.01]	-.50	-.02 [-.02,-.01]	-.50
BPD → NA	—	—	.02 [.01, .02]	.56
BPD → suicidal ideation	.01 [.00, .01]	.25	—	—
BPD x (sleep → suicidal ideation)	.00 [-.00, .00]	-.10	—	—
BPD x (sleep → NA)	—	—	-.00 [-.00, .00]	-.15
BPD x (NA → suicidal ideation)	—	—	—	—

Note. $N=201$ (between), $N= 4076$ (within); → indicates regression; β parameter estimates are standardized. 95% credibility intervals of unstandardized parameter estimates are in parentheses. Bolded values indicate the credibility interval does not contain zero; NA = negative affect.