Vicarious Ratings of Self vs. Other-directed Social Touch in Women with and Recovered from Anorexia Nervosa

Ashleigh Bellard^{1*}, Paula Trotter¹, Francis McGlone¹ & Valentina Cazzato¹

¹School of Psychology, Faculty of Health, Liverpool John Moores University,
Liverpool, UK

Correspondence to: Miss Ashleigh Bellard, School of Psychology, Faculty of Health,
Liverpool John Moores University, Liverpool, e: A.M.Bellard@2020.ljmu.ac.uk

Abstract

Anorexia Nervosa (AN) is an eating pathology characterized by restricted eating, body image distortions and impaired socio-cognitive abilities. Altered responses to affective touch – a pleasant interoceptive stimulus hypothesised to involve activation of the C-Tactile (CT) system, may contribute to the aetiology and maintenance of this disorder. Here, we investigated whether third-party social touch vicarious ratings of different body sites at CT-optimal vs. non-CT optimal velocities differed in women with and recovered from AN (RAN) and healthy controls (HCs). Thirty-five HCs, 27 AN and 29 RAN provided pleasantness ratings for two different tasks designed to probe expectations of how touch is perceived by self (self-directed touch) vs. others (other-directed touch). Findings revealed that both clinical groups, compared to HCs, did not differ in their pleasantness ratings to touch for another but when evaluating touch for self, both clinical groups rated CT-optimal touch as less pleasant than HCs. These findings suggest that AN and RAN women demonstrate an atypical vicarious pleasantness response to affective touch involving self, but not others. Therefore, as atypical responses persist even after recovery, treatment interventions should focus on overcoming an impairment in differentiating between self and other affective touch experience, which could help prevent post-recovery relapsing.

Introduction

Anorexia Nervosa (AN) is an eating disorder typically characterised by body image disturbances, including overestimations in body shape and size [1, 2, 3], atypical eating and dieting behaviours, including maintaining an extremely low and unhealthy Body Mass Index (BMI) [4]. Individuals with AN also show disturbances in interoceptive processing, which is defined as the understanding of internal bodily states such as taste, hunger, stomach distention, pain, heartbeat, and gut attention [5, 6, 7, 8, 9]. They also abnormally interpret the internal physiological condition of their body such as sensations of hunger and pain [10]. A mismatch when combining information from internal and external bodily sensations may contribute towards the aetiology of AN [9, 10, 11].

Prior investigations examining the link between interoceptive deficits and AN have mainly concentrated on unrewarding interoceptive signalling, including pain (thermal), neutral interoceptive signalling (heart rate), and interoceptive signals specific to the symptoms of AN (hunger and taste), with only few studies focussing instead on rewarding interoceptive stimuli, such as a gentle, caress-like affective touch [12, 13, 14, 15]. Affective touch processing is hypothesised to be facilitated by activation of low-threshold mechanoreceptive C-fibres, known as C-Tactile afferents (CTs) [16, 17, 18, 19]. CTs optimally respond to gentle stroking in velocities ranging from 1-10cm/s [20], which is generally perceived as pleasant and rewarding in neurotypical populations, with ~3cm/s being perceived as most pleasant [16, 18, 20]. A plethora of investigations have highlighted that the Insula Cortex, specifically the posterior and anterior and posterior regions, is activated when an individual is experiencing affective touch to the hairy skin of the arm [21, 22, 23]. The posterior Insula is also thought to be the centre for processing and evaluating the hedonic properties of interoceptive cues, and receives information regarding internal bodily states, thus contributing to interoceptive processing [24, 25]. Henceforth, affective touch might be considered a sub-modality of interoception, supported by the Insula Cortex, which conveys affective and affiliative aspects of social touch in agreement with the ‘Social Touch Hypothesis’ [26, 27].

To the best of our knowledge, only four independent studies have provided so far evidence for an impairment in the affective touch system in individuals with AN [12, 13, 14, 15, but also see Carey et al [28, 29] and Cazzato et al [30] for evidence on subthreshold EDs]. A seminal study by Crucianelli et al [14], reported that AN patients (restrictive-type)

perceived gentle skin stroking touch at CT-optimal velocities as less pleasant relative to healthy controls, while there was no such difference for CT-suboptimal velocities, thus suggesting that individuals with AN have abnormalities in the perception of the pleasantness of CT-mediated, affective touch. These results were corroborated by a (brain imaging) study by Davidovic et al. [13] which showed that at behavioural level, AN rated CT-optimal touch as less pleasant than a group of matched HCs. On the other hand, neuroimaging results were controversial. No differences in the activation of the Insula were observed when comparing conditions of gentle touch and no tactile stimulation amongst the two groups of AN and HCs. A further brain imaging study by Bischoff-Grethe and colleagues [12] assessed brain correlates of anticipation and perception of gentle touch on the forearm and the palm and specifically in individuals who have recovered from AN. The findings of this study show that RAN did not rate the pleasantness of CT-touch as less pleasant than controls and that there were not any CT-specific neuroimaging findings at the group level, thus not confirming the findings of the previous two studies. Finally, a recent study by Crucianelli et al. [15] showed that clinical groups of AN and RAN anticipated and rated delivered tactile stimuli as less pleasant than HCs. However, the latter difference was not related to the CT optimality of the stimulation. On the contrary, differences in the perception of CT-optimal touch were best explained by individual differences in alexithymia and interoceptive sensibility. All these studies offer an important theoretical advancement in the understanding of affective touch experiences in disordered eating. Yet, the results are inconclusive and leave this research question still unanswered. Furthermore, the studies above have only focused on directly experienced touch (with touch delivered to the self mainly on the forearm and palm) thus, neglecting the importance of investigating vicariously experienced touch.

In the general population, it has been reported that viewing others being touched induces vicarious touch experiences, which is based on the recruitment of brain regions, which originally were responsive to first-hand somatosensation, including the somatosensory cortex, the parietal operculum and the insular cortex [31, 32, 33, 34]. Furthermore, we have recently demonstrated that ratings of vicarious touch have the same relationship between velocity and anticipated pleasantness as directly felt touch [35]. However, there are several reasons to expect differences in ratings of vicariously experienced touch in the case of AN and particularly when this is self- compared to when this is other-directed. Indeed, interpreting the affective state of two persons exchanging interpersonal touch requires social cognition, such as Theory of Mind (ToM). One important aspect of ToM is cognitive

perspective taking which is one's capacity to compare state of self (1st person perspective) with another individual (3rd person perspective), with both perspectives involving different neural and cognitive processes [36, 37]. It is well known that anorexics have difficulties in attributing mental states (intentions, feelings, beliefs) to others and in explaining and predicting others' behaviour, which involves the decoding of affective stimuli [38, 39, 40, 41, 42, 43, 44]. Further, fMRI studies have revealed that even after recovery, anorexics have reduced functioning in social and cognitive networks, compared to healthy controls, which again can account for their difficulty in ToM perspective taking [38, 45]. Given that previous studies have found reduced pleasantness in individuals with AN during real experience of touch [13, 14, 15], and that observing touch activates touch experiences and corresponding brain regions [46, 47], it is reasonable to expect that when anorexics are asked to rate the perceived pleasantness of others receiving affective touch, they will use their own experiences to interpret the feelings generated in another. In doing so, one might expect that reduced pleasantness observed during real tactile experience should also extend to vicarious responses to social touch, a hypothesis which has not been investigated yet, therefore leaving an important gap in the literature of vicarious touch experience in AN.

The current study

With the present study we aimed to fill this gap by investigating whether third-party vicarious ratings of social touch delivered at CT-optimal vs. CT non-optimal velocities differ in women reporting current diagnoses of AN. Participants were shown a series of videos with affective touch being delivered to various body sites of actors, after which participants answered two questions: "How much would you like to be touched like that?" (self-directed touch) and "How pleasant do you think that action was for the person being touched?" (other-directed touch). We expected patients with AN to perceive vicarious self-directed tactile stimuli as less pleasant than other-directed touch and compared to healthy controls. Furthermore, we explored whether specific dimensions of body awareness and of social touch experiences, may predict patients' third-party vicarious pleasantness ratings of touch when delivered to self, compared to when delivered to others [48]. Accordingly, we focused on the 'Trusting' subscale of the Multidimensional Assessment of Interoceptive Awareness (MAIA) [49] given that a previous work by Brown et al. [50] reported that lower trust in one's body signals was most robustly associated with ED psychopathology including higher restraint,

eating concern, weight and shape concern, and binge eating and purging symptoms. Furthermore, a recent study confirmed that items from the MAIA Trusting subscale were the most central in bridging interoceptive awareness and ED symptoms [51].

On the other hand, we focused on ‘Attitude to intimate touch’ of The Touch Experiences and Attitudes Questionnaire (TEAQ) [52] because previous research reported that AN patients experience intimate stimuli with lower valence and dominance than healthy controls [53]. Furthermore, women suffering from disordered eating report difficulties experiencing closeness with a partner and low levels of satisfaction in relationships [54], they also express a fear of intimacy [55] and report avoidance of interpersonal relationships with men [56]. Importantly, we also recruited recovered AN (RAN) individuals to examine whether the differences in self- vs. other-directed third-party ratings hypothesized in AN as explained above may also apply to recovered patients, in comparison to HCs. Indeed, by testing women who have recovered from AN, it is possible to draw more solid conclusions about whether any potential abnormality observed in processing tactile pleasure are a cause or a consequence of starvation, thus ruling out the impact of malnutrition in the acute stages of illness which can cause cognitive and emotional deficits per se.

In line with previous evidence of difficulties in attributing mental states (intentions, feelings, beliefs) to others in both AN and RAN, it was hypothesized that both clinical groups will provide lower pleasantness ratings for self- compared to other-directed touch. However, given that results of a reduction in the perceived tactile pleasantness in terms of specificity to CT-optimal stroking velocities are to date controversial, we did not make any specific prediction with these regards. Finally, we hypothesized that scores obtained to the MAIA Trusting and to the TEAQ Attitude to intimate touch scales would predict pleasantness ratings for self-directed social touch, for both AN and RAN, but not for HCs.

Materials and Methods

Participants

A total sample of 91 female participants, assigned to three groups (Current Anorexics vs. Remitted Anorexics vs. Healthy controls) based on self-reported Anorexia Nervosa Diagnosis (Current AN diagnosis, previous AN diagnosis and no AN diagnosis) were recruited for this online study. The total sample size was calculated based on the results of a

power analysis using G*Power 3.0.10 [57], which indicated that a minimum sample of 81 participants was needed to detect a medium effect ($f = .25$) with 95% power, using a Mixed ANOVA with alpha at .05 (two tailed).

Only females were used in the current investigation, as females are typically more sensitive to touch discrimination and respond more positively to touch than males [58], and because the incidence of eating disorders is greater in women than men [59]. Twenty-seven were current anorexics aged 18-47 ($M = 25.56$, $SD = 6.95$) and with a mean BMI of 17.58 ($SD = 1.64$) (see Table 1). Information regarding type of treatment and numbers of years diagnosed for this group is reported in Table 2. An additional 29 remitted anorexics aged 18-47 ($M = 27.31$, $SD = 7.12$), with a mean BMI of 22.59 ($SD = 3.03$) were recruited (see Table 1). Information regarding number of years recovered is reported in Table 2. A total sample of 35 female control participants ($M_{age} = 27.20$, $SD = 8.81$; $M_{BMI} = 24.77$, $SD = 4.47$), were recruited for the healthy control group (see Table 1).

Participants for the healthy control group were recruited through external contacts, via social media, the University SONA participant recruitment scheme and through those known to the researchers. Healthy controls were eligible to take part if they self-reported no history or current diagnosis of any psychiatric or neurological disorders (including eating disorders and body dysmorphic disorder) when questioned. No other psychometric constraints were applied because we wanted to recruit women who are representative of the non-eating disordered female population, many of whom have concerns about body image [60].

Current and remitted AN were recruited by contacting eating disorder charities such as the Beating Eating Disorders Organisation (BEAT) and MQ Mental Health Research, in addition to contacting clinicians known to the researchers, who work directly with these patient groups. Gatekeeper permission was granted to assist with the recruitment of individuals with and recovered from AN. As well as this, patient groups were also recruited through Anorexia support groups on social media. All patients were eligible if they had a primary diagnosis of AN according to the DSM-5 [4]. Furthermore, we used scores obtained from the Eating Disorder Examination Questionnaire (EDE-Q) which is a self-report version of the Eating Disorder Examination (EDE) structured interview [61], as a screening questionnaire for eating disordered behaviour.

All participants had normal or corrected to normal vision (individuals with contact lenses or glasses were able to take part), were not pregnant, did not suffer from any skin

conditions such as eczema or any chronic pain conditions such as fibromyalgia. Eligibility for all participants was checked through an online participant screening questionnaire. All participants gave their implied informed consent to take part in the study and were debriefed through the debrief sheet presented at the end of the study. Participants were offered compensation for their time taking part. They were offered the chance to voluntarily enter themselves into a prize draw, with two chances of winning a £25 Amazon voucher and level 4 BSc Psychology students were awarded with course credits. The study was carried out in accordance with the Helsinki declaration of ethical standards. The study protocol was approved by Liverpool John Moores University (LJMU)'s University Research Ethics Committee (UREC, protocol: 20/NSP/025).

Measures

Self-report Questionnaires

Participant Screening

An initial set of seven screening questions were used to determine participants' eligibility. The questionnaire asked participants to answer "true" or "false" to a series of statements. These statements included: "I am 18 years old or over.", "I am female.", "I do not suffer from any form of skin condition.", "I do not suffer from any chronic pain condition.", "I am not pregnant.", "I have normal or corrected to normal vision". A final question asked participants to confirm they had read the participant information sheet and they agreed to take part in the study by clicking an "I agree" option. For the healthy control group only, an additional question ascertained that they did not have a current or previous diagnosis of AN. If a participant answered "false" to any of the statements, or did not agree to take part, an "if then" function was applied, so that ineligible participants were directed to the end of the survey, thanking them for their time in taking part.

To classify clinical populations into AN and RAN groups, patients were required to answer to the following statements: 'Please specify your current diagnosis of Anorexia Nervosa' with the options of identifying either: 'Current diagnosis of Anorexia Nervosa', 'Previous diagnosis of Anorexia Nervosa' or 'No diagnosis/history of Anorexia Nervosa'. For AN, this group was asked to specify how many years they have had a formal diagnosis, and to declare any treatment they are currently undergoing, through answering the following

questions: ‘If you have a current diagnosis of Anorexia Nervosa, for how many years have you had this?’ and ‘Are you currently undergoing any treatment for Anorexia Nervosa (e.g., psychiatric treatments, SSRIs, tranquilizers, and or CBT)?’. The RAN group were asked to identify the number of years they have been recovered from AN: ‘If you have a previous diagnosis of Anorexia Nervosa, for how many years have you been recovered?’. Questions specific to both AN populations were not displayed to individuals with no AN diagnosis.

Demographics Questionnaire

The demographic information taken from participants included their age, gender, biological sex, ethnicity, relationship status and education level. Participants were also asked to state their height (cm/ft) and weight (Kg/lbs) which was used to calculate their BMI.

Eating Disorder Examination Questionnaire (EDE-Q)

The Eating Disorder Examination Questionnaire (EDE-Q) [61] consists of 28 items, which asked participants to self-report any eating disorder symptoms they may have experienced over the last 4 weeks. This questionnaire comprised of four subscales: Restraint, Eating Concern, Weight Concern and Shape Concern. Participants responded to items using a 7-point Likert scale, for the first 12 questions using a scale of 0= No Days, 6= Every day. Items for these questions included: “Have you had a definite desire to have a totally flat stomach?” and “Have you had a definite fear that you might gain weight?”. For the next 6 questions participants had to state the number of days (up to 28 days) that each statement corresponded to, statements included: “Over the past 28 days, how many times have you eaten what other people would regard as an usually large amount of food (given the circumstances)?” and “Over the past 28 days, how many times have you made yourself sick (vomit) as a means of controlling your shape or weight?”. For the next 6 questions, the scale used ranged from 0= Not at all to 6= Markedly, items for these questions included “Has your weight influenced how you think about (judge) yourself as a person?” and “How dissatisfied have you been with your shape?”. The final 4 questions asked participants to self-report their current weight, height, if they have missed any menstrual periods and how many and if they are taking the pill. A sum of scores from all four subscales gave an indication of the severity of eating disorder symptoms. This questionnaire has good internal consistency in clinical populations, with Cronbach α = 0.70-0.83 and in healthy populations. with Cronbach α =0.78-

0.93 [62, 63]. Individual subscales have also shown to have good internal consistency: Restraint (Cronbach α =0.70–0.85), Eating Concern (Cronbach α =0.73–0.86), Shape Concern (Cronbach α =0.83–0.93), and Weight Concern (Cronbach α =0.72–0.89) [64]. This questionnaire was administered to assess eating disorder diagnosis and symptom severity.

Multidimensional Assessment of Interoceptive Awareness Questionnaire

The Multidimensional Assessment of Interoceptive Awareness (MAIA) [49] is a 32-item questionnaire which was administered to investigate eight dimensions of interoceptive bodily awareness: Noticing (4 items), Not Distracting (3 items), Not Worrying (3 items), Attention Regulation (7 items), Emotional Awareness (5 items), Self-regulation (4 items), Body Listening (3 items) and Trusting (3 items). Items were answered using a 6-point Likert scale ranging from 0 = Never to 5 = Always. Questions included: “When I am tense I notice where the tension is located in my body.” and “I notice when I am uncomfortable in my body”. Each individual dimension is scored by the average of scores from questions corresponding to that subscale, with some questions being reversed scored. This questionnaire is a reliable measure, as it has been previously used in research measuring interoceptive awareness in both healthy [49] and with an eating disorder population [50]. The MAIA questionnaire was found to have good internal consistency, with Cronbach α = .90 [65]. For this investigation, we were particularly interested on the MAIA Trusting subscale because previous studies by Brown and colleagues [50, 51] reported that this subscale is the most central in bridging interoceptive awareness and ED symptoms.

Touch Experiences and Attitudes Questionnaire

The Touch Experiences and Attitudes Questionnaire (TEAQ) [52] is a 57-item questionnaire which was administered to examine current experiences of positive touch and positive experience of touch during childhood, as well as an individual’s attitude towards positive touch. Questions were answered using a 5-point Likert scale ranging from 1 = “Disagree strongly”, 2 = “Disagree a little”, 3 = “Neither agree nor disagree”, 4 = “Agree a little”, 5 = “Agree strongly”. Questions included: “I dislike people being very physically affectionate towards me.” and “There was a lot of physical affection during my childhood.” A mean score was calculated for each of the six subscales; friends and family touch (FFT) (11

items), current intimate touch (CIT) (14 items), childhood touch (ChT) (9 items), attitude to self-care (ASC) (5 items), attitude to intimate touch (AIT) (13 items) and attitude to unfamiliar touch (AUT) (5 items), with negatively worded questions reversed scored. The TEAQ questionnaire was found to have a good internal consistency with Cronbach $\alpha = 0.78$ – 0.92 [52, 66]. For this investigation, we were particularly interested on the TEAQ Attitude to intimate touch (AIT) subscale because previous investigations suggest that AN patients experience intimate stimuli with lower valence and dominance than healthy controls [53]. They also report difficulties experiencing closeness with a partner and low levels of satisfaction in relationships [54], they show a fear of intimacy [55] and report avoidance of interpersonal relationships with men [56].

Self- and Other-directed affective touch video clips

Affective touch task consisted of 15, 6 s long video clips depicting actors being touched by another person, at different body sites. The videos displayed a female actor receiving touch by a male actor. This choice was made because only females were recruited in the current investigation and we wanted to avoid any confound due to using different genders receiving touch, to those observing this touch. Moreover, we chose to display videos showing only males delivering touch and females receiving touch as generally males appear to initiate touch more than females do [67, 68]. All videos were taken from a database of previously standardised video clips [66]. Touch was delivered across five different body regions: non-CT-innervated body site i.e., palm vs. CT-innervated body sites i.e., back, ventral forearm, cheek and upper arm) with 3 different stroking velocities: static (0 cm/s), slow (5cm/s) and fast (30 cm/s). The order in which the videos were presented was counterbalanced amongst participants. To avoid any confound linked to emotional expression of the agent delivering the touch [69], all videos were displayed zoomed up to the body part of the receiver and the hand of the deliverer of touch, which ensured that the actors faces were not included in the video (see Figure 1).

Immediately after viewing each video, participants were probed to respond to two questions, one asking: “How much would you like to be touched like that?” (self-directed touch) using a VAS scale ranging from 0 = not at all to 100 = extremely. This question was related to the participant themselves in their level of wanting the touch they are viewing. Participants also responded to a second question asking: “How pleasant do you think that

action was for the person being touched?” (3rd person perspective) using a 100-point VAS scale ranging from 0 = very unpleasant to 100 = extremely pleasant [66]. This question concerns the ability of participants to determine mental states of another and evaluate how pleasant the touch received was for the actor in the video (other-directed touch). Overall, there was a total of 15 videos randomly presented, with each video being presented once to participants, each displayed in 240 p YouTube quality in full HD (1920×1080 pixels) at 25 fps rate [66].

-----Insert Figure 1 here-----

General procedure

The study was conducted online using Qualtrics software, Version 60,939 of the Qualtrics Research Suite (Copyright © 2015 Qualtrics., Provo, UT, USA. <http://www.qualtrics.com>). Participants were provided with an invitation email containing a brief description of the study and the hyperlink to the online study if participants were happy to take part. Once clicked, participants were provided with the participant information sheet, screening questionnaire and gave their implied consent to take part. Once consent was obtained, participants engaged with the affective touch videos task, during which they were asked to rate using a 100-point VAS scale: “How pleasant do you think that action was for the person being touched?” and “How much would you like to be touched like that?” using a VAS scale ranging from 0 = not at all to 100 = extremely. The order participants viewed the videos was counterbalanced, as well as the order they were shown the two questions. After completion of the videos, participants filled out questionnaires which assessed body image and eating concerns, body awareness, experiences and attitudes to touch, as well as social isolation and longing for touch since the COVID-19 pandemic respectively, by means of the EDE-Q [61], MAIA [49], TEAQ [52] and the COVID-19 touch experiences and eating behaviour Questionnaire (see supplementary information). The order participants filled out these questionnaires was counterbalanced. Those participants who were not eligible were taken to the end of the survey, in which they were thanked for their time. Overall, the online study lasted approximately 35 minutes. Data collection began on 10th July 2020 and ended on 27th June 2021.

Data Handling and Statistical Analysis

Data were analysed using STATISTICA 8.0 (StatSoftInc, Tulsa, Oklahoma). Inspection of model residuals indicated data were normally distributed. Assumptions of sphericity were not violated. Third-party pleasantness ratings of social touch were analysed by conducting a 4-way Mixed ANOVA with a between-subjects factor of Group (3 levels: AN, RAN and HCs) and within-subjects factors of velocity (3 levels: 0cm/s, 5cm/s and 30cm/s), body sites (5 levels: Upper arm, ventral forearm, back, cheek and palm) and Task (2 levels: Other-directed touch and self-directed touch). All demographic information and scores for the self-reports are reported as Mean (M) and Standard Deviation of the mean (SD). A significance threshold of $p < .05$ was used as the significance threshold for each of the effects. All pairwise comparisons were assessed using the Duncan's post-hoc test correction for multiple comparisons, which reduces the size of the critical difference depending on the number of steps separating the ordered means. This procedure is optimal for testing in the same design effects that may have different sizes [70, 71, 72], as it is expected in our case for differences between patients AN, RAN and HCs participants and, within the patient group, between AN and RAN patients, as well as for the size of the effects of the different within-subject variables (i.e., Self- vs. Other-directed touch effects).

Independent samples t-tests, with Bonferroni Correction were conducted to assess group differences for subscales of the EDE-Q, MAIA and TEAQ questionnaires, by examining significant differences in mean scores across each group. A significance threshold of $p < .05$ was used as the significance threshold for each of the effects.

In keeping with previous studies [73-74], a Pleasant Touch Awareness (PTA) index was calculated as the difference in pleasantness ratings between CT-optimal (5 cm/s) and CT non-optimal stroking (30 cm/s), weighted by the average scores calculated separately for each participant and location ($PTA = (\text{pleasantness ratings at 5cm/s} - \text{pleasantness ratings at 30 cm/s}) / \text{overall touch pleasantness}$). PTA measures the degree to which an individual prefers CT-optimal as opposed to CT non-optimal velocities [75], and it was used in the current study to compare preference for CT-optimal touch across the three groups (AN, RAN and HCs) and the two tasks. Furthermore, an Overall Touch Pleasantness (OTP index was calculated as the average across CT-optimal velocity (5cm/s) and CT-non optimal velocities (0cm/s and 30cm/s). This index was calculated separately for each group and task [73, 74].

OTP refers to the extent to which an individual finds touch pleasant, which is not CT-specific. Thus, in the current study we wanted to examine overall touch pleasantness amongst the three groups (AN, RAN and HCs) and self- vs. other-directed social touch.

For exploratory purposes, we performed a series of exploratory Multiple Linear Regression analyses to investigate the predictive role of TEAQ AIT and MAIA Trusting subscales on both PTA and OTA, across the three groups and tasks.

Results

Univariate Statistics

Table 1 reports means and standard deviations for the demographic and psychometric questionnaire subscales, which has been calculated separately for AN, RAN and HCs. The far-right column shows the output of pairwise comparisons between the three groups, which have been adjusted for multiple comparisons (Bonferroni-corrected). The three groups were matched for age. However, as expected, all groups differed regarding their self-reported BMI, with AN having a significantly lower BMI than HCs and RAN, and RAN had significantly lower (but still in the normal-weight range) BMI than HCs. Furthermore, as expected, AN had significantly higher EDE-Q global scores than RAN and HCs, whilst no difference was observed between RAN and HCs.

Between AN and HCs, there was a significant difference observed for the subscale MAIA Self-regulation, with HCs demonstrating higher ability to self-regulate distress when paying attention to inner body sensations, compared to AN. HCs had also significantly higher MAIA Trusting scores and thus reported greater experiences of their body as safe and trustworthy compared to both AN and RAN. No other group differences were observed for any other interoceptive facet, as measured by the MAIA.

For touch experiences and attitudes, HCs currently experience more Family and Friends touch and current intimate touch compared to AN. During childhood, HCs reported experiencing significantly more childhood touch compared to both AN and RAN. HCs also reported more positive attitudes towards self-care, and intimate touch compared to both AN and RAN and reported higher attitudes to unfamiliar touch compared to AN. Both AN and RAN did not differ in their current and previous touch experiences, and both demonstrated similar attitudes towards touch and self-care.

-----Insert Table 1 here-----

Additional demographics such as ethnicity, relationship status and education level were also collected and are reported in Table 2. We carried out Chi-Square analysis between all 3 groups to investigate whether there were any differences in these demographics between groups. Importantly, there was no significant differences in ethnicity ($\chi^2_6 = 3.28$; $p = .773$), relationship status ($\chi^2_{16} = 13.18$; $p = .659$) and education level ($\chi^2_{10} = 5.33$; $p = .868$) between groups.

-----Insert Table 2 here-----

Self-directed vs Other-directed affective touch ratings

The repeated measures ANOVA revealed a significant main effect of Body sites [$F(4,352) = 19.10$, $p < .001$, $\eta p^2 = .18$], with the back being rated as most pleasant overall (all $ps < .001$). The upper arm, ventral forearm and palm were all rated as more pleasant than the cheek (all $ps < .001$). No other significant differences in pleasant ratings for the other body sites were observed (all $ps > .351$). There was also a significant main effect of velocity [$F(2,176) = 11.99$, $p < .001$, $\eta p^2 = .12$], with CT-optimal touch (5cm/s) rated as significantly more pleasant than touch applied at CT non-optimal velocities (0cm/s and 30cm/s) (all $ps < .001$). No significant differences were observed between the two CT non-optimal velocities ($p = .682$). There was a significant main effect of Task [$F(1,88) = 67.57$, $p < .001$, $\eta p^2 = .43$], with overall touch being rated as more pleasant for other-directed touch compared to self-directed touch ($p < .001$). A significant main effect of Group was also observed [$F(2, 88) = 6.43$, $p = .002$, $\eta p^2 = .13$]. Overall, there were significant differences in pleasantness ratings between both HCs and AN ($p = .022$) and HCs and RAN ($p = .002$), with HCs providing higher overall pleasantness ratings compared to AN and RAN. No significant differences were observed in pleasantness ratings provided by AN and RAN ($p = .301$).

A significant 2-way interaction of body sites $\times$ velocity was revealed [$F(8, 704) = 8.64$, $p < .001$, $\eta p^2 = .09$] with a velocity of 5cm/s to the ventral forearm being rated as significantly

more pleasant than 30cm/s to the ventral forearm ($p < .001$) and 0cm/s to the ventral forearm ($p < .001$). Touch delivered at 5cm/s to the cheek was rated as significantly more pleasant than touch delivered at both 30cm/s ($p < .001$) and 0cm/s ($p < .001$). Also, touch delivered at 5cm/s to the palm was rated as significantly more pleasant than that delivered at 30cm/s ($p = .007$) and 0cm/s ($p = .005$). All other body sites and velocities were not significant ($p > .224$).

Crucially, a 3-way interaction of velocity $\times$ task $\times$ group was also observed [$F(4, 176) = 2.99$, $p = .020$, $\eta p^2 = .06$; See Figure 2]. For self-directed touch, post-hoc comparisons of the three groups of AN, RAN and HCs at each level of velocity revealed that AN and RAN rated touch delivered at 5cm/s significantly lower than HCs (all $p < .05$). No difference in pleasantness ratings was observed between AN and RAN for touch delivered at CT-optimal velocity ($p = .946$). A significant difference was also observed between pleasantness ratings provided by HCs and RAN for 30cm/s ($p = .007$), with HCs providing higher preference ratings for 30cm/s compared to RAN. There were no significant differences amongst the three groups in their ratings of 0cm/s touch (all $ps > .429$). For the other-directed task, there was no significant differences between groups for any of the velocities (all $ps > .05$).

In summary, regardless of whether touch involved the self or another person, all groups demonstrated a preference for CT-optimal touch (5cm/s) compared to CT-non optimal touch (0cm/s and 30cm/s). For all groups, touch evaluated for another person (other-directed touch) was perceived as more pleasant compared to when participants were asked to judge receiving the same touch to themselves (self-directed touch). Crucially, both AN and RAN demonstrated lower preference for both self- and other-directed touch when delivered at CT-optimal velocity (5cm/s) than HCs.

-----Insert Figure 2 here-----

Pleasant touch awareness and overall touch perception related to attitudes in intimate touch and the trusting of one's own body

We performed a series of exploratory (given the relatively small sample size for such analyses) Multiple Linear Regression analyses to explore whether individual differences in experiences to intimate touch and of one's body as safe and trustworthy could predict AN, RAN and HCs' preference for CT-optimal touch (PTA) or for overall touch (regardless of CT-Optimal specificity, OTP). Accordingly, we included TEAQ AIT and MAIA Trusting as predictors of PTA and OTP respectively for self- and other-directed touch. We repeated these regressions for each group separately.

For self-directed touch, we found a significant regression equation for the overall touch pleasantness for HCs [$F(2,34) = 4.359, p = .021$] and for AN [$F(2,26) = 6.365, p = .006$]. For RAN, the regression equation was only marginally significant [$F(2,28) = 3.219, p = .056$]. Specifically, the TEAQ AIT was a significant predictor for overall touch preference for both HCs and AN, suggesting higher positive attitude towards intimate touch to predict higher pleasantness ratings to social touch (regardless of CT optimality of the stimulation) in both HCs and AN, but not in RAN (see Table 3). For other-directed touch, we did not find significant regression equations for the overall touch pleasantness for HCs [$F(2,34) = .839, p = .441$], AN [$F(2,26) = 1.635, p = .216$] or RAN [$F(2,28) = .638, p = .537$] (See Table 3).

We then conducted a separate Multiple Linear Regression analysis for each group to investigate whether TEAQ AIT and MAIA Trusting would predict women' PTA for self-directed touch. Again, we did not find significant regression equations for neither of the three groups (HCs [$F(2,34) = .110, p = .896$], AN [$F(2,26) = .262, p = .772$], RAN [$F(2,28) = .715, p = .499$]). Similarly, we did not find significant regression equations for PTA for other-directed touch in any of the three groups (HCs [$F(2,34) = .432, p = .653$], AN [$F(2,26) = .356, p = .704$] and RAN [$F(2,28) = .089, p = .915$], See Table 3).

-----Insert Table 3 here-----

Discussion

To the best of our knowledge, this is the first investigation to assess whether third-party vicarious ratings of self- vs. other-directed social touch, delivered at CT-optimal vs. CT non-optimal velocities differ in women with a current or previous diagnosis of AN, compared to healthy individuals with no AN diagnosis. Our analysis also included covariates such as

interoceptive awareness and attitudes to touch with the aim to explore whether potential factors which are known to largely contribute towards the aetiology and development of EDs [1, 9, 10, 11] might also explain subjective ratings of pleasantness for social touch.

When taking into consideration both self- and other-directed touch tasks, both groups of patients and healthy comparison subjects showed the expected velocity dependent pleasantness curve for brush stroking, indicating that C-tactile targeted stroking velocities were preferred over slower and faster velocities [20, 76]. Although the case, both AN and RAN demonstrated significantly lower pleasantness ratings for CT-optimal touch compared to HCs. This finding is in line with those of real experiences to touch by Crucianelli et al. [14] as they found that anorexics rated CT-optimal touch (delivered to the self) as less pleasant than healthy controls, which was not evident for CT-non optimal velocities. Additionally, this finding supports research by Davidovic et al. [13], whereby current anorexics rated touch as less pleasant than matched HCs. Yet, this finding contradicts the results obtained by Crucianelli et al. [15] as they found that a reduction in tactile pleasantness was not CT-specific. A possible explanation for this finding is that atypical responses to affective touch is due to an error in bottom-up processing, i.e., an error in the sensation and perception of touch, which could, as suggested by Crucianelli et al. [14], be due to a dysfunctional CT afferent system, particularly as this finding is more evident in the self-directed task.

Both current and remitted anorexics demonstrated atypical and clear differences in vicarious affective touch ratings specifically for self-directed touch, compared to healthy controls, with both clinical groups demonstrating reduced pleasantness ratings compared to healthy controls. This is in line with Crucianelli et al. [15] who found that both women with, and recovered from, AN rated affective touch as less pleasant compared to healthy controls. A possible mutual but not exclusive explanation concerning atypical responses to affective touch to the self could be due to anorexics' anxiety and discomfort towards touch in general [77]. Accordingly, we observed significant differences in the ratings to experience and attitude to social touch across all domains of touch experience and amongst the three groups. Indeed, AN reported significantly less positive experience to familiar and unfamiliar touch, as well as to intimate touch compared to HCs. Interestingly, a significant regression equation for self-reports of touch experiences and overall touch pleasantness for AN, particularly for vicarious self-directed touch lend support to this idea. With these regards, it was found that positive attitude to intimate touch was positively associated with overall touch pleasantness

for current anorexics (but also for HCs), when concerning touch to the self. This is in line with previous findings from Zucker et al. [44], who investigated sensitivity and avoidance of touch in women with current and a previous diagnosis of AN, in comparison to HCs. Both clinical groups demonstrated higher sensitivity and avoidance of touch compared to HCs, with AN demonstrating higher touch sensitivity. Furthermore, another possible explanation for the association between attitude to intimate touch and overall touch perception to the self-directed touch is that typically anorexics tend to struggle and find closeness discomforting and have difficulty in maintaining close intimate relationships with a romantic partner, family, and friends [54, 77]. The more severe the body image disturbances in these populations, the less likely they are to engage and feel comfortable in physical intimacy i.e., touch from a romantic partner or feel comfortable with touch from a loved one [78]. This could explain the reduction in pleasantness of touch to the self for both current anorexics, as low exposure to touch was found to result in lower pleasantness of touch [79]. Yet, our findings are not conclusive and should be interpreted with caution given they are exploratory in their nature. As such, further studies with larger clinical samples should be conducted to better investigate the contribution of social touch experiences and attitudes in predicting anorexics' sensitivity to CT-optimal affective touch.

Contrary to our prediction, we did not find evidence that experience of one's body as safe and trustworthy (MAIA Trusting) might predict women' preference for CT-Optimal touch nor for the overall touch perception, in the three groups. Previous works by Brown et al. [51] reported that items from the MAIA Trusting subscale were the most central in bridging interoceptive awareness and ED symptoms [51]. However, the study by Crucianelli et al. [15] reported that individual differences in the perceived pleasantness of CT-optimal in the AN group were predicted by interoceptive sensibility scores (as measured by the Body Awareness Questionnaire [80]), thus suggesting that other facets of interoception awareness might play a role in the perception of social touch in AN. Thus, future studies in AN are needed to disentangle the exact contribution of multidimensions of interoceptive awareness to preference to social touch to draw firm conclusions about the mechanisms that cause blunted responses to vicarious experience of social touch in this population.

Overall, as expected, we observed group differences for third party ratings of pleasantness of touch for self but not for another. We capitalized on prior research which has reported that individuals with AN have impaired ToM which involves the decoding of affective stimuli [38] both at self and other perspectives [36, 37] and it was found to manifest

even after recovery [39, 40, 41, 42, 43, 44, 81]. In this study, it was expected that anorexics would use their own experiences to interpret the feelings generated in another, which would be less pleasant compared to neurotypical populations. However, both current and remitted anorexics did not perceive CT-targeted touch as less pleasant than healthy controls for other-directed touch. Instead, as found previously, no differences were observed between anorexics and control participants regarding perspective taking when engaging in ratings for other-directed touch [38, 82, 83, 84]. A possible explanation for this finding could be related to ‘learned experiences’ of touch as pleasant when delivered to another, which conversely does not apply to the self. With these regards, it might be the case that anorexics may observe how others respond to the receiving of touch and use this learned experience when making judgments about touch for others, regardless of the fact that they perceive touch to themselves as less pleasant [85]. In doing so, they may have boosted their ToM abilities, which is a process which enhances based on an individual’s experience of that specific interaction. Another possible but not mutually exclusive explanation is that, like found previously, anorexics have an intact theory of mind functioning, comparable to healthy controls [38, 82, 83, 84, 86]. Thus, as suggested by Sedgewick et al. [86], the social-emotional difficulties anorexics face may not be specific to inferring and recognising emotions and mental states but rather this would reflect a bias in the interpretation of their social situations that they are more likely to avoid. Furthermore, it also could be that AN struggle with more complex social situations involving a dynamic of various interactions, as opposed to less complex social situations such as receiving pleasant touch from another individual [86].

Although this study was the first to assess third-party ratings of social touch in current and remitted anorexics, limitations have been identified. First, self-report diagnosis may have posed an issue with the variability of individuals in both clinical samples and also there is a risk of some undiagnosed in the healthy control group. However, these concerns are mitigated by several factors. First, the online survey was anonymous, which should encourage honest responding. Furthermore, participants’ ability to continue to take part to the study depended on their responses, which encouraged the provision of high-quality data. Most importantly, the manipulation check indicated that individuals reporting AN endorsed significantly greater eating pathology on the EDE-Q which is a gold standard measure for the assessment of symptom severity and AN diagnosis in replacement of structured interviews (i.e., Eating disorder examination interview [87, 88]). Future studies should however attempt to replicate

these findings by recruiting AN inpatients whose diagnosis has been established by a qualified clinician.

Although the study controlled for various variables that may affect touch pleasantness, other factors such as touch avoidance, sexuality, mood and empathic traits may have an impact as well. For example, a recent study by Sailer and Ackerley [79] found participants with low touch exposure scores, which on the Social Touch Questionnaire (STQ) [89] is indicative of unpleasantness of social touch and avoidance, rated touch differently, as they could not differentiate various stroking velocities. Furthermore, mood may have influenced social touch perception. Although participants were asked to state if they had any neurological or psychiatric conditions, a form of anhedonia measurement such as the Fawcett-Clark Pleasure Scale (FCPS) [90] or the Snaith-Hamilton Pleasure Scale (SHAPS) [91] should have been administered to account for participants mood on their testing day and those above the cut-off point excluded. This is because anhedonia, a characteristic for depression associated with the reduced ability to experience pleasure, can negatively affect attitudes to social touch [92]. Furthermore, in Crucianelli et al. [15] study, tactile anhedonia was found to be present in RAN as well as AN, suggesting that this could be a trait which manifests even post-recovery which could account for their lack of pleasure towards touch.

Another factor which may have impacted the third-party vicarious ratings of social touch is individual differences in empathic abilities. This factor influences participants' ability during embodiment tasks and vicarious experiences [93, 94, 95]. For example, Peled-Avron et al. [96] found levels of empathy modified ratings and EEG responses to photos of social touch. Participants classified as highly empathetic, had higher mu suppression and rated social touch as more pleasant, as opposed to participants reporting lower empathy levels. Therefore, it would be prudent in future studies that involve third-party ratings of touch, rather than actual touch, to consider empathy measures, as responses to observed touch are modulated by empathy levels and previous research has found that current anorexics have lower levels of empathy compared to RAN and HCs [41, 97].

Conclusions

To conclude, our findings add to previous results of Crucianelli et al. [14] and Davidovic and colleagues [13] in that they show that despite atypical, blunted responses to affective touch, women with AN and RAN still demonstrate preference for CT-optimal touch.

Most importantly, we provide initial evidence that third-party ratings of touch pleasantness for another individual do not differ regardless of eating disorder diagnosis. On the contrary, both RAN and AN groups rated affective touch as less pleasant compared to healthy controls when touch involved the self. Thus, altered responses to vicarious experience of affective touch in AN and RAN might be self-specific and intact when making judgements of touch for someone else. This could be due to a learning mechanism according to which AN have learned that the touch they are viewing would be pleasant for another individual, with multiple facets of interoceptive awareness and social touch experiences possibly mediating this effect. Given that atypical responses persist even after recovery, treatment interventions should focus on overcoming an impairment in differentiating between self and other affective touch experience, which could help prevent post-recovery relapsing.

Data Availability

The data that support the findings of this study are available on request from the corresponding author. The data are not publicly available due to privacy or ethical restrictions.

References

1. Beilharz, F. *et al.* (2019). Dysmorphic concern in anorexia nervosa: Implications for recovery. *Psychiatry Research*, 273, 657-661.
2. Cazzato, V. *et al.* (2016). The effects of body exposure on self-body image and esthetic appreciation in anorexia nervosa. *Experimental Brain Research*, 234(3), 695-709.
3. Urgesi, C. *et al.* (2014). Impaired configural body processing in anorexia nervosa: Evidence from the body inversion effect. *British Journal of Psychology*, 105(4), 486–508.
4. American Psychiatric Association. (2013). Diagnostic and statistical manual of mental disorders (5th ed.).
5. Kerr, K. L. *et al.* (2016). Altered insula activity during visceral interoception in weight-restored patients with anorexia nervosa. *Neuropsychopharmacology*, 41(2), 521-528.

6. Monteleone, P., & Maj, M. (2013). Dysfunctions of leptin, ghrelin, BDNF and endocannabinoids in eating disorders: beyond the homeostatic control of food intake. *Psychoneuroendocrinology*, 38(3), 312-330.
7. Oberndorfer, T. A. *et al.* (2013). Altered insula response to sweet taste processing after recovery from anorexia and bulimia nervosa. *American Journal of Psychiatry*, 170(10), 1143-1151.
8. Perez, M. E., Coley, B., Crandall, W., Di Lorenzo, C., & Bravender, T. (2013). Effect of nutritional rehabilitation on gastric motility and somatization in adolescents with anorexia. *The Journal of Pediatrics*, 163(3), 867-872.
9. Strigo, I. A. *et al.* (2013). Altered insula activation during pain anticipation in individuals recovered from anorexia nervosa: evidence of interoceptive dysregulation. *International Journal of Eating Disorders*, 46(1), 23-33.
10. Kaye, W. H., Fudge, J. L., & Paulus, M. (2009). New insights into symptoms and neurocircuit function of anorexia nervosa. *Nature Reviews Neuroscience*, 10(8), 573-584.
11. Nandrino, J. L. *et al.* (2012). Cognitive and physiological dissociations in response to emotional pictures in patients with anorexia. *Journal of Psychosomatic Research*, 72(1), 58-64.
12. Bischoff-Grethe, A. *et al.* (2018). Neural hypersensitivity to pleasant touch in women remitted from anorexia nervosa. *Translational Psychiatry*, 8(1), 1-13.
13. Davidovic, M. *et al.* (2018). Abnormal brain processing of gentle touch in Anorexia Nervosa. *Psychiatry Research: Neuroimaging*, 281(1), 53-60.
14. Crucianelli, L., Cardi, V., Treasure, J., Jenkinson, P. M., & Fotopoulou, A. (2016). The perception of affective touch in anorexia nervosa. *Psychiatry Research*, 239(1), 72-78.
15. Crucianelli, L. *et al.* (2021). The anticipation and perception of affective touch in women with and recovered from anorexia nervosa. *Neuroscience*, 464, 143-155.
16. Ackerley, R., Carlsson, I., Wester, H., Olausson, H., & Backlund Wasling, H. (2014). Touch perceptions across skin sites: Differences between sensitivity, direction discrimination and pleasantness. *Frontiers in Behavioral Neuroscience*, 54(8), 1-10.
17. Liu, Q. *et al.* (2007). Molecular genetic visualization of a rare subset of unmyelinated sensory neurons that may detect gentle touch. *Nature Neuroscience*, 10(8), 946-948.
18. McGlone, F., Wessberg, J., & Olausson, H. (2014). Discriminative and affective touch: sensing and feeling. *Neuron*, 82(4), 737-755.

19. Woodbury, C. J., Ritter, A. M., & Koerber, H. R. (2001). Central anatomy of individual rapidly adapting low-threshold mechanoreceptors innervating the “hairy” skin of newborn mice: early maturation of hair follicle afferents. *Journal of Comparative Neurology*, 436(3), 304-323.
20. Löken, L. S., Wessberg, J., McGlone, F., & Olausson, H. (2009). Coding of pleasant touch by unmyelinated afferents in humans. *Nature Neuroscience*, 12(5), 547-548.
21. Björnsdotter, M., Löken, L., Olausson, H., Vallbo, Å., & Wessberg, J. (2009). Somatotopic organization of gentle touch processing in the posterior insular cortex. *Journal of Neuroscience*, 29(29), 9314-9320.
22. Gordon, I. *et al.* (2013). Brain mechanisms for processing affective touch. *Human Brain Mapping*, 34(4), 914-922.
23. Morrison, I. (2016). ALE meta-analysis reveals dissociable networks for affective and discriminative aspects of touch. *Human Brain Mapping*, 37(4), 1308-1320.
24. Craig, A. D. (2002). How do you feel? Interoception: the sense of the physiological condition of the body. *Nature Reviews Neuroscience*, 3(8), 655-666.
25. Kirsch, L. P. *et al.* (2020). Damage to the right insula disrupts the perception of affective touch. *Elife*, 9, 1-17.
26. Morrison, I., Löken, L. S., & Olausson, H. (2010). The skin as a social organ. *Experimental Brain Research*, 204(3), 305-314.
27. Olausson, H., Wessberg, J., McGlone, F., & Vallbo, Å. (2010). The neurophysiology of unmyelinated tactile afferents. *Neuroscience & Biobehavioral Reviews*, 34(2), 185-191.
28. Carey, M., Crucianelli, L., Preston, C., & Fotopoulou, A. (2019). The effect of visual capture towards subjective embodiment within the full body illusion. *Scientific Reports*, 9(1), 1-12.
29. Carey, M., Crucianelli, L., Preston, C., & Fotopoulou, A. (2021). The role of affective touch in whole-body embodiment remains equivocal. *Consciousness and Cognition*, 87, 1-15.
30. Cazzato, V., Sacchetti, S., Shelby, S., Makdani, A., Trotter, P. D., & McGlone, F. (2021). Affective touch topography and body image. *PloS One (In Press)*.
31. Blakemore, S. J., Bristow, D., Bird, G., Frith, C., & Ward, J. (2005). Somatosensory activations during the observation of touch and a case of vision–touch synaesthesia. *Brain*, 128(7), 1571-1583.

32. Ebisch, S. J. *et al.* (2008). The sense of touch: embodied simulation in a visuotactile mirroring mechanism for observed animate or inanimate touch. *Journal of Cognitive Neuroscience*, 20(9), 1611-1623.
33. Schaefer, M., Heinze, H. J., & Rotte, M. (2012). Embodied empathy for tactile events: interindividual differences and vicarious somatosensory responses during touch observation. *Neuroimage*, 60(2), 952-957.
34. Morrison, I. *et al.* (2011). Reduced C-afferent fibre density affects perceived pleasantness and empathy for touch. *Brain*, 134(4), 1116-1126.
35. Walker, S. C., Trotter, P. D., Woods, A., & McGlone, F. (2017). Vicarious ratings of social touch reflect the anatomical distribution & velocity tuning of C-tactile afferents: A hedonic homunculus?. *Behavioural Brain Research*, 320, 91-96.
36. Cooper, K., & Mohr, C. (2012). Former eating disorder impairs 3rd person but not 1st person perspective taking: does dance training help?. *Comprehensive Psychology*, 1, 02-06.
37. Ruby, P., & Decety, J. (2003). What you believe versus what you think they believe: a neuroimaging study of conceptual perspective-taking. *European Journal of Neuroscience*, 17(11), 2475-2480.
38. Bora, E., & Köse, S. (2016). Meta-analysis of theory of mind in anorexia nervosa and bulimia nervosa: A specific impairment of cognitive perspective taking in anorexia nervosa?. *International Journal of Eating Disorders*, 49(8), 739-740.
39. Gál, Z., Egyed, K., Pászthy, B., & Németh, D. (2011). Impaired theory of mind in anorexia nervosa. *Psychiatria Hungarica: A Magyar Pszichiatriai Tarsasag Tudományos Folyoirata*, 26(1), 12-25.
40. Hamatani, S. *et al.* (2016). Impaired social cognition in anorexia nervosa patients. *Neuropsychiatric Disease and Treatment*, 12, 2527-2531.
41. Konstantakopoulos, G., Ioannidi, N., Patrikelis, P., & Gonidakis, F. (2020). Empathy, mentalizing, and cognitive functioning in anorexia nervosa and bulimia nervosa. *Social Neuroscience*, 15(4), 477-488.
42. Russell, T. A., Schmidt, U., Doherty, L., Young, V., & Tchanturia, K. (2009). Aspects of social cognition in anorexia nervosa: affective and cognitive theory of mind. *Psychiatry Research*, 168(3), 181-185.
43. Tchanturia, K. *et al.* (2004). 'Theory of mind' in anorexia nervosa. *European Eating Disorders Review: The Professional Journal of the Eating Disorders Association*, 12(6), 361-366.

44. Zucker, N. L. *et al.* (2007). Anorexia nervosa and autism spectrum disorders: guided investigation of social cognitive endophenotypes. *Psychological Bulletin*, 133(6), 976-1006.
45. McAdams, C. J., & Krawczyk, D. C. (2011). Impaired neural processing of social attribution in anorexia nervosa. *Psychiatry Research: Neuroimaging*, 194(1), 54-63.
46. Lee Masson, H., Wallraven, C., & Petit, L. (2017). “Can touch this”: Cross-modal shape categorization performance is associated with microstructural characteristics of white matter association pathways. *Human Brain Mapping*, 38(2), 842-854.
47. Lee Masson, H. L., Van De Plas, S., Daniels, N., & de Beeck, H. O. (2018). The multidimensional representational space of observed socio-affective touch experiences. *Neuroimage*, 175, 297-314.
48. Badoud, D., & Tsakiris, M. (2017). From the body’s viscera to the body’s image: Is there a link between interoception and body image concerns?. *Neuroscience & Biobehavioral Reviews*, 77, 237-246.
49. Mehling, W. E. *et al.* (2012). The multidimensional assessment of interoceptive awareness (MAIA). *PloS One*, 7, e48230.
50. Brown, T. A. *et al.* (2017). Psychometric evaluation and norms for the Multidimensional Assessment of Interoceptive Awareness (MAIA) in a clinical eating disorders sample. *European Eating Disorders Review*, 25(5), 411-416.
51. Brown, T. A. *et al.* (2020). Body mistrust bridges interoceptive awareness and eating disorder symptoms. *Journal of Abnormal Psychology*, 129(5), 445-456.
52. Trotter, P. D., McGlone, F., Reniers, R. L. E. P., & Deakin, J. F. W. (2018). Construction and validation of the Touch Experiences and Attitudes Questionnaire (TEAQ): a self-report measure to determine attitudes toward and experiences of positive touch. *Journal of Nonverbal Behavior*, 42(4), 379-416.
53. Maier, S. *et al.* (2019). Neurobiological signature of intimacy in anorexia nervosa. *European Eating Disorders Review*, 27(3), 315-322.
54. Evans, L., & Wertheim, E. H. (1998). Intimacy patterns and relationship satisfaction of women with eating problems and the mediating effects of depression, trait anxiety and social anxiety. *Journal of Psychosomatic Research*, 44(3-4), 355-365.
55. Pruitt, N. A., Kappius, R. E., & Gorman, P. W. (1992). Bulimia and fear of intimacy. *Journal of Clinical Psychology*, 48(4), 472-476.
56. Thelen, M. H., Farmer, J., Mann, L. M., & Pruitt, J. (1990). Bulimia and interpersonal relationships: A longitudinal study. *Journal of Counseling Psychology*, 37(1), 85-90.

57. Faul, F., Erdfelder, E., Lang, A. G., & Buchner, A. (2007). G* Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior Research Methods*, 39(2), 175-191.
58. Russo, V., Ottaviani, C., & Spitoni, G. F. (2020). Affective touch: A meta-analysis on sex differences. *Neuroscience & Biobehavioral Reviews*, 108, 445-452.
59. Striegel-Moore, R. H. *et al.* (2009). Gender difference in the prevalence of eating disorder symptoms. *International Journal of Eating Disorders*, 42(5), 471-474.
60. Mond, J. M., Hay, P. J., Rodgers, B., & Owen, C. (2006). Eating Disorder Examination Questionnaire (EDE-Q): norms for young adult women. *Behaviour Research and Therapy*, 44(1), 53-62.
61. Fairburn, C. G., & Beglin, S. J. (1994). Assessment of eating disorders: Interview or self-report questionnaire?. *International Journal of Eating Disorders*, 16(4), 363-370.
62. Luce, K. H., & Crowther, J. H. (1999). The reliability of the eating disorder examination—Self-report questionnaire version (EDE-Q). *International Journal of Eating Disorders*, 25(3), 349-351.
63. Peterson, C. B. *et al.* (2007). Psychometric properties of the eating disorder examination-questionnaire: Factor structure and internal consistency. *International Journal of Eating Disorders*, 40(4), 386-389.
64. Berg, K. C., Peterson, C. B., Frazier, P., & Crow, S. J. (2012). Psychometric evaluation of the eating disorder examination and eating disorder examination-questionnaire: A systematic review of the literature. *International Journal of Eating Disorders*, 45(3), 428-438.
65. Valenzuela-Moguillansky, C., & Reyes-Reyes, A. (2015). Psychometric properties of the multidimensional assessment of interoceptive awareness (MAIA) in a Chilean population. *Frontiers in Psychology*, 120(6), 1-12.
66. Trotter, P., Belovol, E., McGlone, F., & Varlamov, A. (2018a). Validation and psychometric properties of the Russian version of the Touch Experiences and Attitudes Questionnaire (TEAQ-37 Rus). *PloS One*, 13(12), e0206905.
67. Henley, N. M. (1973). Status and sex: Some touching observations. *Bulletin of the Psychonomic Society*, 2(2), 91-93.
68. Stier, D. S., & Hall, J. A. (1984). Gender differences in touch: An empirical and theoretical review. *Journal of Personality and Social Psychology*, 47(2), 440-459.

69. Harjunen, V. J., Spapé, M., Ahmed, I., Jacucci, G., & Ravaja, N. (2017). Individual differences in affective touch: Behavioral inhibition and gender define how an interpersonal touch is perceived. *Personality and Individual Differences*, 107, 88-95.
70. Duncan, D. B. (1955). Multiple range and multiple F tests. *Biometrics*, 11(1), 1-42.
71. Dunnett, C. W. (1970). 272. Query: Multiple Comparison Tests. *Biometrics*, 26(1), 139-141.
72. McHugh, M. L. (2011). Multiple comparison analysis testing in ANOVA. *Biochemia Medica*, 21(3), 203-209.
73. Croy, I. *et al.* (2016). Interpersonal stroking touch is targeted to C tactile afferent activation. *Behavioural Brain Research* *SreeTestContent1*, 297, 37-40.
74. Jönsson, E. H. *et al.* (2017). The relation between human hair follicle density and touch perception. *Scientific reports*, 7(1), 1-10.
75. Croy, I., Sehlstedt, I., Wasling, H. B., Ackerley, R., & Olausson, H. (2019). Gentle touch perception: From early childhood to adolescence. *Developmental Cognitive Neuroscience*, 35, 81-86.
76. Essick, G. K. *et al.* (2010). Quantitative assessment of pleasant touch. *Neuroscience & Biobehavioral Reviews*, 34(2), 192-203.
77. Arcelus, J., Witcomb, G. L., & Mitchell, A. (2014). Prevalence of eating disorders amongst dancers: a systemic review and meta-analysis. *European Eating Disorders Review*, 22(2), 92-101.
78. Cash, T. F., Theriault, J., & Annis, N. M. (2004). Body image in an interpersonal context: Adult attachment, fear of intimacy and social anxiety. *Journal of Social and Clinical Psychology*, 23(1), 89-103.
79. Sailer, U., & Ackerley, R. (2019). Exposure shapes the perception of affective touch. *Developmental Cognitive Neuroscience*, 35, 109-114.
80. Shields, S. A., Mallory, M. E., & Simon, A. (1989). The body awareness questionnaire: reliability and validity. *Journal of Personality Assessment*, 53(4), 802-815.
81. de Sampaio, F. T. P. *et al.* (2013). Theory of mind and central coherence in eating disorders: Two sides of the same coin?. *Psychiatry Research*, 210(3), 1116-1122.
82. Adenzato, M., Todisco, P., & Ardito, R. B. (2012). Social cognition in anorexia nervosa: Evidence of preserved theory of mind and impaired emotional functioning. *PLoS One*, 7(8), e44414.

83. Calvo, R. S. *et al.* (2014). Emotional processing in adult vs adolescent patients with eating behavior disorders; emotional recognizing and the mental theory. *Nutricion Hospitalaria*, 29(4), 941-952.
84. Medina-Pradas, C., Navarro, J. B., Álvarez-Moya, E. M., Grau, A., & Obiols, J. E. (2012). Emotional theory of mind in eating disorders. *International Journal of Clinical and Health Psychology*, 12(2), 189-202.
85. Korkmaz, B. (2011). Theory of mind and neurodevelopmental disorders of childhood. *Pediatric Research*, 69(8), 101-108.
86. Sedgewick, F. *et al.* (2019). Similarities and differences in theory of mind responses of patients with anorexia nervosa with and without autistic features. *Frontiers in Psychiatry*, 10(318), 1-10.
87. Fairburn, C. G., Cooper, Z., & O'Connor, M. (1993). The eating disorder examination. *International Journal of Eating Disorders*, 6, 1-8.
88. Fairburn, C. G., Cooper, Z., & O'connor, M. (2008). Eating disorder examination (Edition 16.0 D). *Cognitive Behavior Therapy and Eating Disorders*, 265-308.
89. Wilhelm, F. H., Kochar, A. S., Roth, W. T., & Gross, J. J. (2001). Social anxiety and response to touch: incongruence between self-evaluative and physiological reactions. *Biological Psychology*, 58(3), 181-202.
90. Fawcett, J., Clark, D. C., Scheftner, W. A., & Gibbons, R. D. (1983). Assessing anhedonia in psychiatric patients: The Pleasure Scale. *Archives of General Psychiatry*, 40(1), 79-84.
91. Snaith, R. P. *et al.* (1995). A scale for the assessment of hedonic tone the Snaith–Hamilton Pleasure Scale. *The British Journal of Psychiatry*, 167(1), 99-103.
92. Triscoli, C., Croy, I., & Sailer, U. (2019). Depression predicts interpersonal problems partially through the attitude towards social touch. *Journal of Affective Disorders*, 246, 234-240.
93. Kaplan, J. T., & Iacoboni, M. (2006). Getting a grip on other minds: Mirror neurons, intention understanding, and cognitive empathy. *Social Neuroscience*, 1(3-4), 175-183.
94. Minio-Paluello, I., Baron-Cohen, S., Avenanti, A., Walsh, V., & Aglioti, S. M. (2009). Absence of embodied empathy during pain observation in Asperger syndrome. *Biological Psychiatry*, 65(1), 55-62.

95. Rueda, P., Fernández-Berrocal, P., & Baron-Cohen, S. (2015). Dissociation between cognitive and affective empathy in youth with Asperger Syndrome. *European Journal of Developmental Psychology*, 12(1), 85-98.
96. Peled-Avron, L., Levy-Gigi, E., Richter-Levin, G., Korem, N., & Shamay-Tsoory, S. G. (2016). The role of empathy in the neural responses to observed human social touch. *Cognitive, Affective, & Behavioral Neuroscience*, 16(5), 802-813.
97. Morris, R., Bramham, J., Smith, E., & Tchanturia, K. (2014). Empathy and social functioning in anorexia nervosa before and after recovery. *Cognitive Neuropsychiatry*, 19(1), 47-57.

Acknowledgements

We would like to thank all the participants who have taken part to the study. We would also like to thank BEAT charity, Kel O'Neill and MQ Mental Health for their assistance with data collection through advertising our study to anorexic populations.

Author Contributions

AB and VC constructed the experiment. AB, VC, PT and FM contributed towards the concept and design of the study. AB performed data collection under the supervision of VC, PT and FM. Data analysis was performed by VC and AB. The manuscript was written by AB, with input from VC, PT and FM. AB, VC, PT and FM approved the manuscript before submission.

Additional Information

Competing interests

The authors declare no competing interests.

Figure Legends

Figure 1. Visual illustration of the 5 body sites (CT-innervated body regions: Ventral Forearm, Upper Arm, Cheek and Back vs. the non-CT innervated palm) from the affective touch videos used in this study.

Figure 2. Pleasantness ratings for each velocity, which have been collapsed across body sites, with each bar representing one of three groups (AN, RAN and HCs). Graphs are separated for each task: (a) self-directed touch and (b) other-directed touch.

Tables

Table 1. Mean and standard deviation (in brackets) of demographics and self-report questionnaires scores for AN ($n=27$), RAN ($n=29$) and HCs ($n=35$).

	AN ($n=27$) M (SD)	RAN ($n=29$) M (SD)	HCs ($n=35$) M (SD)	AN vs. RAN p	AN vs HCs P	RAN vs. HCs p
Age (years)	25.56 (6.95)	27.31 (7.12)	27.20 (8.81)	ns	ns	ns
BMI (kg/cm ²)	17.58 (1.64)	22.59 (3.03)	24.77 (4.47)	<.001	<.001	.029
EDE-Q						
Restraint (max 6)	4.15 (1.54)	2.14 (1.93)	1.75 (1.68)	<.001	<.001	ns
Eating Concern (max 6)	3.30 (1.39)	1.83 (1.82)	1.13 (1.23)	.001	<.001	ns
Weight Concern (max 6)	4.16 (1.35)	2.96 (1.97)	2.45 (1.87)	.011	<.001	ns
Shape Concern (max 6)	4.16 (1.33)	3.07 (1.90)	2.77 (1.68)	.016	.001	ns
Global Score (max 6)	3.95 (1.24)	2.49 (1.77)	2.03 (1.47)	.001	<.001	ns
MAIA						
Noticing (max 5)	3.55 (1.01)	3.31 (1.06)	3.34 (1.07)	ns	ns	ns
Not Distracting (max 5)	2.58 (1.14)	2.41 (1.23)	2.80 (0.96)	ns	ns	ns
Not Worrying (max 5)	2.85 (1.29)	3.05 (1.16)	2.71 (0.99)	ns	ns	ns
Attention Regulation (max 5)	2.80 (0.96)	2.85 (1.20)	2.91 (0.98)	ns	ns	ns

Emotional Awareness (max 5)	3.24 (1.30)	3.32 (1.09)	3.46 (1.08)	ns	ns	ns
Self-regulation (max 5)	2.04 (1.25)	2.36 (1.38)	2.83 (1.14)	ns	.012	ns
Body Listening (max 5)	2.00 (1.31)	2.65 (1.24)	2.37 (1.27)	ns	ns	ns
Trusting (max 5)	1.46 (1.32)	2.25 (1.61)	3.14 (1.09)	ns	<.001	.011
TEAQ						
Friends and family touch (FFT; max 5)	2.68 (0.72)	2.99 (1.30)	3.51 (0.91)	ns	<.001	ns
Current intimate touch (CIT; max 5)	2.51 (0.89)	2.72 (0.97)	3.09 (0.90)	ns	.014	ns
Childhood touch (ChT; max 5)	3.09 (1.13)	3.02 (1.21)	3.85 (0.83)	ns	.003	.002
Attitude to self-care (ASC; max 5)	2.85 (1.07)	3.10 (0.87)	3.77 (0.78)	ns	<.001	.002
Attitude to intimate touch (AIT; max 5)	3.33 (1.02)	3.39 (1.08)	3.93 (0.85)	ns	0.14	.030
Attitude to unfamiliar touch (AUT; max 5)	2.30 (0.78)	2.36 (0.83)	2.78 (0.96)	ns	.040	ns

BMI Body Mass Index; *EDE-Q* Eating Disorder Examination Questionnaire; *MAIA*

Multidimensional Assessment of Interoceptive Awareness; *TEAQ* Touch Experiences and Attitudes Questionnaire; *ns* Not Significant

Table 2. Demographic characteristics of current anorexics ($n=27$) recovered anorexics ($n=29$) and HCs ($n=35$), which have been analysed by Chi-square.

	AN <i>n</i> (%)	RAN <i>n</i> (%)	HCs <i>n</i> (%)
Ethnicity			
Caucasian	21 (77.8%)	24 (82.8%)	29 (82.9%)
Mixed	4 (14.8%)	2 (6.9%)	2 (5.7%)
Black	1 (3.7%)	2 (6.9%)	1 (2.9%)
Asian	1 (3.7%)	1 (3.4%)	3 (8.6%)
Relationship Status			
Single	11 (40.7%)	14 (48.3%)	14 (40%)
In a Relationship	10 (37%)	10 (34.5%)	10 (28.6%)
Engaged	1 (3.7%)	0 (0%)	1 (2.9%)
Married	3 (11.1%)	4 (13.8%)	8 (22.9%)
Separated	0 (0%)	0 (0%)	1 (2.9%)
Divorced	0 (0%)	1 (3.4%)	0 (0%)
Widowed	1 (3.7%)	0 (0%)	0 (0%)
Not Specified	0 (0%)	0 (0%)	1 (2.9%)
Prefer not to say	1 (3.7%)	0 (0%)	0 (0%)
Education Level			
High School	9 (33.3%)	6 (20.7%)	7 (20%)
College	4 (14.8%)	5 (17.2%)	8 (22.9%)
Foundation	3 (11.1%)	1 (3.4%)	2 (5.7%)
Bachelors	6 (22.2%)	9 (31%)	11 (31.4%)
Masters	3 (11.1%)	5 (17.2%)	6 (17.1%)
Doctoral/Professional	2 (7.4%)	3 (10.3%)	1 (2.9%)
Treatment			
Medication	3 (11.1%)	-	-
<i>Fluoxetine</i>	1 (3.7%)	-	-
<i>SSRI</i>	2 (7.4%)	-	-

<i>Bupropion</i>	1 (3.7%)	-	-
Therapy	6 (22.2%)	-	-
<i>CBT</i>	1 (3.7%)	-	-
<i>Online Therapy</i>	1 (3.7%)	-	-
<i>Therapy not specified</i>	2 (7.4%)	-	-
Psychological/Psychiatric Treatment	2 (7.4%)	-	-
Day Patient Treatment	1 (3.7%)	-	-
Medication and Therapy	3 (11.1%)	-	-
No Treatment	12 (44.4%)	-	-
	AN	RAN	HCs
	M (SD)	M (SD)	M (SD)
Number of Years diagnosed	6.52 (8.62)	-	-
Number of Years Recovered	-	4.26 (3.60)	-

Table 3. Unstandardized coefficients of the multiple linear regression model for MAIA Trusting and TEAQ Attitude to intimate touch (AIT) for both PTA and OTP separated for each group (AN, RAN and HCs) and task (self-directed and other-directed touch).

				B	SE	β	<i>t-value</i>	<i>p-level</i>
HCs	Self-directed	PTA	MAIA_ Trusting	.040	.087	.082	.464	.646
			TEAQ_AIT	-.011	.111	-.018	-.102	.920
AN			MAIA_ Trusting	.063	.100	.129	.624	.538
			TEAQ_AIT	.030	.129	.048	.233	.818
RAN			MAIA_ Trusting	-.002	.117	-.003	-.014	.989
			TEAQ_AIT	-.188	.175	-.227	-1.072	.293
HCs	Other-directed		MAIA_ Trusting	.057	.062	.161	.918	.365
			TEAQ_AIT	-.017	.079	-.037	-.214	.832
AN			MAIA_ Trusting	.019	.066	.059	.290	.775
			TEAQ_AIT	.061	.085	.147	.718	.480
RAN			MAIA_ Trusting	.033	.078	.091	.422	.677
			TEAQ_AIT	-.020	.116	-.037	-.173	.864
HCs	Self-directed	OTP	MAIA_ Trusting	1.237	2.416	.080	.512	.612
			TEAQ_AIT	8.798	3.074	.450	2.862	.007*
AN			MAIA_ Trusting	3.135	2.041	.259	1.536	.138
			TEAQ_AIT	7.481	2.632	.479	2.843	.009*
RAN			MAIA_ Trusting	4.347	2.078	.407	2.092	.046
			TEAQ_AIT	1.218	3.108	.076	.392	.698
HCs	Other-directed		MAIA_ Trusting	.349	2.152	.028	.162	.872
			TEAQ_AIT	3.477	2.738	.219	1.270	.213
AN			MAIA_ Trusting	2.126	2.239	.186	.950	.352
			TEAQ_AIT	3.796	2.887	.257	1.315	.201
RAN			MAIA_ Trusting	2.345	2.215	.225	1.059	.299
			TEAQ_AIT	-.341	3.314	-.022	-.103	.919

Notes: AN Current Anorexics, RAN Remitted Anorexics, HCs Healthy Controls, PTA Pleasant Touch Awareness, OTP Overall Touch Pleasantness, MAIA Multidimensional Assessment of Interoceptive Awareness, TEAQ Touch Experiences and Attitudes Questionnaire, AIT Attitudes to Intimate Touch. *Indicates $p < 0.01$